

Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

A NOVEL AND CONVENIENT SYNTHESIS OF 4-HALOBUTYRALDEHYDE ACETALS

John A. Hyatt ^a

^a Research Laboratories, Eastman Chemical Company, Kingsport, TN, 37662

Published online: 18 Feb 2009.

To cite this article: John A. Hyatt (2004) A NOVEL AND CONVENIENT SYNTHESIS OF 4-HALOBUTYRALDEHYDE ACETALS, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 36:5, 487-490, DOI: [10.1080/00304940409356636](https://doi.org/10.1080/00304940409356636)

To link to this article: <http://dx.doi.org/10.1080/00304940409356636>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>


10. K. M. Davies, M. J. S. Dewar and P. Rona, *J. Am. Chem. Soc.*, **89**, 6294 (1967).
11. T. Jan, D. Floner and C. Moinet, *Electrochimica Acta*, **42**, 2073 (1997). Partial spectral data were also given in this reference.
12. R. G. R. Bacon and A. Karim, *J. Chem. Soc., Perkin Trans. 1*, 272 (1973).
13. A. Arcoria, J. Barassin and H. Lumbroso, *Bull. Soc. Chim. Fr.*, 2509 (1963).
14. D. L. Boger, H. Zarrinmeyer, *J. Org. Chem.*, **55**, 1379 (1990).
15. C. Yu, B. Liu and L. Hu, *J. Org. Chem.*, **66**, 919 (2001).

A NOVEL AND CONVENIENT SYNTHESIS OF 4-HALOBUTYRALDEHYDE ACETALS

Submitted by John A. Hyatt[†]
(08/02/04)

*Research Laboratories, Eastman Chemical Company
Kingsport, TN 37662*


The dimethyl and diethyl acetals of 4-chlorobutyraldehyde and 4-bromobutyraldehyde (1,1-dimethoxy- and 1,1-diethoxy-4-chlorobutane and 1,1-dimethoxy- and 1,1-diethoxy-4-bromobutane) are valuable synthetic intermediates. In particular, these acetals are useful in the Fischer indole synthesis of substituted tryptophols and tryptamines and in the synthesis of natural products.¹ Despite their utility and structural simplicity, there appears to be only one commercial source which makes available small quantities of one of the acetals.² The published chemistry for the preparation of these compounds, illustrated for chlorobutyraldehyde diethyl acetal, is rather lengthy and involves a poorly reproducible Rosenmund reduction.³ Furthermore, in our hands the



intermediate 4-chlorobutyraldehyde showed a strong tendency to form a tenaciously stable cyclic trimer upon storage. The overall yield of chloroacetal from this process is typically about 22%.

Other published routes are shorter but involve low-temperature⁴ or heavy-metal steps⁵ which are problematical to run at large scale. The commercialization of new process technology⁶ for air oxidation of butadiene to the corresponding mono-epoxide has made available, in large quantity, cyclopropanecarboxaldehyde (CPCA) (*via* a series of gas-phase conversions).⁷ We now report that CPCA can serve as the starting material for a very efficient and convenient new synthesis of 4-halobutyraldehyde acetals.

Reaction of CPCA (**1**) with acetyl chloride in the presence of a catalytic amount of zinc chloride provided an *E/Z* mixture of 4-chloro-1-acetoxy-1-butenes (**2a**) in 75% yield. This cyclopropane ring-opening reaction is reminiscent of the titanium-catalyzed process reported for cyclopropyl ketones by Oshima and coworkers.⁸ Compound **2a** has been previously reported only as a low-yield product from the thermal decomposition of a chromium complex.⁹ Although Hirsh and coworkers¹⁰ had reported the use of mercuric acetate to catalyze the conversion of enol esters to acetals, we found that the use of a simple resin gave comparable yields. Treatment of **2a** with excess methanol or ethanol in the presence of Amberlyst-15 acidic ion-exchange resin catalyst provided dimethyl and diethyl acetals **3a** and **3c** in 74% and 65% yield, respectively. The corresponding reaction of **1** with acetyl bromide produced the hitherto unreported bromo enol esters **2b** in 79% yield. The corresponding acetal **3b** was formed in 71% yield.



This work provides a short, convenient route to the methyl and ethyl acetals of both 4-chloro- and 4-bromobutyraldehyde.¹¹

EXPERIMENTAL SECTION

Proton NMR spectra were recorded in CDCl_3 solvent with TMS internal standard on a Varian Gemini 300 instrument. Gas chromatography (GC) was carried out on an HP 5890 instrument fitted with a 25 m DB-5 column operated from 40°C to 220°C at 10°C/min. Solvents and reagents were used as received from Aldrich Chemical Co. CPCA (**1**) is an Eastman Chemical Company commercial product.

4-Chloro-1-acetoxy-1-butene (2a).— To a stirred solution of 62.4 g (0.80 mole) of acetyl chloride in 750 mL of ethyl acetate at room temperature under nitrogen atmosphere was added in one portion 2.0 g (0.015 mole, 1.8 mol%) of anhydrous zinc chloride. There was then added 56.0 g (0.80 mole) of cyclopropanecarboxaldehyde at such a rate that the temperature of the reaction mixture remained between 40°C and 50°C without external cooling (1-2 hrs). The mixture was

then stirred at 40–45°C for 5 h and allowed to cool to rt. The reaction mixture was washed with aqueous 5% NaHCO₃, dried over MgSO₄, filtered and the solvent removed under reduced pressure. Distillation of the crude product through an 8" Vigreux column gave 89.1 g (75%) of a colorless liquid, bp 75–77°C/10 mm Hg. The product was found to consist of a roughly 50/50 *E/Z* mixture by NMR: analysis. IR: (neat) 1760 cm⁻¹ (*lit.*⁹ IR: 1759 cm⁻¹); ¹H NMR: (δ) 7.1–7.2 (m, 1 H), 5.45 (sextet, 0.5 H), 4.95 (q, 0.5 H), 3.5–3.6 (m, 2 H), 2.63 (q, 1 H), 2.42 (q, 1 H), 2.16 (s, 1.5 H), 2.12 (s, 1.5 H) (*lit.*⁹ ¹H NMR: for *Z* isomer: (δ) 7.13 (d, 1 H), 4.93 (q, 1 H), 3.54 (t, 2 H), 2.63 (q, 2 H), 2.16 (s, 3 H)). ¹³C NMR: (δ) 168.0 (*E*), 167.7 (*Z*), 137.5 (*E*), 136.0 (*Z*), 110.5 (*E*), 109.2 (*Z*), 44.1 (*E*), 43.6 (*Z*), 30.8 (*E*), 28.0 (*Z*), 20.7 (*Z*), 20.6 (*E*) (*lit.*⁹ ¹³C NMR: for *Z* isomer (δ) 167.1, 135.9, 109.2, 43.5, 27.9, 20.7).

4-Bromo-1-acetoxy-1-butene (2b).— A reaction performed as above using 56.3 g (0.46 mole) of acetyl bromide, 2.0 g (0.015 mole) of zinc chloride, and 32.2 g (0.46 mole) of **1** in 350 mL of ethyl acetate provided 70.6 g (80%) of **2b** as a colorless liquid, bp 70–73°C/6 mm Hg. NMR: δ 7.1–7.25 (m, 1 H), 5.42 (m, sextet, 0.5 H), 4.92 (q, 0.5 H), 3.38 (m, 2 H), 2.75 (q, 1 H), 2.56 (q, 1 H), 2.19 (s, 1.5 H), 2.13 (s, 1.5 H).

Anal. Calcd for C₆H₉BrO₂: C, 37.33; H, 4.70. Found, C, 37.47, H 4.66

4-Chlorobutyraldehyde Dimethyl Acetal (3a).— A solution of 72 g (0.49 mole) of **2a** in 300 mL of anhydrous methanol (240 g, 7.5 moles) was treated with 15 g of Amberlyst-15 acidic ion-exchange resin and stirred under reflux overnight, at which time GC analysis indicated complete conversion of **2a** to a single major product accompanied by several minor impurities. The catalyst was filtered off and the reaction mixture was treated with 2 g of solid sodium bicarbonate. The volume of the mixture then was reduced to about 75 mL on the rotary evaporator, and the residual bicarbonate was filtered off. Distillation of the filtrate through a 6" Vigreux column gave 55.0g (74%) of a colorless liquid, bp 85–90°C/25mm Hg, *lit.*³ bp 76–78°C/20 mm. ¹H NMR: (δ) 4.40 (t, 1 H), 3.58 (t, 2 H), 3.32 (s, 6 H), 1.65–1.9 (m, 4 H). (*Lit.*³ ¹H NMR: (δ) 4.38 (t, 1 H), 3.60 (t, 2 H), 3.30 (s, 6 H), 1.67–1.9 (m, 4 H)).

4-Chlorobutyraldehyde Diethyl Acetal (3c).— Reaction of 13.7 g (0.092 mole) of **2a** with 100 mL of 95% ethanol, 5 g of triethyl orthoformate (added as water scavenger) and 1.40 g of Amberlyst-15 provided 10.8 g (65%) of **4a** as a colorless liquid, bp 45–50°C/1mm Hg. ¹H NMR: (δ) 4.51 (t, 1 H), 3.4–3.8 (m, 6 H), 1.7–1.9 (m, 4 H), 1.10 (t, 6 H). ¹H NMR: of a commercial sample:² 4.51 (t, 1 H), 3.4–3.8 (m, 6 H), 1.7–1.9 (m, 4 H), 1.10 (t, 6 H).

4-Bromobutyraldehyde Dimethyl Acetal (3b).— This compound was prepared from **2b** in a manner exactly analogous to the preparation of **3a**. There was obtained a 71% yield of **3b**, bp 64–66°C/6 mm Hg. ¹H NMR: (δ) 4.42 (t, 1 H), 3.46 (t, 2 H), 3.38 (s, 6 H), 1.98 (m, 2 H), 1.80 (m, 2 H) (*lit.*¹² ¹H NMR (CCl₄) (δ) 4.25 (t, 1 H), 3.50 (t, 2 H), 3.20 (s, 6 H), 1.55–2.75 (m, 4 H)).

REFERENCES

[†] Address inquiries to: Department of Chemistry, East Tennessee State University, Box 70695 Johnson City, TN 37614-1710

1. a) C. Chen, C. H. Senanayake, T. J. Bill, R. D. Larsen, T. R. Verhoeven, and P. J. Reider, *J. Org. Chem.*, **59**, 3738 (1994). b) R. C. Glen, G. R. Martin, A. P. Hill, R. M. Hyde, P. M. Wollard, J. A. Salmon, J. Buckingham, and A. D. Robertson, *J. Med. Chem.*, **38**, 3566 (1995). c) A. D. Robertson, A. P. Hill, R. C. Glen, and G. R. Martin, *US Patent 5,466,699* (1995); CA: 116:174146. d) J. H. van Maarseveen, S. J. E. Mulders, R. W. H. Aben, C. G. Kruse, and H. W. Scheeren, *Tetrahedron*, **51**, 4841 (1995). e) W. R. Roush and S. E. Hall, *J. Am. Chem. Soc.*, **103**, 5200 (1981). f) B. M. Trost and T. A. Grese, *J. Org. Chem.*, **57**, 686 (1992).
2. Avocado Research Chemicals Ltd., Shore Rd., Heysham, Lancs, UK.
3. T. Athar, *Indian J. Chem.*, **37B**, 1037 (1998).
4. L. Crombie and D. Fisher, *Tetrahedron Lett.*, **26**, 2477 (1985).
5. C. P. Forbes, G. L. Wenteler, and A. Wiechers, *J. Chem. Soc. Perkin Trans. I*, 2353 (1977)
6. a) B. Venepalli, *Pharmaceutical Manufacturing International*, 93 (1996). b) D. Denton, S. Falling, J. Monnier, J. Stavinoha, and W. Watkins, *Chimica Oggi/Chemistry Today*, 17 (1996).
7. S. Liang and T. Price, *US Patent 5,633,419* (1997); CA: 127:4858.
8. Z. Han, S. Uehira, T. Tsuritani, H. Shinokubo, and K. Oshima, *Tetrahedron*, **57**, 987 (2001).
9. B. C. Soderberg, J. Liu, T. W. Ball, and M. J. Turbeville, *J. Org. Chem.*, **62**, 5945 (1997).
10. D. H. Hirsh, R. I. Hoaglin, and D. G. Kubler, *J. Org. Chem.*, **23**, 1083 (1958).
11. J. A. Hyatt, US Patent Application filed 2003.
12. M. Ihara, T. Takahashi, N. Shimizu, Y. Ishida, I. Sudow, K. Fukumoto, and T. Kametani, *J. Chem. Soc., Perkin Trans. I*, 529 (1989).
