

- constants for deuterated media are substantially higher than the true values. Moreover, the amount of protium is likely to change in the course of reaction.
- (25) Wojcicki, A., personal communication.
- (26) Christian, S. D.; Stevens, T. L. *J. Phys. Chem.* **1972**, *76*, 2039.
- (27) Bell, R. P. "The Proton in Chemistry", 2nd ed.; Methuen: London, 1975; Chapter 11.
- (28) Whitesides, T. H.; Neilan, J. P. *J. Am. Chem. Soc.* **1973**, *95*, 907.
- (29) Flood, T. C.; Miles, D. L. *J. Organomet. Chem.* **1977**, *127*, 33.

- (30) Flood, T. C.; Rosenberg, E.; Sarhangi, A. *J. Am. Chem. Soc.* **1977**, *99*, 4334, have described related hydrido species with other metal complexes in acid media.
- (31) Anderson, S. N.; Fong, C. W.; Johnson, M. D. *J. Chem. Soc., Chem. Commun.* **1973**, 163.
- (32) The formation of hydroperoxide was confirmed by its preparation through reaction of benzylcobaloxime with oxygen at 60 °C and then with 2% trifluoroacetic acid.³³
- (33) Bied-Charreton, C.; Gaudemer, A. *Tetrahedron Lett.* **1976**, 4153.

Lewis Acid Mediated Reactions of Organocopper Reagents. A Remarkably Enhanced Regioselective γ -Attack of Allylic Halides and Direct Alkylation of Allylic Alcohols via $\text{RCu}\cdot\text{BF}_3$ System¹

Yoshinori Yamamoto,* Shinichi Yamamoto, Hidetaka Yatagai, and Kazuhiro Maruyama

Contribution from the Department of Chemistry, Faculty of Science, Kyoto University, Kyoto 606, Japan. Received August 29, 1979

Abstract: Chemical reactivities and selectivities of a new class of organocopper reagent, RCu -Lewis acid, are described. Regioselective γ -attack of allylic halides is realized irrespective of the degree of substitution at the two ends of the allylic systems, and of the structural factors (cyclic or acyclic) involved. Among the Lewis acids examined, $\text{BF}_3\cdot\text{OEt}_2$ is the most effective with respect to the selectivity and total yield. Propargyl chloride and acetate are converted into 1,2-heptadiene by $n\text{-BuCu}\cdot\text{BF}_3$. Allylic alcohols react with 3 equiv of $\text{RCu}\cdot\text{BF}_3$ to produce the corresponding alkylation products in high yield. The stereochemistry of the reactions of $\text{RCu}\cdot\text{BF}_3$ is examined by using Goering's system, that is, 5-methyl-2-cyclohexenyl chloride, acetate, and alcohol. The substitution proceeds through a formal anti $\text{S}_{\text{N}}2'$ in the case of *cis*-5-methyl-2-cyclohexenyl acetate and through a formal syn $\text{S}_{\text{N}}2'$ in *trans*-5-methyl-2-cyclohexen-1-ol. On the other hand, the stereochemical integrity disappears in the reaction of *cis*-5-methyl-2-cyclohexen-1-ol and the chloride (1). It is proposed that the "ate" complex between RCu and BF_3 is involved as a reactive intermediate.

An important and yet frequently elusive goal of synthetic chemistry is the selective synthesis of a desired stereo- or regioisomer. Examples illustrating the importance of regioselective syntheses are manifold, control of the regiochemistry in the reaction of allylic carbanions with electrophiles being one example² while regiochemical control of the reaction of allylic substrates with nucleophiles is another.³ In general, substitution reactions of allylic substrates without (or with) complete allylic rearrangement via organometallics are still unpredictable processes.^{4,5} Among such reactions, several promising results to direct nucleophiles to the α position are obtained. Regioselective α -attack is achieved by using allylic substrates containing oxygen in the leaving group (OAc, OTos, etc.),⁶⁻¹¹ although isomerization, rearrangement, and formation of side products in many cases are reported.^{8,10,11} On the other hand, in the case of the reaction of allylic halides⁷ with organolithium and magnesium compounds, there are difficulties on the points of allylic transposition, geometrical isomerization, and cyclization.¹²⁻¹⁶ Several literature sources refer to prenyl halides, cyclic allylic halides, or branched allylic compounds in which cases the problem of partial rearrangement either does not exist or is overshadowed by attendant factors.^{4,17-19} Recently it was reported that acyclic allylic halides undergo cross coupling at the α position via dialkylcuprate- Me_2S system.²⁰

The efforts to direct nucleophiles to the γ position are relatively scarce. Although γ -unsubstituted allylic substrates undergo a facile γ -alkylation,^{10,21-23} the substitution appears to be controlled by steric factors.^{19g-i} Recently exclusive syn γ -attack on the allylic system of 5-phenyl-2-cyclohexenyl carbamates was reported.²⁴ Consequently, the regiochemistry

is highly system dependent²⁵ and the influencing factors are the nature of the leaving group, the degree of substitution of the two ends of the allylic system, the steric factors, the solvent system, and the nature of the nucleophile.

Previously we reported preliminary results dealing with regioselective γ -attack toward allylic halides by $\text{RCu}\cdot\text{BF}_3$ and direct alkylation of allylic alcohols.¹ We now report the full details of that work as well as the stereochemical aspects via $\text{RCu}\cdot\text{BF}_3$. The results reported here indicate that (1) regioselective γ -attack is realized irrespective of the nature of the leaving group, the degree of substitution at the two ends of the acyclic allylic systems,²⁶ and the structural factors (cyclic or acyclic), (2) direct displacement of the OH group of allylic alcohols is achieved via $\text{RCu}\cdot\text{BF}_3$, (3) the substitution proceeds through a formal anti $\text{S}_{\text{N}}2'$ pathway in the case of *cis*-5-methyl-2-cyclohexenyl acetate and syn $\text{S}_{\text{N}}2'$ in *trans*-5-methyl-2-cyclohexenol, and (4) the stereochemical integrity disappears in the reaction of *cis*- and *trans*-5-methyl-2-cyclohexenyl chloride and *cis*-5-methyl-2-cyclohexen-1-ol (eq 1).


Results and Discussion

Regiochemistry of the Reaction of Allylic Halides with Organocopper Reagents in the Presence of Lewis Acids. As we previously reported, certain observations suggest that the regiochemistry in the substitution reaction of allylic substrates with organocopper reagents must be influenced by the presence of organoboranes.^{1a,27} Therefore, systematic investigation on the regiochemistry of the reaction of cinnamyl halides with organocopper reagents in the presence of various Lewis acids was carried out (Table I). As evident, an enhanced γ regiose-

Table I. Reaction of Cinnamyl Chloride with Organocopper Reagents^a

entry	reagent (RM)	solvent	products, % ^b	
			PhCH(R)CH=CH ₂ (γ)	PhCH=CHCH ₂ R (α)
1	MeCu	ether	77 (74) ^c	23 (26) ^c
2	Me ₂ CuLi	ether	(30) ^c	(70) ^c
3	MeCu·BEt ₃	ether	87	13
4	MeCu·B(<i>n</i> -Bu) ₃	ether	89 (90) ^c	11 (10) ^c
5	MeCu·B(OCH ₃) ₃	ether	79	21
6	MeCu·BF ₃	ether	89 (90) ^c	11 (10) ^c
7	MeCu·BF ₃	THF	99.5	0.5
8	MeCu·BCl ₃	THF	99	1
9	<i>n</i> -BuCu	ether	52	48
10	<i>n</i> -BuCu	ether, LiI free ^d	56	44
11	<i>n</i> -BuCu·B(<i>n</i> -Bu) ₃	ether	87	13
12	<i>n</i> -BuCu·BF ₃	ether	91	9
13	<i>n</i> -BuCu·BCl ₃ ^e	ether	88	12
14	<i>n</i> -BuCu·TiCl ₄ ^e	ether	94	6
15	<i>n</i> -BuCu·AlCl ₃	ether	67	33
16	<i>n</i> -BuCu·AlEt ₃	ether	70	30
17	<i>n</i> -BuCu·B(<i>n</i> -Bu) ₃	THF	99.5	0.5
18	<i>n</i> -BuCu·BF ₃	THF	99.7	0.3
19	<i>n</i> -BuCu·BCl ₃	THF	99	1
20	<i>n</i> -BuCu·TiCl ₄	THF	<i>f</i>	<i>f</i>
21	<i>n</i> -BuCu from <i>n</i> -BuMgCl	THF	42	58
22	<i>n</i> -BuCu·BF ₃ from <i>n</i> -BuMgCl	THF	64	36
23	<i>n</i> -Bu ₂ CuLi	THF	82	18
24	<i>n</i> -BuMgCl	THF	1	99
25	PhCu	ether	84 (86) ^c	16 (14) ^c
26	PhCu·BF ₃	ether	84 (81) ^c	16 (19) ^c
27	PhCu·BEt ₃	ether	75	25
28	PhCu·BCl ₃ ^e	ether	79	21
29	PhCu·BCl ₃ ^e	THF	50	50

^a All reactions were carried out on a 1-mmol scale with the same procedure as described in the Experimental Section. Normally, cinnamyl chloride was used except where otherwise indicated. ^b By GLC. Total yields were usually excellent except where otherwise indicated. Only small amounts of byproducts (<4%), such as reduction products, the dimer of cinnamyl unit, cinnamyl iodide, and/or RI (alkyl iodide), were detected in some cases. ^c Cinnamyl bromide was used. ^d *n*-BuCu·P(*n*-Bu)₃ complex free from LiI was prepared as previously described: Whitesides, G. M.; Casey, C. P.; Krieger, J. K. *J. Am. Chem. Soc.* **1971**, *93*, 1379. ^e Total yields were ca. 60%. ^f Cinnamyl chloride was recovered.



lectivity is observed in alkylcopper–Lewis acid systems. $RCu \cdot BF_3$ is the most effective with respect to the selectivity and total yield among the Lewis acids examined (entries 6, 7, 12, and 18). THF is superior to ether for directing to the γ position, though titanium tetrachloride complex in THF gives quite an unsatisfactory result (entry 20). It is noteworthy that both *n*-BuCu and *n*-BuCu·BF₃ from *n*-BuMgCl are less reg-

ioselective even in THF than those from *n*-BuLi (entries 21 and 22). The result of entry 24 suggests a simple and very promising procedure for directing alkyl groups to the α position; actually such a reaction is frequently carried out in an ether solvent.¹⁶ Unfortunately, the regiochemical control of phenylcopper systems (entries 25–29) is not successful. This may be due to the complex formation between BF₃ and the π electrons of the aromatic ring. However, utilization of 2 equiv of BF₃ does not exert any appreciable influence on the regiochemistry.

Of particular interest is the question whether the enhanced γ regioselectivity will be a reflection of the complex formation between Lewis acids and LiI present in the reaction mixture. It seems that the difference between entries 9 and 10 is negligible, though the comparison is not completely adequate because of the presence of P(*n*-Bu)₃ in the latter. The effect of excess amounts of LiI upon the regioselectivity can be ignored (see also Table II); the dimeric product of the cinnamyl unit slightly increased in the presence of LiI. Further, if such a complex formation is an important factor, PhCu·BF₃ system should be equally effective. Since I[−] is a soft base, a relatively soft acid (BR₃) will form a stronger complex with LiI than a hard acid (BF₃).²⁸ This leads to an expectation that $RCu \cdot BR'_3$ must direct the R group to the γ position with greater regioselectivity than $RCu \cdot BF_3$. However, the experiment gives a reversed result (entry 11 vs. 12). These considerations suggest that other factors, except (or in addition to) complex formation, play a major role in the control of the regiochemistry.

Reaction of various allylic halides with organometallics was examined to establish a generality of the enhanced γ regioselectivity of the $RCu \cdot BF_3$ system (Table II). In contrast to cinnamyl chloride, a remarkable salt effect of LiI is observed

Table II. Reaction of Allylic Halides with Organometallic Reagents^a

entry	allylic halide/RM	solvent	γ product, % ^b	α product, % ^b
	$\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{CH}_2\text{Cl} \end{array}$		$\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{CH} \\ \diagup \\ \text{R} \end{array} \text{CH}=\text{CH}_2$	$\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{CH}_2\text{R} \end{array} \quad (t/c)^f$
1	<i>n</i> -BuLi ^d	ether	46	54
2	<i>n</i> -BuMgCl	THF	10	90 (100/~0)
3	<i>n</i> -Bu ₂ CuLi	ether	4	96 (99/1)
4	<i>n</i> -BuCu	ether	68	32
5	<i>n</i> -BuCu	THF	79	21
6	<i>n</i> -BuCu, 2 equiv of LiI ^e	THF	30	70 (85/15)
7	<i>n</i> -BuCu·B(<i>n</i> -Bu) ₃	ether	73	27
8	<i>n</i> -BuCu·BF ₃	ether	94	6
9	<i>n</i> -BuCu·BF ₃	THF	98	2
10	<i>n</i> -BuCu·TiCl ₄ ^f	ether	98	2
11	PhCu	ether	84	16 (97/3)
12	PhCu·BF ₃ ^d	ether	84	16
	$\begin{array}{c} \text{CH}_2=\text{CHCHCH}_3 \\ \\ \text{Cl} \end{array}$		$\begin{array}{c} \text{RCH}_2 \\ \diagdown \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{H} \quad \text{CH}_3 \end{array} \quad (t/c)^f$	$\begin{array}{c} \text{CH}_2=\text{CHCHCH}_3 \\ \\ \text{R} \end{array}$
13	<i>n</i> -BuLi ^d	ether	92 (82/18)	8
14	<i>n</i> -BuMgCl	THF	79 (40/60)	21
15	<i>n</i> -Bu ₂ CuLi	ether	78 (54/46)	22
16	<i>n</i> -BuCu	ether	90 (42/58)	10
17	<i>n</i> -BuCu·BF ₃	ether	96 (55/45)	4
18	<i>n</i> -BuCu·BF ₃	THF	~100 (80/20)	
19	<i>n</i> -BuCu·TiCl ₄ ^f	ether	97 (70/30)	3
20	Ph ₂ CuLi	ether	81 (58/42)	19
21	PhCu	ether	94 (57/43)	6
22	PhCu·BF ₃	ether	98 (59/41)	2
	CH ₃ CH=CHCH=CHCH ₂ Cl		CH ₃ CH=CHC(R)HCH=CH ₂	CH ₃ CH=CHCH=CHCH ₂ R
23	<i>n</i> -BuCu·BF ₃	THF	~100	CH ₃ C(R)HCH=CHCH=CH ₂ ^g
	$\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{CH}_2\text{Br} \end{array}$		$\begin{array}{c} (\text{CH}_3)_2\text{CCH}=\text{CH}_2 \\ \\ \text{R} \end{array}$	$\begin{array}{c} \text{CH}_3 \\ \diagdown \\ \text{C}=\text{C} \\ \diagup \quad \diagdown \\ \text{CH}_3 \quad \text{CH}_2\text{R} \end{array}$
24	MeLi ^d	ether	12	88
25	Me ₂ CuLi	ether	trace	~100
26	MeCu	ether	60	40
27	MeCu·B(<i>n</i> -Bu) ₃	ether	80	20
28	MeCu·BF ₃ (1:0.5) ^h	ether	74	26
29	(1:1) ^h	ether	94	6
30	(1:2) ^h	ether	95	5
31	MeCu·BF ₃	THF	95	5
32	MeCu·BCl ₃ ^d	THF	90	10
33	<i>n</i> -Bu ₂ CuLi	ether	trace	~100
34	<i>n</i> -BuCu	ether	79	21
35	<i>n</i> -BuCu·B(<i>n</i> -Bu) ₃	ether	90	10
36	<i>n</i> -BuCu·BF ₃	ether	90	10
37	<i>n</i> -BuCu·BCl ₃	THF	67	33
38	PhLi	ether	trace	~100
39	PhCu	ether	80	20
40	PhCu·BF ₃	ether	80	20

^a All reactions were carried out on a 1-mmol scale. Total yields were usually excellent except where otherwise indicated. ^b By GLC. ^c Trans/cis ratio. ^d Total yields were about 50–60%. ^e See footnotes to Table I. ^f Considerable amounts of byproducts were detected. ^g Not detected. ^h The ratio of MeCu to BF₃.

in crotyl chloride (entries 5 and 6). Importance of the ratio of RCu to BF₃ is clearly indicated in entries 28–30; the ratio of 1:1 is essential. Here also, RCu·BF₃ in THF is the most effective and such an enhanced regioselectivity is not detected or is negligible in the case of PhCu (entries 11 and 12, 21 and 22, and 39 and 40). Consequently, it is now clear that highly regioselective γ -attack is accomplished generally via RCu·BF₃ irrespective of the degree of substitution of the two ends of the allylic system.

Regiochemistry of the Reaction of Propargyl Chloride and Acetate. The similar α/γ problem arises in the reaction of

propargyl derivatives with organometallics. Although the regiochemistry is also system dependent, it is generally recognized that the reaction of organocopper reagents proceeds through γ -attack (allene formation).^{29–31} In some cases, direct substitution (S_N2 type) to produce acetylenic derivatives is realized by blocking the terminal position of the acetylene with the Me₃Si group.³² On the other hand, widely divergent results are reported in the reaction with Grignard reagents or organolithium.³³ Accordingly, the reaction of propargyl chloride, acetate, and alcohol with RCu·BF₃ was briefly examined; the chloride and acetate were converted into the corresponding

Table III. Reaction of Allylic Alcohols and Acetate with RCu -Lewis Acids^a

entry	substrate (1 mmol)/RM (mmol)	γ product, % ^b	α product, % ^b	total yield, % ^b
1	$\text{C}_6\text{H}_5\text{CH}=\text{CHCH}_2\text{OH}$ $n\text{-BuCu}\cdot\text{BF}_3$ (1)	$\text{C}_6\text{H}_5\text{CH(R)CH}=\text{CH}_2$ 83	$\text{C}_6\text{H}_5\text{CH}=\text{CHCH}_2\text{R}$ 17	17
2	$n\text{-BuCu}\cdot\text{BCl}_3$ (1)	<i>c</i>	<i>c</i>	<i>c</i>
3	$n\text{-BuCu}\cdot\text{BF}_3$ (2)	91	9	66
4	$n\text{-BuCu}\cdot\text{BF}_3$ (3)	91	9	97
5	$n\text{-BuCu}\cdot\text{TiCl}_4$ (3)	trace	trace	trace
6	$\text{MeCu}\cdot\text{BF}_3$ (3)	95	5	96
7	$n\text{-BuCu}$ (2)	<i>c</i>	<i>c</i>	<i>c</i>
8	$n\text{-Bu}_2\text{CuLi}$ (2)	<i>c</i>	<i>c</i>	<i>c</i>
9	$\text{CH}_3\text{CH}=\text{CHCH}_2\text{OH}$ $n\text{-BuCu}\cdot\text{BF}_3$ (3)	$\text{CH}_3\text{CH(R)CH}=\text{CH}_2$ 86	$\text{CH}_3\text{CH}=\text{CHCH}_2\text{R}$ 14	99
10	$\text{CH}_2=\text{CHCH}(\text{OH})\text{CH}_3$ $n\text{-BuCu}\cdot\text{BF}_3$ (2)	$\text{RCH}_2\text{CH}=\text{CHCH}_3$ 90 (55/45) ^d	$\text{CH}_3\text{CH(R)CH}=\text{CH}_2$ 10	44
11	$n\text{-BuCu}\cdot\text{BF}_3$ (3)	94 (55/45) ^d	6	86
12	$\text{CH}_3\text{CH}=\text{CHCH}=\text{CHCH}_2\text{OH}$ $n\text{-BuCu}\cdot\text{BF}_3$ (2)	$\text{CH}_3\text{CH}=\text{CHC(R)HCH}=\text{CH}_2$ 100	$\text{CH}_3\text{CH}=\text{CHCH}=\text{CHCH}_2\text{R}$ $\text{CH}_3\text{CH(R)CH}=\text{CHCH}=\text{CH}_2$ <i>e</i>	70
13	$(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{OH}$ $n\text{-BuCu}\cdot\text{BF}_3$ (3)	$(\text{CH}_3)_2\text{C(R)CH}=\text{CH}_2$ 98	$(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{R}$ 2	60
14	$(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}=\text{CH}_2$ $n\text{-BuCu}\cdot\text{BF}_3$ (3)	$(\text{CH}_3)_2\text{C}=\text{CHCH}_2\text{R}$ 20	$(\text{CH}_3)_2\text{C(R)CH}=\text{CH}_2$ 80	85
15	$\text{C}_6\text{H}_5\text{CH}=\text{CHCH}_2\text{OAc}$ $n\text{-Bu}_2\text{CuLi}$ (1)	$\text{C}_6\text{H}_5\text{CH(R)CH}=\text{CH}_2$ trace	$\text{C}_6\text{H}_5\text{CH}=\text{CHCH}_2\text{R}$ 99	50
16	$n\text{-BuCu}\cdot\text{BCl}_3$ (1)	60	40	23
17	$n\text{-BuCu}\cdot\text{BF}_3$ (2)	75	25	78
18	$n\text{-BuCu}\cdot\text{BF}_3$ (3)	62	38	92
19	$n\text{-BuCu}\cdot\text{AlCl}_3$ (3)	68	32	85

^a All reactions were performed on a 1-mmol scale in ether. ^b By GLC analysis. The stereochemistry of the minor product was not determined.^c Not detected. ^d Trans/cis ratio. ^e The α - and ω -alkylated products were not detected.

allene ($\text{R} = n\text{-Bu}$) in high yields and 1-heptyne was not detected. In contrast to the allylic alcohols, the reaction of propargyl alcohol was quite sluggish even in the presence of excess $n\text{-BuCu}\cdot\text{BF}_3$ and only trace amounts of the allene were detected with GLC.

Reaction of Allylic Alcohols and Their Derivatives with RCu -Lewis Acids. The reaction of allylic acetates with RCu -Lewis acids was examined as a logical extension of the work on allylic halides. During this study, a new reaction was encountered, involving the replacement of the hydroxy group of allylic alcohols by an alkyl function in the reaction of $\text{RCu}\cdot\text{BF}_3$ with such alcohols. The results are summarized in Table III. Although the reaction of allylic alcohols with 1 or 2 equiv of $n\text{-BuCu}\cdot\text{BF}_3$ did produce the alkylation products (entries 1, 3, and 10), the best result was obtained when 3 molar equiv of the reagent was used. As expected, the corresponding alkylcopper and cuprate reagents did not give the desired products at all (entries 7 and 8). Use of other Lewis acids such as BCl_3 (entry 2), $\text{B}(n\text{-Bu})_3$, and TiCl_4 (entry 5) gave quite unsatisfactory results. It was essential to use ether as a solvent; utilization of THF led to poor results. The reaction of cinnamyl acetate with $n\text{-BuCu}\cdot\text{BF}_3$ or $n\text{-BuCu}\cdot\text{AlCl}_3$ also gave the alkylation products (entries 16–19), but the regioselectivity was relatively low.

Some other features of the data in Table III deserve comment. Crotyl alcohol and its isomer, 1-buten-3-ol, are converted into 3-methyl-1-heptene and 2-octene, respectively (entries 9–11). Therefore, the reaction of these alcohols does not involve a common intermediate such as 1-methylallyl cation or radical. A similar result is obtained in the case of crotyl chloride and its isomer (Table II, entries 7–9 vs. 17–19). In contrast to these alcohols, both prenyl alcohol and its isomer, 3-methyl-1-buten-3-ol, gave 3,3-dimethyl-1-heptene as a major product (entries 13 and 14). This suggests that the latter alcohol undergoes isomerization to prenyl alcohol under the reaction conditions. Another possibility is that the latter alcohol, sub-

stituted with an allyl and two methyl groups, undergoes facile ionization under the influence of BF_3 (or $\text{RCu}\cdot\text{BF}_3$) and the resulting 1,1-dimethylallyl cation is immediately trapped with RCu (or $\text{RCu}\cdot\text{BF}_3$).³⁴

In connection with these observations, the role of BF_3 was investigated. It is known that the complex formation between alcohols and BF_3 is a facile process.³⁵ Therefore, it might be thought that complexes can react with an alkylating reagent. However, this possibility was eliminated, though not completely, by a control experiment.^{1b} Consequently, as mentioned later, the reaction must involve a cyclic transition state since $\text{S}_{\text{N}}2'$ -type substitution occurs even in a positionally unbiased system. Although the reaction of allylic alcohols proceeds with less regioselectivity than that of allylic halides, direct alkylation of allylic alcohols via an organocopper reagent is the first example³⁶ and synthetically useful; allylic alcohols are easily handled in comparison with allylic halides, which often are highly reactive.


The reaction of other allylic substrates with $\text{RCu}\cdot\text{BF}_3$ was also examined. Allyl phenyl ether was converted into 1-heptene (26% yield) under the influence of 3 equiv of $n\text{-BuCu}\cdot\text{BF}_3$. Similar reaction of allylamine, allylammonium bromide, and allyl sulfide gave trace amounts of 1-heptene. Phenyl iodide, benzyl chloride, cyclohexyl bromide, 1-bromoadamantane, and benzyl alcohol did not react with excess amounts of $n\text{-BuCu}\cdot\text{BF}_3$, while n -amyl iodide and diphenylchloromethane gave the corresponding butylated products in 10–20% yields. Consequently, it is possible to carry out a chemoselective alkylation of the allylic and propargyl derivatives in the presence of these functional groups.

Stereochemical Aspects. Stereochemistry of the $\text{S}_{\text{N}}2'$ reaction has been of synthetic and mechanistic interest for many years.^{3,37} In this connection, stereochemistry of the formal $\text{S}_{\text{N}}2'$ reaction via an organometallic is of current interest. The mode of substitution is shown to be exclusively anti (trans) in both cyclohexenyl systems^{19g,h,23c} and acyclic allylic ethers,^{23b} while

Table IV. Reaction of 5-Methyl-2-cyclohexenyl Derivatives with Organocopper Reagents^a

entry	cyclohexenyl derivative (cis/trans ratio)	product (<i>cis</i> -4/ <i>trans</i> -4 ratio)		
		<i>n</i> -BuCu-BF ₃	<i>n</i> -Bu ₂ CuLi	<i>n</i> -BuCu
1	<i>cis</i> -1 (97/3)	60/40	65/35	51/49
2	<i>cis</i> - and <i>trans</i> -1 (60/40)	67/33		
3	<i>cis</i> - and <i>trans</i> -1 (37/63)	67/33		
4	<i>cis</i> -2 (90/10)	23/77	12/88	
5	<i>cis</i> -3 (97/3)	59/41		
6	<i>trans</i> -3 (17/83)	22/78		
7	<i>cis</i> -2- γ -d	4- α -d		
8	<i>cis</i> -3- γ -d	4- α -d		
9	<i>cis</i> -3- α -d	4- γ -d		

^a All reactions were carried out as described in the Experimental Section. Three equivalents of *n*-BuCu-BF₃ was used in the reaction of 3; the product (4) was obtained in 47–52% yields. The reaction of 1 proceeded smoothly and the product was obtained in high yield. Deuterium was labeled as indicated (>98% d). *cis*-2- γ -d indicates that deuterium is substituted at γ position of *cis*-2, and analogous notation is used for other systems.



exclusive syn (*cis*) γ -attack on the cyclohexenyl carbamate systems is reported.²⁴ Therefore, we examined the stereochemistry of the formal S_N2' reaction via RCu-BF₃ by using Goering's cyclohexenyl system. The results are listed in Table IV.


The reaction of the allylic halide (1) proceeds with complete loss of the stereochemistry (entries 1–3) regardless of the reagent types. This may be due to a labile characteristic of chloride as a leaving group, which leads to an allylic intermediate or which causes a facile isomerization of the starting materials under the reaction conditions. Although it is now clear that the stereochemical integrity disappears in the reaction of 1, the regiochemical integrity (S_N2' type) must be presumably retained as is apparent from the results of reactions of crotyl chloride and its isomer (Table II). Unfortunately, 1- α -d or 1- γ -d could not be prepared in reasonably pure form. The reaction of *cis*-2 with *n*-BuCu-BF₃ or *n*-Bu₂CuLi proceeds via anti S_N2' as would be expected from the literature^{19g,h,23b,c} (entries 4 and 7).

The reaction of 3 is a puzzling system; the stereochemical integrity of *cis*-3 disappears in a similar manner as that of 1 (entry 5), while *trans*-3 retains its configuration (entry 6). In both cases, the regiochemical integrity (S_N2' type) is retained (entries 8 and 9). Examination of the composition of the recovered alcohols revealed that the *trans* isomer reacted faster

than the *cis* isomer. These observations are tentatively explained in terms of steric hindrance during the attack of the reagent toward *cis*-3.


For the case of a cyclohexenyl derivative disposed in a half-chair conformation, the bonds undergoing rupture and formation are both quasi-axial. Further, it is reasonable to assume a cyclic transition state since a formal S_N2' is observed in this positionally unbiased system (entries 8 and 9). Therefore, *trans*-3 may be expected to be less sterically hindered and more energetically favored than *cis*-3, so that it would react faster than *cis*-3. Thus, *trans*-3 leads readily to *trans*-4 via a syn S_N2' type reaction. On the other hand, *cis*-3 is more sterically hindered and less energetically favored and thus may be energetically similar to or higher than *cis*-3, *quasi-equatorial*, leading to the mixture of *cis*- and *trans*-4. Consequently, the stereochemistry is more system dependent than the regiochemistry, at least in the present system, and will be determined by the nature of the leaving group and conformational and configurational factors.^{38,39}

Concerning the Reactive Intermediate. The enhanced γ regioselectivity, the direct alkylation of allylic alcohols, and the stereochemical aspects make it desirable to obtain better insight into the reactive intermediate. Treatment of cinnamyl chloride with BF₃·OEt₂ followed by the addition of *n*-BuCu exhibited similar regioselectivity as the reaction of normal *n*-BuCu free from BF₃, though total yields of butylated products became low. Interaction between BF₃ and LiI is not a major factor for the high regioselectivity as already mentioned. Consequently, complex formation between RCu and BF₃ is most probable, and the present observations seem to be a reflection of such a complex. Although the coordination of BF₃ with donor atoms, such as N, O, P, and S, is well known,^{35,40} the complexes between BF₃ and carbon atom as a donor are relatively scarce. To our knowledge, the complexes reported so far are [Me₃Sn]⁺[CF₃BF₃][−]⁴¹ and K[CF₃BF₃],⁴² the structure of which is determined by X-ray diffraction methods. Complexes between basic transition metal organometallics and BF₃ are also reported.⁴³ Since it is widely accepted that lithium, sodium, and magnesium borates are formed from the reaction of alkylolithium, sodium, and magnesium compounds with trialkylboranes,⁴⁴ it is reasonable that the similar ate complexes of copper are produced from the reaction between alkylcoppers and trialkylboranes.^{27,45} Actually, it is reported that copper methyltrialkylborates, prepared from the reaction of lithium trialkylborates with copper halides, exhibit high γ regioselectivity (96%) toward cinnamyl




chloride.⁴⁶

Therefore, it is proposed that the complex $RCu \cdot BF_3$ possesses an ate complex type structure, $RBF_3^-Cu^+$, at low temperature. Since CuF is unknown,⁴⁷ this ate complex presumably does not give RBF_2 easily.



The reaction mixture of n -BuCu with BF_3 at $-78^\circ C$ was allowed to warm to room temperature. GLC examination of the mixture revealed the formation of only trace amounts of tri- n -butylborane. This is a marked contrast to the reaction of alkylmagnesium and lithium compounds with BF_3 , which gives excellent yields of trialkylboranes.⁴⁸ The oxidation of the mixture with H_2O_2 -NaOH gave only trace amounts of n -BuOH.⁴⁹ These results indicate that the compounds having n -Bu-boron bonds, such as n -Bu₃B and n -BuBF₂, are not involved in the mixture. Instead, $RCu \cdot BF_3$ or more preferably $RBF_3^-Cu^+$ presumably dissociates into RCu and BF_3 upon heating. Preparation of n -BuCu \cdot BF₃ at $-30^\circ C$ and subsequent reaction with cinnamyl chloride exhibited the same regioselectivity as that at $-70^\circ C$. Thus, the complex is stable at least at $-30^\circ C$. Without further knowledge of the exact structure of $RCu \cdot BF_3$, it is difficult to postulate any convincing mechanism for the reaction. However, the following results strongly suggest involvement of the ate complex as an intermediate.

Dark green ate complex **5**, prepared from 1-octenyl-9-BBN and methylcopper in ether at $0^\circ C$, is stable at $0^\circ C$ left overnight.⁵⁰ This complex does not dissociate into 1-octenylcopper and methyl-9-BBN or into 1-octenyl-9-BBN and methylcopper, while copper bis(1-hexenyl)di- n -butylborate undergoes a facile dissociation into 1-hexenylcopper.^{27,45} The regiochemical behavior of **5** to crotyl chloride was compared with



that of other related copper derivatives. The results are summarized in Table V.⁵¹ The ate complex **5** exhibited higher regioselectivity than free 1-octenylcopper (entry 1 vs. 5). The mixture between 1-octenylcopper and n -butyl-9-BBN showed a similarly high regioselectivity (entry 3). On the other hand, the regioselectivity of **6** was close to that of free 1-octenylcopper (entry 4 vs. 5), suggesting a facile dissociation of **6** into 1-hexenylcopper. Consequently, it is now clear that formation of ate complexes causes an enhanced γ regioselectivity. Finally, the postulation of the cyclic transition state, involving the pentavalent aluminate in the reduction of certain allylic halides with $LiAlH_4$,^{39,52} should be recalled when the mechanism of the present reactions is considered.

Experimental Section

¹H NMR spectra were recorded on a JEOL JNM-MH-100 instrument; chemical shifts (δ) are expressed in parts per million relative to Me₄Si. IR spectra were recorded on a JASCO IRA-1 spectrophotometer. Mass spectra were recorded on a Hitachi GC-M-52 instrument (22 eV). Elemental analyses were performed by the Kyoto University Microanalytical Laboratories, and the results are within the accepted limits ($\pm 0.3\%$). All temperatures were uncorrected.

Reagents. Reagent grade solvents were purified by standard techniques and kept over a drying agent. Methylolithium in ether was prepared by standard procedure,^{53a} and n -butyllithium in hexane was

Table V. Reaction of Alkenylcopper Derivatives with Crotyl Chloride^a

entry	copper derivative	γ -attack, %	α -attack, %
1	5	87.5	12.5
2	5 + 3-chloro-1-butene ^b	87.5	12.5
3	n -Hex-CH=CH-Cu + n -Bu-BBN	84	16
4	n -Bu-CH=CH-Cu + $Cu^+ 6^-$	73	28
5	n -Hex-CH=CH-Cu	77.5	22.5

^a All reactions were carried out in ether at $-30^\circ C$. The ratio was determined by GLC. ^b Instead of crotyl chloride, 3-chloro-1-butene was used. ^c Prepared from bis(1-hexenyl)chloroborane with 3 equiv of MeCu.²⁷

a commercial product. Phenyllithium^{53b} and n -butylmagnesium chloride^{53c} were prepared. The titrations were performed by Gilman's⁵⁴ and Eastham's⁵⁵ methods. Cuprous iodide was purchased and purified.⁵⁶ Organoboranes, such as B(n -Bu)₃, n -Bu-9-BBN, and 1-octenyl-9-BBN, were prepared by the reported procedures.⁵⁷ Other simple chemicals were purchased and used as such. LiI was prepared in situ from the reaction of n -BuLi with MeI since it was highly hygroscopic.

Reaction of RCu -Lewis Acids with Allylic Halides and Related Compounds. The following procedure for the reaction of cinnamyl chloride with n -BuCu \cdot BF₃ is representative. In a 200-mL flask, equipped with a magnetic stirrer and maintained under N₂, were placed 1.9 g (10 mmol) of CuI and 20 mL of dry THF. n -BuLi in hexane (1.3 M, 10 mmol) was added at $-30^\circ C$ and the resulting mixture (black) was stirred at this temperature for 5 min. The mixture was then cooled to $-70^\circ C$ and BF₃ \cdot OEt₂ (47%, 1.3 mL, 10 mmol) was slowly added. The color changed from black to very black. After the mixture was stirred for a few minutes, cinnamyl chloride (1.53 g, 10 mmol) was slowly added by a syringe, and the mixture was allowed to warm to room temperature by stirring. The mixture was directly analyzed by GLC or was filtered through the column of alumina using light petroleum ether. The olefin thus obtained in an essentially pure form was distilled under reduced pressure through Kugelrohr: 1.64 g (3-phenyl-1-heptene), 94%, bp $65-66^\circ C$ (5 mmHg).

The same procedure was employed for other Lewis acids and allylic halides. Boron trichloride and aluminum trichloride were dissolved in ether and these solutions were added to RCu at $-70^\circ C$. Triethylaluminum in hexane solution was purchased and used as such. Although the reaction of PhCu-Lewis acids was carried out as described above, a slightly different condition was used in the reaction of MeCu-Lewis acids; MeCu was prepared at $0^\circ C$ (yellow) and then cooled to $-70^\circ C$. The product ratio was determined by GLC using a column of DC550 (2 m). In some cases, the GC-mass instrument was used for the analysis (SE-30, 1 m). Prenyl bromide was prepared from the reaction of isoprene with HBr. 2,4-Hexadienyl chloride was prepared by Dr. Y. Naruta in our laboratories from the reaction of 2,4-hexadienyl-1-ol, which was purchased from Tokyo Kasei Co., Ltd., with Ph₃P-CCl₄.⁵⁸ The chloride was contaminated with the regioisomers even by using this procedure and purified by fractional distillations. The procedure via SOCl₂⁵⁹ or via PBr₃,⁶⁰ which had been used for the synthesis of the corresponding bromide, was less effective for the selective synthesis of the desired chloride: bp $55-58^\circ C$ (10 mmHg); n_D^{20} 1.5010 (lit.⁵⁹ 1.5000); ¹H NMR (CCl₄) δ 1.76 (3 H, d, J = 6 Hz), 3.96 (2 H, d, J = 6 Hz), 5.0-6.4 (4 H, m); m/e (M⁺) 116, 118. The reaction of 2,4-hexadienyl chloride with n -BuCu \cdot BF₃ was carried out as described above. GLC examination of the reaction mixture revealed that the reaction was very clean and a single peak appeared at the region expected for the butylated product. The usual workup followed by distillation gave 3- n -butyl-1,4-hexadiene (cis and trans mixture) in 83% yield; bp $100-103^\circ C$ (20 mmHg); IR (cm⁻¹, in CCl₄) 1640, 995, 970, 909; ¹H NMR (CCl₄) δ 0.90 (3 H, t, J = 7 Hz), 1.23 (6 H, bs), 1.62 and 1.63 (3 H, d, J = 6 Hz), 2.50 (1 H, m), 4.70-5.90 (5 H, m); m/e (M⁺) 138. Anal. Calcd for C₁₀H₁₈: C, 86.88; H, 13.12. Found: C, 86.68; H, 13.31.

The same procedure was used for the reaction of allylic alcohols and acetate. In some cases, excess amounts of RCu-Lewis acids were used (Table III). 3,3-Dimethyl-1-butene, 2-methyl-2-pentene, *cis*- and *trans*-2-octene, and *trans*-1-phenyl-2-butene were commercially available. Authentic samples of 3-phenyl-1-butene, *trans*-1-phenyl-1-butene, 3-phenyl-1-heptene, *trans*-1-phenyl-1-heptene, 3,3-diphenyl-1-propene, and 3-methyl-1-heptene were kindly given by Dr. Y. Tanigawa, which were prepared according to the reported methods (ref 11b and 23c). *trans*-1,3-Diphenylpropene,⁶¹ 2-methyl-2-octene,⁶² 2-methyl-4-phenyl-2-butene,⁶³ 3,3-dimethyl-1-heptene,⁶⁴ and 3-methyl-3-phenyl-1-butene⁶⁵ were prepared according to the previous literature. The spectroscopic characteristics of these authentic samples were compared with those of the reaction products listed in Tables I-III. Both materials were completely identical in all respects.

Stereochemical Aspects. The synthesis of the cyclohexenyl derivatives has been described in an earlier paper.⁶⁶ 5-Methyl-1,3-cyclohexadione was prepared from the reaction of ethyl crotonate with ethyl acetoacetate in the presence of sodium ethoxide,⁶⁷ mp 126–128 °C (50% yield) (lit.⁶⁷ 126–127 °C). Treatment of this dione with ethanol in the presence of *p*-toluenesulfonic acid produced 5-methyl-3-ethoxy-2-cyclohexenone in 90% yield, bp 108 °C (4 mmHg) (lit.⁶⁸ 83 °C (0.1 mmHg)). Reduction of ethoxycyclohexenone with LiAlH_4 in ether gave 5-methyl-2-cyclohexenone in 69% yield,⁶⁹ bp 87 °C (20 mmHg) (lit.⁶⁹ 74 °C (8 mmHg)). Further reduction with LiAlH_4 produced 5-methyl-2-cyclohexenol in 82% yield,⁷⁰ bp 82 °C (20 mmHg) (lit.⁷⁰ 74 °C (8 mmHg)). GLC examination using CW-20M (2 m) revealed that the ratio of *cis* to *trans* isomer was 93:7. On the other hand, the reduction with $\text{Al}(\text{O}-i\text{-Pr})_3$ gave the corresponding alcohol, in which *cis*:*trans* = 60:40, in 76% yield.⁷¹ Resolution of partially resolved *cis*- and *trans*-5-methyl-2-cyclohexenol was carried out via their nitrobenzoates as previously described.⁷⁰ 5-Methyl-2-cyclohexenyl chloride was prepared from the reaction of resolved or partially resolved *cis*- and *trans*-5-methyl-2-cyclohexenol with SOCl_2 ,⁶⁶ bp 71 °C (50 mmHg) (42% yield) (lit.⁶⁵ 65 °C (25 mmHg)).⁶⁶ The configurations of the isomeric chlorides were related to those of the isomeric alcohols; they were readily distinguished by the IR bands at 7.81, 10.93, and 13.03 μ for *cis* isomer and at 8.00, 9.52, and 13.4 μ for *trans* isomer.⁶⁶ Both isomers exhibited essentially identical ^1H NMR spectra. 5-Methyl-2-cyclohexenyl acetate was prepared from the corresponding alcohol and acetyl chloride in pyridine in 73% yield, bp 77–78 °C (20 mmHg) (lit.⁶⁶ 83–84 °C (25 mmHg)). The ratio of *cis* to *trans* isomer was determined by GLC using a DC 550 column (1 m). *cis*-2- γ -d, *cis*-3- γ -d, and *cis*-3- α -d were prepared from 5-methyl-3-ethoxy-2-cyclohexenone as described in an earlier paper.^{19a} Total deuterium contents were determined from mass spectra. The deuterium distribution was determined from the total deuterium and relative peak areas for the C_5 and the γ (or α) proton signals in expanded 220-MHz NMR spectra (Varian HR-220), and was found to be discretely labeled as indicated in Table IV. Preparation of deuterium-labeled *trans*- or *cis*-1 was unsuccessful; deuterium was scrambled at the α and γ position of 1 when *cis*-3- γ -d or *cis*-3- α -d was treated with SOCl_2 . Butylated product, 5-methyl-3-butylcyclohexene [bp 95 °C (20 mmHg); ^1H NMR (CCl_4) δ 0.95 (3 H, d, J = 7 Hz), 0.95 (3 H, t, J = 7 Hz), 1.28 (6 H, bs), 1.60–1.80 (3 H, d), 1.95 (3 H, m), 5.52 (2 H, m)], was reduced to 3-methyl-1-butylcyclohexane via the hydroboration-protonolysis procedure.⁵⁷ Authentic material of this hydrocarbon was prepared according to the reported procedure.⁷² The ratio of *cis* to *trans* isomer was determined by GLC using a capillary column (squalane, 0.25 i.d., 45 m) as previously reported.⁷³

Concerning the Reactive Intermediate. *trans*-1-Octenyl-9-borabicyclo[3.3.1]nonane was prepared as previously reported.^{57,74} bp 92–96 °C (0.03 mmHg); ^1H NMR (CCl_4) δ 0.96 (3 H, t, J = 6 Hz), 1.00–2.00 (22 H, m), 2.10–2.30 (2 H, m), 6.08 (1 H, d, J = 18 Hz), 6.68 (1 H, d-t, J = 18 and 6 Hz). *trans*-1-Octenylcopper was prepared from the corresponding lithium and CuI in ether. The reaction of these copper derivatives with crotyl chloride and its isomer was carried out as described above. 3-Methyl-(4*E*)-1,4-undecadiene (γ -attack product of 5 to crotyl chloride): bp 115–125 °C (20 mmHg); IR (cm^{-1} , direct) 1640, 990, 965, 910; ^1H NMR (CCl_4) δ 0.90 (3 H, t, J = 6 Hz), 1.04 (3 H, d, J = 6 Hz), 1.20–1.44 (8 H, m), 1.84–2.10 (2 H, m), 2.64–2.88 (1 H, m), 4.88 (1 H, d-d, J = 18 and 2 Hz), 5.20–5.36 (2 H, m), 5.70 (1 H, d-d-d, J = 18, 10, and 6 Hz). Anal. Calcd for $\text{C}_{12}\text{H}_{22}$: C, 86.66; H, 13.34. Found: C, 86.37; H, 13.33. (2*E*,5*E*)-2,5-Dodecadiene (α -attack product of 5 to crotyl chloride and the γ -attack product to 3-chloro-1-butene in which case (2*Z*,5*E*)

isomer was also produced):⁵¹ bp 125–130 °C (20 mmHg); IR (cm^{-1} , direct) 965; ^1H NMR (CCl_4) δ 0.90 (3 H, t, J = 6 Hz), 1.24–1.36 (8 H, m), 1.60–1.68 (3 H, m), 1.94–2.02 (2 H, m), 2.58–2.72 (2 H, m), 5.28–5.40 (4 H, m). Anal. Calcd for $\text{C}_{12}\text{H}_{22}$: C, 86.66; H, 13.34. Found: C, 86.63; H, 13.10. 3-Methyl-(4*E*)-1,4-nonadiene (γ -attack product of 6 to crotyl chloride): bp 80–90 °C (30 mmHg); IR (cm^{-1} , direct) 1640, 990, 965, 910; ^1H NMR (CCl_4) δ 0.93 (3 H, t, J = 6 Hz), 1.10 (3 H, d, J = 7 Hz), 1.16–1.78 (4 H, m), 1.83–2.16 (2 H, m), 2.66–3.16 (1 H, m), 5.00–5.40 (2 H, m), 5.56–5.71 (2 H, m), 5.83–6.50 (1 H, m). Anal. Calcd for $\text{C}_{10}\text{H}_{18}$: C, 86.88; H, 13.12. Found: C, 86.60; H, 13.22. (2*E*,5*E*)-2,5-Decadiene (α -attack product of 6 to crotyl chloride): bp 80–90 °C (30 mmHg); IR (cm^{-1} , direct) 965; ^1H NMR (CCl_4) δ 0.93 (3 H, t, J = 6 Hz), 1.16–1.56 (4 H, m), 1.66–1.83 (3 H, m), 1.96–2.25 (2 H, m), 2.63–2.90 (2 H, m), 5.50–5.83 (4 H, m). Anal. Calcd for $\text{C}_{10}\text{H}_{18}$: C, 86.88; H, 13.12. Found: C, 86.71; H, 13.10.

Acknowledgment. We gratefully acknowledge support by the Ministry of Education, Science and Culture (Grant-in-Aid 247020, 321709, 334029, 411108, and 484027), the Kurata Foundation, and the JSPS (Basic Chemical Research Foundation).

References and Notes

- (1) For a preliminary report on some aspects of the present study, see: (a) Yamamoto, Y.; Maruyama, K. *J. Am. Chem. Soc.* **1977**, *99*, 8068. (b) *J. Organomet. Chem.* **1978**, *156*, C9.
- (2) For a leading reference, see: Seebach, D.; Geiss, K. H. In "New Applications of Organometallic Reagents in Organic Synthesis", Seyferth, D. Ed.; Elsevier: Amsterdam, 1976; p 1.
- (3) For a leading reference, see: March, J. "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure", 2nd ed.; McGraw-Hill: New York, 1977; p 304. See also the most recent article: Magid, R. M.; Fruchey, O. S. *J. Am. Chem. Soc.* **1979**, *101*, 2107.
- (4) Posner, G. H. *Org. React.* **1975**, *22*, 253.
- (5) Normant, J. F. In ref 2, p 219.
- (6) Böhlmann, F.; Bonnet, H.; Jente, R. *Chem. Ber.* **1968**, *101*, 855.
- (7) Sood, R.; Nagasawa, M.; Sih, C. J. *Tetrahedron Lett.* **1974**, 423.
- (8) Commercon, A.; Bourgain, M.; Delaunay, M.; Normant, J. F.; Villieras, J. *Tetrahedron Lett.* **1975**, 3837.
- (9) Henrick, C. A.; Labovitz, J. N.; Corbin, V. L. *Life Sci.* **1975**, *17*, 699.
- (10) Anderson, R. J.; Henrick, C. A.; Siddall, J. B. *J. Am. Chem. Soc.* **1970**, *92*, 735; **1972**, *94*, 5379.
- (11) (a) Fouquet, G.; Schlosser, M. *Angew. Chem., Int. Ed. Engl.* **1974**, *13*, 82. (b) Tanigawa, Y.; Kanamaru, H.; Sonoda, A.; Murahashi, S.-I. *J. Am. Chem. Soc.* **1977**, *99*, 2361. (c) Mukaiyama, T.; Imaoka, M.; Izawa, T. *Chem. Lett.* **1977**, 1257.
- (12) Henne, A. L.; Chanan, H.; Turk, A. J. *Am. Chem. Soc.* **1941**, *63*, 3474.
- (13) Czernecki, S.; Georgoulis, C.; Gross, B.; Prevost, C. *Bull. Soc. Chim. Fr.* **1968**, *9*, 3713.
- (14) Cristol, S. J.; Overhulst, W. C.; Meek, J. S. *J. Am. Chem. Soc.* **1951**, *73*, 813.
- (15) Wawzonek, S.; Studnicka, B. J.; Zigman, A. R. *J. Org. Chem.* **1969**, *34*, 1316.
- (16) Magid, R. M.; Nieh, E. C.; Gandour, R. D. *J. Org. Chem.* **1971**, *36*, 2099.
- (17) Jukes, A. E. *Adv. Organomet. Chem.* **1974**, *12*, 215.
- (18) For a review article, see: Henrick, C. A. *Tetrahedron Lett.* **1977**, 33, 1845.
- (19) (a) Kuwajima, I.; Doi, Y. *Tetrahedron Lett.* **1972**, 1163. (b) Colonge, J.; Falcot, R. *Bull. Soc. Chim. Fr.* **1957**, 1166. (c) Boumechal, F. D.; Lorne, R.; Linstrumelle, G. *Tetrahedron Lett.* **1977**, 1181. (d) Yamamoto, Y.; Maruyama, K. *J. Am. Chem. Soc.* **1978**, *100*, 6282. (e) Corey, E. J.; Mann, J. *Ibid.* **1973**, *95*, 6832. (f) Gream, G. E.; Pincombe, C. F. *Aust. J. Chem.* **1974**, *27*, 543. (g) Goering, H. L.; Singleton, Jr., V. D. *J. Am. Chem. Soc.* **1976**, *98*, 7854. (h) Kreft, A. *Tetrahedron Lett.* **1977**, 1035. (i) Levisalles, J.; R-Chauvin, M.; Rudler, H. *J. Organomet. Chem.* **1977**, *136*, 103.
- (20) vanMourik, G. L.; Pabon, H. J. *J. Tetrahedron Lett.* **1978**, 2705.
- (21) Rona, P.; Tokes, L.; Tremble, J.; Crabbé, P. *Chem. Commun.* **1969**, 43.
- (22) Normant, J. F.; Commercon, A.; Bourgain, M.; Villieras, J. *Tetrahedron Lett.* **1975**, 3833.
- (23) (a) Claesson, A.; Tärnnefors, I.; Olsson, L. I. *Tetrahedron Lett.* **1975**, 1509. (b) Claesson, A.; Olsson, L. I. *J. Chem. Soc., Chem. Commun.* **1978**, 621. (c) Tanigawa, Y.; Ohta, H.; Sonoda, A.; Murahashi, S.-I. *J. Am. Chem. Soc.* **1978**, *100*, 4610. (d) Cassani, G.; Massardo, P.; Piccardi, P. *Tetrahedron Lett.* **1979**, 633.
- (24) Gallina, C.; Ciattini, P. G. *J. Am. Chem. Soc.* **1979**, *101*, 1035.
- (25) For the coupling of allylic organometallic reagents with allylic halides in which additional complications arise in terms of the possibilities for symmetrical vs. unsymmetrical coupling, see: (a) Courtols, G.; Miglinac, L. *J. Organomet. Chem.* **1974**, *69*, 1. (b) Benkeser, R. A. *Synthesis* **1971**, 347. (c) Reference 19d.
- (26) The reaction of 3-methyl-1-buten-3-ol is an exception among the substrates examined (see Table III).
- (27) Yamamoto, Y.; Yatagai, H.; Maruyama, K.; Sonoda, A.; Murahashi, S.-I. *J. Am. Chem. Soc.* **1977**, *99*, 5652.
- (28) Ho, T.-L. "Hard and Soft Acids and Bases Principle in Organic Chemistry"; Academic Press: New York, 1977.
- (29) Pasto, D. J.; Chou, S.-K.; Fritzen, E.; Shultz, R. H.; Waterhouse, A.; Hennion, G. F. *J. Org. Chem.* **1978**, *43*, 1389, and references cited therein.

- (30) Vermeer, P.; Meijer, J.; Brandsma, L. *Recl. Trav. Chim. Pays-Bas* **1975**, *94*, 112. Tadema, G.; Vermeer, P.; Meijer, J.; Brandsma, L. *Ibid.* **1976**, *95*, 66. Vermeer, P.; Westmijze, H.; Kleijn, H.; van Dijk, L. A. *Ibid.* **1978**, *97*, 56. Tadema, G.; Everhardus, R. H.; Westmijze, H.; Vermeer, P. *Tetrahedron Lett.* **1978**, 3935.
- (31) Luche, J. L.; Barreiro, E.; Dollat, J. M.; Crabbé, P. *Tetrahedron Lett.* **1975**, 4615. Descolins, C.; Henrick, C. A.; Siddall, J. B. *Ibid.* **1972**, 3777. Moreau, J. L.; Gaudemar, M. J. *Organomet. Chem.* **1976**, *108*, 159. Crabbé, P.; Carpio, H. J. *Chem. Soc., Chem. Commun.* **1972**, 904. Crabbé, P.; Barreiro, E.; Dollat, J.-M.; Luche, J. L. *Ibid.* **1976**, 183. Rona, P.; Crabbé, P. *J. Am. Chem. Soc.* **1969**, *91*, 3289. Landor, P. D.; Landor, S. R. *J. Chem. Soc., Perkin Trans. 1* **1973**, 1347.
- (32) Brikmeyer, R. S.; Macdonald, T. L. *J. Chem. Soc., Chem. Commun.* **1978**, 876. For a related system, see: Kleijn, H.; Westmijze, H.; Kruthof, K.; Vermeer, P. *Recl. Trav. Chim. Pays-Bas* **1979**, *98*, 27.
- (33) (a) Pasto, D. J.; Shults, R. H.; McGrath, J. A.; Waterhouse, A. J. *Org. Chem.* **1978**, *43*, 1382. (b) Pasto, D. J.; Chou, S.-K.; Waterhouse, A.; Shults, R. H.; Hennion, G. F. *Ibid.* **1978**, *43*, 1385. (c) Gore, J.; Roumestant, M. L. *Tetrahedron Lett.* **1970**, 891. (d) Bush, C. N.; Applequist, D. E. *J. Org. Chem.* **1977**, *42*, 1076. (e) Jacobs, T. L.; Meyers, R. A. *J. Am. Chem. Soc.* **1964**, *86*, 5244; **1967**, *89*, 6177.
- (34) Although it is of interest to examine the reaction of 3-chloro-3-methyl-1-butene with $\text{RCu}\cdot\text{BF}_3$, it is very difficult to prepare the chloride in pure form.
- (35) Steinberg, H. "Organoboron Chemistry", Vol. 1; Interscience: New York, 1964; p 793.
- (36) Felkin, H.; Swierczewski, G. *Tetrahedron Lett.* **1972**, 1433. Felkin, H.; Costa, E. J.; Swierczewski, G. *J. Organomet. Chem.* **1977**, *134*, 265. Buckwalter, B. L.; Burflitt, I. R.; Felkin, H.; J-Goudket, M.; Naemura, K.; Salomon, M. F.; Wenkert, E.; Wovkulich, P. M. *J. Am. Chem. Soc.* **1978**, *100*, 6445.
- (37) Stork, G.; Kreft III, A. F. *J. Am. Chem. Soc.* **1977**, *99*, 3850, 3851. Cane, D. E.; Murthy, P. P. N. *Ibid.* **1977**, *99*, 8327. Magid, R. M.; Fruchey, O. S. *Ibid.* **1977**, *99*, 8368. Ikota, N.; Ganem, B. *Ibid.* **1978**, *100*, 351. Dobbie, A. A.; Overton, K. H. *J. Chem. Soc., Chem. Commun.* **1977**, 722.
- (38) For importance of the stereochemistry of starting materials, see: Crabbé, P.; Dollat, J.-M.; Gallina, J.; Luche, J.-L.; Velarde, E.; Maddox, M. L.; Tökes, L. *J. Chem. Soc., Perkin Trans. 1* **1978**, 730.
- (39) For the stereochemistry of $\text{S}_{\text{N}}2'$ reaction of metal hydrides, see: Jefford, C. W.; Sweeney, A.; Delay, F. *Helv. Chim. Acta* **1972**, *55*, 2214.
- (40) Booth, H. S.; Martin, D. R. "Boron Trifluoride and Its Derivatives"; Wiley: New York, 1949; p 61.
- (41) Chambers, R. D.; Clark, H. C.; Willis, C. J. *J. Am. Chem. Soc.* **1960**, *82*, 5298. *Proc. Chem. Soc., London* **1960**, 114.
- (42) Brauer, D. J.; Bürger, H.; Pawelke, G. *Inorg. Chem.* **1977**, *16*, 2305.
- (43) Schmid, G. *Angew. Chem., Int. Ed. Engl.* **1970**, *9*, 819.
- (44) For review articles, see: Tochtermann, W. *Angew. Chem., Int. Ed. Engl.* **1966**, *5*, 351. Wittig, G. *Q. Rev., Chem. Soc.* **1966**, 191. Negishi, E. J. *Organomet. Chem.* **1976**, *108*, 281.
- (45) Yamamoto, Y.; Yatagai, H.; Maruyama, K.; Sonoda, A.; Murahashi, S.-I. *Bull. Chem. Soc. Jpn.* **1977**, *50*, 3427.
- (46) Miyaura, N.; Itoh, M.; Suzuki, A. *Bull. Chem. Soc. Jpn.* **1977**, *50*, 2199. For copper tetraarylborates via the cation exchange reaction, see: Nesmeyanov, A. N.; Sazonova, V. A.; Liberman, G. S.; Yemelyanova, L. I. *Izv. Akad. Nauk SSSR, Otd. Khim. Nauk* **1955**, 48.
- (47) Cotton, F. A.; Wilkinson, G. "Advanced Inorganic Chemistry. A Comprehensive Text"; Interscience: New York, 1966; p 895.
- (48) Onak, T. "Organoborane Chemistry"; Academic Press: New York, 1975; p 19.
- (49) Previously we reported that considerable amounts of *n*-BuOH were obtained under this condition.^{1a} However, detailed investigation revealed that most of the *n*-BuOH had its origin in lithium butylate present in *n*-BuLi as an impurity. Use of *n*-BuLi stored for long periods increased the formation of the *n*-BuOH, presumably because of the slow oxidation of *n*-BuLi by air: Panek, E. J.; Kaiser, L. R.; Whitesides, G. M. *J. Am. Chem. Soc.* **1977**, *99*, 3708.
- (50) Since the complex is considerably stable, the structural information is obtained without difficulties and will be published soon. This is the first example of a stable alkenylcopper-trialkylborane ate complex.^{4b}
- (51) Detailed studies concerning these complexes will be published elsewhere by H. Yatagai.
- (52) Hatch, L. F.; Gardner, P. D.; Gilbert, R. E. *J. Am. Chem. Soc.* **1959**, *81*, 5943.
- (53) (a) "Organic Syntheses", Collect Vol. V; Wiley: New York, 1973; p 859. (b) Gilman, H.; Morton Jr., J. W. *Org. React.* **1954**, *8*, 258. (c) Reference 53a, p 762.
- (54) Gilman, H.; Hauben, A. H. *J. Am. Chem. Soc.* **1944**, *66*, 1515.
- (55) Watson, S. C.; Eastham, J. F. *J. Organomet. Chem.* **1967**, *9*, 165.
- (56) Posner, G. H.; Whitten, C. E.; Sterling, J. J. *J. Am. Chem. Soc.* **1973**, *95*, 7788.
- (57) Brown, H. C. "Organic Syntheses via Boranes"; Wiley-Interscience: New York, 1975.
- (58) *Org. Synth.* **1974**, *54*, 63.
- (59) Stepanova, O. S.; Ganin, E. V. *Ukr. Khim. Zh.* **1977**, *43*, 168.
- (60) Mori, K. *Tetrahedron* **1974**, *30*, 3807.
- (61) Bokadia, M. M.; Brown, B. R.; Coburn, D.; Roberts, A.; Somerfield, G. A. *J. Chem. Soc.* **1962**, 1658.
- (62) Normant, J. *Bull. Soc. Chim. Fr.* **1963**, 1888.
- (63) Bateman, L.; Cunneen, J. I. *J. Chem. Soc.* **1951**, 2289.
- (64) Levina, R. Y.; Skvarchenko, R.; Ushakova, T. M. *Zh. Obshch. Khim.* **1956**, *26*, 398.
- (65) Bingham, A. J.; Dyall, L. K.; Norman, R. O.; Thomas, C. B. *J. Chem. Soc. C* **1970**, 1879.
- (66) Goering, H. L.; Neritt, T. D.; Silversmith, E. F. *J. Am. Chem. Soc.* **1955**, *77*, 4042.
- (67) Crossley, A. W.; Ronouf, N. *J. Am. Chem. Soc.* **1915**, *37*, 602.
- (68) Frank, R. L.; Hal Jr., H. K. *J. Am. Chem. Soc.* **1950**, *72*, 1645.
- (69) Nystrom, R. F.; Borwn, W. G. *J. Am. Chem. Soc.* **1947**, *69*, 1197.
- (70) Goering, H. L.; Blanchard, J. P. *J. Am. Chem. Soc.* **1954**, *76*, 5405.
- (71) Macbeth, A. K.; Mills, J. A. *J. Chem. Soc.* **1949**, 2646.
- (72) Liberman, A. L.; Vasina, T. V. *Neftekhimiya* **1962**, *2*, 129.
- (73) Kiselev, A. V.; Migunova, I. A.; Khudyakov, V. L.; Yashin, Y. I. *Zh. Fiz. Khim.* **1966**, *40*, 2910.
- (74) Brown, H. C.; Scouten, C. G.; Liotta, R. *J. Am. Chem. Soc.* **1979**, *101*, 96.

Tunneling Model for Hydrogen Abstraction Reactions in Low-Temperature Solids. Applications to Reactions in Alcohol Glasses and Acetonitrile Crystals¹

Robert J. Le Roy,^{*2} Hisao Murai,^{3,4} and Ffrancon Williams³

Contribution from the Guelph-Waterloo Centre for Graduate Work in Chemistry, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1, and the Department of Chemistry, University of Tennessee, Knoxville, Tennessee 37916. Received August 2, 1979

Abstract: A simple and internally consistent quantitative model for hydrogen-abstraction reactions in low-temperature solids, which implicitly incorporates zero point energy effects which allow for finite reaction rates at $T = 0$ K, is derived and applied to new measurements of H-abstraction rate constants by methyl radicals in methanol and ethanol glasses at $T = 13$ –99 K and in acetonitrile and methyl isocyanide crystals at 69–128 K. Nonlinear least-squares fits of the model to the experimental data yield effective one-dimensional barriers to reaction whose heights are virtually independent of the analytic form used for the potential energy barrier, and are somewhat larger than the activation energies measured for the corresponding reactions in the gas phase. This model predicts that values of the isotopic rate constant ratio $k_{\text{H}}/k_{\text{D}}$ will be larger than 10^{12} at $T = 0$ K.

Quantum-mechanical tunneling has long been believed to make a dominant contribution to the rates of many chemical reactions at low temperatures.^{5,6} However, it is only relatively recently that convincing experimental evidence of this behavior

has begun to appear. In most studies,^{7–17} the evidence that the reactions considered proceeded mainly by tunneling consisted of (1) pronounced curvature in Arrhenius plots, (2) anomalously small activation energies, and (3) anomalously large^{10,14}