Research Paper

# N-acylation of amides through internal nucleophilic catalysis

# Meng Yao, Xia Chen, Yuling Lu, Zhenhua Guan,

# Abstract

Journal of Chemical Research © The Author(s) 2020 Article reuse guidelines: sagepub.com/journals-permissions DOI: 10.1177/1747519820925755 journals.sagepub.com/home/chl

(\$)SAGE

Zengwei Luo and Yonghui Zhang

# An efficient method for N-acylation of amides is described using a pyridine ring as the internal nucleophilic catalyst to give imides in moderate to excellent yields. The methodology provides a facile, air insensitive, and environmentally friendly route to form diversified imide scaffolds, which exist widely in natural products and biologically active materials.

#### **Keywords**

imides, internal nucleophilic catalysis, Mumm rearrangement, N,N-diisopropylethylamine, pyridine

Date received: 28 February 2020; accepted: 20 April 2020



# Introduction

Imides are well represented as valuable intermediates in synthetic organic chemistry, and these structural motifs also occur in natural products and pharmaceuticals such as the antibiotic fumaramidmycin,<sup>1,2</sup> palauimide,<sup>3</sup> thalidomide,<sup>4</sup> the antifeedant ypaoamide,<sup>5</sup> and the platelet-activating factor (PAF) antagonist CV-62096,7 (Figure 1). To date, various methods for imides have been developed. Imides can be prepared by the palladium-catalyzed three-component reaction between terminal alkynes, isonitriles, and sodium carboxylates.8 Yamaguchi reported a copper-catalyzed aerobic oxidative acylation of amides with alcohols for the synthesis of imides.9 Also, Nicolaou reported the oxidation of secondary amides to the corresponding imides using Dess-Martin periodinane.10

In spite of the above-mentioned methods, imides are usually prepared between amides and an excess of activated forms of carboxylic acids, such as acyl chlorides, anhydrides, and esters, under strong basic or acidic conditions (Scheme 1(a)).<sup>11-14</sup> However, for some acid- or basesensitive or biologically related compounds, metal-free and versatile conditions are in demand. Although there are several examples described using mild conditions with a weak base,<sup>15–19</sup> for instance, the conversion of NHBoc to N(Boc)<sub>2</sub> with 4-dimethylaminopyridine (DMAP),<sup>20</sup> the development of versatile and efficient methods for the preparation of imides is still challenging.



Figure 1. Compounds containing an imide core.

Hubei Key Laboratory of Natural Medicinal Chemistry and Resource Evaluation, School of Pharmacy, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, P.R. China

#### **Corresponding authors:**

Zengwei Luo, Hubei Key Laboratory of Natural Medicinal Chemistry and Resource Evaluation, School of Pharmacy, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, P.R. China.

Email: luozengwei@hust.edu.cn

Yonghui Zhang, Hubei Key Laboratory of Natural Medicinal Chemistry and Resource Evaluation, School of Pharmacy, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, P.R. China.

Email: zhangyh@mails.tjmu.edu.cn



Scheme I. Strategies relevant to imides synthesis: (a) traditional work, (b) Unsworth's work, and (c) this work.

Using an internal nucleophilic catalyst may be the solution to this problem. Acylation is one type of basic organic transformation to form amides or esters, which can be accelerated using a nucleophilic catalyst. Pyridine can act as a nucleophile for activated carboxylic acids and is often used as a catalyst in acylation reactions. Recently, Unsworth demonstrated a strategy for the synthesis of medium-sized rings readily from linear precursors mediated by an internal nucleophilic catalyst, a pyridine ring, which can capture the carboxylic acid to provide a reactive intermediate (Scheme 1(b)).<sup>21-23</sup> Inspired by this work, we have taken advantage of internal catalysis to achieve mild N-acylation reactions of amides. Herein, we used the amides possessing a pyridine ring as substrates, the pyridine ring can form an active acylammonium salt with an acyl chloride. Thus, the original intermolecular reaction will be transformed into an intramolecular reaction. The imide will be obtained following a Mumm rearrangement of the intermediate isoimide (Scheme 1(c)).

# **Results and discussion**

With *N*-(pyridin-2-ylmethyl)acetamide (1a) as the acyl acceptor and benzoyl chloride (2a) as the acyl donor, an initial study was performed with *N*,*N*-diisopropylethylamine (DIPEA) as the base in  $CH_2Cl_2$  at room temperature for 3 h (Table 1, entry 1). To our delight, the desired product 3a was obtained in 51% yield. It was found that a higher yield (83%) was achieved when using 1.5 equiv. of DIPEA (entries 2 and 3). Furthermore, the yield was improved to 94% when screening the amount of acyl chloride (entries 4 and 5). Three other bases were also tested (entries 6–8) but all led to a reduction in the yield of 3a.

Next, other aromatic substrates as internal nucleophilic catalysts were examined (Scheme 2). Thiophene and furan were less efficient leading to **3b** and **3c** in moderate yields. The 2-methylpyridine was found to be optimal when altering the carbon chain number between the pyridine ring and the amide (**3d** and **3e**). Moreover, replacing the pyridine ring with a phenyl ring did not give any product **3f**. Thus, the optimized reaction conditions with a pyridine ring as

Table 1. Optimization of the reaction conditions.<sup>a</sup>

	$ \begin{array}{c}                                     $	Base CH <sub>2</sub> Cl <sub>2</sub>	0 Эн <sub>3</sub>
Entry	Base (equiv.)	<b>2a</b> (equiv.)	Yield <sup>b</sup> (%)
1	DIPEA (1.0)	1.0	51
2	DIPEA (1.5)	1.0	83
3	DIPEA (2.0)	1.0	82
4	<b>DIPEA</b> (1.5)	1.3	94
5	DIPEA (1.5)	1.5	93
6	Et <sub>3</sub> N (1.5)	1.3	48
7	Na <sub>2</sub> CO <sub>3</sub> (1.5)	1.3	23
8	CH <sub>3</sub> COOK (1.5)	1.3	35

DIPEA: N,N-diisopropylethylamine.

 $^{a}Conditions:$  1 a (0.2 mmol),  $CH_{2}Cl_{2}$  (2.0 mL), room temperature, 3 h.  $^{b}$  Isolated yields.

Significance for bold value in table I was illustrated as the highlighted sentence.



Scheme 2. Acyl transfer catalyst optimization.

the internal nucleophilic catalyst were found to be 1.5 equiv. of DIPEA and 1.3 equiv. of 2a in  $CH_2Cl_2$  at room temperature for 3 h, from which compound 3a was isolated in 94% yield (entry 4).

After determining the optimized reaction conditions, we then set out to investigate the substrate scope of the amides and the acid chlorides. The results are summarized in Scheme 3.

Aliphatic and  $\alpha,\beta$ -unsaturated amides, especially hindered t-butyryl amide, reacted smoothly to produce the target imides 5a-5d in moderate to excellent yields. Amides bearing either electron-donating or electron-withdrawing groups on the phenyl ring successfully delivered the desired products 5ea-5ej in good to excellent yields. Combined with the above results, the transformation still ran smoothly regardless of the presence of substituents at ortho, meta, or para positions on the phenyl ring (5f-5j). This transformation was also applicable to heteroaryl amides (5k and 5l). Next, the scope of the acyl chlorides was examined. Acetyl chloride and 2-methoxyacetyl chloride reacted with cinnamamide to deliver the products 5m and 5n in 90% and 82% yields, respectively. As in the case of benzamide and isobutyramide, conjugated acyl chlorides and various functional-group-substituted alkyl acyl chlorides also reacted smoothly to provide the corresponding imides 50-5u. It was noteworthy that when a diamide was used as the substrate, product 5v was obtained as the sole product.



Scheme 3. Substrate scope of the amides and acyl chlorides. Reaction conditions: 4 (0.2 mmol), 2 (0.26 mmol), DIPEA (0.3 mmol),  $CH_2CI_2$  (2.0 mL), room temperature, 3 h. Isolated yields are given.

# Conclusion

In conclusion, we have developed a mild and robust *N*-acylation tactic for the preparation of imides from simple amides with acyl chlorides via internal nucleophilic catalysis. The method employs a mild readily available DIPEA as the base and thus provides an inexpensive, environmentally friendly, and easy to operate route toward diverse imide derivatives with good substrate scope. Regarding the widespread distribution of imide fragments, the utility of this method in synthetic chemistry is being explored in our laboratory.

# **Experimental analysis**

Reagents and solvents were purchased from commercial suppliers unless otherwise specified. All reactions were carried out under an air atmosphere. Anhydrous solvents were purified and dried following standard procedures. Purification was generally done by flash column chromatography on brand silica gel (200–300 mesh size). Thinlayer chromatography (TLC) analysis was performed on brand precoated, glass-backed silica gel plates. Nuclear magnetic resonance (NMR) spectra were recorded on a 400-MHz Bruker spectrometer (400 MHz for <sup>1</sup>H NMR, 100 MHz for <sup>13</sup>C NMR). Chemical shifts (<sup>1</sup>H and <sup>13</sup>C) are given in ppm relative to the residual solvent peak (CDCl<sub>3</sub>, 7.26 ppm, 77.0 ppm, respectively). High-resolution mass spectra

(HRMS) were obtained on a Thermo Fisher LC-LTQ-Orbitrap XL spectrometer. For more information about chemical spectra, please see the supplemental material.

General procedure for the synthesis of imides 5. To a mixture of amide 4 (0.2 mmol) and DIPEA (0.3 mmol) in  $CH_2Cl_2$  (2 mL) was added acyl chloride 2 (0.26 mmol) dropwise at 0 °C. The reaction mixture was stirred at room temperature for 3 h. After completion of the reaction, the mixture was poured into water (20 mL) and extracted with  $CH_2Cl_2$  (3 × 20 mL). The combined organic layers were dried with anhydrous  $Na_2SO_4$ . After removal of the solvents in vacuo, the obtained crude product was further purified by column chromatography on silica gel, eluting with a mixture of petroleum and ethyl acetate (5:1) to give the desired products.

*N-Acetyl-N-(pyridin-2-ylmethyl)benzamide* (**3a**): colorless oil; yield: 47.8 mg, (94%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.51 (d, *J*=4.7 Hz, 1H), 7.70 (d, *J*=8.1 Hz, 2H), 7.62 (t, *J*=7.1 Hz, 1H), 7.52 (t, *J*=7.4 Hz, 1H), 7.42 (t, *J*=7.6 Hz, 2H), 7.22 (d, *J*=7.8 Hz, 1H), 7.17–7.11 (m, 1H), 5.09 (s, 2H), 2.26 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.3, 173.5, 156.5, 149.2, 136.6, 135.7, 132.2, 128.6, 128.5, 122.1, 121.4, 50.8, 26.2; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>15</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>Na: 277.0947; found: 277.0944.

*N-Acetyl-N-(thiophen-2-ylmethyl)benzamide* (3b): colorless oil; yield: 33.7 mg, (65%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.60–7.53 (m, 3H), 7.45 (t, *J*=7.6 Hz, 2H), 7.21 (dd, *J*=5.0, 1.4 Hz, 1H), 6.93–6.87 (m, 2H), 5.15 (s, 2H), 2.10 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.9, 173.0, 139.3, 135.7, 132.6, 128.9, 128.5, 127.3, 126.5, 125.8, 44.0, 26.5; HRMS (ESI<sup>+</sup>): *m/z* [M+Na]<sup>+</sup> calcd for C<sub>14</sub>H<sub>13</sub>NO<sub>2</sub>SNa: 282.0559; found: 282.0569.

*N-Acetyl-N-(furan-2-ylmethyl)benzamide* (**3c**): colorless oil; yield: 33.5 mg, (69%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.59 (d, *J*=8.0 Hz, 2H), 7.54 (t, *J*=7.4 Hz, 1H), 7.44 (t, *J*=7.6 Hz, 2H), 7.29 (s, 1H), 6.27 (s, 1H), 6.20 (d, *J*=3.1 Hz, 1H), 4.96 (s, 2H), 2.17 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.0, 172.8, 150.4, 142.1, 135.5, 132.5, 128.8, 128.4, 110.4, 108.4, 42.3, 26.1; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>14</sub>H<sub>13</sub>NO<sub>3</sub>Na: 266.0788; found: 266.0782.

*N*-Acetyl-*N*-(2-(pyridin-2-yl)ethyl)benzamide (3d): colorless oil; yield: 36.5 mg, (68%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.43 (d, *J*=4.2 Hz, 1H), 7.57 (td, *J*=7.65,1.32 Hz, 1H), 7.51–7.47 (m, 3H), 7.40 (t, *J*=7.6 Hz, 2H), 7.15–7.07 (m, 2H), 4.19 (t, *J*=6.9 Hz, 2H), 3.11 (t, *J*=6.9 Hz, 2H), 2.15 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.4, 173.3, 158.5, 149.1, 136.6, 135.4, 132.3, 128.7, 128.5, 123.8, 121.6, 46.3, 37.1, 25.9; HRMS (ESI<sup>+</sup>): *m*/*z* [M + Na]<sup>+</sup> calcd for C<sub>16</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>Na: 291.1104; found: 291.1121.

*N-Pivaloyl-N-(pyridin-2-ylmethyl)benzamide* (5a): colorless oil; yield: 44.4 mg, (75%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.54 (d, *J*=4.7 Hz, 1H), 7.80 (d, *J*=7.1 Hz, 2H), 7.62 (t, *J*=6.9 Hz, 1H), 7.51 (t, *J*=6.7 Hz, 1H), 7.41 (t, *J*=7.4 Hz, 2H), 7.22–7.12 (m, 2H), 4.92 (s, 2H), 1.31 (s, 9H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  187.4, 174.8, 156.5, 149.2, 136.6, 134.6, 132.3, 129.0, 128.6, 122.2, 121.6, 52.6, 43.4, 28.5; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>18</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub>Na: 319.1417; found: 319.1424.

*N-(Cyclopropanecarbonyl)-N-(pyridin-2-ylmethyl)benzamide* (**5b**): colorless oil; yield: 42.6 mg, (76%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.53 (d, *J*=4.4 Hz, 1H), 7.80–7.75 (m, 2H), 7.64 (td, J=7.7, 1.7Hz, 1H), 7.52 (t, J=7.4Hz, 1H), 7.43 (t, J=7.5Hz, 2H), 7.28 (d, J=7.85, 1H), 7.15 (dd, J=7.2, 5.2Hz, 1H), 5.21 (s, 2H), 1.59 (ddd, J=12.4, 7.8, 4.6Hz, 1H), 1.07–1.00 (m, 2H), 0.65 (m, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  177.9, 173.9, 156.9, 149.3, 136.6, 136.2, 132.2, 129.0, 128.5, 122.1, 121.4, 50.7, 18.3, 11.7; HRMS (ESI<sup>+</sup>): m/z [M + Na]<sup>+</sup> calcd for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>Na: 303.1104; found: 303.1115.

*N-Acryloyl-N-(pyridin-2-ylmethyl)benzamide* (5c): colorless oil; yield: 47.9 mg, (90%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, *J*=4.7 Hz, 1H), 7.72 (d, *J*=8.2 Hz, 2H), 7.64 (t, *J*=7.7 Hz, 1H), 7.53 (t, *J*=7.5 Hz, 1H), 7.41 (t, *J*=7.6 Hz, 2H), 7.29 (d, *J*=7.8 Hz, 1H), 7.18–7.12 (m, 1H), 6.31–6.17 (m, 2H), 5.51 (dd, *J*=9.1, 2.7 Hz, 1H), 5.21 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.7, 169.2, 156.4, 149.3, 136.6, 135.8, 132.5, 130.8, 129.1, 128.8, 128.6, 122.2, 121.5, 50.6; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>Na: 289.0947; found: 289.0951.

*N*-(3-Phenylpropioloyl)-*N*-(pyridin-2-ylmethyl)benzamide (**5d**): colorless oil; yield: 60.5 mg, (89%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.55 (d, *J*=4.9 Hz, 1H), 7.88 (d, *J*=7.5 Hz, 2H), 7.67 (t, *J*=7.7 Hz, 1H), 7.53 (t, *J*=7.4 Hz, 1H), 7.45 (t, *J*=7.5 Hz, 2H), 7.35 (t, *J*=7.8 Hz, 2H), 7.24 (m, 2H), 7.21–7.13 (m, 1H), 7.09 (d, *J*=7.7 Hz, 2H), 5.34 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 173.4, 156.1, 155.0, 149.5, 136.6, 136.0, 132.7, 132.6, 130.6, 129.7, 128.5, 128.3, 122.3, 121.5, 119.5, 95.9, 82.7, 49.9; HRMS (ESI<sup>+</sup>): *m*/*z* [M + Na]<sup>+</sup> calcd for C<sub>22</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>Na: 363.1104; found: 363.1113.

*N-Benzoyl-N-(pyridin-2-ylmethyl)benzamide* (5ea): colorless oil; yield: 55.6 mg, (88%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, *J*=4.7 Hz, 1H), 7.66 (td, *J*=7.7, 1.4 Hz, 1H), 7.54 (d, *J*=7.3 Hz, 4H), 7.39 (d, *J*=7.8 Hz, 1H), 7.25 (t, *J*=7.20, 3H), 7.14 (t, *J*=7.5 Hz, 5H), 5.34 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.2, 156.6, 149.3, 136.6, 136.5, 131.7, 129.1, 128.1, 122.3, 121.8, 51.4; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>20</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>Na: 339.1104; found: 311.1111.

*N-Benzoyl-4-methyl-N-(pyridin-2-ylmethyl)benzamide* (**5eb**): colorless oil; yield: 59.3 mg, (90%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.53 (d, *J*=4.9 Hz, 1H), 7.67 (t, *J*=7.7 Hz, 1H), 7.56 (d, *J*=7.4 Hz, 2H), 7.47 (d, *J*=7.9 Hz, 2H), 7.40 (d, *J*=7.8 Hz, 1H), 7.25 (d, *J*=7.0 Hz, 1H), 7.17 (t, *J*=7.9 Hz, 3H), 6.96 (d, *J*=7.9 Hz, 2H), 5.34 (s, 2H), 2.23 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.3, 174.2, 156.7, 149.3, 142.5, 136.6, 136.5, 133.6, 131.6, 129.3, 129.0, 128.8, 128.0, 122.2, 121.8, 51.6, 21.4; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>21</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>Na: 353.1260; found: 353.1262.

*N-Benzoyl-4-(tert-butyl)-N-(pyridin-2-ylmethyl)benzamide* (**5ec**): colorless oil; yield: 63.3 mg, (85%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.54 (d, *J*=4.4Hz, 1H), 7.67 (t, *J*=7.7Hz, 1H), 7.53 (d, *J*=7.5Hz, 2H), 7.49 (d, *J*=8.2Hz, 2H), 7.40 (d, *J*=7.8Hz, 1H), 7.22 (t, *J*=7.2Hz, 1H), 7.17– 7.12 (m, 5H), 5.35 (s, 2H), 1.19 (s, 9H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.4, 174.2, 156.7, 155.3, 149.4, 136.6, 133.5, 131.4, 129.1, 129.0, 128.0, 125.0, 122.2, 121.8, 51.4, 34.9, 30.9; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>24</sub>H<sub>24</sub>N<sub>2</sub>O<sub>3</sub>Na: 395.1730; found: 395.1725. *N-Benzoyl-4-methoxy-N-(pyridin-2-ylmethyl)benzamide* (**5ed**): colorless oil; yield: 55.4 mg, (80%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.59 (d, *J*=4.9 Hz, 1H), 7.72 (t, *J*=7.6 Hz, 1H), 7.63 (dd, *J*=8.4, 2.5 Hz, 4H), 7.46 (d, *J*=7.8 Hz, 1H), 7.33 (t, *J*=7.2 Hz, 1H), 7.23 (q, *J*=7.7 Hz, 3H), 6.73 (d, *J*=8.6 Hz, 2H), 5.40 (s, 2H), 3.79 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.1, 173.7, 162.4, 156.7, 149.3, 136.5, 131.5, 131.4, 128.9, 128.6, 128.1, 122.1, 121.7, 113.4, 55.3, 51.6; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>21</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub>Na: 369.1210; found: 369.1218.

*N-Benzoyl-4-chloro-N-(pyridin-2-ylmethyl)benzamide* (**5ee**): colorless oil; yield: 63.0 mg, (90%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.51 (d, *J*=4.9 Hz, 1H), 7.67 (t, *J*=7.7 Hz, 1H), 7.56 (d, *J*=7.6 Hz, 2H), 7.52 (d, *J*=8.3 Hz, 2H), 7.37 (d, *J*=7.9 Hz, 1H), 7.30 (t, *J*=7.4 Hz, 1H), 7.22–7.11 (m, 5H), 5.33 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.0, 173.3, 156.3, 149.4, 137.8, 136.6, 136.3, 135.0, 132.0, 130.4, 129.1, 128.4, 128.3, 122.3, 121.8, 51.4; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>20</sub>H<sub>15</sub>CIN<sub>2</sub>O<sub>2</sub>Na: 373.0714; found: 373.0712.

*N-Benzoyl-4-bromo-N-(pyridin-2-ylmethyl)benzamide* (**5ef**): colorless oil; yield: 71.7 mg, (91%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.51 (d, *J*=4.4 Hz, 1H), 7.67 (td, *J*=7.7, 1.8 Hz, 1H), 7.58–7.53 (m, 2H), 7.47–7.42 (m, 2H), 7.37 (d, *J*=7.8 Hz, 1H), 7.30 (m, 3H), 7.23–7.14 (m, 3H), 5.33 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.0, 173.4, 156.2, 149.4, 136.5, 136.2, 135.4, 132.0, 131.3, 130.5, 129.1, 128.3, 126.4, 122.3, 121.7, 51.4; HRMS (ESI<sup>+</sup>): *m/z* [M+Na]<sup>+</sup> calcd for C<sub>20</sub>H<sub>15</sub>BrN<sub>2</sub>O<sub>2</sub>Na: 417.0209; found: 417.0208.

*N-Benzoyl-N-(pyridin-2-ylmethyl)-4-(trifluoromethyl)* benzamide (**5eg**): colorless oil; yield: 63.0 mg, (82%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, *J*=4.9 Hz, 1H), 7.69 (d, *J*=7.8 Hz, 3H), 7.56 (d, *J*=7.5 Hz, 2H), 7.43 (d, *J*=7.7 Hz, 2H), 7.38 (d, *J*=7.6 Hz, 1H), 7.33–7.25 (m, 1H), 7.19 (t, *J*=6.9 Hz, 3H), 5.35 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.0, 173.0, 156.0, 149.4, 139.9, 136.6, 136.0, 132.8 (q, *J*=32.0 Hz), 132.09, 129.2, 129.1, 128.3, 125.3 (q, *J*=4.0 Hz), 123.4 (d, *J*=271.0 Hz), 122.4, 121.8, 51.2; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>21</sub>H<sub>15</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>Na: 407.0978; found: 407.0970.

*N-Benzoyl-4-cyano-N-(pyridin-2-ylmethyl)benzamide* (**5eh**): colorless oil; yield: 53.2 mg, (78%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, *J*=4.7 Hz, 1H), 7.72–7.65 (m, 3H), 7.58–7.53 (m, 2H), 7.46 (d, *J*=8.3 Hz, 2H), 7.37 (d, *J*=7.8 Hz, 1H), 7.32 (t, *J*=7.4 Hz, 1H), 7.23–7.17 (m, 3H), 5.35 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.7, 172.5, 155.7, 149.3, 140.6, 136.8, 135.8, 132.3, 131.8, 129.3, 129.1, 128.5, 122.5, 121.8, 117.8, 114.6, 51.1; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>21</sub>H<sub>15</sub>N<sub>3</sub>O<sub>2</sub>Na: 364.1056; found: 364.1060.

*N-Benzoyl-4-(methylsulfonyl)-N-(pyridin-2-ylmethyl)benzamide* (**5ei**): colorless oil; yield: 63.1 mg, (80%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.53 (d, *J*=4.7 Hz, 1H), 7.79–7.67 (m, 5H), 7.57 (d, *J*=7.3 Hz, 2H), 7.38 (d, *J*=7.8 Hz, 1H), 7.31 (t, *J*=7.4 Hz, 1H), 7.20 (t, *J*=7.8 Hz, 3H), 5.36 (s, 2H), 2.93 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.8, 172.5, 155.7, 149.3, 142.4, 141.5, 136.8, 135.9, 132.3, 129.7, 129.2, 128.4, 127.2, 122.5, 121.8, 51.1, 44.2; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>21</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>SNa: 417.0879; found: 417.0881.

*N-Benzoyl-4-nitro-N-(pyridin-2-ylmethyl)benzamide* (**5ej**): colorless oil; yield: 65.7 mg, (91%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, *J*=4.5 Hz, 1H), 8.02 (d, *J*=8.6 Hz, 2H), 7.74 (d, *J*=8.6 Hz, 2H), 7.69 (t, *J*=7.7 Hz, 1H), 7.58 (d, *J*=7.8 Hz, 2H), 7.37 (d, *J*=7.8 Hz, 1H), 7.31 (t, *J*=7.4 Hz, 1H), 7.20 (t, *J*=7.7 Hz, 3H), 5.36 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.7, 172.3, 155.7, 149.4, 148.9, 142.3, 136.7, 135.8, 132.4, 129.8, 129.2, 128.5, 123.2, 122.5, 121.8, 51.2; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>20</sub>H<sub>15</sub>N<sub>3</sub>O<sub>4</sub>Na: 384.0955; found: 384.0956.

*N-Benzoyl-N-(pyridin-2-ylmethyl)-[1,1'-biphenyl]-2carboxamide* (**5f**): colorless oil; yield: 40.5 mg, (80%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.47 (d, *J*=4.5 Hz, 1H), 7.56 (td, *J*=7.7, 1.7 Hz, 1H), 7.43–7.40 (m, 3H), 7.39–7.33 (m, 3H), 7.27–7.22 (m, 4H), 7.21 (d, *J*=11.7 Hz, 1H), 7.18– 7.16 (m, 1H), 7.15–7.12 (m, 1H), 7.12–7.07 (m, 3H), 4.87 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.4, 172.9, 156.4, 149.3, 139.7, 139.4, 136.6, 136.0, 135.8, 131.6, 130.3, 129.8, 129.8, 128.7, 128.6, 128.5, 127.9, 127.2, 122.2, 121.7, 51.0; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>26</sub>H<sub>20</sub>N<sub>2</sub>O<sub>2</sub>Na: 415.1417; found: 415.1429.

*N-Benzoyl-2-bromo-N-(pyridin-2-ylmethyl)benzamide* (**5g**): colorless oil; yield: 63.9 mg, (81%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.55 (d, *J*=4.8 Hz, 1H), 7.67 (t, *J*=7.8 Hz, 1H), 7.57 (d, *J*=7.6 Hz, 2H), 7.46 (d, *J*=7.6 Hz, 1H), 7.39 (d, *J*=7.9 Hz, 1H), 7.28 (t, *J*=8.45, 2H), 7.23– 7.16 (m, 3H), 7.10 (t, *J*=7.5 Hz, 1H), 7.01 (t, *J*=7.8 Hz, 1H), 5.35 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.0, 171.1, 156.2, 149.4, 137.8, 136.5, 136.4, 133.3, 131.6, 131.3, 130.3, 128.6, 128.1, 126.8, 122.3, 121.9, 120.8, 50.5; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>20</sub>H<sub>15</sub>BrN<sub>2</sub>O<sub>2</sub>Na: 417.0209; found: 417.0205.

*N-Benzoyl-3-methyl-N-(pyridin-2-ylmethyl)benzamide* (**5h**): colorless oil; yield: 59.4 mg, (90%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.53 (d, *J*=4.7 Hz, 1H), 7.67 (t, *J*=7.6 Hz, 1H), 7.54 (d, *J*=7.7 Hz, 2H), 7.40 (d, *J*=7.8 Hz, 1H), 7.35 (d, *J*=4.5 Hz, 1H), 7.33 (s, 1H), 7.24 (d, *J*=7.3 Hz, 1H), 7.16 (t, *J*=7.6 Hz, 3H), 7.04 (d, *J*=4.9 Hz, 2H), 5.35 (s, 2H), 2.21 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.4, 174.3, 156.7, 149.4, 137.9, 136.6, 136.6, 136.3, 132.4, 131.6, 129.7, 129.0, 128.1, 128.0, 126.3, 122.3, 121.8, 51.5, 21.1; HRMS (ESI<sup>+</sup>): *m*/*z* [M + Na]<sup>+</sup> calcd for C<sub>21</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>Na: 353.1260; found: 353.1257.

*N-Benzoyl-3-methoxy-N-(pyridin-2-ylmethyl)benzamide* (**5i**): colorless oil; yield: 54.7 mg, (79%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.54 (d, *J*=4.6 Hz, 1H), 7.67 (t, *J*=7.6 Hz, 1H), 7.56 (d, *J*=7.7 Hz, 2H), 7.40 (d, *J*=7.9 Hz, 1H), 7.30–7.24 (m, 1H), 7.21–7.13 (m, 4H), 7.07 (dd, *J*=14.7, 6.7 Hz, 2H), 6.79 (d, *J*=8.1 Hz, 1H), 5.35 (s, 2H), 3.70 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.2, 174.1, 159.2, 156.6, 149.3, 137.7, 136.7, 136.5, 131.7, 129.2, 129.1, 128.1, 122.3, 121.8, 121.6, 118.4, 113.7, 55.4, 51.4; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>21</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub>Na: 369.1210; found: 269.1214.

*N-Benzoyl-3-bromo-N-(pyridin-2-ylmethyl)benzamide* (5j): colorless oil; yield: 67.0 mg, (85%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.53 (d, *J*=4.5Hz, 1H), 7.69 (m, 2H), 7.55 (d, *J*=7.5 Hz, 2H), 7.49 (d, *J*=7.4 Hz, 1H), 7.36 (t, *J*=8.3 Hz, 2H), 7.29 (t, *J*=7.29, 1H), 7.23–7.15 (m, 3H), 7.02 (t, *J*=7.8 Hz, 1H), 5.34 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.0, 172.7, 156.2, 149.4, 138.4, 136.6, 136.3, 134.3, 132.0, 131.9, 129.6, 129.0, 128.3, 127.5, 122.3, 122.1, 121.7, 51.3; HRMS (ESI<sup>+</sup>): m/z [M + Na]<sup>+</sup> calcd for C<sub>20</sub>H<sub>15</sub>BrN<sub>2</sub>O<sub>2</sub> 417.0209; found: 417.0214.

*N-Benzoyl-N-(pyridin-2-ylmethyl)thiophene-2-carboxamide* (**5k**): colorless oil; yield: 57.3 mg, (89%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, *J*=4.7 Hz, 1H), 7.71–7.61 (m, 3H), 7.44–7.36 (m, 3H), 7.32 (t, *J*=7.3 Hz, 1H), 7.24 (t, *J*=7.32, 2H), 7.20–7.14 (m, 1H), 6.81 (t, *J*=4.4 Hz, 1H), 5.35(s, 2H), <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.5, 167.5, 156.5, 149.1, 139.5, 136.8, 136.2, 133.3, 133.0, 131.8, 129.0, 128.3, 127.2, 122.3, 121.8, 51.8; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>18</sub>H<sub>14</sub>N<sub>2</sub>O<sub>2</sub>SNa: 345.0668; found: 345.0678.

*N-Benzoyl-N-(pyridin-2-ylmethyl)furan-2-carboxamide* (51): colorless oil; yield: 53.3 mg, (87%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.51 (d, *J*=4.7 Hz, 1H), 7.66 (dd, *J*=15.9, 7.7 Hz, 3H), 7.40 (d, *J*=7.7 Hz, 1H), 7.33 (t, *J*=6.1 Hz, 1H), 7.29–7.19 (m, 3H), 7.17–7.11 (m, 1H), 6.95 (s, 1H), 6.22 (s, 1H), 5.32 (s, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.3, 162.9, 156.6, 149.4, 147.9, 145.3, 136.6, 136.0, 131.8, 128.7, 128.1, 122.2, 121.4, 118.8, 112.3, 51.3; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>18</sub>H<sub>14</sub>N<sub>2</sub>O<sub>3</sub>Na: 329.0897; found: 329.0902.

*N*-Acetyl-*N*-(pyridin-2-ylmethyl)cinnamamide (5m): colorless oil; yield: 50.4 mg, (90%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.56 (d, *J*=4.9 Hz, 1H), 7.76 (d, *J*=15.5 Hz, 1H), 7.67 (d, *J*=6.36 Hz, 1H), 7.52 (s, 2H), 7.37 (s, 3H), 7.28 (d, *J*=13.6 Hz, 2H), 7.20 (s, 1H), 5.14 (s, 2H), 2.53 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.8, 169.4, 156.7, 149.4, 145.2, 136.9, 134.7, 130.4, 128.8, 128.3, 122.5, 121.7, 120.5, 49.5, 26.3; HRMS (ESI<sup>+</sup>): *m*/*z* [M + Na]<sup>+</sup> calcd for C<sub>17</sub>H<sub>16</sub>N<sub>2</sub>O<sub>3</sub>Na: 303.1104; found: 303.1115.

*N*-(2-*Methoxyacetyl*)-*N*-(*pyridin*-2-*ylmethyl*)*cinnamamide* (**5n**): colorless oil; yield: 50.9 mg, (82%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.57 (d, *J*=4.7 Hz, 1H), 7.76 (d, *J*=15.5 Hz, 1H), 7.67 (td, *J*=1.9, 7.6 Hz, 1H), 7.51 (dd, *J*=6.6, 2.9 Hz, 2H), 7.37 (dd, *J*=5.1, 1.8 Hz, 3H), 7.32 (d, *J*=7.9 Hz, 1H), 7.26 (d, *J*=15.5 Hz, 2H), 7.21 (dd, *J*=7.1, 5.3 Hz, 1H), 5.18 (s, 2H), 4.66 (s, 2H), 3.49 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 174.0, 169.1, 156.3, 149.4, 146.3, 137.0, 134.4, 130.6, 128.9, 128.4, 122.7, 122.2, 119.3, 74.6, 59.3, 48.8; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for  $C_{18}H_{18}N_2O_3Na$ : 333.1210; found: 333.1214.

*N*-(*Pyridin-2-ylmethyl*)-*N*-(3,3,3-trifluoropropanoyl) benzamide (**50**): colorless oil; yield: 55.4 mg, (86%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, *J*=4.6 Hz, 1H), 7.71– 7.67 (m, 2H), 7.64 (td, *J*=7.7, 1.6 Hz, 1H), 7.54 (t, *J*=7.5 Hz, 1H), 7.43 (t, *J*=7.6 Hz, 2H), 7.18 (t, *J*=6.4 Hz, 2H), 5.05 (s, 2H), 3.74 (q, *J*=9.9 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  174.1, 166.9 (t, *J*=3.0 Hz), 155.3, 149.2, 136.8, 134.4, 132.5, 128.8, 128.4, 123.7 (d, *J*=276.0 Hz), 122.5, 121.6, 51.3, 41.9 (q, *J*=30.0 Hz); HRMS (ESI<sup>+</sup>): *m*/*z* [M + Na]<sup>+</sup> calcd for C<sub>16</sub>H<sub>13</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>Na: 345.0821; found: 345.0825.

*Propyl Benzoyl(pyridin-2-ylmethyl)carbamate* (**5p**): colorless oil; yield: 48.8 mg, (82%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.56 (d, *J*=4.2 Hz, 1H), 7.70–7.62 (m, 3H), 7.48 (t, *J*=7.3 Hz, 1H), 7.41 (t, *J*=7.5 Hz, 2H), 7.30 (d, *J*=7.8 Hz, 1H), 7.20–7.13 (m, 1H), 5.18 (s, 2H), 3.92 (t, *J*=6.6 Hz, 2H), 1.27 (h, *J*=7.1 Hz, 2H), 0.58 (t, *J*=7.4 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  173.0, 156.9, 155.1, 149.4, 136.9, 136.6, 131.3, 128.1, 127.8, 122.2, 121.0, 68.7, 50.4, 21.5, 10.0; HRMS (ESI<sup>+</sup>): m/z [M + Na]<sup>+</sup> calcd for C<sub>17</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub>Na: 321.1210; found: 321.1225.

(*E*)-*N*-*Isobutyryl*-*N*-(*pyridin*-2-*ylmethyl*)*but*-2-enamide (**5q**): colorless oil; yield: 48.3 mg, (98%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52(d, *J*=4.8 Hz, 1H), 7.63 (t, *J*=7.6 Hz, 1H), 7.17 (d, *J*=7.3 Hz, 2H), 7.07–6.98 (m, 1H), 6.48 (d, *J*=14.9, 1H), 5.04 (s, 2H), 3.32 (p, *J*=6.1, 5.6 Hz, 1H), 1.89 (d, *J*=6.9 Hz, 3H), 1.15 (d, *J*=6.6 Hz, 6H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  181.5, 169.5, 156.9, 149.3, 144.9, 136.8, 125.2, 122.3, 121.2, 49.2, 34.8, 19.5, 18.3; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>14</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>Na: 269.1260; found: 269.1256.

*Methyl* 3-Oxo-3-(*N*-(*pyridin-2-ylmethyl*)*isobutyramido*) propanoate (**5r**): colorless oil; yield: 51.7 mg, (93%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, *J*=4.8 Hz, 1H), 7.67 (t, *J*=7.6 Hz, 1H), 7.29 (d, *J*=7.8 Hz, 1H), 7.19 (t, *J*=5.0 Hz, 1H), 5.12 (s, 2H), 3.93 (s, 2H), 3.73 (s, 3H), 3.05 (p, *J*=6.6 Hz, 1H), 1.10 (d, *J*=6.6 Hz, 6H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$ 181.1, 169.2, 168.0, 156.3, 149.3, 137.1, 122.5, 121.2, 52.3, 48.7, 46.0, 34.2, 19.2; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>14</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>Na: 301.1159; found: 301.1158.

*N*-(2-*Methoxyacetyl*)-*N*-(*pyridin*-2-*ylmethyl*)*isobutyramide* (**5s**): colorless oil; yield: 45.0 mg, (90%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.50 (d, *J*=4.7 Hz, 1H), 7.64 (td, *J*=7.7, 1.7 Hz, 1H), 7.23–7.15 (m, 2H), 5.08 (s, 2H), 4.56 (s, 2H), 3.45 (s, 3H), 3.13 (hept, *J*=6.7 Hz, 1H), 1.12 (d, *J*=6.7 Hz, 6H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 181.0, 174.1, 156.2, 149.3, 136.9, 122.5, 121.5, 74.7, 59.2, 48.2, 34.2, 19.3; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>13</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub>Na: 273.1210; found: 273.1218.

*N-Isobutyryl-N-(pyridin-2-ylmethyl)cyclopropanecarboxamide* (**5**t): colorless oil; yield: 41.8 mg, (85%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.53 (d, *J*=3.8 Hz, 1H), 7.66 (td, *J*=7.7, 1.7 Hz, 1H), 7.18 (dd, *J*=7.6, 4.0 Hz, 2H), 5.16 (s, 2H), 3.41 (p, *J*=6.7 Hz, 1H), 2.21–2.15 (m, 1H), 1.17 (d, *J*=6.7 Hz, 6H), 1.10–1.06 (m, 2H), 0.92–0.87 (m, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  181.5, 177.8, 157.1, 149.2, 137.0, 122.4, 121.1, 49.2, 35.1, 19.6, 15.8, 10.5; HRMS (ESI<sup>+</sup>): *m/z* [M + Na]<sup>+</sup> calcd for C<sub>14</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>Na: 269.1260; found: 269.1259.

4-Chloro-N-isobutyryl-N-(pyridin-2-ylmethyl)butanamide (**5u**): colorless oil; yield: 54.2 mg, (96%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.52 (d, J=33.8 Hz, 1H), 7.65 (t, J=7.6 Hz, 1H), 7.17 (d, J=7.3 Hz, 2H), 5.06 (s, 2H), 3.60 (t, J=6.2 Hz, 2H), 3.27 (p, J=6.6 Hz, 1H), 2.98 (t, J=6.8 Hz, 2H), 2.13 (p, J=6.56 Hz, 2H), 1.15 (d, J=6.6 Hz, 6H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  181.3, 175.7, 156.6, 149.3, 136.9, 122.5, 121.2, 48.8, 44.2, 35.1, 34.9, 27.7, 19.4; HRMS (ESI<sup>+</sup>): m/z [M + Na]<sup>+</sup> calcd for C<sub>14</sub>H<sub>19</sub>ClN<sub>2</sub>O<sub>2</sub>Na: 307.0998; found: 307.0997.

*Benzyl*(*2-(Benzoyl(pyridin-2-ylmethyl)carbamoyl)-6-methylphenyl)carbamate* (**5v**): colorless oil; yield: 92.0 mg (96%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.28 (d, *J*=4.9 Hz, 1H), 7.72 (dd, *J*=8.2, 1.3 Hz, 2H), 7.43 (td, *J*=7.7, 1.6 Hz, 1H), 7.34 (d, *J*=7.4 Hz, 1H), 7.29–7.21 (m, 5H), 7.16–7.07 (m, 4H), 7.04–6.96 (m, 2H), 6.86 (d, *J*=6.6 Hz, 2H), 5.06 (s, 2H), 4.91 (s, 2H), 2.24 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 171.2, 166.2, 155.8, 154.2, 148.7, 137.0, 135.7, 134.8, 134.5, 133.3, 133.2, 131.5, 128.5, 128.4, 128.2, 128.2, 127.7, 126.0, 122.1, 121.6, 69.2, 49.8, 19.0; HRMS (ESI<sup>+</sup>): m/z [M + Na]<sup>+</sup> calcd for C<sub>29</sub>H<sub>25</sub>N<sub>3</sub>O<sub>4</sub>Na: 502.1737; found: 502.1746.

#### **Declaration of conflicting interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this paper.

#### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this paper: This work was financially supported by the National Natural Science Foundation of China (No. 81102334, 31370372, 31170323), the Program for New Century Excellent Talents in University, State Education Ministry of China (NCET-2008-0224), and the Fundamental Research Funds for the Central Universities (2017KFYXJJ152).

#### Supplemental material

Supplemental material for this paper is available online.

#### References

- Maruyama HB, Suhara Y, Suzuki-Watanabe J, et al. J Antibiot 1975; 28: 636.
- Maruyama HB, Suhara Y, Suzuki-Watanabe J, et al. J Antibiot 1975; 28: 648.
- Bai W-J, Jackson SK and Pettus TRR. Org Lett 2012; 14: 3862.
- 4. Somers GF. Br J Pharmacol Chemother 1960; 15: 111.
- Nagle DG, Paul VJ and Ann Roberts M. *Tetrahedron Lett* 1996; 37: 6263.
- Takatani M, Yoshioka Y, Tasaka A, et al. J Med Chem 1989; 32: 56.
- Trova MP, Wissner A, Carroll ML, et al. *J Med Chem* 1993; 36: 580.
- 8. He Y, Wang Y, Liang X, et al. Org Lett 2018; 20: 7117.
- 9. Kataoka K, Wachi K, Jin X, et al. Chem Sci 2018; 9: 4756.
- Nicolaou KC and Mathison JN. Angew Chem Int Ed 2005; 117: 6146.
- Giovannini A, Savoia D and Umani-Ronchi A. J Org Chem 1989; 54: 228.
- Savoia D, Concialini V, Roffia S, et al. J Org Chem 1991; 56: 1822.
- 13. Andrus MB, Li W and Keyes RF. Tetrah Let 1998; 39: 5465.
- 14. Ke D, Zhan C, Li X, et al. Synlett 2009; 1506.
- Yamada S, Yaguchi S and Matsuda K. *Tetrahedron Lett* 2002; 43: 647.
- Schindler CS, Forster PM and Carreira EM. Org Lett 2010; 12: 4102.
- Tsunoda T, Sasaki O, Takeuchi O, et al. *Tetrahedron* 1991; 47: 3925.
- El-Saghier AMM, Makhlouf MA and Farhat MF. *Phosphorus Sulfur Silicon Relat Elem* 2006; 181: 1569.
- 19. Jiang Y-Y, Dou G-Y, Xu K, et al. *Org Chem Front* 2018; 5: 2573.
- Cink RD, Lukin KA, Bishop RD, et al. Org Process Res Dev 2020; 24: 183.
- 21. Lawer A, Rossi-Ashton JA, Stephens TC, et al. *Angew Chem Int Ed* 2019; 58: 13942.
- 22. Stephens TC, Lodi M, Steer AM, et al. *Chem Eur J* 2017; 23: 13314.
- 23. Stephens TC, Lawer A, French T, et al. *Chem Eur J* 2018; 24: 13947.