

Featured Article

# DIRECT SYNTHESIS OF N,N-DISUBSTITUTED FORMAMIDES BY OXIDATION OF IMINES USING HFIP/UHP SYSTEM

Natalia Llopis, Patricia Gisbert, and Alejandro Baeza

*J. Org. Chem.*, **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.0c01579 • Publication Date (Web): 12 Aug 2020

Downloaded from pubs.acs.org on August 13, 2020

## Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

# DIRECT SYNTHESIS OF *N,N*-DISUBSTITUTED FORMAMIDES BY OXIDATION OF IMINES USING HFIP/UHP SYSTEM

*Natalia Llopis, Patricia Gisbert and Alejandro Baeza\**

AUTHOR ADDRESS. Departamento de Química Orgánica and Instituto de Síntesis Orgánica (ISO), Facultad de Ciencias, Universidad de Alicante. Apdo. 99, E-03080 Alicante, Spain.

KEYWORDS: Green oxidation, Formamides, Fluorinated alcohol, Imines, Water peroxide

## ABSTRACT

The straightforward synthesis of *N,N*-disubstituted formamides using 1,1,1,3,3,3-hexafluoroisopropanol (HFIP) and H<sub>2</sub>O<sub>2</sub> combination is described. The unique features of HFIP allowed the utilization of a green oxidant such as H<sub>2</sub>O<sub>2</sub> and the products, arising from an oxidation-rearrangement sequence, were obtained in good to high yields under smooth reaction conditions.


## INTRODUCTION

Formamides are considered valuable intermediates in organic synthesis which have been extensively utilized in the synthesis of pharmaceutically relevant compounds such as fluoroquinolones,<sup>1</sup> substituted aryl imidazoles<sup>2</sup> or 1,2-dihydroquinolines.<sup>3</sup> Besides, during the last years, formamides have been also used as Lewis bases being employed as catalysts in numerous organic transformations.<sup>4</sup> Further to this, these organic compounds play an important role as reagents in the Vilsmeier formylation reaction.<sup>5</sup>


On the other hand, imines are considered very accessible and versatile compounds. Thus, it is not surprising that oxidation of imines has been addressed in a myriad of articles giving rise to the corresponding oxaziridines<sup>6</sup> or *N*-substituted amides<sup>7</sup> (Scheme 1, eq. a). However, to the best of our knowledge, there are only a couple of precedents in the literature describing the oxidative rearrangement of imines in order to get access to *N,N*-disubstituted formamides (Scheme 1, eq. b).<sup>8</sup> In addition, in both routes, the authors use organic peroxyacids, requiring the presence of a strong Lewis or Brønsted acid; hence generating a stoichiometric amount of non-degradable waste in a low atom-economy process. Therefore, the development of a more environmentally benign oxidative route for the synthesis of such formamides from readily available imines, which can overcome such problems, would be highly desirable. In this regard, continuing with an ongoing project about the use of fluorinated alcohols in organic transformations as solvents and reaction mediators,<sup>9,10</sup> we envisioned the employment of H<sub>2</sub>O<sub>2</sub> or the more stable form UHP (urea-hydrogen peroxide adduct) as a green alternative oxidants to those previously described ones in combination with 1,1,1,3,3,3-hexafluoroisopropanol (HFIP) for such purpose (Scheme 1, eq. c).<sup>11</sup> The use of this system would considerably reduce the amount and the environmental impact of the generated waste.<sup>12</sup> The results of this research are herein reported.

### Scheme 1. Imines oxidation


a) Well-known procedures<sup>6,7</sup>



b) 2 reported procedures<sup>8</sup>



c) Our proposal




## RESULTS AND DISCUSSION

The optimization of the reaction conditions was tackled, using *N*-benzylideneaniline (**1a**) as model substrate (Table 1). Firstly, different organic polar protic solvents (namely HFIP, H<sub>2</sub>O, MeOH and TFE) were tested using 2 equiv. of H<sub>2</sub>O<sub>2</sub> (30% v/v) as oxidizing agent at 45 °C (Table 1, entries 1-4). As depicted in Table 1, only HFIP gave a good conversion towards the formation of **2a** (Table 1, entry 1). It is noteworthy that, although in poor conversion, MeOH gave rise to the formation of the regioisomer **3a** (Table 1, entry 3). Other commonly employed polar aprotic and apolar organic solvents were also essayed (DMSO, MeCN, EtOAc, THF, acetone, CHCl<sub>3</sub>, PhMe) remaining the imine unaltered or slightly hydrolysed. Attempts to ameliorate the conversion of **2a** were performed by increasing or lowering the amount of H<sub>2</sub>O<sub>2</sub> employed using HFIP as solvent (Table 1, entries 5 and 7). However, whereas the use of 3 equiv. barely changed the result, the employment of 1.5 equiv. resulted in lower conversion. The same experiments were performed with MeOH, trying to favor the formation of **3a** (Table 1, entries 6 and 8). Unfortunately, the same trend was observed and no significant improvement was accomplished. It is also noteworthy that, at this point, different solvent mixtures HFIP/MeOH

and HFIP/H<sub>2</sub>O were also tested without further success. Focused on the formation of compound **2a**, refining the reaction conditions in order to increase the conversion was next addressed. Thus, the concentration was examined and it was observed that changing the concentration from 1 M to 0.6 M improved the conversion of **2a** reaching up to 91% (Table 1, compare entry 9 and 10). The reaction at room temperature was also essayed but the conversion was not as high as at 45 °C (Table 1, entry 11). Finally, from the presence of benzaldehyde and aniline in the GC-MS analysis, arising from the partial hydrolysis of imine, it was decided to switch to an anhydrous H<sub>2</sub>O<sub>2</sub> source in order to minimize such background reaction. In this regard, when UHP was used instead, an excellent conversion towards the formation of **2a** was achieved (Table 1, entry 12). Further changes in concentration or temperature did not produce any substantial improvement (Table 1, entries 13-15).

**Table 1. Optimization of the reaction conditions<sup>a</sup>**



Entry	H <sub>2</sub> O <sub>2</sub> source (x equiv.)	Solvent	Temp (° C)	Conv. ( <b>2a</b> / <b>3a</b> ratio) <sup>b</sup>
1	H <sub>2</sub> O <sub>2</sub> (2 equiv.)	HFIP	45	85% (98/2)
2	H <sub>2</sub> O <sub>2</sub> (2 equiv.)	H <sub>2</sub> O	45	<10%
3	H <sub>2</sub> O <sub>2</sub> (2 equiv.)	MeOH	45	35% (4/96)
4	H <sub>2</sub> O <sub>2</sub> (2 equiv.)	TFE	45	<15%
5	H <sub>2</sub> O <sub>2</sub> (1.5 equiv.)	HFIP	45	77% (95/5)

6	H <sub>2</sub> O <sub>2</sub> (1.5 equiv.)	MeOH	45	<15%
7	H <sub>2</sub> O <sub>2</sub> (3 equiv.)	HFIP	45	82% (98/2)
8	H <sub>2</sub> O <sub>2</sub> (3 equiv.)	MeOH	45	39% (4/96)
9	H <sub>2</sub> O <sub>2</sub> (2 equiv.)	HFIP <sup>c</sup>	45	91% (>99/1)
10	H <sub>2</sub> O <sub>2</sub> (2 equiv.)	HFIP <sup>d</sup>	45	68% (95/5)
11	H <sub>2</sub> O <sub>2</sub> (2 equiv.)	HFIP <sup>c</sup>	25	86% (99/1)
<b>12</b>	<b>UHP (2 equiv.)</b>	<b>HFIP</b>	<b>45</b>	<b>97% (&gt;99/1)</b>
13	UHP (2 equiv.)	HFIP <sup>c</sup>	45	92% (>99/1)
14	UHP (2 equiv.)	HFIP <sup>d</sup>	45	87% (>99/1)
15	UHP (2 equiv.)	HFIP	25	93% (>99/1)


<sup>a</sup> Reaction conditions: all the reactions were carried out using 0.15 mmol of imine **1a** and the corresponding amount of oxidant in 150  $\mu$ L of the solvent at the given temperature for 15 h. <sup>b</sup> Conversion and **2a**/**3a** ratio, determined by GC-MS. <sup>c</sup> 250  $\mu$ L of HFIP were used (0.6 M). <sup>d</sup> 50  $\mu$ L of HFIP were used (3 M).





Once the optimal reaction conditions were selected, the scope of the methodology was performed by using different imines (Table 2). As previously commented, imine **1a** produced the corresponding *N,N*-diphenylformamide (**2a**) in high yield (Table 2, entry 1). Good yield was also achieved when imine **1b** was tested (Table 2, entry 2). However, imine derived from *para*-anisidine **1c** gave only decomposition and hydrolysis products regardless the reaction conditions employed, probably due to its more basic character, which makes the substrate more prone to be protonated by HFIP and hence more easily hydrolysed (Table 2, entry 3). Aldimine **1d** bearing an electron-withdrawing group in the aniline moiety, produced the desired product in a 62% yield (Table 2, entry 4). Next, aldimines synthesized from different *para*-substituted benzaldehydes were essayed, obtaining high yields in the corresponding disubstituted formamides regardless the electronic nature of such substituent (Table 2, entries 5-8). It is worth mentioning that in the case of imine **1h** bearing an electron-withdrawing group (CF<sub>3</sub>) produced a

substantial 10% of the corresponding isomer **3h** arising from the benzylic oxidation (Table 2, entry 8). Imine derived from salicylaldehyde was also explored but unfortunately turned out to be unreactive, probably due to an internal hydrogen bond, and low conversion towards the desired product was detected even when increasing the amount of UHP (Table 2, entry 9). This assumption was further reinforced in view of the good yield achieved for the product **2j** when *o*-anisaldehyde imine derivative **1j** was essayed (Table 2, entry 10). Imine **1k** bearing *para*-methyl substituents in both aromatic rings produced formamide **2k** in good yield (Table 2, entry 11). Next, heteroaromatic aldehyde imine derivatives were employed and different behaviors were observed depending on the electronic properties of the aromatic ring. Thus, whereas imine **1l** containing an electron-rich heteroaromatic ring such as furan, gave rise to decomposition and hydrolysis products, imine **1m** bearing a pyridine ring, considered as electron poor aromatic ring, produced compound **3m** in good yield as main product (Table 2, entries 12 and 13). Imines derived from aliphatic amines were next evaluated. When imine **1n** was the substrate of choice, the formation of a mixture of both *N*-butylbenzamide (**3n**) and the corresponding formamide **2n** was observed in moderate yield (Table 2, entry 14). However, the reaction with imine **1o** bearing a cyclohexane ring resulted in the formation of the oxaziridine **4o** (Table 2, entry 15). Unexpectedly, *N,N*-diphenylformamide **2a** was obtained in high yield when benzylamine imine derivative **1p** was employed (Table 2, entry 16). Next, imines derived from cyclohexanecarbaldehyde were examined. Good results were obtained with imine **1q** (Table 2, entry 17). Contrariwise, oxaziridine **4r** was obtained in excellent yields from imine **1r** (Table 2, entry 18). Finally, cinnamaldehyde derivative imine **1s** was also tested and to our surprise a complex mixture of products was obtained (Table 2, entry 19). We were able to distinguish (by GC-MS) among them benzaldehyde and formamide **2a** as major products (together with lower
















amounts of *N*-phenylformamide). At this point we speculate about a possible hydrolysis and retroaldol reaction, which would form benzaldehyde and aniline, hence the *in situ* formation of imine **1a** (see Scheme 2) which would lead to the formation of **2a**. Finally, a few ketimines were also tested. Firstly, imine derived from acetophenone (**1t**) was essayed, but unfortunately only 20% conversion towards the formation of the corresponding *N,N*-diphenylacetamide **2t** (Table 2, entry 20) was observed. Imine derived from cyclohexyl methyl ketone **1u** underwent hydrolysis under the reaction conditions (Table 2, entry 21). Finally, all alkylic ketimine **1v** produced the corresponding oxaziridine **4v** in excellent yields (Table 2, entry 22).
















**Table 2. Scope of the reaction<sup>a</sup>**









Entry	Imine	Product	Yield <sup>b</sup> (2/3 ratio) <sup>c</sup>
1	 <b>1a</b>	 <b>2a</b>	89 (>99/1)
2	 <b>1b</b>	 <b>2b</b>	82 (>99/1)



3	 <b>1c</b>	—	—
4	 <b>1d</b>	 <b>2d</b>	62 (>99/1)
5	 <b>1e</b>	 <b>2e</b>	87 (>99/1)
6	 <b>1f</b>	 <b>2b</b>	80 (>99/1)
7	 <b>1g</b>	 <b>2g</b>	88 (96/4)
8	 <b>1h</b>	 <b>2h</b>	91 (90/10)
9	 <b>1i</b>	 <b>2i</b>	<20% conv. <sup>d</sup>
10	 <b>1j</b>	 <b>2j</b>	90 (98/2)

11	 <b>1k</b>	 <b>2k</b>	79 (>99/1)
12	 <b>1l</b>	—	—
13	 <b>1m</b>	 <b>3m</b>	72 (15/85)
14	 <b>1n</b>	 <b>3n</b>	57 (35/65) <sup>e</sup>
15	 <b>1o</b>	 <b>4o</b>	98
16	 <b>1p</b>	 <b>2a</b>	83 (>99/1)
17	 <b>1q</b>	 <b>2q</b>	84 (>99/1)
18	 <b>1r</b>	 <b>4r</b>	95


19	 <b>1s</b>	Complex Mixture	>95% conv. <sup>d</sup>
20	 <b>1t</b>	 <b>2t</b>	~20% conv. <sup>d</sup>
21	 <b>1u</b>	—	—
22	 <b>1v</b>	 <b>4v</b>	95

<sup>a</sup> Reaction conditions: all the reactions were carried out using 0.15 mmol of imine **1** and 2 equiv. of UHP in 150  $\mu$ L of HFIP (1 M) at 45  $^{\circ}$ C for 15 h. <sup>b</sup> Isolated yield of **2** or the **2/3** mixture after preparative TLC or chromatography column. <sup>c</sup> **2/3** ratio determined by GC-MS from the crude mixture. <sup>d</sup> Determined by GC-MS. <sup>e</sup> Not purely isolated; yield determined by <sup>1</sup>H NMR from the crude.

After the scope of the reaction was tested, we decided to explore the *one-pot* imine formation-oxidative rearrangement sequence (Scheme 2). Thus, onto *in situ* formed benzylideneaniline **1a**, synthesised by mixing benzaldehyde and aniline for 20 minutes in the absence of solvent, UHP (2 equiv.) and HFIP was added in one portion. After 15 hours and subsequent purification the formamide **2a** was obtained in 81% yield. It is also worth mentioning that when this *one-pot* was implemented in a larger scale reaction (5 mmol), the corresponding formamide **2a** was obtained in a notable 73% yield. In addition, since benzylidene amine it is not a very stable compound, we decided to perform the same *one-pot* sequence reaction but using ammonium hydroxide. Using a slightly modified procedure (longer reaction time to form the imine and 5 equiv. of both


ammonium hydroxide and UHP) we were able to isolate benzamide **3w** in a good 59% yield, being in addition, the only regioisomer observed.

**Scheme 2. One-pot imine formation-oxidation sequence.**



Finally, based on experimental evidences, a possible reaction mechanism was proposed as depicted in Scheme 3. Firstly, the corresponding oxaziridine **B** would be formed thanks to the electrophilic activation of hydrogen peroxide by HFIP<sup>13</sup> which allows the nucleophilic attack of the iminic nitrogen and subsequent oxaziridine formation from the oxime derivative **A**. The formation of intermediate **B** is seemingly plausible from the fact that this kind of products have been isolated, for example in compound **4o**, **4r** and **4v**. From this intermediate, a Meinwald-type rearrangement mediated by HFIP could occur rendering the expected amide. Although less probable, a Beckmann-type rearrangement from intermediate **A** cannot be ruled out.

**Scheme 3. Proposed reaction mechanism.**



**CONCLUSIONS**

In summary, we have developed an efficient and environmentally benign methodology for the synthesis of formamides by means of the oxidation of aldimines using UHP as green oxidant. The use of this safe and non-toxic oxidant in absence of any Lewis acid has been possible because the presence of HFIP as solvent and reaction mediator. The peculiar properties of such fluorinated alcohols allow the electrophilic activation of hydrogen peroxide which is crucial for the reaction to proceed. The corresponding formamides **2** were obtained generally in good yields under smooth reaction conditions. However, the formation of the corresponding aryl amides **3** or oxaziridines **4** was also observed for imines derived from electron-poor aldehydes or cyclohexanamine, respectively. The *one-pot* imine formation-oxidative rearrangement sequence has been also described obtaining good results even when a big-scale reaction was performed. In this regard this *one-pot* sequence allowed the synthesis of benzamide in good yields using benzaldehyde and aqueous ammonia which to the best of our knowledge represents the first example of this transformation in absence of metal or Lewis acid catalysts. Finally, the mechanism was proposed to proceed *via* formation of the corresponding oxaziridine and subsequent rearrangement.

## EXPERIMENTAL SECTION

### General Remarks:

All reagents and solvents were obtained commercially and used without further purification. Substrates that were not commercially available were synthesized according to known literature procedures.

NMR spectra were performed on a Bruker AV-300 or Bruker AV-400 (Bruker Corporation) using CDCl<sub>3</sub> as solvent and TMS as internal standard unless otherwise stated.

Low resolution mass spectra (MS) were recorded in the electron impact mode (EI, 70 eV, He as carrier phase) using an Agilent GC/MS 5973 Network Mass Selective Detector spectrometer apparatus equipped with a HP-5MS column (Agilent technologies, 30 m  $\times$  0.25 mm) and giving fragment ions in  $m/z$  with relative intensities (%) in parentheses.

Analytical TLC was performed on Merck silica gel plates and the spots visualized with UV light at 254 nm. Flash chromatography employed Merck silica gel 60 (0.040-0.063 mm). Silica gel 60 F<sub>254</sub> containing gypsum was employed for preparative layer chromatography.

### ***General Procedure for the Oxidation Reaction of Imines***

In a capped tube, onto the corresponding imine (0.15 mmol), HFIP (150  $\mu$ L) and UHP (2.0 equiv.) were added in one portion. The reaction was then stirred at 45  $^{\circ}$ C in a sand bath for 15 hours, until the reaction was judged to be completed by GC-MS. After this time, the reaction mixture was filtered over silica/celite plug and then the solvent was evaporated and the crude material was directly purified by flash chromatography or preparative TLC using hexane/ethyl acetate mixtures.

Spectroscopic and analytical data for isolated compounds are given below:

*N,N-Diphenylformamide (2a)*:<sup>14</sup> Slightly yellow solid; purification by preparative TLC (hexane/ethyl acetate 8.0/2.0), 29.0 mg, 89% yield;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{H}}$  = 8.69 (s, 1H), 7.46 – 7.39 (m, 4H), 7.37 – 7.27 (m, 4H), 7.23 – 7.16 (m, 2H) ppm;  $^{13}\text{C}\{\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{C}}$  = 161.8, 141.7, 139.6, 129.7, 129.2, 127.1, 126.9, 126.1, 125.1 ppm; MS (EI):  $m/z$  197 ( $\text{M}^+$ , 100%), 169 (58), 168 (79), 167 (45), 104 (12), 77 (17), 66 (19), 51 (13).

*N*-Phenyl-*N*-(*p*-tolyl)formamide (**2b**):<sup>15</sup> Dark orange oil; purification by preparative TLC (hexane/ethyl acetate 8.0/2.0), 25.3 mg, 82% yield, mixture of rotamers; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ<sub>H</sub> = 8.69 (s, 1H), 8.64 (s, 1H), 7.47 – 7.35 (m, 5H), 7.35 – 7.26 (m, 5H), 7.24 (s, 2H), 7.20 – 7.14 (m, 4H), 7.13 – 7.05 (m, 2H), 2.39 (s, 3H), 2.37 (s, 3H) ppm; <sup>13</sup>C{H} NMR (75 MHz, CDCl<sub>3</sub>): δ<sub>C</sub> = 161.8, 141.9, 139.8, 139.2, 137.2, 136.9, 130.3, 129.8, 129.6, 129.1, 126.8, 126.6, 126.2, 125.8, 125.3, 124.7, 29.7, 21.1, 20.9 ppm; MS (EI): *m/z* 211 (M<sup>+</sup>, 100%), 183 (68), 182 (75), 180 (14), 168 (11), 167 (38), 108 (13), 91 (18), 80 (13), 77 (17).

*N*-(4-Acetylphenyl)-*N*-phenylformamide (**2d**):<sup>16</sup> White solid; purification by preparative TLC (hexane/ethyl acetate 8.5/1.5), 22.2 mg, 62% yield; mixture of rotamers; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ<sub>H</sub> = 8.83 (s, 1H), 8.66 (s, 1H), 8.17 – 8.09 (m, 2H), 8.06 – 7.94 (m, 7H), 7.95 – 7.86 (m, 2H), 7.80 (d, *J* = 8.8 Hz, 1H), 7.60 (ddd, *J* = 8.8, 5.0, 1.6 Hz, 1H), 7.56 – 7.35 (m, 11H), 7.25 – 7.20 (m, 4H), 2.62 (s, 4H), 2.61 (s, 4H) ppm; <sup>13</sup>C{H} NMR (101 MHz, CDCl<sub>3</sub>): δ<sub>C</sub> = 161.8, 161.3, 132.3, 130.0, 129.9, 129.8, 129.5, 129.4, 129.3, 128.9, 128.4, 127.9, 127.7, 127.1, 126.8, 126.1, 124.9, 123.4, 123.2, 119.3, 26.8, 26.6 ppm; MS (EI): *m/z* 239 (M<sup>+</sup>, 100%), 196 (72), 167 (58), 121 (23), 77 (12).

*N*-(4-Methoxyphenyl)-*N*-phenylformamide (**2e**):<sup>16</sup> Dark yellow solid; purification by preparative TLC (hexane/ethyl acetate 8.5/1.5), 36.1 mg, 87% yield, mixture of rotamers; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ<sub>H</sub> = 8.69 (s, 1H), 8.59 (s, 1H), 7.46 – 7.35 (m, 4H), 7.34 – 7.30 (m, 3H), 7.26 – 7.19 (m, 3H), 7.19 – 7.12 (m, 4H), 6.95 (dd, *J* = 8.9, 1.3 Hz, 4H), 6.79 (d, *J* = 1.0 Hz, 1H), 3.85 (s, 3H), 3.83 (s, 3H) ppm; <sup>13</sup>C{H} NMR (101 MHz, CDCl<sub>3</sub>): δ<sub>C</sub> = 161.9, 161.9, 129.6, 129.1, 127.7, 127.2, 126.7, 126.5, 125.4, 124.4, 114.9, 114.5, 55.5, 55.4 ppm; MS (EI): *m/z* 277 (M<sup>+</sup>, 100%), 206 (32), 199 (17), 184 (76), 154 (20), 124 (13), 76 (11).

*N*-(4-Chlorophenyl)-*N*-phenylformamide (**2g**):<sup>17</sup> Brown solid; 30.4 mg, 88% yield (without further purification); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ<sub>H</sub> = 8.65 (d, *J* = 2.6 Hz, 1H), 7.49 – 7.36 (m, 5H), 7.30 – 7.24 (m, 2H), 7.19 (dd, *J* = 5.3, 3.3 Hz, 1H), 7.13 (d, *J* = 8.7 Hz, 1H) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>): δ<sub>C</sub> = 161.8, 161.5, 141.2, 140.3, 139.2, 138.1, 132.9, 132.4, 129.9, 129.4, 129.3, 127.5, 127.3, 127.2, 126.1, 126.1, 125.3 ppm; MS (EI): *m/z* 231 (M<sup>+</sup>, 100%), 205 (20), 204 (11), 203 (60), 168 (49), 167 (84), 166 (17), 77 (11), 66 (12).

*N*-Phenyl-*N*-[4-(trifluoromethyl)phenyl]formamide (**2h**): white sticky solid; purification by preparative TLC (hexane/ethyl acetate 8.0/2.0), 36.1 mg, 91% yield, mixture of rotamers (and regioisomer **3h**); only the signals for the major isomer **2h** are given: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ<sub>H</sub> = 8.78 (s, 1H), 8.66 (s, 2H), 8.01 (d, *J* = 8.0 Hz, 1H), 7.75 (d, *J* = 8.1 Hz, 1H), 7.66 (dd, *J* = 11.1, 8.7 Hz, 8H), 7.51 – 7.35 (m, 15H), 7.27 (dd, *J* = 7.6, 3.9 Hz, 5H), 7.21 (d, *J* = 7.6 Hz, 5H) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>): δ<sub>C</sub> = 164.4, 161.7, 161.2, 144.9, 142.8, 140.8, 138.7, 138.3, 137.6, 133.6, 130.0, 129.5, 129.1, 128.5, 128.1, 127.9, 127.7, 127.6, 126.9, 126.8, 126.7, 126.2, 126.2, 125.9, 125.8, 125.7, 125.4, 124.9, 124.0, 122.5, 120.4 ppm; MS (EI): *m/z* 265 (M<sup>+</sup>, 100%), 237 (73), 236 (34), 216 (17), 168 (16), 167 (42), 77 (11), 66 (13); HRMS (GC/MS-EI/Q-TOF): *m/z* calcd. for C<sub>14</sub>H<sub>10</sub>F<sub>3</sub>NO 265.0714, found 265.0717.

*N*-(2-Methoxyphenyl)-*N*-phenylformamide (**2j**): Dark orange oil; purification by preparative TLC (hexane/ethyl acetate 8.0/2.0), 31.1 mg, 90% yield, mixture of rotamers; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ<sub>H</sub> = 8.57 (s, 1H), 8.26 (s, 1H), 7.31 – 7.21 (m, 4H), 7.17 – 7.01 (m, 6H), 6.93 (d, *J* = 8.3 Hz, 3H), 3.69 (s, 3H), 3.69 (s, 3H) ppm; <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>): δ<sub>C</sub> = 163.42, 162.41, 155.6, 155.1, 141.6, 140.0, 129.9, 129.8, 129.7, 129.6, 129.5, 129.4, 128.8, 127.5, 126.4, 126.2, 124.8, 123.1, 121.2, 112.5, 112.4, 55.7 ppm; MS (EI): *m/z* 227 (M<sup>+</sup>, 53%), 200 (11), 199



(71), 185 (14), 184 (100), 183 (23), 167 (16), 166 (14), 156 (24), 129 (25), 128 (17), 77 (15), 51 (11); HRMS (GC/MS-EI/Q-TOF):  $m/z$  calcd. for  $C_{14}H_{13}NO_2$  227.0946; found 227.0939.

*N,N-Di-p-tolylformamide (2k)*:<sup>18</sup> Slightly orange solid; purification by preparative TLC (hexane/ethyl acetate 8.0/2.0), 26.8 mg, 79% yield;  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta_H$  = 8.63 (s, 1H), 7.21 (t,  $J$  = 5.5 Hz, 6H), 7.07 (d,  $J$  = 8.3 Hz, 2H), 2.39 (s, 3H), 2.37 (s, 3H) ppm;  $^{13}C\{H\}$  NMR (101 MHz,  $CDCl_3$ ):  $\delta_C$  = 161.8, 139.4, 137.2, 136.9, 136.7, 130.2, 129.7, 125.8, 124.9, 21.0, 20.9 ppm; MS (EI):  $m/z$  225 ( $M^+$ , 100%), 197 (65), 196 (69), 181 (23), 180 (29), 108 (20), 91 (19), 80 (20), 65 (13).

*N-Phenyl-3-pyridinecarboxamide (3m)*:<sup>19</sup> Dark orange solid; purification by preparative TLC (hexane/ethyl acetate 8.0/2.0), 20.5 mg, 72% yield (90:10 isomeric mixture), only the major isomer signals are given;  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta_H$  = 9.10 (s, 1H), 8.75 (d,  $J$  = 3.6 Hz, 1H), 8.39 (s, 1H), 8.23 (dt,  $J$  = 7.9, 1.8 Hz, 1H), 7.66 (d,  $J$  = 7.9 Hz, 2H), 7.47 – 7.35 (m, 3H), 7.19 (t,  $J$  = 7.4 Hz, 1H) ppm;  $^{13}C\{H\}$  NMR (101 MHz,  $CDCl_3$ ):  $\delta_C$  = 163.9, 152.2, 147.8, 137.5, 135.6, 130.9, 129.2, 125.1, 123.7, 120.5 ppm; MS (EI):  $m/z$  198 ( $M^+$ , 61%), 197 (16), 106 (100), 78 (54), 51 (17).

*N-Butylbenzamide (3n)*:<sup>20</sup> White solid; purification by preparative TLC (hexane/ethyl acetate 8.0/2.0), 15.5 mg, 57% yield (not pure isolated);  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta_H$  = 7.81 – 7.73 (m, 2H), 7.50 (dd,  $J$  = 5.0, 3.7 Hz, 1H), 7.47 – 7.42 (m, 2H), 6.10 (s, 1H), 3.54 – 3.44 (m, 2H), 1.71 – 1.58 (m, 2H), 1.45 (dd,  $J$  = 15.1, 7.4 Hz, 2H), 0.99 (t,  $J$  = 7.3 Hz, 3H) ppm; MS (EI):  $m/z$  177 ( $M^+$ , 11%), 135 (20), 134 (19), 105 (100), 77 (33).

*2-Cyclohexyl-3-phenyl-1,2-oxaziridine (4o)*:<sup>21</sup> White solid; purification by preparative TLC (hexane/ethyl acetate 8.5/1.5), 26.8 mg, 98% yield;  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta_H$  = 7.47 –

7.43 (m, 1H), 7.42 – 7.38 (m, 1H), 4.55 (s, 1H), 2.16 – 2.05 (m, 1H), 1.90 – 1.80 (m, 1H), 1.78 – 1.72 (m, 1H), 1.70 – 1.56 (m, 1H), 1.47 (ddd,  $J = 23.0, 12.2, 3.8$  Hz, 1H), 1.39 – 1.21 (m, 2H) ppm;  $^{13}\text{C}\{\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{C}} = 135.3, 129.9, 128.5, 127.4, 79.8, 70.2, 31.6, 29.2, 25.7, 24.6, 24.1$  ppm; MS (EI):  $m/z$  203 ( $\text{M}^+$ , 40%), 122 (78), 105 (100), 77 (42).

*N-Cyclohexyl-N-phenylformamide (2q)*:<sup>16</sup> Orange oil; purification by preparative TLC (hexane/ethyl acetate 8.5/1.5), 25.9 mg, 84% yield;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{H}} = 8.17$  (s, 1H), 7.51 – 7.36 (m, 3H), 7.23 – 7.12 (m, 2H), 4.54 – 4.27 (m, 1H), 1.88 (d,  $J = 12.2$  Hz, 2H), 1.78 (d,  $J = 13.8$  Hz, 2H), 1.66 – 1.57 (m, 1H), 1.47 – 1.25 (m, 5H), 1.01 (qt,  $J = 12.9, 3.7$  Hz, 1H) ppm;  $^{13}\text{C}\{\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{C}} = 162.6, 138.6, 129.3, 129.1, 128.2, 53.7, 31.4, 25.7, 25.3$  ppm; MS (EI):  $m/z$  203 ( $\text{M}^+$ , 15%), 160 (16), 132 (17), 122 (13), 121 (100), 93 (34), 77 (15).

*2,3-Dicyclohexyl-1,2-oxaziridine (4r)*: White solid; purification by preparative TLC (hexane/ethyl acetate 8.0/2.0), 28.5 mg, 95% yield; m.p. 48-49 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{H}} = 3.48$  (d,  $J = 6.6$  Hz, 1H), 1.99 (d,  $J = 13.4$  Hz, 1H), 1.86 – 1.71 (m, 8H), 1.71 – 1.57 (m, 4H), 1.35 – 1.12 (m, 8H) ppm;  $^{13}\text{C}\{\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{C}} = 85.3, 69.3, 40.2, 31.6, 290, 27.7, 27.5, 26.3, 25.7, 25.2, 24.5, 23.9$  ppm; MS (EI):  $m/z$  209 ( $\text{M}^+$ , 30%), 192 (19), 154 (23), 150 (18), 138 (42), 128 (98), 127 (18), 126 (48), 125 (44), 110 (42), 98 (18), 95 (28), 84 (22), 83 (79), 82 (33), 81 (18), 68 (14), 67 (52), 56 (39), 55 (100), 54 (55); HRMS (GC/MS-EI/Q-TOF):  $m/z$  calcd. for  $\text{C}_{13}\text{H}_{23}\text{NO}$  209.1780, found 209.1778.

*2-Cyclohexyl-1-oxa-2-azaspiro[2.5]octane (4v)*: Slightly yellow oil; purification by flash chromatography (hexane/ethyl acetate 8.0/2.0), 25.7 mg, 95% estimated yield (not pure isolated);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{H}} = 2.55 – 2.27$  (m, 1H), 1.97 – 1.77 (m, 4H), 1.75 – 1.55 (m, 6H), 1.54 – 1.38 (m, 3H), 1.34 – 1.16 (m, 6H) ppm;  $^{13}\text{C}\{\text{H}\}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta_{\text{C}} = 85.1, 60.2,$

36.6, 31.9, 29.7, 29.2, 27.8, 25.7, 25.5, 25.4, 24.9, 24.4, 24.2 ppm; MS (EI):  $m/z$  195 ( $M^+$ , 15%), 179 (20), 178 (34), 152 (15), 138 (12), 136 (25), 114 (100), 113 (11), 98 (47), 97 (12), 96 (20), 83 (16), 82 (15), 81 (21), 69 (18), 67 (24), 56 (13), 55 (52), 54 (20); HRMS (GC/MS-EI/Q-TOF):  $m/z$  calcd. for  $C_{12}H_{21}NO$  195.1623, found 195.1625.

*Benzamide (3w)*:<sup>22</sup> White solid; purification by preparative TLC (hexane/ethyl acetate 9.0/1.0), 12.3 mg, 59% yield;  $^1H$  NMR (400 MHz, DMSO- $d_6$ ):  $\delta_H$  = 7.99 (s, 1H), 7.88 (dd,  $J$  = 5.3, 3.4 Hz, 2H), 7.51 (ddd,  $J$  = 6.3, 3.7, 1.3 Hz, 1H), 7.48 – 7.41 (m, 2H), 7.38 (s, 1H) ppm;  $^{13}C\{^1H\}$  NMR (101 MHz, DMSO- $d_6$ ):  $\delta_C$  = 168.4, 134.7, 131.7, 128.6, 127.9 ppm; MS (EI):  $m/z$  121 ( $M^+$ , 84%), 105 (100), 77 (83), 51 (25).

## ASSOCIATED CONTENT

### Supporting Information.

The following files are available free of charge.

Copies of NMR spectra for all isolated compounds.

## AUTHOR INFORMATION

### Corresponding Author

\*E-mail: alex.baeza@ua.es.

ORCID: 0000-0002-6230-1162.

### Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENT

We would like to thank the Spanish Ministerio de Economía, Industria y Competitividad (CTQ2015-66624-P and CTQ2017-88171-P) and Spanish Ministerio de Ciencia, Innovación y Universidades (PGC2018-096616-B-I00) and the University of Alicante (VIGROB-316/19, UADIF19-106) for the financial support.

## REFERENCES

- (1) Jackson, A.; Meth-Cohn, O. A New Short and Efficient Strategy for the Synthesis of Quinolone Antibiotics. *J. Chem. Soc., Chem. Commun.* **1995**, 1319-1320.
- (2) Chen, B. C.; Bednarz, M. S.; Zhao, R.; Sundeen, J. E.; Chen, P.; Shen, Z.; Skoumbourdis, A. P.; Barrish, J. C. A New Facile Method for the Synthesis of 1-Arylimidazole-5-carboxylates. *Tetrahedron Lett.* **2000**, *41*, 5453-5456.
- (3) Kobayashi, K.; Nagato, S.; Kawakita, M.; Morikawa, O.; Konishi, H. Synthesis of 1-Formyl-1,2-dihydroquinoline Derivatives by a Lewis Acid-Catalyzed Cyclization of *o*-(1-Hydroxy-2-alkenyl)phenyl Isocyanides. *Chem. Lett.* **1995**, 575-576.
- (4) (a) Kobayashi, S.; Nishio, K. Facile and Highly Stereoselective Synthesis of Homoallylic Alcohols Using Organosilicon Intermediates. *J. Org. Chem.* **1994**, *59*, 6620-6628. (b) Kobayashi, S.; Yasuda, M.; Hachiya, I. Trichlorosilane-Dimethylformamide (Cl<sub>3</sub>SiH-DMF) as an Efficient Reducing Agent. Reduction of Aldehydes and Imines and Reductive Amination of Aldehydes under Mild Conditions Using Hypervalent Hydridosilicates. *Chem. Lett.* **1996**, 407-408.
- (5) Downie, I. M.; Earle, M. J.; Heaney, H.; Shuhaibar, K. F. Vilsmeier Formylation and Glyoxylation Reactions of Nucleophilic Aromatic Compounds Using Pyrophosphoryl Chloride. *Tetrahedron.* **1993**, *49*, 4015-4034.

(6) (a) Tanaka, N.; Tsutsumi, R.; Uraguchi, D.; Ooi, T. *N*-Sulfonyl  $\alpha$ -Imino Ester-Derived Chiral Oxaziridines: Catalytic Asymmetric Synthesis and Application as a Modular Chiral Organic Oxidant. *Chem. Commun.* **2017**, 53, 6999-7002. (b) Ji, N.; Yuan, J.; Xue, S.; Zhang, J.; He, W. Novel Chiral Thiourea Organocatalysts for the Catalytic Asymmetric Oxaziridination. *Tetrahedron.* **2016**, 72, 512-517. (c) Kraïem, J.; Ghedira, D.; Ollevier, T. Hydrogen Peroxide/Dimethyl Carbonate: a Green System for Epoxidation of *N*-Alkylimines and *N*-Sulfonylimines. One-pot Synthesis of *N*-Alkyloxaziridines from *N*-Alkylamines and (Hetero)aromatic Aldehydes. *Green Chem.* **2016**, 18, 4859-4864. (d) Jin, Y.; Zhang, T.; Zhang, W.; Chang, S.; Feng, B. Enantioselective Organocatalytic Oxaziridination of *N*-Tosyl Aldimine Catalyzed by Cinchona Alkaloid-Ester Derivatives. *Chirality*, **2014**, 26, 150-154. (e) Tsutsumi, R.; Kim, S.; Uraguchi, D.; Ooi, T. The Practical Preparation of Chiral *N*-Sulfonyl Oxaziridines via Catalytic Asymmetric Payne Oxidation. *Synthesis.* **2014**, 46, 871-878. (f) Zhang, T.; He, W.; Zhao, X.; Jin, Y. Asymmetric Oxaziridination Catalyzed by Cinchona Alkaloid Derivatives Containing Sulfide. *Tetrahedron.* **2013**, 69, 7416-7422. (g) Uraguchi, D.; Tsutsumi, R.; Ooi, T. Catalytic Asymmetric Oxidation of *N*-Sulfonyl Imines with Hydrogen Peroxide–Trichloroacetonitrile System. *J. Am. Chem. Soc.* **2013**, 135, 8161-8164 and references cited therein.

(7) (a) De Souza, G. F. P.; von Zuben, T. W.; Salles Jr, A. G. “On Water” Metal-Catalyst-Free Oxidative Coupling–Amidation of Amines To Access Imines and Amides. *ACS Sustainable Chem. Eng.* **2017**, 5, 8439-8446. (b) Rostamnia, S.; Doustkhah, E.; Golchin-Hosseini, H.; Zeynizadeh, B.; Xin, H.; Luque, R. Efficient Tandem Aqueous Room Temperature Oxidative Amidations Catalysed by Supported Pd Nanoparticles on Graphene Oxide. *Catal. Sci. Technol.* **2016**, 6, 4124-4133. (c) Liang, J.; Lv, J.; Shang, Z.-C. Metal-Free Synthesis of Amides by Oxidative Amidation of Aldehydes with Amines in PEG/Oxidant System. *Tetrahedron.* **2011**, 67, 8532-8535. (d) Mohamed, M. A.; Yamada, K.-I.; Tomioka, K. Accessing the Amide Functionality by the Mild and Low-Cost Oxidation of Imine. *Tetrahedron Lett.* **2009**, 50, 3436-3438. (e) Leung, C. H.; Voutchkova, A. M.; Crabtree, R. H.; Balcells, D.; Eisenstein, O. Atom Economic Synthesis of Amides *via* Transition Metal Catalyzed Rearrangement of Oxaziridines. *Green Chem.* **2007**, 9, 976-979. (f) Larsen, J.; Jørgensen, K. A.; Christensen, D. Duality of the Permanganate Ion in the Oxidation of Imines. Oxidation of Imines to Amides *J. Chem. Soc., Perkin Trans. 1*, **1991**, 5, 1187-1190.

- (8) (a) An, G.-I.; Kim, M.; Kim, J. Y.; Rhee, H. Oxidation of Aldimines to Amides by *m*-CPBA and  $\text{BF}_3 \cdot \text{OEt}_2$ . *Tetrahedron Lett.* **2003**, *44*, 2183-2186. (b) Nongkunsarn, P.; Ramsden, C. A. Oxidative Rearrangement of Imines to Formamides Using Sodium Perborate. *Tetrahedron*. **1997**, *53*, 3805-3830. (c) Nongkunsarn, P.; Ramsden, C. A. An Oxidative Rearrangement of *C,N*-Diaryldimines to Formamides Using Sodium Perborate. *Tetrahedron Lett.* **1993**, *34*, 6773-6776.
- (9) (a) Pérez, J. M.; Maquilón, C.; Ramón, D. J.; Baeza, A. Hexafluoroisopropanol-Promoted Metal-Free Allylation of Silyl Enol Ethers with Allylic Alcohols. *Asian J. Org. Chem.* **2017**, *6*, 1440-1444. (b) Trillo, P.; Baeza, A.; Nájera, C. Fluorinated Alcohols As Promoters for the Metal-Free Direct Substitution Reaction of Allylic Alcohols with Nitrogenated, Silylated, and Carbon Nucleophiles. *J. Org. Chem.* **2012**, *77*, 7344-7354.
- (10) For selected reviews about the use of fluorinated alcohols as promoters in organic transformations, see: (a) An, X.-D.; Xiao, J. Fluorinated Alcohols: Magic Reaction Medium and Promoters for Organic Synthesis. *Chem. Rec.* **2019**, *19*, 142-161. (b) Colomer, I.; Chamberlain, A. E. R.; Haughey, M. B.; Donohoe, T. J. Hexafluoroisopropanol as a Highly Versatile Solvent. *Nat. Rev. Chem.* **2017**, *1*, 0088. (c) Bégué, J.-P. ; Bonnet-Delpon, D.; Crousse, B. Fluorinated Alcohols: A New Medium for Selective and Clean Reaction. *Synlett* **2004**, 18-29.
- (11) Llopis, N.; Baeza, A. Oxidation of Electron-Rich Arenes Using HFIP-UHP System. *J. Org. Chem.* **2020**, *85*, 6159-6164.
- (12) Berkessel, A. "Catalytic Oxidations with Hydrogen Peroxide in Fluorinated Alcohol Solvents" *Modern Oxidation Methods*; Bäckvall, J.-E., Ed; Wiley-VCH: Weinheim, 2010, p 117.
- (13) (a) Neimann, K.; Neumann, R. Electrophilic Activation of Hydrogen Peroxide: Selective Oxidation Reactions in Perfluorinated Alcohol Solvents. *Org. Lett.* **2000**, *2*, 2861-2863. (b) Berkessel, A.; Adrio, J. A. Dramatic Acceleration of Olefin Epoxidation in Fluorinated Alcohols: Activation of Hydrogen Peroxide by Multiple H-Bond Networks. *J. Am. Chem. Soc.* **2006**, *128*, 13412-13420. (c) Berkessel, A.; Adrio, J. A.; Hüttenhain, D.; Neudörfl, J. M. Unveiling the "Booster Effect" of Fluorinated Alcohol Solvents: Aggregation-Induced Conformational Changes and Cooperatively Enhanced H-Bonding. *J. Am. Chem. Soc.* **2006**, *128*, 8421-8426.

- (14) Pathare, S. P.; Jain, A. K. H.; Akamanchi, K. G. Sulfated Tungstate: a Highly Efficient Catalyst for Transamidation of Carboxamides with Amines. *RSC Adv.* **2013**, *3*, 7697-7703.
- (15) Nongkunsarn, P.; Ramsden, C. A. Oxidative Rearrangement of Imines to Formamides using Sodium Perborate. *Tetrahedron* **1997**, *53*, 3805-3830.
- (16) Jiang, L. Copper/*N,N*-Dimethylglycine Catalyzed Goldberg Reactions Between Aryl Bromides and Amides, Aryl Iodides and Secondary Acyclic Amides. *Molecules* **2014**, *19*, 13448-13460.
- (17) Phillips, D. P.; Zhu, X. – F. Lau, T. L.; He, X.; Yang, K.; Liu, H. Copper-Catalyzed C–N Coupling of Amides and Nitrogen-Containing Heterocycles in the Presence of Cesium Fluoride. *Tetrahedron* **2009**, *50*, 7293-7296.
- (18) Yang, S.; Li, P.; Wang, Z.; Wang, L. Photoinduced Oxidative Formylation of *N,N*-Dimethylanilines with Molecular Oxygen without External Photocatalyst. *Org. Lett.* **2017**, *19*, 3386-3389.
- (19) Martinelli, J. R.; Clark, T. P.; Watson, D. A.; Munday, R. H.; Buchwald, S. L. Palladium-Catalyzed Aminocarbonylation of Aryl Chlorides at Atmospheric Pressure: The Dual Role of Sodium Phenoxide. *Angew. Chem. Int. Ed.* **2007**, *46*, 8460 –8463.
- (20) Sureshbabu, P.; Azeez, S.; Chaudhary, P.; Kandasamy, J. *tert*-Butyl Nitrite Promoted Transamidation of Secondary Amides Under Metal and Catalyst Free Conditions. *Org. Biomol. Chem.* **2019**, *17*, 845-850.
- (21) Singhal, S.; Jain, S. L.; Prasad, V. V. D. N.; Sain, B. An Environmentally Friendly Oxidation System for the Selective Oxygenation of Aldimines to Oxaziridines with Anhydrous TBHP and Alumina-Supported MoO<sub>3</sub> as a Recyclable Heterogeneous Catalyst. *Eur. J. Org. Chem.* **2007**, 2051–2054.
- (22) Chen, J.; Xia, Y.; Lee, S. Transamidation for the Synthesis of Primary Amides at Room Temperature. *Org. Lett.* **2020**, *22*, 3504–3508.

