

CHEM MED CHEM

CHEMISTRY ENABLING DRUG DISCOVERY

Accepted Article

Title: Synthesis and evaluation of new nifurtimox-adamantane adducts with trypanocidal activity

Authors: Angeliki-Sofia Foscolos, Ioannis Papanastasiou, Andrew Tsotinis, Martin C. Taylor, and John M. Kelly

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *ChemMedChem* 10.1002/cmdc.201900165

Link to VoR: <http://dx.doi.org/10.1002/cmdc.201900165>

WILEY-VCH

www.chemmedchem.org

A Journal of



COMMUNICATION

Synthesis and evaluation of new nifurtimox-adamantane adducts with trypanocidal activity

Angeliki-Sofia Foscolos,^[a] Ioannis Papanastasiou,^{[a]*} Andrew Tsotinis,^[a] Martin C. Taylor,^[b] and John M. Kelly^[b]

In memory of Professor George B. Foscolos, Department of Pharmacy, National and Kapodistrian University of Athens

- [a] Ms. A.-S. Foscolos, 0000-0001-8123-5060; Asst. Prof. I. Papanastasiou, 0000-0001-6974-2561; Prof. A. Tsotinis, 0000-0002-4334-4810
School of Health Sciences, Department of Pharmacy, Division of Pharmaceutical Chemistry
National and Kapodistrian University of Athens
Panepistimioupoli-Zografou, 157 84 Athens, GREECE
E-mail: papanastasiou@pharm.uoa.gr
- [b] Assoc. Prof. M.C. Taylor, 0000-0003-4147-0693; Prof. J.M. Kelly, 0000-0003-4305-5258
Department of Pathogen Molecular Biology, London School of Hygiene and Tropical Medicine
Keppel Street, London WC1 E7HT, UK

Supporting information for this article is given via a link at the end of the document.

Abstract: The synthesis and pharmacological evaluation of the C-1 substituted adamantane hydrazones **1a-d**, their C-2 substituted isomers **2a-d** and the C-1 substituted adamantane furanico carboxamides **3a-c** is described. The new adamantane derivatives exhibited an interesting pharmacological profile, in terms of trypanocidal activity and selectivity. Of the compounds tested, the phenylacetoxhydrazone **1b** showed the most promising profile against African trypanosomes ($EC_{50}=11 \pm 0.9$ nM; $SI_{TD}=770$).

Vector-borne kinetoplastid diseases, such as Chagas disease (CD),^[1] leishmaniasis and human African trypanosomiasis (HAT)^[2] threaten almost a billion people worldwide.^[3] The available drugs against these neglected tropical diseases (NTDs) are characterized by toxicity, limited efficacy and increasing resistance.^[4] For example, nifurtimox and benznidazole which are used against CD, can cause severe side-effects and treatment is often unsuccessful.^[5] This has led the World Health Organization (WHO) to coordinate public sector and private partnerships as part of a global effort to develop new and safer drugs.^[3] Fexinidazole, which was recommended by the European Medicines Agency in November 2018, is a successful example of the collaboration between the Drugs for Neglected Diseases initiative (DNDi) and pharmaceutical chemistry sector.^[6]




Figure 1. Nifurtimox, benznidazole, fexinidazole and adamantane carbohydrazone derivatives.

We have been interested in adamantane chemistry^[7-21] and have prepared a large number of analogues in an attempt to exploit adamantane's role in bioactivity. We prepared a series of adamantane carbohydrazones,^[22] and showed that these derivatives are very potent trypanocidals. Following this work, we now describe the preparation of a series of phenylhydrazone analogues, **1a-d** and **2a-d**, which in general have very promising antitrypanosomal activity. The new derivatives share common structural features with nifurtimox and also contain a phenyl

substituted adamantane ring. The adamantane core of the novel adducts accommodates aromatic substitutions at the C-1 and C-2 positions. In addition, the 5-nitro-2-furanyl pharmacophore group is attached to the phenyladamantane by a hydrazone bond with a linker, consisting of one or two methylene groups and an oxygen atom. This bond is altered to carboxamide in compounds **3a-c**, with compound **3c** also bearing a benzyl group on the amide nitrogen, to mimic the benznidazole skeletal arrangement (**Figure 2**).





Figure 2. New nifurtimox-adamantane adducts with trypanocidal activity.


The 4-(adamant-1-yl)phenylhydrazides, **1a-d**, were prepared by previously described methods,^[22] using as starting materials ethyl 4-(adamant-1-yl)benzoate (**2**),^[18] ethyl 4-(adamant-1-yl)phenyl acetate (**4**),^[18] ethyl 4-(adamant-1-yl)phenylpropionate (**6**)^[18] and ethyl 4-(adamant-1-yl)phenoxyacetate (**8**)^[23] as shown in **Scheme 1**.



Scheme 1. Reagents and conditions: (a) hydrazine hydrate, EtOH, **3**: 160 °C, autoclave, 6 d, 81%; **5**: 130 °C, autoclave, 5 d, 85%; **7**: reflux, 2 d, 80%; **9**: reflux, 1 d, 98%; (b) 5-nitro-2-furaldehyde, EtOH, r.t., 12 h, 55-80%.

The preparation of analogues **2a-c** was realised by the reaction sequence illustrated in **Scheme 2**.


COMMUNICATION



Scheme 2. Reagents and conditions: (a) Jones reagent 8N in acetone, 3 h, 40 %; (b) i. SOCl₂, gentle reflux, 60 min ii. EtOH (abs.), gentle reflux, 90 min, 79 %; (c) hydrazine hydrate, EtOH, 13 : 160 °C, autoclave, 6 d, 55%, 15 : 160 °C, autoclave, 6 d, 72%, 18 : reflux, 1 d, 79%; d) 5-nitro-2-furaldehyde, EtOH, r.t., 12 h, 39-82 %; (e) i. triethyl phosphonoacetate and NaH, anh. THF, 0 °C, Ar, 30 min and then at r.t., 60 min ii. saturated ammonium chloride solution, 0 °C, 66 %; (f): H₂/PtO₂, EtOH, 40 psi, r.t., 3 h, 91 %.


4-(2-Adamantyl)benzaldehyde (**10**)^[17] was oxidised by the Jones reagent to the corresponding benzoic acid **11**, which was esterified to the respective benzoate **12**. Under Emmons-Horner reaction conditions benzaldehyde **10** afforded the *trans*-ethyl cinnamate **16**, which was then hydrogenated to the corresponding saturated propionate **17**. Ethyl 4-(adamant-2-yl)benzoate (**12**), ethyl 4-(adamant-2-yl)phenylacetate (**14**)^[17] and ethyl 4-(adamant-2-yl)phenylpropionate (**17**) gave the desired hydrazones **2a-c** via standard procedures.

The synthesis of the 2-(4-(adamant-2-yl)phenoxy)-*N*-((5-nitrofuran-2-yl)methylene)acetohydrazide (**2d**) is shown in **Scheme 3**.



Scheme 3. Reagents and conditions: (a) ClCH₂CO₂Et, K₂CO₃, dry DMF, 83 %; (b) hydrazine hydrate, EtOH, reflux, 1 d, 98 %; (c) 5-nitro-2-furaldehyde, EtOH, r.t., 12 h, 92 %.


The 4-(2-adamantyl)phenol (**19**),^[24,25] was ethoxycarbonyl-methylated in the presence of potassium carbonate in dry DMF to afford the respective phenoxyacetate **20**, which led to the corresponding hydrazone **2d** following the above methods. Due to the inefficiency of the literature methods for the preparations of 4-(2-adamantyl)phenol (**19**), we developed the alternative synthetic routes to **19**, shown in **Scheme 4**.



Scheme 4. Reagents and conditions: (a) i. 4-methoxyphenylmagnesium bromide, Et₂O/THF, Ar, r.t., 3 h, ii. HCl 10%, 0 °C, 75 %; (b) i. TFA, DCM, Ar, r.t., 15 min, ii. Et₃SiH, Ar, r.t., 1 h, iii. H₂O, 0 °C, 24:72 %, 27:56 %; (c) i. BBBr₃, DCM, r.t., ii. H₂O, 0 °C, 97 %; (d) *n*-BuLi, THF, -80 °C; (e) i. adamantanone, -80 °C ii. H₂O, 0 °C, 45 %; (f) H₂/10% Pd-C, EtOAc, 2 h, almost quant. yield.

The first route involves the reaction of 4-methoxyphenylmagnesium bromide with 2-adamantanone (**22**) to give the carbinol **23**, which was then reduced to the 2-(4-methoxyphenyl)adamantane (**24**). O-demethylation of **24** led to the desired phenol **19** in 52% overall yield (from adamantanone **22**). The second reaction pathway involved the lithiation of the aryl bromide **25**,^[26] which was added to the 2-adamantanone (**22**) to give the 2-(4-(benzyloxy)phenyl)adamantan-2-ol (**26**). This was sequentially reduced and hydrogenated to give 4-(2-adamantyl)phenol (**19**) in 25 % overall yield (from the aryl bromide **30**).

The *N*-substituted-5-nitrofuran-2-carboxamides **3a-c** were prepared as shown in **Scheme 5**.



Scheme 5. Reagents and conditions: (a) i. SOCl₂, reflux, 1 h, ii. benzylamine, THF, r.t., 2 h, 83 %; (b) i. LiAlH₄, THF, reflux, 2 h ii. EtOH, H₂O, NaOH 10%, 0 °C, 83 %; (c) i. 5-nitro-2-furoic acid, SOCl₂, reflux, 1 h, ii. appropriate amine **28**, **29**, **32**, acetone/pyridine, dropwise, r.t., 1 d, 43-53 %.

4-(1-Adamantyl)benzoic acid (**30**)^[18] was converted to the 4-(adamant-1-yl)-*N*-benzylbenzamide (**31**), which was then reduced to the respective benzylamine **32**. The methanamine **28**,^[27] the ethanamine **29**^[27] and the benzylamine **32** were coupled with the intermediate, 5-nitro-2-furoic chloride, to afford the desired amides **3a-c**, as shown in **Scheme 5**.

The new nifurtimox-adamantane adducts were tested for their activity against the bloodstream form *Trypanosoma brucei* and *Trypanosoma cruzi* epimastigotes and the results are shown in **Tables 1** and **2**.

Table 1. Anti-*T. brucei* screening of nifurtimox-adamantane derivatives.

Cmpd	<i>T. brucei</i> EC ₅₀ (nM) ^[a]	<i>T. brucei</i> EC ₉₀ (nM) ^[a]	L6 cells EC ₅₀ (μM) ^[a]	S.I. EC ₅₀ L6/Tb ^[b]
1a	19.0 ± 1.00	28.0 ± 4.00	1.85 ± 0.08	95
1b	11.0 ± 0.90	23.9 ± 0.60	8.49 ± 0.17	770
1c	15.9 ± 1.80	36.9 ± 12.4	8.28 ± 0.24	520
1d	29.0 ± 1.00	35.0 ± 1.00	3.04 ± 0.21	105
2a	41.0 ± 3.00	81.0 ± 18.0	-	-
2b	12.4 ± 0.40	25.2 ± 1.90	2.90 ± 0.40	235
2c	17.4 ± 1.80	41.2 ± 5.90	3.74 ± 0.36	215
2d	87.0 ± 5.00	172 ± 9.00	-	-
a	881 ± 66.0	1230 ± 20.0	14.5 ± 0.40	16
3b	353 ± 38.0	488 ± 10.0	4.06 ± 0.23	12
3c	322 ± 17.0	653 ± 51.0	1.38 ± 0.15	4.3
Nifurtimox	2510 ± 90 ^[28]			

[a] EC₅₀ and EC₉₀; concentration that inhibits growth by 50% and 90%, respectively. [b] S.I.; selectivity index, the ratio of EC₅₀ values obtained with L6 cells and *T. brucei*.

Table 2. Anti-*T. cruzi* screening of nifurtimox-adamantane derivatives.

Cmpd	<i>T. cruzi</i> EC ₅₀ (nM) ^[a]	<i>T. cruzi</i> EC ₉₀ (nM) ^[a]	L6 cells EC ₅₀ (μM) ^[a]	S.I. EC ₅₀ L6/Tc ^[b]
1a	178 ± 13.0	404 ± 81.0	-	-
1b	33.9 ± 2.00	63.0 ± 5.00	8.49 ± 0.17	250
1c	70.7 ± 13.8	193 ± 7.00	8.28 ± 0.24	115
1d	85.0 ± 9.00	196 ± 7.00	-	-
2a	170 ± 5.00	330 ± 5.00	-	-
2b	55.4 ± 1.20	112 ± 3.00	2.90 ± 0.40	52
2c	95.2 ± 19.0	245 ± 14.0	3.74 ± 0.36	39
2d	330 ± 4.00	873 ± 9.00	-	-
3a	16200 ± 500	26600 ± 200	14.5 ± 0.40	0.9
3b	770 ± 89.0	1880 ± 43.0	4.06 ± 0.23	5.3
3c	1550 ± 140	5340 ± 190	1.38 ± 0.15	0.9
Nifurtimox	3100 ± 500 ^[29]			

[a] EC₅₀ and EC₉₀; concentration that inhibits growth by 50% and 90%, respectively. [b] S.I.; selectivity index, the ratio of EC₅₀ values obtained with L6 cells and *T. cruzi*.

It is apparent that the hydrazone nifurtimox-adamantane adducts **1a-d** and **2a-d** are more potent trypanocidals than the parent drug

COMMUNICATION

(I) and 3–9 fold more effective against *T. brucei* than *T. cruzi*. The most active adduct, with the best selectivity, was the phenylacetoxo hydrazone **1b** ($EC_{50}=11\pm0.9$ nM and $SI_{Tb}=770$). The spacer between the phenyl ring and the carbonyl group seems to have a significant impact on activity and cytotoxicity. It seems that the present structural modification comprising of a phenyl ring insertion between the adamantane core and the hydrazone side chain has improved the pharmacological characteristics of the new molecules, in terms of activity and toxicity, compared to the adamantane carbohydrazones (IV), we previously reported.^[22] The direct attachment of the hydrazone linker to the phenyl ring decreased the potency, and a one or two methylene spacer was associated with enhanced activity. Adducts **1b**, **1c** and **2b**, **2c** exhibited higher trypanocidal activity than analogues **1a** and **2a**, respectively. Conversely, the replacement of one methylene by an oxygen atom had a detrimental impact on activity. The position of substitution on the adamantane core (C-1, C-2), influenced cytotoxicity and the C-1 substituted hydrazones **1b** ($SI_{Tb}=770$) and **1c** ($SI_{Tb}=520$) were found to be more selective than the corresponding C-2 substituted adducts **2b** ($SI_{Tb}=235$) and **2c** ($SI_{Tb}=215$). The same pattern was also observed in the *T. cruzi* results. The third series of derivatives, the 5-nitro-furonic carboxamides **3a-c**, exhibited reduced activity in comparison to the hydrazones, which implies that these analogues may follow the same mechanistic pathway as nifurtimox.^[30–32] This observation is in agreement with our previous test results on adamantane carbohydrazones (IV).^[22] In conclusion, the new nifurtimox-adamantane hydrazone adducts show higher trypanocidal potency than both the parent drug (I) and the adamantane carbohydrazones (IV). The optimum potency arose from the combination of 4-(adamant-1-yl)phenyl substitution and a two methylene spacer, whilst their 4-(adamant-2-yl)phenyl substituted isomers were almost equipotent but less selective.

Experimental Section

General procedure for the preparation of hydrazones 1a-d and 2a-d.

A solution of the respective hydrazide (0.96 mmol) in ethanol (5 mL) was added to a solution of 5-nitro-2-furaldehyde (200 mg, 1.42 mmol) in ethanol (5 mL). The mixture was stirred under Argon in darkness for 12 h. The resulting precipitate was filtered, washed with a small amount of ethanol and dried to give the corresponding product as a yellow solid, which was recrystallized from chloroform.

General procedure for the preparation of carboxamides 3a-c.

5-Nitro-2-furoic acid (240 mg, 1.52 mmol) was treated with $SOCl_2$ (10 mL) at 60–65 °C for 60 min. Excess $SOCl_2$ was removed under reduced pressure and subsequently by azeotropic distillation with benzene (5 mL). The residue obtained was dissolved in anhydrous acetone (10–15 mL) and a solution of the respective amine (1.44 mmol) in anhydrous pyridine (2 mL) was added dropwise under stirring. The reaction mixture was stirred at room temperature for 1 d, the solvent removed *in vacuo* and the residue was treated with water. The resulting mixture was extracted with CH_2Cl_2 and the organic extracts washed with water, 10% sol. HCl and water. The combined organic phases were dried over Na_2SO_4 and the solvent evaporated. The crude product was purified by gradient flash column chromatography, using as eluent a mixture of CH_2Cl_2 /MeOH, to give the corresponding carboxamide as an orange viscous semi-solid, which was recrystallized from chloroform.

Conflicts of interest

The authors have no conflicts to declare.

Acknowledgements

We thank Dr. Dimitra Benaki, Department of Pharmacy, Division of Pharmaceutical Chemistry, National and Kapodistrian University of Athens for NMR experiments.



Keywords: 4-(adamant-1-yl)phenyl substitution • 4-(adamant-2-yl)phenyl substitution • hydrazone • nifurtimox • trypanocidal activity

References:

- [1] L. E. Echeverria, C. A. Morillo, *Infect. Dis. Clin. North Am.* **2019**, 33 (1), 119–134. DOI: 10.1016/j.idc.2018.10.015.
- [2] E. Bottieau, J. Clerinx, *Infect. Dis. Clin. North Am.* **2019**, 33 (1), 61–77. DOI: 10.1016/j.idc.2018.10.003.
- [3] S. P. S. Rao, M. P. Barrett, G. Dranoff, C. J. Faraday, C. R. Gimpelewicz, A. Hailu, C. L. Jones, J. M. Kelly, J. K. Lazdins-Helds, P. Mäser, et al., *ACS Infect. Dis.* **2019**, 5 (2), 152–157. DOI: 10.1021/acsinfecdis.8b00298.
- [4] A. F. Francisco, S. Jayawardhana, M. D. Lewis, M. C. Taylor, J. M. Kelly, *Parasitology* **2017**, 144 (14), 1871–1880. DOI: 10.1017/S0031182017001469.
- [5] M. Keenan, J. H. Chaplin, *Prog. Med. Chem.* **2015**, 54, 185–230. DOI: 10.1016/bs.pmch.2014.12.001.
- [6] "Fexinidazole (HAT) – DNDi," available at <https://www.dndi.org/diseases-projects/portfolio/fexinidazole/>, accessed on 20 February 2018.
- [7] I. Papanastasiou, A. Tsotinis, G. B. Foscolos, S. R. Prathalingam, J. M. Kelly, *J. Heterocycl. Chem.* **2008**, 45 (5), 1401–1406. DOI: 10.1002/jhet.5570450524.
- [8] I. Papanastasiou, A. Tsotinis, N. Kolocouris, S. R. Prathalingam, J. M. Kelly, *J. Med. Chem.* **2008**, 51 (5), 1496–1500. DOI: 10.1021/jm7014292.
- [9] I. Papanastasiou, G. B. Foscolos, A. Tsotinis, J. Ol, J. Ov, S. R. Prathalingam, J. M. Kelly, *Heterocycles* **2008**, 75 (8), 2043–2061.
- [10] G. Foscolos, I. Papanastasiou, A. Tsotinis, *Lett. Org. Chem.* **2008**, 5 (1), 57–59. DOI: 10.2174/157017808783330117.
- [11] I. Papanastasiou, K. C. Prousis, K. Georgikopoulou, T. Pavlidis, E. Scoulica, N. Kolocouris, T. Calogeropoulou, *Bioorg. Med. Chem. Lett.* **2010**, 20 (18), 5484–5487. DOI: 10.1016/j.bmcl.2010.07.078.
- [12] I. Papanastasiou, G. Foscolos B., A. Tsotinis, J. Kelly M., *African Trypanosomiasis: Clinical Symptoms, Diagnosis and Treatment*, Nova Science Publishers, NY **2016**.
- [13] S. Riganas, I. Papanastasiou, G. B. Foscolos, A. Tsotinis, K. Dimas, V. N. Kourafalos, A. Eleutheriades, V. I. Moutsos, H. Khan, P. Margarita, et al., *Med. Chem. Shariqah United Arab Emir.* **2012**, 8 (4), 569–586.
- [14] S. Riganas, I. Papanastasiou, G. B. Foscolos, A. Tsotinis, J. J. Bourguignon, G. Serin, J. F. Mirjolet, K. Dimas, V. N. Kourafalos, A. Eleutheriades, et al., *Bioorg. Med. Chem.* **2012**, 20 (10), 3323–3331. DOI: 10.1016/j.bmc.2012.03.038.
- [15] S. Riganas, I. Papanastasiou, G. B. Foscolos, A. Tsotinis, G. Serin, J.-F. Mirjolet, K. Dimas, V. N. Kourafalos, A. Eleutheriades, V. I. Moutsos, et al., *J. Med. Chem.* **2012**, 55 (22), 10241–10261. DOI: 10.1021/jm3013008.
- [16] A. Koperniku, I. Papanastasiou, G. B. Foscolos, A. Tsotinis, M. C. Taylor, J. M. Kelly, *MedChemComm.* **2013**, 4 (5), 856–859. DOI: 10.1039/C3MD00081H.
- [17] I. Papanastasiou, S. Riganas, G. B. Foscolos, A. Tsotinis, S. A. Akhtar, M. A. Khan, K. M. Rahman, D. E. Thurston, *Lett. Org. Chem.* **2014**, 12, 319–323.
- [18] A. Koperniku, A.-S. Foscolos, I. Papanastasiou, G. B. Foscolos, A. Tsotinis, D. Schols, *Lett. Org. Chem.* **2016**, 13, 171–176.
- [19] A.-S. Foscolos, I. Papanastasiou, A. Tsotinis, N. Kolocouris, G. B. Foscolos, A. Vocat, S. T. Cole, *Med. Chem.* **2017**, 13 (7), 670–681. DOI: 10.2174/1573406413666170125112709.
- [20] I. Papanastasiou, in *Ration. Drug Des.* (Eds: T. Mavromoustakos, T. F. Kellici), Vol. 1824, Springer New York, New York, NY **2018**.
- [21] M.-O. Georgiadis, V. Kourbeli, V. Ioannidou, E. Karakitsios, I. Papanastasiou, A. Tsotinis, D. Komiotis, A. Vocat, S. T. Cole, M. C. Taylor, et al., *Bioorg. Med. Chem. Lett.* **2019**, 29 (11), 1278–1281. DOI: 10.1016/j.bmcl.2019.04.010.
- [22] A.-S. Foscolos, I. Papanastasiou, G. B. Foscolos, A. Tsotinis, T. F. Kellici, T. Mavromoustakos, M. C. Taylor, J. M. Kelly, *MedChemComm.* **2016**, 7 (6), 1229–1236. DOI: 10.1039/C6MD00035E.
- [23] K. Lee, J.-I. Goo, H. Y. Jung, M. Kim, S. K. Boovanahalli, H. R. Park, M.-O. Kim, D.-H. Kim, H. S. Lee, Y. Choi, *Bioorg. Med. Chem. Lett.* **2012**, 22 (24), 7456–7460. DOI: 10.1016/j.bmcl.2012.10.046.
- [24] Y. Arredondo, M. Moreno-Mañas, R. Pleixats, *Synth. Commun.* **1996**, 26 (21), 3885–3895. DOI: 10.1080/00397919608003808.
- [25] N. Basarić, I. Žabčić, K. Mlinarić-Majerski, P. Wan, *J. Org. Chem.* **2010**, 75 (1), 102–116. DOI: 10.1021/jo902004n.
- [26] W.-J. Hu, X.-L. Zhao, M.-L. Ma, F. Guo, X.-Q. Mi, B. Jiang, K. Wen, *Eur. J. Org. Chem.* **2012**, 2012 (7), 1448–1454. DOI: 10.1002/ejoc.201101599.
- [27] C. Chyi Tseng, I. Handa, A. N. Abdel-Sayed, L. Bauer, *Tetrahedron.* **1988**, 44 (7), 1893–1904. DOI: 10.1016/S0040-4020(01)90332-1.
- [28] B. S. Hall, C. Bot, S. R. Wilkinson, *J. Biol. Chem.* **2011**, 286 (15), 13088–13095. DOI: 10.1074/jbc.M111.230847.
- [29] M. C. O. Campos, L. L. Leon, M. C. Taylor, J. M. Kelly, *Mol. Biochem. Parasitol.* **2015**, 201 (1), 83. DOI: 10.1016/j.molbiopara.2015.05.006.
- [30] S. R. Wilkinson, M. C. Taylor, D. Horn, J. M. Kelly, I. Cheeseman, *Proc. Natl. Acad. Sci.* **2008**, 105 (13), 5022–5027. DOI: 10.1073/pnas.0711014105.
- [31] S. R. Wilkinson, J. M. Kelly, *Expert Rev. Mol. Med.* **2009**, 11. DOI: 10.1017/S1462399409001252.
- [32] S. Patterson, S. Wyllie, *Trends Parasitol.* **2014**, 30 (6), 289–298. DOI: 10.1016/j.pt.2014.04.003.

COMMUNICATION

Entry for the Table of Contents



The insertion of a phenyl ring between the adamantane core and the hydrazone side chain of the new nifurtimox-adamantane adducts led to a higher trypanocidal activity and lower toxicity than the parent drug. The most active adduct with the best selectivity is the phenylacetoxo hydrazone **1b** ($EC_{50}=11 \pm 0.9$ nM and $SI_{Tb}=770$).