[-1

## Mizoroki–Heck-Type Reaction Mediated by Potassium *tert*-Butoxide\*\*

Eiji Shirakawa,\* Xuejing Zhang, and Tamio Hayashi\*

The Mizoroki–Heck reaction is a versatile method used to prepare styrene derivatives from readily available aryl halides and alkenes.<sup>[1]</sup> A major drawback of this method is the necessary use of an expensive palladium complex as a catalyst.<sup>[2]</sup> Addition of an aryl radical, which is generated from an aryl halide, to an alkene and subsequent elimination of a hydrogen radical in a similar way to homolytic aromatic substitution<sup>[3]</sup> is a possible approach for a new pathway leading to Mizoroki–Heck-type reactions. However, attack of aryl radicals to alkenes usually results in an addition reaction but not a substitution reaction.<sup>[4]</sup> Herein, we report that the substitution reaction of aryl halides with styrene derivatives takes place by just using KOtBu, EtOH, and DMF, possibly through a radical mechanism.<sup>[5,6]</sup>

Treatment of 4-iodotoluene (1a) with styrene (2o: 5 equiv), KOtBu<sup>[7]</sup> (3 equiv), and EtOH (20 mol %) in DMF at 80 °C for 2 hours gave 73 % yield of 4-methylstilbene (3ao) with 17 % yield of toluene (Table 1, entry 1).<sup>[8]</sup> Use of DMF as a solvent is essential: the reaction in MeCN, 1,4-dioxane, or toluene did not give 3ao at all (Table 1, entries 2–4).<sup>[9]</sup> The reaction took place also in the absence of EtOH but the yield of 3ao was slightly lower (Table 1, entry 5). A low conversion of 1a was observed when either sodium or lithium *tert*-butoxide was used, both of which have lower basicity than KOtBu (Table 1, entries 6 and 7). A decrease in the amount of styrene severely affected the yield (Table 1, entry 8).

The present *tert*-butoxide-promoted arylation of alkenes was applied to various aryl halides and styrene derivatives (Table 2).<sup>[10]</sup> Aryl and heteroaryl iodides underwent the coupling with styrene (**2o**), where substitution with a methoxy, phenyl, or methyl group on the benzene ring is compatible (Table 2, entries 1–5). In contrast, the reaction of phenyl iodides having an electron-withdrawing CF<sub>3</sub> group gave no coupling product with **2o** but reduction product **4g** 

[*] Prof. E. Shirakawa, Dr. X. Zhang, Prof. T. Hayashi
Department of Chemistry, Graduate School of Science
Kyoto University
Sakyo, Kyoto, 606-8502 (Japan)
Fax: (+81) 75-753-3988
E-mail: shirakawa@kuchem.kyoto-u.ac.jp

- [\*\*] This work was supported financially by a Grant-in-Aid for Scientific Research (the Global COE Program "Integrated Materials Science" on Kyoto University) from Ministry of Education, Culture, Sports, Science and Technology (Japan). We are grateful to Ms. Yuki Yamamoto, Dr. Kenji Kitayama, and Mr. Ikuo Takahashi (Daicel Chemical Industries, Ltd.) for ICP analysis. We thank Dr. Kazufumi Kohno (Tokyo University of Agriculture and Technology) for carrying out duplicate experiments.
- Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201008220.

$\begin{array}{c} \text{MOtBu} \\ \text{(3 equiv)} \\ \text{EtOH} \\ \text{1a (1 equiv)} & \text{2o (5 equiv)} \end{array}$							
Entry	Solvent	MOtBu	Conv. of <b>1 a</b> [%] <sup>[b]</sup>	Yield of <b>3 ao</b> [%] <sup>[b]</sup>	Yield of <b>4a</b> [%] <sup>[b]</sup>		
1	DMF	KOtBu	>99	73	17		
2	MeCN	KOtBu	4	<1	<1		
3	1,4-dioxane	KOtBu	20	<1	6		
4	toluene	KOtBu	6	<1	_		
5 <sup>[c]</sup>	DMF	KO <i>t</i> Bu	>99	58	15		
6	DMF	NaOtBu	5	<1	4		
7	DMF	LiOtBu	8	<1	<1		
8 <sup>[d]</sup>	DMF	KOtBu	> 99	39	32		

[a] The reaction was carried out in DMF (0.50 mL) at 80 °C for 2 h under nitrogen using 4-iodotoluene (**1**a: 0.23 mmol), styrene (**2**o: 1.2 mmol), MOtBu (0.69 mmol), and EtOH (0.046 mmol). [b] Determined by GC analysis. [c] In the absence of EtOH. [d] Styrene (0.28 mmol) was used. DMF = N,N-dimethylformamide.

with a high conversion (Table 2, entry 6). Almost no reaction took place with p-tolyl bromide or chloride (Table 2, entries 7 and 8), whereas not only iodide but also bromide and even chloride underwent coupling when the organic moiety was a 1- or 2-naphthyl group (Table 2, entries 9–13). Substitution by a conjugating group such as phenyl and alkenyl also enhanced the reactivity of aryl bromides and chlorides (Table 2, entries 14-17). Styrenes having an electron-donating or electron-withdrawing substituent were arylated in the reaction with 1a (Table 2, entries 18 and 19). Notably, 4-chlorostyrene reacted with styrene (20) at the chloroarene moiety (Table 2, entry 17) and with aryl halide 1a at the alkene moiety (Table 2, entry 19). Almost no conversion of 1a was observed in the reaction with a styrene having a strong electron-withdrawing  $CF_3$  group (Table 2, entry 20). The reaction was applicable to an ortho-substituted styrene and to the synthesis of a polymethoxylated stilbene (Table 2, entries 21 and 22). Benzofuran accepted arylation at the 2position under the same reaction conditions (Table 2, entry 23). Unfortunately, the reaction of alkyl-substituted alkenes, acrylic acid derivatives, or vinyl ethers did not give synthetically significant yields of products under the present conditions.

Formation of the reduction product **4** of aryl halide **1** suggests involvement of aryl radical intermediates, which are generated through single-electron transfer (SET) to Ar-X to give the radical anion  $[Ar-X]^{-}$ , and subsequent facile

Communications

Table 2: Coupling of aryl halides with styrene derivatives mediated by  $KOtBu.^{[a]}$ 

Ar	$Ar^{2} + 20-2t$	KO <i>t</i> Bu (3 equiv) EtOH (20 mol%) DMF, 80 °C	Ar <sup>1</sup>	Ar <sup>2</sup>
<b>1</b> or <b>2u</b> (1 equiv) (5 equiv)				3
Entry	Aryl halide 1	Ar <sup>2</sup> (2)	<i>t</i> [h]	Yield [%] <sup>[b]</sup>
1	Ph—I ( <b>1 b</b> )	Ph ( <b>2 o</b> )	2	66
2	4-MeOC <sub>6</sub> H <sub>4</sub> I ( <b>1 c</b> )	Ph ( <b>2o</b> )	2	65
3	4-PhC <sub>6</sub> H <sub>4</sub> —I ( <b>1 d</b> )	Ph ( <b>2o</b> )	2	71
4	2-MeC <sub>6</sub> H <sub>4</sub> —I ( <b>1 e</b> )	Ph ( <b>2o</b> )	2.5	61
5	2-pyridyl–I ( <b>1 f</b> )	Ph ( <b>2o</b> )	2	57
6	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub> -1 (1g)	Ph ( <b>2o</b> )	2	<1
7	4-MeC <sub>6</sub> H <sub>4</sub> -Br ( <b>1 a'</b> )	Ph ( <b>2o</b> )	5	2 <sup>[c]</sup>
8	4-MeC <sub>6</sub> H <sub>4</sub> Cl ( <b>1 a</b> ″)	Ph ( <b>2o</b> )	17	$< 1^{[d]}$
9	1-naphthyl—I ( <b>1h</b> )	Ph ( <b>2 o</b> )	2	68
10	1-naphthyl—Br ( <b>1 h'</b> )	Ph ( <b>2o</b> )	6	65
11	1-naphthyl–Cl ( <b>1 h″</b> )	Ph ( <b>2o</b> )	6	63
12	2-naphthyl—Br ( <b>1 i'</b> )	Ph ( <b>2o</b> )	3	76
13	2-naphthyl–Cl ( <b>1 i</b> ")	Ph ( <b>2o</b> )	3	83
14	4-PhC <sub>6</sub> H <sub>4</sub> —Br ( <b>1 d'</b> )	Ph ( <b>2o</b> )	2	62
15	4-styrylC <sub>6</sub> H <sub>4</sub> -Cl ( <b>1 j</b> ")	Ph ( <b>2o</b> )	2	69
16	4-vinylC <sub>6</sub> H <sub>4</sub> -Br ( <b>1 k</b> ')	Ph ( <b>2 o</b> )	2	62
17	4-vinylC <sub>6</sub> H <sub>4</sub> Cl ( <b>1 k</b> ")	Ph ( <b>2o</b> )	6	61
18	4-MeC <sub>6</sub> H <sub>4</sub> -I ( <b>1 a</b> )	4-MeOC <sub>6</sub> H <sub>4</sub> ( <b>2 p</b> )	5.5	56
19	4-MeC <sub>6</sub> H <sub>4</sub> -I ( <b>1 a</b> )	$4-ClC_6H_4$ ( <b>2q</b> = <b>1k''</b> )	2	67
20	4-MeC <sub>6</sub> H <sub>4</sub> -I ( <b>1 a</b> )	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub> (2r)	2	$< 1^{[d]}$
21	4-MeC <sub>6</sub> H <sub>4</sub> -I ( <b>1 a</b> )	2-MeC <sub>6</sub> H <sub>4</sub> ( <b>2 s</b> )	2	74
22	4-MeOC <sub>6</sub> H <sub>4</sub> -I ( <b>1 c</b> )	3,5-(MeO) <sub>2</sub> C <sub>6</sub> H <sub>3</sub> (2t)	3	56
23	4-MeC <sub>6</sub> H <sub>4</sub> -1 (1a)	benzofuran ( <b>2 u</b> )	3	55 <sup>[e]</sup>

[a] The reaction was carried out in DMF (0.50 mL) at 80 °C under nitrogen using an aryl halide (1: 0.23 mmol), an alkene (2: 1.2 mmol), KOtBu (0.69 mmol), and EtOH (0.046 mmol). [b] Yield of isolated product based on 1. Unless otherwise noted, conversion of 1 was >99%. [c] 46% conversion of 1, where 3- and 4-(*tert*-butoxy)toluene compounds were the main products. [d] Less than 5% conversion of 1. [e] 2-(4-Tolyl)benzofuran was obtained in 55% yield.

elimination of X<sup>-</sup>.<sup>[11]</sup> As metal alkoxides are known to act as single-electron donors towards aromatic compounds, alkyl iodides, and ketones,<sup>[12]</sup> SET from KOtBu to aryl halides is possibly in operation here. Treatment of 4-iodobiphenvl (1d: 1 equiv) with KOtBu (3 equiv) and EtOH (20 mol %) in DMF at 80°C for 0.25 hours gave reduction product 4d in 24% yield with 79% conversion of 1d (Table 3, entry 1).<sup>[13]</sup> The yield of 4d was largely improved by addition of 1,4-cyclohexadiene (5; 4.6 equiv), which is an efficient hydrogen radical donor, thus strongly suggesting the existence of aryl radical intermediates (Table 3, entry 2). The yield was almost the same in the absence of EtOH (Table 3, entry 3). In contrast, a low yield with a low conversion was observed in the reaction of *p*-tolyl chloride (**1**a''; Table 3, entry 4), which was not consumed at all also in the reaction with styrene (20; Table 1, entry 8). The correlation, which was observed also in the solvent effect (Table 3, entry 5), implies that aryl radical generation is a critical step in the coupling reaction.

Radical anions are known to be generated more easily when the starting arenes have lower absolute values of reduction potential and/or half-wave potential, which are lowered with substitution by an electron-withdrawing group including halogen (lowering effect: I > Br > Cl) and/or con**Table 3:** Reaction of aryl halides with KOtBu in the presence of 1,4-cyclohexadiene.<sup>[a]</sup>



[a] The reaction was carried out in DMF (0.50 mL) at 80 °C for 0.25 h under nitrogen using an aryl halide (1: 0.23 mmol), KOtBu (0.69 mmol), 1,4-cyclohexadiene (5: 1.1 mmol), and EtOH (0.046 mmol). [b] Determined by GC analysis. [c] In the absence of 5. [d] In the absence of EtOH.

jugating group such as phenyl and alkenyl as well as extended aromaticity with fused rings such as naphthalene.<sup>[14]</sup> The reactivity order of aryl halides here is consistent with aryl radical generation through radical anion intermediates, if we consider the following: 1) Aryl iodides are easily reduced to radical anions and thus fully consumed irrespective of their substituents in the reaction with styrene, though those having an electron-withdrawing group are converted exclusively into arene 4 (Table 2, entry 6), probably through further reduction of aryl radicals to aryl anions. 2) Substitution by a conjugating group or extended conjugation by introduction of a naphthalene ring makes aryl bromides and chlorides sufficiently reactive (compare Table 2, entries 7-8 vs. 10-17). 3) p-Tolyl iodide (1a) fails to react when the coupling partner is an electron-deficient styrene, which accepts an electron more easily than 1a (Table 2, entry 20).

Thus far, generated aryl radicals must add to styrene (2o), but the subsequent steps and the role of EtOH are unclear at present (Scheme 1). As adduct 7 has to lose a proton and an



Scheme 1. A plausible reaction mechanism.

Angew. Chem. Int. Ed. 2011, 50, 4671-4674



electron to be converted into coupling product **3**, the subsequent pathways can be diverted into three types (a–c) according to the timing of elimination of these species. In path a, a proton is first removed to give radical anion **8**, which may give an electron to **1** to therefore give **3** and regenerate the radical anion of **1** through  $S_{RN}$ 1 mechanism (a1)<sup>[11]</sup> or is oxidized by *t*BuO<sup>•</sup> (a2). Path b includes elimination of both species at once as H<sup>•</sup>, whereas oxidation of **7** is followed by deprotonation in path c. Considering the high oxidation ability of *t*BuO<sup>•</sup>, which is generated upon SET, it can possibly act as an oxidant as in paths a2, b, or c. As shown in Scheme 2,



 $\textit{Scheme 2.}\ Reaction of $\alpha$-cyclopropylstyrene that gives ring-opening products.$ 

the reaction of  $\alpha$ -cyclopropylstyrene (9) with *p*-tolyl iodide (1a) gave a mixture of (*E*)- and (*Z*)-1-(*p*-tolyl)-2-phenyl-2pentenes (10) in addition to their double bond isomers 11, albeit in a moderate combined yield (32%). Products 10 are likely to be derived from ring-opening products 13 of addition product 12 (corresponds to 7 in Scheme 2). As cyclopropyl-methyl radicals such as 12 are known to readily undergo ring opening,<sup>[15]</sup> this result supports the reaction pathway that includes addition of aryl radicals to styrenes. Base-catalyzed isomerization of 10 is likely to give 11.

In conclusion, we have disclosed transition-metal-free Mizoroki–Heck-type reaction, which proceeds through aryl radical intermediates and gives various stilbene derivatives by simply using KOtBu, EtOH, and DMF.

Received: December 28, 2010 Revised: February 25, 2011 Published online: April 6, 2011

**Keywords:** alkenes  $\cdot$  aryl radicals  $\cdot$  C–C coupling  $\cdot$  single-electron transfer  $\cdot$  synthetic methods

- For reviews, see: a) I. P. Beletskaya, A. V. Cheprakov, *Chem. Rev.* 2000, 100, 3009-3066; b) S. Braese, A. de Meijere in *Metal-Catalyzed Cross-Coupling Reactions*, 2nd ed. (Eds: A. de Meijere, F. Diederich), Wiley-VCH, Weinheim, 2004, pp. 217-315; c) M. Larhed, A. Hallberg in *Handbook of Organopalladium Chemistry for Organic Synthesis*, Vol. 1 (Eds.: E. Negishi, A. de Meijere), Wiley-Interscience, New York, 2002, chap. IV.2, pp. 1133-1178.
- [2] The Mizoroki-Heck reaction catalyzed by a metal other than palladium is also known but is rare. For a review, see: L.

Ackermann, R. Born in *Mizoroki–Heck Reaction* (Ed.: M. Oestreich), Wiley, Chichester, **2009**, pp. 383–403.

- [3] For reviews of homolytic aromatic substitution with aryl radicals, see: a) R. Bolton, G. H. Williams, *Chem. Soc. Rev.* 1986, *15*, 261–289; b) J. Fossey, D. Lefort, J. Sorba, *Free Radicals in Organic Chemistry*, Wiley, Chichester, 1995, Chapter 14, pp. 166–180; c) A. Studer, M. Bossart in *Radicals in Organic Synthesis, Vol. 2* (Eds.: P. Renaud, M. P. Sibi), Wiley-VCH, Weinheim, 2001, chap. 1.4, pp. 62–80.
- [4] For a recent review, see: M. R. Heinrich, *Chem. Eur. J.* 2009, 15, 820–833.
- [5] KOtBu in combination with or without a bidentate nitrogencontaining ligand is used in the coupling of aryl halides with (hetero)arenes, where generation of aryl radicals from aryl halides is considered to be involved; a) S. Yanagisawa, K. Ueda, T. Taniguchi, K. Itami, Org. Lett. 2008, 10, 4673-4676; b) W. Liu, H. Cao, H. Zhang, H. Zhang, K. H. Chung, C. He, H. Wang, F. Y. Kwong, A. Lei, J. Am. Chem. Soc. 2010, 132, 16737-16740; c) C.-L. Sun, H. Li, D.-G. Yu, M. Yu, X. Zhou, X.-Y. Lu, K. Huang, S.-F. Zheng, B.-J. Li, Z.-J. Shi, Nat. Chem. 2010, 2, 1044-1049. We have reported that coupling of aryl halides with arenes mediated by NaOtBu proceeds through a homolytic aromatic substitution pathway, where a catalytic amount of a phenanthroline derivative is essential for generation of the aryl radical from aryl halides; d) E. Shirakawa, K. Itoh, T. Higashino, T. Hayashi, J. Am. Chem. Soc. 2010, 132, 15537-15539. For a discussion on the mechanism of these reactions, see e) A. Studer, D. P. Curran Angew. Chem. 2011, in press; Angew. Chem. Int. Ed. 2011, in press.
- [6] The coupling of iodobenzene with styrene in the absence of any transition-metal catalysts is reported to take place at a high temperature and a high pressure in supercritical water, thus giving stilbene in a moderate yield. R. Zhang, F. Zhao, M. Sato, Y. Ikushima, *Chem. Commun.* 2003, 1548–1549.
- [7] For the reactions shown in Table 1–Table 3, KOtBu, which was purchased from Wako Pure Chemicals Industries, Ltd. (product number 163–08425), was used as received. Its ICP-AES and ICP-MS analysis showed that the contents of Co, Ni, Ru, Rh, Pd, Ag, Ir, Pt, and Au are less than 0.05 ppm and those of Fe and Cu are 1.0 ppm and 1.1 ppm, respectively. See also Ref. [8].
- [8] The reaction using a sublimed grade KOtBu (99.99% purity, Aldrich Co., product number 659878) or a KOtBu purified through sublimation by us gave coupling product **3ao** in a similar yield (80% or 79% yield, respectively). The contents of Fe and Cu in the sublimed KOtBu are < 0.2 ppm (under detection limit) and 0.25 ppm, respectively (ICP-AES analysis). These results amply indicate that the coupling is promoted by KOtBu itself rather than catalyzed by contaminated transition metals. We asked Dr. Kazufumi Kohno to repeat the reaction and he confirmed that the reaction under the same conditions gave **3ao** in 71% yield.
- [9] N,N-Dimethylacetamide and dimethylsulfoxide also were effective as solvents but addition of their enolates to styrene and the coupling products took place and thus the yield of **3ao** was slightly lower than that obtained when DMF was used.
- [10] The products are all E isomers. The corresponding Z isomer or 1,1-disubstitued product was not observed in any reaction in Table 2.
- [11] J. F. Bunnett, Acc. Chem. Res. 1978, 11, 413-420. See also Ref. [3b].
- [12] a) E. C. Ashby, A. B. Goel, R. N. DePriest, J. Org. Chem. 1981, 46, 2429-2431; b) C. G. Screttas, M. Micha-Screttas, C. T. Cazianis, Tetrahedron Lett. 1983, 24, 3287-3288; c) E. C. Ashby, D.-H. Bae, W.-S. Park, R. N. Depriest, W.-Y. Su, Tetrahedron Lett. 1984, 25, 5107-5110; d) E. C. Ashby, J. N. Argyropoulos, J. Org. Chem. 1986, 51, 3593-3597.

## Communications

- [13] A large amount of biphenyls having a *tert*-butoxy, dimethylamino, or hydroxy group at the 3- or 4-positions, all of which are most likely to be produced through 4-phenylbenzyne, was observed as by-products. These by-products were produced also in the coupling reaction but in lower amounts.
- [14] a) J. W. Sease, F. G. Burton, S. L. Nickol, J. Am. Chem. Soc. 1968, 90, 2595–2598; b) K. Meerholz, J. Heinze, J. Am. Chem. Soc.

**1989**, *111*, 2325–2326; c) J. Grimshaw, *Electrochemical Reactions and Mechanisms in Organic Chemistry*, Elsevier, Amsterdam, **2000**, pp. 54–56, pp. 122–126, and pp. 239–243.

[15] For examples, see: a) V. W. Bowry, J. Lusztyk, K. U. Ingold, J. Am. Chem. Soc. 1991, 113, 5687–5698; b) V. W. Bowry, K. U. Ingold, J. Am. Chem. Soc. 1991, 113, 5699–5707; c) A. Ogawa, I. Ogawa, N. Sonoda, J. Org. Chem. 2000, 65, 7682–7685.