## Zinc-Catalyzed Synthesis of Acylsilanes Using Carboxylic Acids and a Silylborane in the Presence of Pivalic Anhydride

Kenta Tatsumi, Sae Tanabe, Yasushi Tsuji,<sup>®</sup> and Tetsuaki Fujihara\*<sup>®</sup>

Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering, Kyoto University, Nishikyo-ku, Kyoto 615-8510, Japan

Supporting Information

ABSTRACT: Zinc-catalyzed synthesis of acylsilanes using carboxylic acids and a silylborane has been achieved in the presence of pivalic anhydride. Various carboxylic acids were converted to the corresponding acylsilanes. The in situ formation of mixed anhydrides was essential in the present reaction.



cylsilanes are carbonyl compounds in which silicone and oxygen atoms are directly bound to the same carbon atom.<sup>1</sup> As a result of this unique feature, acylsilanes are very versatile and can be used in many organic transformations, such as reactions with various nucleophiles via Brook rearrangement,<sup>2</sup> photochemical reactions,<sup>3</sup> and transitionmetal-catalyzed reactions.4

Classically, acylsilanes are synthesized by silvlation of lithiated 1,3-dithianes followed by a deprotection step.<sup>5</sup> Alternative preparation methods include the oxidation of organosilicon compounds.<sup>6,7</sup> Direct synthesis of acylsilanes via silylation of carbonyl compounds using silyl metals such as lithium,<sup>8</sup> magnesium,<sup>9</sup> or aluminum<sup>10</sup> species was also reported. However, functional group tolerance was quite low due to their high reactivity. Although silvl copper species <sup>1</sup> a provided acylsilanes with various functional groups,<sup>1</sup> stoichiometric amount of copper wastes was generated after the reaction.

In contrast to classical synthetic methods, transition-metalcatalyzed preparation of acylsilanes using mild silylation reagents is highly reliable. Yamamoto reported the palladium-catalyzed synthesis of acylsilanes from acid chlorides and disilanes (Scheme 1a).<sup>12</sup> Later, Riant reported that acylsilanes could be synthesized by the reaction of acid anhydrides with silvlboranes in the presence of Cu catalysts (Scheme 1b).<sup>13</sup> These methods result in various acylsilanes; however, the reaction starting materials, such as acid chlorides or acid anhydrides, must be prepared from the corresponding carboxylic acids before use. Herein, the Zn-catalyzed synthesis of acylsilanes using carboxylic acids and a silylborane is reported, where pivalic anhydride was found to be an indispensable additive (Scheme 1c). The key to success of this reaction is the in situ formation of a mixed anhydride.<sup>14</sup>

First, sodium benzoate (1a-Na) was reacted with a silylborane, PhMe<sub>2</sub>Si-B(pin), using pivalic anhydride as an additive in the presence of a zinc catalyst in 1:1 (v/v) ClCH<sub>2</sub>CH<sub>2</sub>Cl/toluene at 90 °C for 18 h (Table 1).<sup>15</sup> Without

## Scheme 1. Transition-Metal-Catalyzed Synthesis of Acylsilanes

(a) Pd-catalyzed reaction using acid chlorides

$$R \rightarrow OR + Me_2PhSi \rightarrow B(pin)$$
 Cu cat.

(c) Zn-catalyzed reaction using carboxylic acids (This Work)

a zinc catalyst, the reaction did not proceed at all (entry 1). Employing  $ZnCl_2$  as the catalyst, an acylsilane (2a) was obtained in 76% GC yield (entry 2).  $Zn(OAc)_2$  was found to be less effective as the catalyst (entry 3), producing the product 2a in 66% yield, while  $Zn(O_2CtBu)_2$  showed high catalytic activity (entry 4). A commercially available ZnEt<sub>2</sub> was also effective for the reaction, giving **2a** in 86% yield (entry 5). Lithium benzoate (1a-Li) and potassium benzoate (1a-K)were not effective substrates for the reaction (entries 6 and 7). In the case of benzoic acid (1a) as the substrate, the desired product was not detected at all (entry 8). On the other hand, in situ deprotonation using NaH dramatically improved the yield of 2a (entry 9). Under the reaction conditions, pure 2a was isolated in 82% yield. The reaction using 2.0 mol of 1a (0.24 g) afforded 3a in 64% yield (0.31 g). Next, other catalysts were tested and  $Pd(OAc)_2$ , Cu(OAc), and  $AlCl_3$  were found to be ineffective (entries 10-12). When other silvl sources such as Et<sub>3</sub>Si–B(pin) or Me<sub>3</sub>Si–SiMe<sub>3</sub> were tested in

Received: November 20, 2019

# Table 1. Optimization of Reaction Conditions for Zn-Catalyzed Synthesis of an Acylsilane $2a^{a}$

1a-M (M = or 1a (M	OM + Me <sub>2</sub> PhSi—B(pin) = Na, K, Li) = H)	Catalyst (5.0 mol %) (tBuCO) <sub>2</sub> O (1.5 equiv) CICH <sub>2</sub> CH <sub>2</sub> Cl/toluene (1:1, 2.0 mL) 90 °C, 18 h	SiPhMe <sub>2</sub>
entry	catalysts	М	yield of $2a (\%)^b$
1	none	Na	0
2	ZnCl2	Na	76
3	$Zn(OAc)_2$	Na	66
4	$Zn(O_2CtBu)_2$	Na	86
5	ZnEt2 (1.0 M in hex	ane) Na	86
6	ZnEt <sub>2</sub>	Li	56
7	ZnEt <sub>2</sub>	K	13
8	ZnEt <sub>2</sub>	H (1a)	0
9 <sup>c</sup>	ZnEt <sub>2</sub>	H (1a)	86 $(82)^d$
10 <sup>c</sup>	$Pd(OAc)_2$	H (1a)	0
11 <sup>c</sup>	Cu(OAc)	H (1a)	0
12 <sup>c</sup>	AlCl <sub>3</sub>	H (1a)	0

<sup>*a*</sup>Reaction conditions: **1a** (0.40 mmol), Me<sub>2</sub>PhSi–B(pin) (0.52 mmol), catalyst (0.020 mmol, 5.0 mol %), (tBuCO)<sub>2</sub>O (0.60 mmol) in ClCH<sub>2</sub>CH<sub>2</sub>Cl/toluene (1:1 (v/v), 2.0 mL) at 90 °C for 18 h. <sup>*b*</sup>GC yield by an internal standard method. <sup>*c*</sup>NaH (1.1 equiv) was added. <sup>*d*</sup>Isolated yield of **2a**.

place of  $Me_2PhSi-B(pin)$ , the desired products were not obtained at all.<sup>15</sup>

Next, the substrate scope was examined using various carboxylic acids in the presence of NaH (Table 2). Aromatic carboxylic acids bearing electron-donating (1b and 1c) and electron-withdrawing groups (1d-g) on the phenyl ring afforded the corresponding acylsilanes (2b-g) in moderateto-good yields (entries 1-6). In the reaction, bromo and iodo groups were intact (entries 4 and 5). A boronic ester moiety was also tolerated (entry 7). A sterically hindered substrate (1i) was smoothly converted to the product (2i) in 80% yield (entry 8). 1-Naphthoic acid (1j) also gave the product (2j) in good yield (entry 9). Aliphatic carboxylic acids (1k-n) also afforded the corresponding products (2k-n) (entries 10–13). The ester functional group in 1m was tolerated (entry 12). Cyclohexanecarboxylic acid (1n) also reacted with silylborane, giving 2n in moderate yield (entry 13). On the other hand, bulky pivalic acid 10 did not afford the corresponding product at all (entry 14). Other carboxylic acids having a chiral center at the  $\alpha$ -position such as *N*,*N*-dibutylalanine were not suitable substrates.

To obtain insights into the reaction mechanism, some control experiments were carried out (Scheme 2). When sodium benzoate (1a-Na) was reacted with PhMe<sub>2</sub>Si-B(pin) in the absence of pivalic anhydride, the product 2a was not obtained at all (Scheme 2a). Thus, carboxylate was not directly silylated, and the formation of a mixed anhydride by the reaction of 1a-Na and pivalic anhydride (in equilibrium) is indispensable. Next, benzoic anhydride (3) as an alternative substrate for the mixed anhydride reacted with PhMe<sub>2</sub>Si-B(pin). Although 2a was obtained, the yield was low (Scheme 2b). In order to generate the mixed anhydride, sodium pivalate was added to the reaction with 3 (Scheme 2c). As a result, the yield of 2a was increased to 52% yield, indicating in situ generated mixed anhydride is a key intermediate for the present reaction.

## Table 2. Scope of Carboxylic Acids for the Synthesis of Acylsilanes $^{a}$



<sup>*a*</sup>Reaction conditions: 1 (0.40 mmol), Me<sub>2</sub>PhSi–B(pin) (0.52 mmol), ZnEt<sub>2</sub> (1.0 M in hexane, 0.020 mmol, 5.0 mol %), (tBuCO)<sub>2</sub>O (0.60 mmol), NaH (0.44 mmol) in ClCH<sub>2</sub>CH<sub>2</sub>Cl/toluene = 1:1 (v/v, 2.0 mL) at 90 °C for 18 h. <sup>*b*</sup>Isolated yield of 2. <sup>*c*</sup>ZnEt<sub>2</sub> (10 mol %) was used.

A plausible catalytic cycle for the reaction discovered in this work is presented in Scheme 3. First,  $ZnEt_2$  reacts with pivalic anhydride, giving zinc carboxylate **A**. Next, transmetalation with silylborane affords silylzinc intermediate **B**, which was recently reported by Uchiyama.<sup>16</sup> Then, **B** reacts with in situ generated mixed anhydride **C** to afford an intermediate **D**. In

Scheme 2. Control Experiments To Elucidate the Reaction Mechanism



#### Scheme 3. Plausible Catalytic Cycle



this step, **B** selectively reacts with a less hindered carbonyl group in **C** due to the bulky *t*Bu group. Such steric difference is important for site selectivity.<sup>14</sup> Finally, an elimination step results in the formation of product 2, and the zinc carboxylate species regenerates. Considering what is outlined in Scheme 2c, an in situ generated sodium carboxylate may support the transmetalation step (step a).

In conclusion, a zinc compound was found to catalyze the reaction of carboxylic acids with a silylborane in the presence of pivalic anhydride. The reactions yielded the corresponding acylsilanes in moderate-to-high yields. Further studies on the reaction mechanism and the application of the principles of this study to other silylation reactions are currently underway.

## ASSOCIATED CONTENT

#### **Supporting Information**

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acs.orglett.9b04151.

Experimental procedures and characterization of the products (PDF)

## AUTHOR INFORMATION

#### **Corresponding Author**

\*E-mail: tfuji@scl.kyoto-u.ac.jp. ORCID <sup>©</sup>

Yasushi Tsuji: 0000-0001-6733-0539 Tetsuaki Fujihara: 0000-0002-6687-2528

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

This work was supported by JSPS KAKENHI Grant Number 18H04257 in Precisely Designed Catalysts with Customized Scaffolding (T.F.) and 17H03096 in Grant-in-Aid for Scientific Research B (Y.T.) from MEXT, Japan. T.F. also acknowledged the Tokuyama Science Foundation for financial support.

## REFERENCES

 For a review, see Zhang, H.-J.; Priebbenow, D. L.; Bolm, C. Chem. Soc. Rev. 2013, 42, 8540-8571 and references cited therein..
 For selected examples, see (a) Reich, H. J.; Rusek, J. J.; Olson, R. E. J. Am. Chem. Soc. 1979, 101, 2225-2227. (b) Kuwajima, I.; Kato, M. J. Chem. Soc., Chem. Commun. 1979, 708-709. (c) Takeda, K.; Fujisawa, M.; Makino, T.; Yoshii, E.; Yamaguchi, K. J. Am. Chem. Soc. 1993, 115, 9351-9352. (d) Takeda, K.; Nakajima, A.; Takeda, M.; Okamoto, Y.; Sato, T.; Yoshii, E.; Koizumi, T.; Shiro, M. J. Am. Chem. Soc. 1998, 120, 4947-4959. (e) Lettan, R. B., II; Galliford, C. V.; Woodward, C. C.; Scheidt, K. A. J. Am. Chem. Soc. 2009, 131, 8805-8814. (f) Unger, R.; Weisser, F.; Chinkov, N.; Stanger, A.; Cohen, T.; Marek, I. Org. Lett. 2009, 11, 1853-1856. (g) Sasaki, M.; Kondo, Y.; Kawahata, M.; Yamaguchi, K.; Takeda, K. Angew. Chem., Int. Ed. 2011, 50, 6375-6378. (h) Priebbenow, D. L. J. Org. Chem. 2019, 84, 11813-11822.

(3) (a) Ito, K.; Tamashima, H.; Iwasawa, N.; Kusama, H. J. Am. Chem. Soc. 2011, 133, 3716–3719. (b) Zhang, H.-J.; Becker, P.; Huang, H.; Pirwerdjan, R.; Pan, F.-F.; Bolm, C. Adv. Synth. Catal. 2012, 354, 2157–2161. (c) Ishida, K.; Tobita, F.; Kusama, H. Chem. - Eur. J. 2018, 24, 543–546.

(4) (a) Obora, Y.; Ogawa, Y.; Imai, Y.; Kawamura, T.; Tsuji, Y. J. Am. Chem. Soc. 2001, 123, 10489–10493. (b) Schmink, J. R.; Krska, S. W. J. Am. Chem. Soc. 2011, 133, 19574–19577.

(5) (a) Brook, A. G.; Duff, J. M.; Jones, P. F.; Davis, N. R. J. Am. Chem. Soc. **1967**, 89, 431–434. (b) Corey, E. J.; Seebach, D.; Freedman, R. J. Am. Chem. Soc. **1967**, 89, 434–436.

(6) (a) Reich, H. J.; Eisenhart, E. K. J. Org. Chem. **1984**, 49, 5282– 5283. (b) Linderman, R. J.; Suhr, Y. J. Org. Chem. **1988**, 53, 1569– 1572. (c) Lipshutz, B. H.; Lindsley, C.; Susfalk, R.; Gross, T. Tetrahedron Lett. **1994**, 35, 8999–9002. (d) Reddy, G. P.; Reddy, J. S.; Das, S.; Roisnel, T.; Yadav, J. S.; Chandrasekhar, S.; Grée, R. Org. Lett. **2013**, 15, 1524–1527.

(7) (a) Hassner, A.; Soderquist, J. A. J. Organomet. Chem. 1977, 131, C1–C4. (b) Brook, A. G. J. Am. Chem. Soc. 1957, 79, 4373–4275.
(c) Corriu, R. J. P.; Masse, J. P. J. Organomet. Chem. 1970, 22, 321–332. (d) Degl'Innocenti, A.; Walton, D. R. M; Seconi, G.; Pirazzini, G.; Ricci, A. Tetrahedron Lett. 1980, 21, 3927–3928. (e) Cossrow, J.; Rychnovsky, S. D. Org. Lett. 2002, 4, 147–150. (f) Inoue, A.; Kondo, J.; Shinokubo, H.; Oshima, K. J. Am. Chem. Soc. 2001, 123, 11109–11110. (g) Kondo, J.; Shinokubo, H.; Oshima, K. Org. Lett. 2006, 8, 1185–1187.

(8) (a) Fleming, I.; Ghosh, U. J. Chem. Soc., Perkin Trans. 1 1994, 1, 257–262.
(b) Clark, C. T.; Milgram, B. C.; Scheidt, K. A. Org. Lett. 2004, 6, 3977–3980.

(9) Picard, J. P.; Calas, R.; Dunogues, J.; Duffaut, N.; Gerval, J.; Lapouyade, P. J. Org. Chem. **1979**, 44, 420–424.

(10) (a) Kang, J.; Lee, J. H.; Kim, K. S.; Jeong, J. U.; Pyun, C. *Tetrahedron Lett.* **1987**, *28*, 3261–3262. (b) Nakada, M.; Nakamura, S.-I.; Kobayashi, S.; Ohno, M. *Tetrahedron Lett.* **1991**, *32*, 4929–4932. (11) (a) Bonini, B. F.; Comes-Franchini, M.; Mazzanti, G.; Ricci, A.; Sala, M. J. Org. Chem. **1996**, *61*, 7242–7243. (b) Bonini, B. F.; Comes-Franchini, M.; Laboroi, F.; Mazzanti, G.; Ricci, A.; Varchi, G. J. Org. Chem. **1999**, *64*, 8008–8013.

(12) (a) Yamamoto, K.; Suzuki, S.; Tsuji, J. *Tetrahedron Lett.* **1980**, 21, 1653–1656. (b) Yamamoto, K.; Hayashi, A.; Suzuki, S.; Tsuji, J. *Organometallics* **1987**, 6, 974–979. (c) Geng, F.; Maleczka, R. E. *Tetrahedron Lett.* **1999**, 40, 3113–3114.

(13) Cirriez, V.; Rasson, C.; Riant, O. Adv. Synth. Catal. 2013, 355, 3137–3140.

(14) For the reactions using pivalic anhydride, see (a) Nagayama, K.; Shimizu, I.; Yamamoto, A. Chem. Lett. 1998, 27, 1143-1144.
(b) Nagayama, K.; Shimizu, I.; Yamamoto, A. Bull. Chem. Soc. Jpn. 2001, 74, 1803-1815. (c) Gooßen, L. J.; Ghosh, K. Angew. Chem., Int. Ed. 2001, 40, 3458-3460. (d) Gooßen, L. J.; Ghosh, K. Chem. Commun. 2002, 836-837. (e) Fujihara, T.; Cong, C.; Terao, J.; Tsuji, Y. Adv. Synth. Catal. 2013, 355, 3420-3424. (f) Pan, F.; Lei, Z.-Q.; Wang, H.; Li, H.; Sun, J.; Shi, Z. - J. Angew. Chem., Int. Ed. 2013, 52, 2063-2067. (g) Maetani, S.; Fukuyama, T.; Ryu, I. Org. Lett. 2013, 15, 2754-3219.

(15) See Supporting Information for details.

(16) Nagashima, Y.; Yukimori, D.; Wang, C.; Uchiyama, M. Angew. Chem., Int. Ed. 2018, 57, 8053–8057.