


Inhibition of 1-Deoxy-D-Xylulose-5-Phosphate Reductoisomerase by Lipophilic Phosphonates: SAR, QSAR, and Crystallographic Studies

Lisheng Deng,^{†,§} Jiasheng Diao,^{†,§} Pinhong Chen,[†] Venugopal Pujari,[¶] Yuan Yao,[†] Gang Cheng,[†] Dean C. Crick,[¶] B. V. Venkataram Prasad,[‡] and Yongcheng Song^{*,†}[†]Department of Pharmacology and [‡]Verna and Marrs McLean Department of Biochemistry and Molecular Biology, Baylor College of Medicine, 1 Baylor Plaza, Houston, Texas 77030, United States[¶]Department of Microbiology, Colorado State University, 1682 Campus Delivery, Ft. Collins, Colorado 80523, United States

S Supporting Information

ABSTRACT: 1-Deoxy-D-xylulose-5-phosphate reductoisomerase (DXR) is a novel target for developing new antibacterial (including antituberculosis) and antimalaria drugs. Forty-one lipophilic phosphonates, representing a new class of DXR inhibitors, were synthesized, among which 5-phenylpyridin-2-ylmethylphosphonic acid possesses the most activity against *E. coli* DXR (*EcDXR*) with a K_i of 420 nM. Structure–activity relationships (SAR) are discussed, which can be rationalized using our *EcDXR*:inhibitor structures, and a predictive quantitative SAR (QSAR) model is also developed. Since inhibition studies of DXR from *Mycobacterium tuberculosis* (*MtDXR*) have not been performed well, 48 *EcDXR* inhibitors with a broad chemical diversity were found, however, to generally exhibit considerably reduced activity against *MtDXR*. The crystal structure of a *MtDXR*:inhibitor complex reveals the flexible loop containing the residues 198–208 has no strong interactions with the 3,4-dichlorophenyl group of the inhibitor, representing a structural basis for the reduced activity. Overall, these results provide implications in the future design and development of potent DXR inhibitors.



INTRODUCTION

Isoprenoid biosynthesis, producing very important substances such as isoprenyl (e.g., farnesyl and undecaprenyl) diphosphates, vitamins, steroids, and terpenoids, is essential to the growth of all organisms. Isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) are two common precursors for isoprenoid biosynthesis. In humans and animals, the mevalonate pathway is used to synthesize IPP and DMAPP (Figure 1). On the other hand, the vast majority of bacteria and apicomplexan parasites such as *Plasmodium falciparum*, the causative agent of malaria, use the non-mevalonate (or methylerythritol phosphate, MEP) isoprene biosynthesis pathway discovered in the 1990s (Figure 1).¹ Since humans do not have the 7 enzymes in the pathway, the non-mevalonate pathway, especially the second enzyme 1-deoxy-D-xylulose-5-phosphate reductoisomerase (DXR), has attracted much interest for its potential to be a new anti-infective drug target.²

DXR catalyzes the isomerization and reduction of 1-deoxy-D-xylulose-5-phosphate (DXP) to 2-C-methyl-D-erythritol-4-phosphate (MEP) with a Mg^{2+} and NADPH being the enzyme cofactors (Figure 1). Fosmidomycin (**1**, Chart 1) (as well as its analogue FR900098 (**2**)) is a very potent DXR inhibitor³ and has strong antibacterial activity against many Gram-negative bacteria such as *Escherichia coli* and *Pseudomonas aeruginosa*.⁴ Particularly remarkable is its profound antimalaria activity in preclinical

studies and recent clinical trials.⁵ However, since **1** is a highly polar molecule with great water solubility, it has a short half-life (~ 1 h) in plasma as well as poor oral availability. In addition, many pathogens, such as *Mycobacterium tuberculosis*, the causative agent of tuberculosis that kills ~ 1.6 million people each year, are completely resistant to **1** and sensitive bacteria can also easily develop drug resistance. This is because **1** is actively transported into cells via glycerol-3-phosphate transporter GlpT.⁶ Pathogens without GlpT or with a mutant GlpT are therefore not affected by **1** due to limited cellular uptake.^{2d,7} Great interest has therefore been generated to develop potent inhibitors of DXR.^{8–10}

A potential solution is to develop lipophilic DXR inhibitors that are able to passively permeate into bacterial cells. Medicinal chemistry studies based on the structures of **1** and **2** have been carried out for the past decade.⁸ Although most modifications resulted in compounds with largely reduced activity, derivatives of **1** and **2** having an α -substituent still possess the majority of the inhibitory activity.^{8j} The best in this series are compounds **3** and **4** with a 3,4-dichlorophenyl group, which have similar enzyme activity as **1** but showed greatly increased activity against *P. falciparum* (EC_{50} of 90 nM for **3** and 250 nM for **4** vs 1100 nM for **1**),^{8j} presumably due to enhanced cellular uptake resulting

Received: March 28, 2011

Published: May 11, 2011





Figure 1. Non-mevalonate and mevalonate isoprene biosynthesis pathways.

Chart 1. Structures of Known DXR Inhibitors 1–10



from their lipophilicity. In addition, only a few DXR inhibitors have been identified that are structurally distinct from **1** and **2**.⁹ Two bisphosphonates (**5** and **6**, Chart 1) bearing a hydrophobic pyridine and isoquinoline side chain are relatively weak inhibitors (IC_{50} : 4 and 7 μM).^{9a} We recently used a coordination chemistry based design to find a lipophilic, drug-like DXR inhibitor **7** (IC_{50} : 1.4 μM) without a phosphonate group, which exhibited good activity against a broad spectrum of pathogenic bacteria.^{9b} Moreover, in order to probe the hydrophobic nature of the DXR active site, we identified three new lipophilic phosphonate inhibitors **8–10** having a pyridine or quinoline group with IC_{50} values as low as 0.84 μM .^{9c}

Our X-ray crystallographic studies of *E. coli* DXR (*EcDXR*) in complex with **8–10** reveal a new inhibitor binding mode as well as an important role of Trp211 in inhibitor recognition.^{9c} The great plasticity of a flexible loop (residues 205–215) of *EcDXR* allows the indole ring of Trp211 to form π – π stacking and charge transfer interactions with the electron-deficient pyridine/

quinoline group of these inhibitors. Previously reported crystal structures of *EcDXR* in complex with **1**,¹¹ **5**, and **6**^{9a} also show favorable hydrophobic interactions between Trp211 and the inhibitors. In addition, our structures show that the conformational change of the flexible loop opens up a mainly hydrophobic pocket, where the pyridine/quinoline ring of **8–10** is located. These structures have been used to model the binding modes of other lipophilic DXR inhibitors, such as compounds **3** and **4**.

Here, we report synthesis, inhibitory activity, structure–activity relationships (SAR) and quantitative SAR (QSAR) of 41 lipophilic phosphonate and related compounds against *E. coli* DXR. In addition, since *M. tuberculosis* is a Gram-positive bacterium (which behaves very differently from Gram-negative bacteria, e.g., *E. coli*) and causes enormous human mortalities, we also report the activity of 48 *EcDXR* inhibitors with a broad range of chemical diversity against *M. tuberculosis* DXR (*MtDXR*). The results show that these inhibitors generally have considerably lowered affinity to *MtDXR*. In addition, to investigate the

possible structural basis for the reduced activity, a crystal structure of *Mt*DXR:inhibitor complex is disclosed, showing the flexible loop of *Mt*DXR, especially the residue Trp203 (the aligned counterpart of *Ec*DXR Trp211) does not have strong interactions with the hydrophobic group of the inhibitor.

RESULTS AND DISCUSSION

SARs of Lipophilic Phosphonates in *Ec*DXR Inhibition. In our previous study,^{9c} pyridine/quinoline containing phosphonate compounds **8–10** were identified to be a new class of *Ec*DXR inhibitors. On the basis of the structure of **8**, we synthesized a total of 41 compounds to analyze the structure–activity relationships (SAR) of this class of compounds. These compounds were tested to inhibit recombinant *Ec*DXR, and their inhibition constants (K_i) are shown in Table 1.

First, compounds **11–13** (Table 1) were synthesized to find the optimal linker length between the pyridine ring and the phosphonate group. Compound **11** without a linker has essentially no activity on *Ec*DXR at 100 μ M, while **12** and **13** having two and three methylenes are active with K_i values of 18.6 and 9.5 μ M, respectively. However, compared to **8** with a K_i of 2.3 μ M, they are $\sim 8\times$ and $4\times$ less potent, suggesting that one methylene linker is best for DXR inhibition.

Second, we optimized the aromatic ring of **8** in order to find more potent DXR inhibitors. Compounds **14** and **15** with a pyridin-3-yl and pyridin-4-yl group were found to be active ($K_i = 5.0$ and 3.5 μ M). In addition, other N-containing aromatic heterocyclic compounds **10** (with quinoline), **16** (with pyrazine), and **17** (with pyrimidine) also exhibit inhibitory activity against *Ec*DXR, with K_i values ranging from 3.5 to 11.2 μ M. However, phenyl ring containing compounds **18–20** with a variety of linker length ($n = 0–2$) are inactive. Since the 3,4-dichlorophenyl group is present in potent *Ec*DXR inhibitors **3** and **4** (Chart 1) with antimalaria activity,^{8j} compound **21** having a 3,4-dichlorophenyl group was prepared, but it was found to have no activity. Compounds **22** and **23** containing electron-donating hydroxyl groups also possess no inhibitory activity. To further probe the electronic effects of the aromatic ring, compounds **24** and **25** were synthesized, which have hydroxyl (or tautomeric carbonyl) substituted pyridine rings that are more electron-rich. Compared to unsubstituted parent compounds **14** and **12**, these two compounds are essentially inactive. These results indicate that a highly electron-deficient aromatic ring, such as pyridine, quinoline, and pyrimidine, is needed for DXR inhibition.


Third, compounds **26–38** were synthesized to see which substituents on the pyridine ring are favored. Compared to the parent compound **8**, compound **26** with a 4-bromo substituent has a slightly weakened inhibitory activity with a K_i of 3.5 μ M. **27** with a 4-phenyl group exhibits better activity ($K_i = 1.1$ μ M), while compound **28** bearing a 4-(4-fluorophenyl) group possesses a reduced activity ($K_i = 3.5$ μ M). For compounds **9** and **29–31** that have a 5-substituent, compound **9** with a 5-phenyl group shows a K_i of 0.42 μ M against *Ec*DXR, which is $5\times$ as potent as **8**, while compound **29** with a 5-bromo group exhibits a weaker inhibition ($K_i = 6.7$ μ M). Compound **30** with a 5-hydroxyl group has strong activity with a K_i of 0.7 μ M, being $\sim 3\times$ as potent as **8**. This is rather unexpected given that 5-hydroxypyridine is more electron-rich and, according to the SAR found above, should lead to a reduced activity. However, the high activity of **30** could be rationalized by molecular modeling using the crystal structure of the *Ec*DXR:**8** complex, as discussed below. Compound **31** is

inactive likely due to its electron-donating 5-benzyloxy group, which is also sterically more demanding. Compounds **32** and **33** bearing a 6-bromo and -phenyl group, respectively, show slightly weaker inhibitory activity ($K_i = 3.6$ and 3.5 μ M) against *Ec*DXR than does **8**. Since compound **9** possesses a greatly improved potency, five derivatives **34–38** were made in an effort to find more active inhibitors. However, neither electron-withdrawing -F/-Cl (for **34** and **35**, $K_i = 1.1$ and 1.6 μ M) nor electron-donating -Me/-OMe (for **36** and **37**, $K_i = 3.9$ and 3.6 μ M) on the 5-phenyl ring provide enhanced potency. **38** with a bulkier naphthyl substituent is also a weak inhibitor with a K_i of 13.9 μ M. These results show that introducing a 5-phenyl group leads to the strongest *Ec*DXR inhibitor **9** among this class of compounds.

Finally, the phosphonate group in compounds **8** and **9** was replaced by an isosteric carboxyl, hydroxamate, or sulfonate group with a rationale that the replacement with a less polar, negatively charged group could result in a higher lipophilicity, thereby improving cellular uptake. However, compounds **39–44** that are analogues of **8** with either a carboxyl or a hydroxamate group exhibit no inhibitory activity against *Ec*DXR at 100 μ M. Similarly, compounds **45–48**, which are analogues of **9** containing a carboxyl, hydroxamate, or sulfonate group, are also inactive. These results show that the phosphonate group is essential to the tight binding of this class of inhibitors.

SAR Rationalization. With the availability of the crystal structures of *Ec*DXR:**8–10** complexes, it is possible to rationalize the structure–activity relationships observed above. Our crystallographic studies show that there are two common interactions between *Ec*DXR and these inhibitors.^{9c} The first one is the π – π stacking and charge transfer interactions between the electron-deficient pyridine/quinoline group of **8–10** and the indole ring of the residue Trp211, with the two parallel rings having a distance of ~ 3.5 Å. To see if compounds **11–13** are potentially able to form such interactions, we docked these compounds into the crystal structure of DXR:**8** complex, using the program Glide¹² in Schrödinger,¹³ and the results are shown in Figure 2a–c. The phosphonate group of **11** is predicted to be located very close to the phosphonate binding site of **8**. However, its pyridine ring cannot form a π – π interaction with the indole ring of Trp211, which might be responsible for its non-activity. The docking results of compounds **12** and **13** indicate their pyridine rings can be parallel to the indole, thanks to their longer, flexible linkers. The modeling studies are in line with the experimental data showing **12** and **13** are active against *Ec*DXR (Table 1). In addition, the importance of Trp211 in inhibitor recognition also determines that an electron-deficient aromatic ring is needed for *Ec*DXR inhibition, because of its stronger electrostatic interaction with the electron-rich indole ring. Moreover, the crystal structure of DXR:**9** complex reveals that the conformation of Trp211 is different from that of DXR:**8**, with the indole ring flipping $\sim 180^\circ$ in order to have more hydrophobic interactions with the 5-phenyl group of **9**.^{9c} This might account for the improved activity of compound **9**.

The second key interaction is the tight binding of the phosphonate group in **8–10**, with all of the three negatively charged O atoms forming a network of electrostatic and H-bond interactions with the residues Lys227, Asn226, Ser221, and Ser185. However, a carboxylate, hydroxamate, or sulfonate group possesses considerably reduced polarity as well as negative charges at the physiological pH. Lack of activity of compounds **39–48** against *Ec*DXR can therefore be rationalized.

Table 1. Structures of Compounds 8–48 and Their K_i Values against *EcDXR* and *MtDXR*

Compound ID	R-	n	-X	<i>EcDXR</i> K_i (μM)	<i>MtDXR</i> K_i (μM)
11	Pyridin-2-yl	0	-P(O)(OH) ₂	NI ^a	NI
8	Pyridin-2-yl	1	-P(O)(OH) ₂	2.3	1.6
12	Pyridin-2-yl	2	-P(O)(OH) ₂	18.6	NI
13	Pyridin-2-yl	3	-P(O)(OH) ₂	9.5	NI
14	Pyridin-3-yl	1	-P(O)(OH) ₂	5.0	NI
15	Pyridin-4-yl	1	-P(O)(OH) ₂	3.5	6.8
10	Quinolin-2-yl	1	-P(O)(OH) ₂	7.9	6.9
16		1	-P(O)(OH) ₂	11.2	NI
17		1	-P(O)(OH) ₂	3.5	12.4
18	Phenyl-	0	-P(O)(OH) ₂	NI	NI
19	Phenyl-	1	-P(O)(OH) ₂	NI	NI
20	Phenyl-	2	-P(O)(OH) ₂	NI	NI
21	3,4-Dichlorophenyl-	1	-P(O)(OH) ₂	NI	NI
22	2-Hydroxyphenyl-	2	-P(O)(OH) ₂	NI	NI
23	2,3-Dihydroxyphenyl-	1	-P(O)(OH) ₂	NI	NI
24		1	-P(O)(OH) ₂	NI	NI
25		2	-P(O)(OH) ₂	NI	NI
26	4-Bromopyridin-2-yl	1	-P(O)(OH) ₂	3.5	32.7
27	4-Phenylpyridin-2-yl	1	-P(O)(OH) ₂	1.1	6.9
28	4-(4-Fluorophenyl)pyridin-2-yl	1	-P(O)(OH) ₂	3.5	NI
29	5-Bromopyridin-2-yl	1	-P(O)(OH) ₂	6.7	45.2
9	5-Phenylpyridin-2-yl	1	-P(O)(OH) ₂	0.42	3.2
30	5-Hydroxypyridin-2-yl	1	-P(O)(OH) ₂	0.7	17.3
31	5-Benzyloxy pyridin-2-yl	1	-P(O)(OH) ₂	NI	NI
32	6-Bromopyridin-2-yl	1	-P(O)(OH) ₂	3.6	NI
33	6-Phenylpyridin-2-yl	1	-P(O)(OH) ₂	3.5	NI
34	5-(4-Fluorophenyl)pyridin-2-yl	1	-P(O)(OH) ₂	1.1	50.7
35	5-(4-Chlorophenyl)pyridin-2-yl	1	-P(O)(OH) ₂	1.6	NI
36	5-(4-Methylphenyl)pyridin-2-yl	1	-P(O)(OH) ₂	3.9	NI
37	5-(3-Methoxyphenyl)pyridin-2-yl	1	-P(O)(OH) ₂	3.6	NI
38	5-(2-Naphthyl)pyridin-2-yl	1	-P(O)(OH) ₂	13.9	NI
39	Pyridin-2-yl	0	-COOH	NI	NI
40	Pyridin-2-yl	1	-COOH	NI	NI
41	Pyridin-3-yl	1	-COOH	NI	NI
42	Pyridin-4-yl	1	-COOH	NI	NI
43	Pyridin-2-yl	0	-CONHOH	NI	NI
44	Pyridin-2-yl	1	-CONHOH	NI	NI
45	5-Phenylpyridin-2-yl	0	-COOH	NI	NI
46	5-Phenylpyridin-2-yl	1	-COOH	NI	NI
47	5-Phenylpyridin-2-yl	1	-CONHOH	NI	NI
48	5-Phenylpyridin-2-yl	1	-SO ₃ H	NI	NI

^a No inhibition observed at 100 μM .




Figure 2. Lowest energy docking structures of compounds (A) 11, (B) 12, (C) 13, and (D) 30 in the *EcDXR*:8 structure, together with superimposed structure of 8 (in yellow).

Table 2. K_i Values of 1–7 against *EcDXR* and *MtDXR*

compound ID	<i>EcDXR</i> K_i (μM)	<i>MtDXR</i> K_i (μM)
1	0.027	0.14
2	0.019	0.13
3	0.058	1.2
4	0.036	1.5
5	6.3	32.7
6	15.3	NI ^a
7	0.70	21.8

^aNo inhibition observed at 100 μM .

Moreover, molecular modeling could be used to explain why compound 30 with the electron-donating 5-hydroxy group possesses unexpected, strong inhibition ($K_i = 0.7 \mu\text{M}$). As shown in Figure 2d, the docking result of 30 in the DXR:8 structure indicates that, in addition to the phosphonate binding and the π - π stacking with Trp211, the 5-hydroxy group of 30 forms a hydrogen bond with the carbonyl of Asn210. This extra H-bond could compensate the weakened π - π /charge transfer interactions between 30 and the protein.

QSAR Study in *EcDXR* Inhibition. The above qualitative SAR analysis and rationalization demonstrate that a complex combination of electronic, steric, hydrophobic, and H-bonding features of the substituents affect the activity of these compounds. In order to understand these observations in a quantitative and predictive manner, we next carried out a 3-D quantitative structure activity relationship (QSAR) study, using the program Phase¹⁴ in Schrödinger. Since compounds 1–4 (Chart 1) are more potent *EcDXR* inhibitors (Table 2, K_i : 19–58 nM) than 8–10 and known to occupy the same binding site, they were also included in an effort to generate a broader and more predictive QSAR model. In addition, for those inactive compounds in Table 1 against *EcDXR*, their K_i values were arbitrarily assigned to be 300 μM for QSAR calculation. The 45 compounds (1–4

and 8–48) were constructed, minimized, and aligned with compound 8 as a template, using the “flexible ligand alignment” module in Maestro¹⁵ in Schrödinger, which recognizes common features within the molecules (e.g., similar partial charge, hydrophobicity, aromaticity, and H-bond donor/acceptor). The aligned compounds were imported into Phase, and a partial least-squares (PLS) method was used to correlate the *EcDXR* activities of these compounds with the Phase field data calculated from their 3-D structures. The QSAR model yielded $r^2 = 0.95$, q^2 (no. of factors) = 0.78 (5), F -test = 158.5, and a $\text{p}K_i$ error of 0.30, as shown in Table 3 and Figure 3a. Five leave-5-out training/test sets were applied to further validate the model, obtaining a good predictivity with $r^2 \geq 0.94$, $q^2 \geq 0.62$, F -test ≥ 123.5 , as shown in detail in Table 3. The compound alignment and the 3-D QSAR virtualization showing negative ionic, electron-withdrawing, hydrophobic, H-bond donor Phase fields are shown in Figure 3b–f. The results are in good agreement with the qualitative SAR observations. For example, the negatively charged phosphonate group (Figure 3c) and the electron-deficient aromatic ring (Figure 3d) are important for the activity. A hydrophobic group (e.g., phenyl) at the 5-position of the pyridine ring of compound 9 is favored (Figure 3e). In addition, the negative charge (due to chelating Mg^{2+}) and the electron-withdrawing feature of the hydroxamate group as well as the hydrophobic nature of the carbon skeleton of compounds 1–4 contribute together to their superior *EcDXR* inhibitory activities (Figure 3c–e).

Inhibition of *MtDXR*. Since the discovery of the non-mevalonate isoprene biosynthesis pathway, DXRs from many species, including bacteria, plants, and parasites, have been cloned and characterized.^{3,5a,7,16} However, inhibition studies of DXRs other than *E. coli* DXR have not been well performed, except that 1 has been tested and shown broad activity on these enzymes. To this end, we are particularly interested in DXR from *M. tuberculosis* (*MtDXR*). This bacterium is a major human pathogen, causing deaths of ~ 1.6 million people a year worldwide. In addition, it has become more and more resistant to first line treatments such as isoniazid and rifampicin during the past decade.¹⁷ Previous studies demonstrated that *M. tuberculosis* exclusively uses the non-mevalonate pathway and DXR is essential for its survival,^{2d,7} suggesting DXR is a target for developing new antituberculosis drugs, but the bacterium is completely resistant to the treatment of 1 due to lack of cellular uptake. Without GlpT (glycerol-3-phosphate transporter), 1 cannot penetrate the *M. tuberculosis* cell membrane and reach the target in cytosol. Nevertheless, 1 was found to be a strong inhibitor of *MtDXR*. However, no other *EcDXR* inhibitors have been tested against *MtDXR*.

We examined the activity of 48 *EcDXR* inhibitors (compounds 1–48) on recombinant DXR from *M. tuberculosis*. It is also noteworthy that these inhibitors represent a wide range of chemical diversity. Compounds 1 and 2 are highly polar molecules containing a hydroxamate and a negatively charged phosphonate group, while 3 and 4 are their derivatives with a hydrophobic 3,4-dichlorophenyl side chain.^{8j} Compounds 5 and 6 are isoquinoline/pyridine containing bisphosphonates,^{9a} which are extremely polar with 4 negative charges at the physiological pH. Distinct from the above compounds, 7 is a lipophilic, drug-like DXR inhibitor without a phosphonate.^{9b} Moreover, all of compounds 8–48 consist of a lipophilic side chain and a polar headgroup (phosphonate, carboxylate, hydroxamate, or sulfonate).

Activities of compounds 1–7 against *MtDXR* as well as *EcDXR* are shown in Table 2. In our study, the reference compound 1

Table 3. QSAR Results for *Ec*DXR Inhibition Using Phase

compound	p <i>K_i</i> (M) ^a	phase p <i>K_i</i> predictions					
		training set	test set 1 ^b	test set 2	test set 3	test set 4	test set 5
1	7.57	7.67	7.58	7.82	7.42	7.78	7.62
2	7.72	7.95	7.84	7.48	7.99	8.07	7.93
3	7.24	7.13	6.94	7.06	7.14	7.15	7.16
4	7.44	7.13	7.10	6.93	7.20	6.97	7.17
8	5.64	5.57	5.51	5.66	5.59	5.62	5.64
9	6.38	5.85	5.81	5.77	5.86	5.91	5.94
10	5.10	4.76	4.75	4.77	4.80	4.65	4.82
11	3.52	3.72	3.65	3.79	3.80	3.62	4.06
12	4.73	4.54	4.57	4.59	4.48	4.54	4.62
13	5.02	5.18	5.22	5.24	5.15	5.18	4.74
14	5.30	4.88	4.88	4.93	4.92	4.83	4.89
15	5.46	5.03	4.97	5.10	5.10	4.93	5.06
16	4.95	5.40	5.34	5.69	5.40	5.46	5.43
17	5.46	6.09	6.07	6.15	6.19	6.05	6.15
18	3.52	3.29	3.50	3.36	3.33	3.27	3.32
19	3.52	4.20	4.12	4.36	4.13	4.18	4.19
20	3.52	3.72	3.74	3.61	3.46	3.62	3.71
21	3.52	3.78	3.72	3.91	3.76	3.68	3.76
22	3.52	3.51	3.58	3.34	3.55	3.47	3.58
23	3.52	3.68	3.59	3.84	3.75	3.64	3.65
24	3.52	3.48	3.45	3.57	4.35	3.46	3.48
25	3.52	3.53	3.55	3.56	3.53	3.50	3.56
26	5.46	5.61	5.66	5.75	5.61	5.64	5.65
27	5.96	5.60	6.09	5.84	5.53	5.57	5.57
28	5.46	5.53	6.06	5.80	5.46	5.49	5.49
29	5.17	5.16	5.10	5.22	5.16	5.18	5.20
30	6.15	5.50	5.47	5.58	5.57	5.52	5.56
31	3.52	3.29	3.22	3.38	3.43	3.29	3.28
32	5.46	5.29	5.25	5.42	5.28	5.27	5.35
33	5.46	5.60	5.56	5.74	5.58	5.61	5.57
34	5.96	5.88	5.86	5.77	5.85	5.90	5.97
35	5.80	5.82	5.82	5.69	5.86	5.76	5.91
36	5.41	5.80	5.79	5.67	5.84	5.78	5.88
37	5.46	5.75	5.71	5.66	5.70	5.88	5.82
38	4.86	4.67	5.09	4.58	4.79	4.66	4.72
39	3.52	3.36	3.29	3.33	3.41	3.30	3.41
40	3.52	3.53	3.52	3.81	3.53	4.05	3.48
41	3.52	3.59	3.59	3.50	3.58	3.63	3.57
42	3.52	3.57	3.59	3.50	3.54	3.57	3.53
43	3.52	3.38	3.33	3.32	3.42	3.36	3.33
44	3.52	3.31	3.33	3.36	3.24	3.50	3.41
45	3.52	3.68	3.70	3.32	3.62	3.83	3.68
46	3.52	3.50	3.48	4.02	3.52	3.80	3.44
47	3.52	3.59	3.58	3.35	3.55	4.19	3.59
48	3.52	3.89	3.80	3.87	3.86	3.95	3.86
	<i>r</i> ²	0.95	0.95	0.94	0.95	0.95	0.96
	<i>q</i> ²	0.78	0.68	0.65	0.62	0.76	0.76
	F-test	158.5	131.5	128.3	123.5	165.8	151.7
	N	5	5	4	5	4	5
	<i>n</i>	45	40	40	40	40	40

^aAn arbitrary p*K_i* value of 3.52 (*K_i* = 300 μM) is given to all inactive compounds. ^bBold values represent predicted activities of compounds that were not included in the training set.

exhibited a *K_i* value of 140 nM against *Mt*DXR, which is very close to that of 150 nM found in a previous study^{2d} (calculated from the IC₅₀ of 310 nM). It is noted that in both assays physiologically relevant Mg²⁺ (4 mM) was used as the cofactor. **1** was also reported to have a lower IC₅₀ value of 80 nM using 2.1 mM of Mn²⁺.¹⁸ However, such a high concentration of Mn²⁺ is not present in any organisms and the *K_m* value for the substrate DXP was not measured in the experiment. This result therefore cannot be used to compare to our data. In addition, we determined the *K_i* value of **1** for *Ec*DXR to be 27 nM, which is also similar to a literature value of 21 nM.¹⁹ Therefore, our study shows that **1** binds to *Mt*DXR ~5× more weakly than *Ec*DXR. Likewise, compound **2**, another extremely potent *Ec*DXR inhibitor,^{8j} was found to possess a *K_i* of 19 nM against *Ec*DXR in our experiment, while the *K_i* of **2** for *Mt*DXR is 130 nM, indicating again the inhibitor has ~6× less affinity to the *M. tuberculosis* enzyme. Even more drastic are the differences between the *K_i* values of compounds **3** and **4** for these two DXR proteins. The two compounds bearing a bulky α-(3,4-dichloro)phenyl group are potent inhibitors of *Ec*DXR with *K_i* values of 58 and 36 nM, approaching the activities of their parent compounds **1** and **2**. However, their inhibition constants against *Mt*DXR are 1.2 and 1.5 μM, being ~10× less active than **1** and **2** (*K_i*^{*Mt*DXR} = 0.14 and 0.13 μM). In addition, compared to *Ec*DXR, the binding of compounds **3** and **4** to *Mt*DXR exhibits enormous 21- and 42-fold affinity losses. These results show that introducing an α-(3,4-dichloro)phenyl group is highly disfavored with respect to targeting *Mt*DXR. Similarly, bisphosphonates (**5** and **6**) and compound **7** (Table 2) were also found to be greater than 5× and 31× less active against the *M. tuberculosis* enzyme, respectively.

The *K_i* values of compounds **8**–**48** against *Mt*DXR are shown in Table 1. Except for compounds **8** and **10**, these compounds possess considerably reduced binding affinities to *Mt*DXR. The best *Ec*DXR inhibitor **9** (*K_i*^{*Ec*DXR} = 0.42 μM) in this series has a 7-fold affinity loss to *Mt*DXR (*K_i*^{*Mt*DXR} = 3.2 μM). In addition, it appears that the *M. tuberculosis* enzyme is very sensitive to any additional substituents on **9**. For example, compound **34** with only a small 4'-fluoro substituent, which is still a strong *Ec*DXR inhibitor (*K_i*^{*Ec*DXR} = 1.1 μM), is much less active against *Mt*DXR with a *K_i*^{*Mt*DXR} of 50.7 μM. Other derivatives of **9**, compounds **35**–**37** with a 4'-chloro, 4'-methyl, and 3'-methoxy substituent, are essentially inactive against *Mt*DXR. On the other hand, compounds **8** and **10** possess slightly improved *K_i* values (1.6 and 6.9 μM, respectively) for *Mt*DXR, compared to their *K_i*^{*Ec*DXR} values of 2.3 and 7.9 μM.

The inhibition study of *M. tuberculosis* DXR is of interest, since it shows most inhibitors, including the reference compounds **1** and **2**, have significantly reduced binding affinities to the enzyme (*K_i*^{*Ec*DXR} ≫ *K_i*^{*Mt*DXR}). More problematic is that the substrate DXP, however, binds to *Mt*DXR twice as tightly as *Ec*DXR (*K_m*^{*Mt*DXR} = 47.1 μM^{2d} vs *K_m*^{*Ec*DXR} = 99 μM²⁰). These two factors render the inhibition of *M. tuberculosis* DXR in the presence of DXP even more difficult and make it a great challenge to design potent *Mt*DXR inhibitors.

Crystallographic Studies of *Mt*DXR. Due to the limited number of active *Mt*DXR inhibitors, our attempts for QSAR studies in *Mt*DXR inhibition did not yield a predictive model. X-ray crystallography was used to investigate the possible structural basis for the generally reduced inhibitor affinity to the protein. DXR is highly conserved across the species using the non-mevalonate pathway.^{9c} *Mt*DXR and *Ec*DXR share an overall




Figure 3. (A) Correlation between the experimental *Ec*DXR inhibitory activity and predicted activity by the phase QSAR model, showing $r^2 = 0.95$. (B) Flexible alignment of compounds (only heavy atoms and polar hydrogens displayed for clarity), with compound **9** shown in a ball and stick model and **1** in a tube model. (C) Phase negative ionic fields, pink favorable, yellow disfavored. (D) Electron-withdrawing fields, red favorable. (E) Hydrophobic fields, cyan favorable, purple disfavored. (F) H-bond donor fields, orange disfavored.

~40% sequence identity and 69% similarity (Supporting Information Figure S1), with the active site showing even higher homology (~50% identity). In fact, despite the 5× affinity difference, **1** binds to the two DXRs in a very similar fashion, as shown in the aligned *Mt*DXR:1 (PDB code: 2JCV)¹⁸ and *Ec*DXR:1 (PDB code: 2EGH)^{11d} structures (Figure S2). Three common interactions are (1) the electrostatic/H-bonds between the phosphonate and the protein; (2) metal ion (Mg^{2+} or Mn^{2+}) chelation by the hydroxamate group; and (3) the hydrophobic interaction of the inhibitor backbone with a conserved Trp residue (Trp211 for *Ec*DXR and Trp203 for *Mt*DXR), which also shields the active site from the solvent. From the two DXR:1 structures, it is still unclear what factor determines the affinity difference.

We turned to compound **4** for its larger affinity difference ($K_i^{EcDXR} = 36$ nM vs $K_i^{MtDXR} = 1500$ nM). In addition, **4** is also 10× less active than **1** against *Mt*DXR, while these two compounds have similar activity on *Ec*DXR. We determined the X-ray crystal structure of *Mt*DXR in complex with **4**, Mn^{2+} , and NADPH, at 2.5 Å. Details of data processing and refinement are shown in Table 4 and the overall structure of the *Mt*DXR:4 complex illustrated in Figure S3a. Although the protein crystallizes as a homodimer, only one monomer (monomer A) contains inhibitor **4** as well as Mn^{2+} and NADPH in the active site. In addition, a flexible loop consisting of the residues 198–208, which acts as a “cap” opening and closing when the substrate or an inhibitor enters the active site, adopts a closed conformation (Figure S3b). The other monomer (monomer B) includes a Mn^{2+} , SO_4^{2-} , and NADPH in the active site, with the flexible loop in a widely open conformation due to crystal packing interactions (Figure S3c).¹⁸ No inhibitor is found in the active site of monomer B, while a SO_4^{2-} ion is located in the phosphonate binding site of **1**. The above phenomena can also be observed in the *Mt*DXR:1 structure.¹⁸ There is one molecule

of **4** found on the surface of monomer B, likely due to nonspecific binding. Each of the two Mn^{2+} ions is anchored to the protein via coordination to three carboxylate groups of the residues Asp151, Glu153, and Glu222. Moreover, the binding of both NADPH molecules in the monomers A and B closely resemble those found in the *Mt*DXR:1 structure. Since monomer A mimics the solution state of the *Mt*DXR:4 complex better, the discussion below is focused on monomer A. In addition, the racemic form of **4** was used in the crystallization experiment and it is not unambiguous to assign the absolute configuration of the α -carbon atom of the inhibitor. As shown in Figure S4, good models for (*R*)- and (*S*)-**4** can be generated, in which the phosphonates, hydroxamates, and 3,4-dichlorophenyl groups are located almost in the same sites, but the conformations of the linker carbon skeletons differ from each other. We therefore assigned 50% occupancy for each enantiomer in the final structure. While (*R*)-**4** is used for the following discussion on inhibitor–protein interactions, the (*S*)-**4** molecule possesses essentially the same properties.

Close-up views of the active site of the *Mt*DXR:4 structure and protein–ligand interactions are shown in Figure 4a and Supporting Information Figure S5, respectively. The phosphonate group of **4** is located in the phosphonate binding site of **1**, with all of the three O atoms forming hydrogen bonds and/or electrostatic interactions with the residues Lys219, Asn218, Ser213, and Ser177. The hydroxamate group of **4** is found to chelate Mn^{2+} , with the distances between the two O atoms and the metal ion being 2.04 and 2.51 Å. These two features are common to all known structures of DXR:1 complexes. The 3,4-dichlorophenyl group of **4** is located in a hydrophobic cavity, surrounded by Pro265, Ser245, and Trp203. However, there is no π – π stacking interaction between the 3,4-dichlorophenyl and the indole ring of Trp203. Moreover, it is remarkable that the

electron density for the residues 198–203 in the flexible loop is relatively weak with high B factor values, showing that this part of the loop flips around and has no strong interactions with the inhibitor **4**. On the contrary, the loop in the *MtDXR*:**1** structure is bound well near the center of the active site, with the indole ring of Trp203 covers on top of the relatively hydrophobic carbon skeleton of **1** (Figure 4b). It can therefore be inferred that the weak interactions between Trp203 and the 3,4-dichlorophenyl ring could account for the reduced activity of compound **4** on *MtDXR*.

Table 4. Data Processing and Refinement Statistics

Data Processing	
Wavelength (Å)	1.542
Space group	$P2_1$
Unit cell dimensions	
a, b, c (Å)	67.25, 65.18, 85.41
α, β, γ (°)	90.0, 101.3, 90.0
Resolution (Å)	100–2.56(2.6–2.56)
Unique reflections	22365 (1144)
Completeness (%)	93.9 (94.5)
Redundancy	3.9 (3.8)
R_{sym} (%)	13.6 (51.8)
I/σ (I)	11.2 (2.2)
Refinement	
Resolution (Å)	33.45–2.55 (2.71–2.55)
Number of reflections used in working set	21222 (3505)
R/R_{free}^a (%)	20.8/27.8
Number of non-hydrogen atoms	5866
Number of solvent waters	119
Mean B-factor from Wilson plot (Å ²)	49.90
Mean B-factor, protein atoms (Å ²)	52.42
Mean B-factor, solvent atoms (Å ²)	44.21
Mean B-factor, inhibitors (Å ²)	59.47
Mean B-factor, NADPH (Å ²)	51.54
Mean B-factor, Mn^{2+} (Å ²)	59.17
Mean B-factor, SO_4^{2-} (Å ²)	56.24
Root mean square deviations from ideality	
Bond length (Å)	0.008
Bond Angle (°)	1.3

^a A subset of the data (5%) was excluded from the refinement and used to calculate R_{free} .

Next, we analyzed the binding differences of compound **4** in *MtDXR* and *EcDXR*. Although the crystal structure of *EcDXR* in complex with **4** has not been available, a model of how **4** binds to the protein has been generated in our previous study^{9c} by using the crystal structure of the *EcDXR*:**8** complex, which is also supported by experimental data. The modeled *EcDXR*:**4** structure was aligned with the *MtDXR*:**4** structure, and as shown in Figure 4c, the two molecules of **4** can almost be superimposed onto each other and possess a similar binding pose. Nevertheless, a notable difference exists with respect to the conformation of the flexible loops of the two proteins. The *EcDXR* loop (in purple) is located closer to the inhibitor, with the Trp211 indole ring forming a π – π stacking interaction with the 3,4-dichlorophenyl group of **4**, affording a tight binding mode. This is also in line with the potent inhibitory activity of **4** ($K_i^{\text{EcDXR}} = 36$ nM). On the other hand, the loosely bound loop of *MtDXR* might be again responsible for the considerably lowered affinity of **4** to the enzyme ($K_i^{\text{MtDXR}} = 1500$ nM).

The exact origin for the weak interaction between Trp203 in *MtDXR* and compound **4** is unclear, but should be attributed to factors from both the protein and the inhibitor. From the side of the protein, the flexible loop of *EcDXR* has one conformationally more rigid proline residue (Pro209), while that of *MtDXR* contains two proline residues (Pro201 and Pro207), which could restrict the conformational change of Trp203 from having a favorable π – π stacking interaction with the 3,4-dichlorophenyl group of inhibitor **4**. However, this might not be the only reason, since Trp203 is still able to move closely to **1** (Figure 4b). From the side of the inhibitor, the relatively weak electron deficiency on the 3,4-dichlorophenyl ring could be another unfavorable factor. For example, phosphonate compound **21** containing this group (Table 1) has essentially no activity against both *EcDXR* and *MtDXR*. The more electron-deficient pyridine analogue **8** is a good DXR inhibitor and, interestingly, shows slightly improved activity on *MtDXR* over *EcDXR*, suggesting a pyridine containing analogue of **4** could be particularly active against *MtDXR*.

Synthesis Aspect. Compounds **1–6** were prepared according to published methods^{8j,9a} and **39–42** are commercially available. Compounds **7–10**, **14**, **15**, **21**, and **23** were from our previous studies.^{9b,c} Scheme 1 shows the general synthetic methods for making the other compounds. 2-Bromopyridine or bromobenzene was converted to their corresponding phosphonate diethyl ester, using either a palladium catalyzed reaction or a lithium–halogen exchange followed by treatment with diethyl chlorophosphate. The esters thus obtained were hydrolyzed with





Figure 4. (A) Active site of the *MtDXR*:**4** structure. The protein backbone is shown as green curved lines, **4** as a ball and stick model, and Mn^{2+} as a brown sphere. (B) Crystal structure of *MtDXR*:**4** (in green) superimposed with the structure of *MtDXR*:**1** (in orange). For clarity, only the flexible loops containing residues 198–208 of the two structures are shown as curved lines. (C) Crystal structure of *MtDXR*:**4** (in green) superimposed with a modeled structure of *EcDXR*:**4** complex (in purple). Only the flexible loops (residues 198–208 for *MtDXR* and 205–215 for *EcDXR*) of the two structures are shown as curved lines.

Scheme 1^a

^a Reagents and conditions: (i) PH(OEt)_2 , $\text{Pd(Ph}_3\text{P)}_4$, Ph_3P , DIPEA, toluene, reflux; (vi) $n\text{-BuLi}$, CIP(O)(OEt)_2 , THF, -78°C ; (iii) TMSBr , CH_3CN , rt, then $\text{MeOH/H}_2\text{O}$; (iv) P(OEt)_3 , 120°C , overnight; (v) RB(OH)_2 , $\text{Pd(Ph}_3\text{P)}_4$, Na_2CO_3 , toluene/ H_2O (v/v 3:1), 90°C ; (vi) $m\text{-CPBA}$, CHCl_3 ; (vii) Ac_2O , 90°C ; (viii) KOH , MeOH ; (ix) SOCl_2 , CH_2Cl_2 ; (x) LDA , 2-bromoethylphosphonate, THF, -78°C ; (xi) NaH , tetraethyl diphosphonate, THF, 0°C ; (xii) H_2 , 5% Pd/C , MeOH ; (xiii) NH_2OBn , HOBT , EDC , TEA , CH_2Cl_2 ; (ixv) Na_2SO_3 , H_2O reflux.

bromotrimethylsilane to afford compounds **11** and **18**. Compounds **12**, **16**, **17**, **19**, **20**, **22**, and **24** with 1 or 2 methylene linkers were synthesized from their corresponding chloride with an Arbuzov reaction, followed by bromotrimethylsilane mediated hydrolysis. For making pyridine containing phosphonates **26–38**, a substituted 2-picoline, which was either commercially available or prepared from a bromo-substituted 2-picoline using a Suzuki coupling, was oxidized with *m*-chloroperbenzoic acid (*m*-CPBA) to give a pyridine oxide. It was then treated with Ac_2O at 90°C followed by hydrolysis (KOH) in MeOH to afford a 2-hydroxymethyl pyridine, which was reacted with thionyl chloride to give the corresponding chloride. It can then be readily converted to **26–38**, using an Arbuzov reaction followed by hydrolysis. To make compound **13** with a three-methylene linker, 2-picoline was treated with lithium diisopropylamide (LDA) followed by alkylation with diethyl 2-bromoethylphosphonate. The resulting diethyl ester was hydrolyzed with bromotrimethylsilane to give **13**. The reaction of the sodium salt of tetraethyl diphosphonate with 2-methoxypyridine-5-carbaldehyde produced an α,β -unsaturated phosphonate, which subjected to hydrogenation (5% Pd/C) and hydrolysis to give compound **25**. Suzuki coupling of methyl 5-bromopyridine-2-carboxylate or -2-acetate with benzenboronic acid afforded,

after hydrolysis, compounds **45** and **46**. Hydroxamates **43**, **44**, and **47** were synthesized from their corresponding acids with an amide forming reaction with *O*-benzylhydroxylamine, followed by hydrogenation. Finally, compound **48** was made from sodium sulfite and 2-chloromethyl-5-phenylpyridine.

CONCLUSION

Our SAR/QSAR and crystallographic studies of *E. coli* and *M. tuberculosis* DXRs have resulted in several novel observations that have implications in the design of potent inhibitors. First, a total of 41 lipophilic phosphonate and related compounds were synthesized and their inhibitory activity tested against *E. coli* DXR, with the best compound **9** (5-phenylpyridin-2-ylmethylphosphonic acid) possessing a K_i^{EcDXR} of 420 nM. Second, SAR studies show that (1) an electron-deficient aromatic ring, especially a pyridine, and a phosphonate headgroup are essential for the inhibition; (2) one methylene linker between these two groups is optimal; and (3) certain substituents (e.g., 5-phenyl or 5-hydroxyl) on the aromatic ring provide enhanced activity. In addition, these SARs can be rationalized using the crystal structures of the *EcDXR*:**8** and *EcDXR*:**9** complexes. We also developed a 3-D QSAR model for *EcDXR* inhibition that had good predictivity. Third, the 41 compounds as well as 7 known *EcDXR* inhibitors with a broad range of chemical diversity were tested against *M. tuberculosis* DXR, in an effort to find new antituberculosis agents. The vast majority of the inhibitors were found to exhibit considerably weakened activity against *MtDXR*, compared to that on *EcDXR*. In particular, fosmidomycin and its derivatives **1–4** with low nM K_i values against *EcDXR* show $5\text{--}40\times$ lower affinity to *MtDXR* (K_i^{MtDXR} : 130–1500 nM). Fourth, the X-ray crystal structure of *MtDXR* in complex with **4** is disclosed. The fosmidomycin moiety of **4** exhibits an expected binding mode, with the phosphonate located in the favored site and the hydroxamate group chelating the central metal ion. However, the flexible loop of *MtDXR* shows relatively weak electron density in the structure, in which Trp203 indole ring has no $\pi\text{--}\pi$ stacking interaction with the 3,4-dichlorophenyl group of **4**. This could represent a structural basis for the considerably lowered activity of **4** against *MtDXR*.

EXPERIMENTAL SECTION

All reagents were purchased from Alfa Aesar (Ward Hill, MA) or Aldrich (Milwaukee, WI). Compounds **1–6** were prepared according to published methods^{8j,9a} and **39–42** are commercially available. Compounds **7–10**, **14**, **15**, **21**, and **23** were from our previous studies.^{9b,c} All compounds were characterized by ^1H , ^{31}P , and/or ^{19}F NMR on a Varian (Palo Alto, CA) 400-MR spectrometer. The purities were monitored by a Shimadzu Prominence HPLC with a Zorbax C18 or C8 column (4.6×250 mm) or ^1H (at 400 MHz) absolute spin-count quantitative NMR analysis with imidazole as an internal standard. The purities of all compounds were found to be $>95\%$.

General Method A. A chloride (1 mmol) was heated with triethyl phosphite (5 mmol) at 120°C under N_2 overnight. Upon removal of excess triethyl phosphite under reduced pressure, the residue was purified with flash column chromatography to give the corresponding phosphonic acid diethyl ester.

General Method B. To a solution of a phosphonic acid diethyl ester (1 mmol) in dry CH_3CN (3 mL) was slowly added bromotrimethylsilane (3 mmol) at 0°C . After stirring overnight at room temperature, the solution was evaporated to dryness and MeOH (5 mL) was added to the residue. The solvent was removed under

reduced pressure again and the residue redissolved in MeOH (5 mL). Neutralization with 2 M NaOH to pH = 8 gave a disodium salt of a phosphonic acid as a white powder, which may be recrystallized in H₂O/acetone if needed.

General Method C. A mixture of a bromide (1 mmol), a boronic acid (1.5 mmol), Na₂CO₃ (424 mg, 4 mmol), and Pd(Ph₃P)₄ (58 mg, 5 mmol %) in toluene/H₂O (v/v 3:1, 9 mL) was heated to 90 °C under N₂ overnight. The product was extracted with EtOAc (3 × 10 mL) and purified with flash column chromatography.

General Method D. To a solution of a 2-methylpyridine (25 mmol) in CHCl₃ (100 mL) was added *m*-CPBA (11.2 g with 50% purity, 32.5 mmol). After stirring for 5 h, the reaction was quenched by adding saturated NaHCO₃ (30 mL) and the aqueous layer extracted with CH₂Cl₂ (15 mL). The combined organic phases were washed with 2 N HCl (15 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give the corresponding pyridine oxide, which was used without further purification. It was then treated with acetic anhydride (50 mL) at 90 °C for 12 h and the resulting dark solution concentrated under reduced pressure. The residue was dissolved in MeOH (30 mL) and KOH (2.1 g, 15 mmol) added at 0 °C. After stirring for 3 h at room temperature, the solvent was removed and the residue treated with saturated NaHCO₃ (10 mL) and extracted with EtOAc (3 × 10 mL). The combined organic layers were dried, filtered, concentrated, and purified with flash column chromatography (silica gel, hexane/EtOAc: 1/1) to give a pyridin-2-ylmethyl alcohol. To a solution of the alcohol (1 mmol) in CH₂Cl₂ (5 mL) was added thionyl chloride (0.22 mL, 3 mmol) solution in CH₂Cl₂ (2 mL) at 0 °C.

The reaction was warmed to room temperature and stirred overnight. Upon removal of the solvent, the residue was treated with saturated Na₂CO₃ (5 mL) and extracted with CH₂Cl₂ (3 × 10 mL). The combined organic layers were dried, filtered, and evaporated to give the corresponding chloride, which was used without purification, following the general methods A and B, to produce a phosphonic acid disodium salt as a white powder.

Pyridin-2-ylphosphonic Acid Disodium Salt (11). To a solution of 2-bromopyridine (790 mg, 5 mmol), diethyl phosphite (967 μL, 7.5 mmol), and diisopropylethylamine (DIPEA, 5 mL, 25 mmol) in toluene (40 mL) were successively added Ph₃P (1.31 g, 5 mmol) and Pd(Ph₃P)₄ (290 mg, 5 mmol %) under N₂. The reaction mixture was refluxed overnight. After removing solvent under reduced pressure, the residue was purified with column chromatography (silica gel, EtOAc) to give the corresponding phosphonate (350 mg), which was hydrolyzed, following the general method A, to give compound **11** as a white powder (320 mg, 31.5% yield). ¹H NMR (400 MHz, D₂O): δ 8.41 (d, *J* = 4.8 Hz, 1H), 7.69–7.67 (m, 2H), 7.23–7.21 (m, 1H). ³¹P NMR (162 MHz, D₂O): δ 6.0.

2-(Pyridin-2-yl)ethylphosphonic Acid Disodium Salt (12). To a solution of 2-(pyridin-2-yl)ethanol (2.46 g, 20 mmol) was added slowly thionyl chloride (5.74 mL, 80 mmol) at 0 °C. The reaction mixture was heated to 80 °C for 3 h. Upon removal of excessive thionyl chloride under reduced pressure, the residue was treated with saturated NaHCO₃ (30 mL) and extracted with CH₂Cl₂ (3 × 20 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered, and evaporated to give the crude chloride, which was used for the next step without purification. Compound **12** was obtained from the chloride following the general methods A and B as a white powder (3.8 g, 82% yield). ¹H NMR (400 MHz, D₂O): δ 8.29 (d, *J* = 4.8 Hz, 1H), 7.77–7.73 (m, 1H), 7.32 (d, *J* = 8.4 Hz, 1H), 7.21 (m, 1H), 2.93–2.68 (m, 2H), 1.80–1.72 (m, 2H). ³¹P NMR (162 MHz, D₂O): δ 22.4.

3-(Pyridin-2-yl)propylphosphonic Acid Disodium Salt (13). To a solution of diisopropylamine (1.51 mL, 10.8 mmol) in THF (12 mL) was added dropwise *n*-BuLi (2.46 mL, 2.93 M, 7.2 mmol) at –10 °C. After 30 min, the solution was cooled to –78 °C and 2-picoline (558 mg, 6 mmol) was added slowly. The reaction mixture

was stirred for 1 h, followed by addition of 2-bromoethylphosphonate (960 μL, 5 mmol). After 2 h, the reaction was warmed to room temperature and stirred for additional 1 h. The reaction was quenched with saturated NH₄Cl (10 mL) and extracted with EtOAc (3 × 25 mL). The combined organic phases were dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified with column chromatography (silica gel, EtOAc) to give the corresponding diethyl phosphonate, which was hydrolyzed, following the general method A, to afford compound **13** as a white powder (516 mg, 33.4% yield). ¹H NMR (400 MHz, D₂O): δ 8.30 (d, *J* = 4.8 Hz, 1H), 7.77–7.71 (m, 1H), 7.27 (d, *J* = 8.4 Hz, 1H), 7.19 (m, 1H), 2.74 (t, *J* = 7.6 Hz, 2H), 1.82–1.75 (m, 2H), 1.41–1.32 (m, 2H). ³¹P NMR (162 MHz, D₂O): δ 21.8.

Pyrazin-2-ylmethylphosphonic Acid Disodium Salt (16). To a solution of 2-methylpyrazine (1 mL, 22 mmol) in the CCl₄ (80 mL) were added *N*-chlorosuccinimide (4.27 g, 31.5 mmol) and benzoyl peroxide (0.26 g, 1.1 mmol). The reaction mixture was stirred overnight. Upon removal of the solvent under reduced pressure, the residue was purified with column chromatography (silica gel, hexane/EtOAc: 6/1) to give the corresponding chloride, which was used following the general methods A and B, to give compound **16** as an off-white powder (1.46 g, 30.4% yield). ¹H NMR (400 MHz, D₂O): δ 8.47 (d, *J* = 1.6, 1H), 8.38 (s, 1H), 8.31 (d, *J* = 1.6 Hz, 1H), 3.10 (d, *J* = 20.4 Hz, 2H). ³¹P NMR (162 MHz, D₂O): δ 15.9.

Pyrimidin-4-ylmethylphosphonic Acid Disodium Salt (17). Compound **17** was prepared from 4-methylpyrimidine, following a similar procedure as **16**, as an off-white powder (158 mg, 37% yield). ¹H NMR (400 MHz, D₂O): δ 8.84 (m, 1H), 8.46 (d, *J* = 4.8 Hz, 1H), 7.44 (s, 1H), 3.03 (d, *J* = 20.4 Hz, 2H). ³¹P NMR (162 MHz, D₂O): δ 13.9.

Phenylphosphonic Acid Disodium Salt (18). To a solution of bromobenzene (468 mg, 3 mmol) in THF (9 mL) was added *n*-BuLi (1.23 mL, 2.93 M, 3.6 mmol) at –78 °C. After stirring 1.5 h, diethyl chlorophosphate (776 mg, 4.5 mmol) was added. The reaction mixture was stirred for additional 3 h at the same temperature, quenched with saturated NH₄Cl (10 mL), and extracted with EtOAc (3 × 10 mL). The combined organic phases were dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified with flash column chromatography (silica gel, hexane/EtOAc: 1/1) to give the diethyl phosphonate, which was hydrolyzed, following the general method B, to afford compound **18** as a white powder (340 mg, 56% overall yield). ¹H NMR (400 MHz, D₂O): δ 7.30–7.21 (m, 5H). ³¹P NMR (162 MHz, D₂O): δ 17.2.

Benzylphosphonic Acid Disodium Salt (19). Compound **19** was prepared from benzyl chloride (250 mg, 2 mmol), following the general methods A and B, as a white powder (293 mg, 68.1% yield). ¹H NMR (400 MHz, D₂O): δ 7.20–7.04 (m, 5H), 2.13 (d, *J* = 20.0 Hz, 2H). ³¹P NMR (162 MHz, D₂O): δ 21.6.

2-Phenylethylphosphonic Acid Disodium Salt (20). Compound **20** was prepared from phenylethylchloride (280 mg, 2 mmol), following the general methods A and B, as a white powder (300 mg, 65% yield). ¹H NMR (400 MHz, D₂O): δ 7.16–7.01 (m, 5H), 2.21 (m, 2H), 2.13 (m, 2H). ³¹P NMR (162 MHz, D₂O): δ 19.5.

2-(2-Hydroxyphenyl)ethylphosphonic Acid Disodium Salt (22). Compound **22** was prepared from 2-chloroethylphenol (160 mg, 1 mmol), following the general methods A and B, as a white powder (94 mg, 40.5% yield). ¹H NMR (400 MHz, D₂O): δ 7.10 (d, *J* = 7.2 Hz, 2H), 6.99 (t, *J* = 7.2 Hz, 1H), 6.79 (t, *J* = 7.2 Hz, 1H), 6.73 (t, *J* = 7.2 Hz, 1H), 2.61 (m, 2H), 1.57 (m, 2H). ³¹P NMR (162 MHz, D₂O): δ 22.5.

5-(1-Hydroxy-pyridin-2-one)methylphosphonic Acid Disodium Salt (24). To a solution of 6-methoxypyridine-3-carbaldehyde (274 mg, 2 mmol) in methanol (6 mL) was slowly added NaBH₄ (114 mg, 3 mmol) at 0 °C. After 1 h at room temperature, the solvent was removed under reduced pressure and the reaction quenched with water

and extracted with EtOAc (3×10 mL). The combined organic phases were dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The resulting crude alcohol was dissolved in CH_2Cl_2 (8 mL) and a solution of SOCl_2 (0.43 mL, 6 mmol) in CH_2Cl_2 (5 mL) added slowly at 0°C . After 3 h, the solvent was removed under vacuum. The residue was treated with saturated Na_2CO_3 (10 mL) and extracted with CH_2Cl_2 (3×10 mL). The combined organic layers were dried, filtered, and evaporated to give the corresponding chloride, which was used following the general method A to give the diethyl phosphonate. It (1 mmol) was dissolved in CHCl_3 (5 mL) and *m*-CPBA (690 mg, 50%, 2 mmol) added. After 5 h, the reaction was quenched by adding saturated NaHCO_3 (10 mL) and the aqueous layer extracted with CH_2Cl_2 (15 mL). The combined organic phases were washed with 2 N HCl (5 mL), dried over Na_2SO_4 , filtered, and concentrated under reduced pressure to give the crude product. It was refluxed in acetyl chloride (5 mL) for 1 h. After excessive acetyl chloride was removed, the residue was washed with diethyl ether and hydrolyzed, following the general Method B, to give compound **24** as a white powder (152 mg, 61% overall yield). ^1H NMR (400 MHz, D_2O): δ 7.98 (s, 1H), 7.42 (d, $J = 8.8$ Hz, 1H), 6.63 (d, $J = 8.8$ Hz, 1H), 2.72 (d, $J = 18.8$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 18.6.

2-(Pyridin-2-one-6-yl)ethylphosphonic Acid Disodium Salt (25). To a solution of tetraethyl diphosphonate (1.1 mL, 5.5 mmol) in THF (15 mL) was added NaH (242 mg, 6.1 mmol, 60% in oil) at 0°C . 6-methoxy-pyridine-2-carbaldehyde (730 mg, 5 mmol) was added to the resulting solution. After 2 h at room temperature, the reaction mixture was quenched with saturated NH_4Cl and extracted with EtOAc (3×15 mL). The combined organic phases were dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The crude product was hydrogenated with 5% Pd/C in methanol (15 mL) for 2 h to afford the saturated ester, which was hydrolyzed, following the general method B, to afford compound **25** as a white powder (816 mg, 66% yield). ^1H NMR (400 MHz, D_2O): δ 7.49–7.41 (m, 1H), 6.32–6.23 (m, 2H), 2.61 (m, 2H), 1.57 (m, 2H). ^{31}P NMR (162 MHz, D_2O): δ 19.3.

(4-Bromopyridin-2-yl)methylphosphonic Acid Disodium Salt (26). It was prepared following the general method D as a white powder in an overall 41% yield from 4-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.07 (d, $J = 7.2$ Hz, 1H), 7.58 (s, 1H), 7.32 (d, $J = 7.2$ Hz, 1H), 2.89 (d, $J = 19.6$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 14.8.

(4-Phenylpyridin-2-yl)methylphosphonic Acid Disodium Salt (27). It was prepared following the general method D as a white powder in an overall 51% yield from 4-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.30 (d, $J = 4.4$ Hz, 1H), 7.72 (s, 1H), 7.70 (s, 1H), 7.66 (s, 1H), 7.45–7.39 (m, 4H), 3.01 (d, $J = 19.6$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 15.6.

4-(4-Fluorophenyl)pyridin-2-ylmethylphosphonic Acid Disodium Salt (28). It was prepared following the general method D as a white powder in an overall 37% yield from 4-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.25 (d, $J = 7.2$ Hz, 1H), 7.74 (t, $J = 8.8$ Hz, 2H), 7.62 (s, 1H), 7.38 (m, 1H), 7.16 (t, $J = 8.8$ Hz, 2H), 2.97 (d, $J = 20$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 15.5. ^{19}F NMR (376 MHz, D_2O): δ -116.2.

(5-Bromopyridin-2-yl)methylphosphonic Acid Disodium Salt (29). It was prepared following the general method D as a white powder in an overall 46% yield from 5-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 7.52–7.46 (m, 1H), 7.32 (s, 1H), 7.30 (s, 1H), 2.93 (d, $J = 19.6$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 15.3.

(5-Hydroxypyridin-2-yl)methylphosphonic Acid Disodium Salt (30). It was prepared from **31** by hydrogenation with 5% Pd/C in 99% yield as a white powder. ^1H NMR (400 MHz, D_2O): δ 7.64 (m, 1H), 7.30–7.28 (m, 2H), 2.96 (d, $J = 19.2$ Hz, 2H); ^{31}P NMR (162 MHz, D_2O): δ 14.9.

(5-Benzyloxy-pyridin-2-yl)methylphosphonic Acid Disodium Salt (31). It was prepared following the general method D as

a white powder in an overall 21% yield from 5-benzyloxy-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.01 (s, 1H), 7.39–7.28 (m, 7H), 5.08 (s, 2H), 2.90 (d, $J = 19.6$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 16.3.

(6-Bromopyridin-2-yl)methylphosphonic Acid Disodium Salt (32). It was prepared following the general method D as a white powder in an overall 45% yield from 6-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 7.52–7.46 (m, 1H), 7.32 (s, 1H), 7.30 (s, 1H), 2.93 (d, $J = 19.6$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 15.3.

(6-Phenylpyridin-2-yl)methylphosphonic Acid Disodium Salt (33). It was prepared following the general method D as a white powder in an overall 40% yield from 6-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 7.77–7.64 (m, 3H), 7.48–7.32 (m, 5H), 3.03 (d, $J = 20$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 15.6;

5-(4-Fluorophenyl)pyridin-2-ylmethylphosphonic Acid Disodium Salt (34). It was prepared following the general method D as a white powder in an overall 32% yield from 5-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.48 (s, 1H), 7.85 (d, $J = 8.0$ Hz, 1H), 7.59–7.56 (m, 2H), 7.44 (d, $J = 8.8$ Hz, 1H), 7.13 (t, $J = 8.8$ Hz, 2H), 2.98 (d, $J = 20$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 15.5. ^{19}F NMR (376 MHz, D_2O): δ -115.8.

5-(4-Chlorophenyl)pyridin-2-ylmethylphosphonic Acid Disodium Salt (35). It was prepared following the general method D as a white powder in an overall 36% yield from 5-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.50 (s, 1H), 7.86 (d, $J = 8.0$ Hz, 1H), 7.56–7.40 (m, 5H), 2.98 (d, $J = 20$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 15.4.

5-(4-Methylphenyl)pyridin-2-ylmethylphosphonic Acid Disodium Salt (36). It was prepared following the general method D as a white powder in an overall 28% yield from 5-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.50 (s, 1H), 7.84 (d, $J = 7.2$ Hz, 1H), 7.51 (d, $J = 8.0$ Hz, 2H), 7.42 (d, $J = 7.2$ Hz, 1H), 7.26 (d, $J = 8.0$ Hz, 2H), 2.97 (d, $J = 20$ Hz, 2H), 2.24 (s, 3H). ^{31}P NMR (162 MHz, D_2O): δ 15.5.

5-(3-Methoxyphenyl)pyridin-2-ylmethylphosphonic Acid Disodium Salt (37). It was prepared following the general method D as a white powder in an overall 32% yield from 5-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.56 (s, 1H), 7.84 (d, $J = 8.0$ Hz, 1H), 7.38–7.35 (m, 2H), 7.20–7.17 (m, 2H), 6.91 (d, $J = 8.0$ Hz, 1H), 3.77 (s, 3H), 3.10 (d, $J = 20$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 16.2.

5-(Naphthalen-2-yl)pyridin-2-ylmethylphosphonic Acid Disodium Salt (38). It was prepared following the general method D as a white powder in an overall 27% yield from 5-bromo-2-methylpyridine. ^1H NMR (400 MHz, D_2O): δ 8.57 (s, 1H), 7.98–7.78 (m, 5H), 7.63–7.61 (m, 1H), 7.44–7.38 (m, 3H), 2.97 (d, $J = 20$ Hz, 2H). ^{31}P NMR (162 MHz, D_2O): δ 15.5.

N-Hydroxy-pyridine-2-carboxamide (43). To a solution of pyridine-2-carboxylic acid (411 mg, 3 mmol), NH_2OBn (525 μL , 4.5 mmol), and triethylamine (1.26 mL, 9 mmol) in the CH_2Cl_2 (20 mL) were added HOBT (405 mg, 3 mmol) and *N*-ethyl-*N'*-(3-amino-propyl)carbodiimide hydrochloride (864 mg, 4.5 mmol), respectively. The reaction mixture was stirred overnight at room temperature. The reaction was quenched with water (10 mL) and extracted with CH_2Cl_2 (3×10 mL). The combined organic phases were washed successively with saturated Na_2CO_3 (10 mL) and brine, dried, filtered, and evaporated. The resulting oil was purified with flash column chromatography (silica gel, hexane/EtOAc: 2/1) to give the amide, which was hydrogenated with 10% Pd/C in methanol (12 mL) for 3 h to afford compound **43** as a pale yellow oil (201 mg, 48.9% overall yield). ^1H NMR (400 MHz, D_2O): δ 8.46–8.44 (m, 1H), 7.90–7.86 (m, 1H), 7.82–7.80 (m, 1H), 7.50–7.47 (m, 1H).

N-Hydroxy-(pyridin-2-yl)acetamide (44). It was prepared from pyridin-2-ylacetic acid following the similar procedure as **112** as

a colorless oil (169 mg, 37% overall yield). ^1H NMR (400 MHz, D_2O): δ 8.32–8.30 (m, 1H), 7.72–7.68 (m, 1H), 3.56 (s, 2H).

5-Phenylpyridine-2-carboxylic Acid (45). The methyl ester of compound **45** was prepared from 5-bromopyridine-2-carboxylate (324 mg, 1.5 mmol), following the general method C. It was dissolved in MeOH (5 mL) and 3 equiv of KOH was added at 0 °C. After stirring 3 h at room temperature, the solvent was removed under vacuum. The residue was dissolved in H_2O (1 mL), acidified with 2 M HCl, and extracted with ethyl acetate (30 mL). The organic layer was concentrated and purified with column chromatography (silica gel, hexane/EtOAc: 5/1) to give **45** as a white solid (214 mg, 72% yield). ^1H NMR (400 MHz, D_2O): δ 8.68 (s, 1H), 8.03–8.02 (m, 1H), 7.82 (d, J = 8.4 Hz, 1H), 7.61 (s, 1H), 7.59 (s, 1H), 7.44–7.35 (m, 3H).

(5-Phenylpyridin-2-yl)acetic Acid (46). It was prepared from 5-bromopyridine-2-acetate (350 mg, 1.5 mmol) following the similar procedure as **45**, as a white solid (257 mg, 81% yield). ^1H NMR (400 MHz, D_2O): δ 8.82 (s, 1H), 8.60 (d, J = 7.2 Hz, 1H), 7.83 (d, J = 7.2 Hz, 1H), 7.62 (s, 1H), 7.61 (s, 1H), 7.47–7.44 (m, 3H), 4.01 (s, 2H).

N-Hydroxy-(5-phenylpyridin-2-yl)acetamide (47). It was prepared from **46** (212 mg, 1 mmol) following the similar procedure as **43**, as a white solid (164 mg, 73% yield). ^1H NMR (400 MHz, d_6 -DMSO): δ 10.73 (br, 1H), 8.86 (s, 1H), 8.75 (s, 1H), 8.00 (d, J = 8.0 Hz, 1H), 7.70 (s, 1H), 7.68 (s, 1H), 7.50–7.46 (m, 2H), 7.42–7.40 (m, 2H), 4.01 (s, 2H).

5-Phenylpyridin-2-ylmethanesulfonic Acid Sodium Salt (48). The mixture of 2-chloromethyl-5-phenylpyridine (203.5 mg, 1 mmol) and sodium sulfite (126 mg, 1 mmol) in the water (3 mL) was heated under reflux for 12 h. The solvent was concentrated under reduced pressure and the solid obtained was filtered and washed with diethyl ether to give **48** as a white powder (240 mg, 97%). ^1H NMR (400 MHz, D_2O): δ 8.76 (s, 1H), 8.10 (d, J = 8.4 Hz, 1H), 7.73–7.48 (m, 6H), 4.37 (s, 2H).

DXR Inhibition Assay. *E. coli* DXR expression, purification, and activity assay were carried out as previously reported.^{9b,c} In brief, the inhibition assay was performed in a 96-well microplate using the purified *EcDXR* (10 nM), 2 mM MgCl_2 , 100 μM DXP, 50 μM NADPH in 50 mM HEPES buffer (pH = 7.5) containing 25 $\mu\text{g}/\text{mL}$ BSA (bovine serum albumin). Inhibitors were preincubated with *EcDXR* for 10 min at 30 °C, before adding DXP to initiate the reaction. The process was monitored at 340 nm with a Beckman DTX-880 microplate reader. The initial velocities of wells containing increasing concentrations of an inhibitor were calculated and input into *Prism* (version 3.0, GraphPad Software, Inc., La Jolla, CA). The IC_{50} values were then obtained by using standard dose–response curve fitting in the software. The IC_{50} s used for calculating K_i values were the mean values of at least three experiments with standard deviations of less than 20%. K_i values were calculated using the formula $K_i = \text{IC}_{50}/(1 + [\text{S}]/K_m)$, where $[\text{S}]$ is the concentration of DXP (100 μM) and K_m is the literature value of 99 μM .²⁰

M. tuberculosis DXR was expressed and purified as described in the literature.¹⁸ BL21-DE3 strain (Lucigen) was transformed and cultured at 37 °C in LB medium containing ampicillin (50 $\mu\text{g}/\text{mL}$). Upon reaching an optical density of ~ 0.6 at 600 nm, DXR expression was induced for 3 h by adding 0.42 mM isopropylthiogalactoside at 37 °C. Cells were then harvested and resuspended in 50 mM NaH_2PO_4 (pH 8.0), 300 mM NaCl (buffer A). After addition of 1 mM phenylmethylsulfonyl fluoride and sonication at 0 °C, the lysate was centrifuged at 20 000 rpm for 20 min and the supernatant was collected and subjected to an affinity column chromatography using the Talon cobalt resin (Clontech). The resin was washed with 5 mM imidazole in buffer A, and then the protein was eluted with 150 mM imidazole in buffer A. After desalting (using HiTrap, GE Healthcare) to 20 mM Tris pH 7.5, 150 mM NaCl, 2% glycerol, the protein was concentrated and stored in small aliquots at -80 °C.

The inhibition assay for *MtDXR* was carried out similarly as the *EcDXR* experiment. It was done in a 96-well microplate using the purified *MtDXR* (50 nM), 4 mM MgCl_2 , 100 μM DXP, 50 μM NADPH in 50 mM HEPES buffer (pH = 7.6) containing 25 $\mu\text{g}/\text{mL}$ BSA. Compounds were incubated with *MtDXR* for 10 min at 30 °C before adding DXP. The initial velocities were calculated and the IC_{50} values for inhibitors obtained by using *Prism*. The IC_{50} s used for calculating K_i values were from at least three independent experiments with standard deviations of less than 20%. K_i values were calculated using the formula $K_i = \text{IC}_{50}/(1 + [\text{S}]/K_m)$, where $[\text{S}]$ is the concentration of DXP (100 μM) and K_m is the literature value of 47.1 μM .²⁴

Molecular Modeling. Docking studies were performed as previously reported^{9b,c} using Schrödinger suite 2010,¹³ which includes all of the programs described below. Crystal structures of *E. coli* DXR were prepared using the module “protein preparation wizard” in *Maestro 9.1*¹⁵ with default protein parameters: water molecules were removed, hydrogens added, inhibitors extracted, and NADPH remained in the protein structure for docking. H-bonds were then optimized and the protein was energy-minimized using OPLS-2005 force field. A receptor grid, which is large enough to contain the whole active site, was generated using Glide¹² without constraints. Compounds were built, minimized using OPLS-2005 force field in *Maestro*, and then docked into the prepared protein structure using Glide (docking parameters: standard-precision and dock flexibly).

QSAR studies were performed with default settings using *Phase*¹⁴ (version 3.2). All compounds were built and geometry- and energy-minimized using the OPLS-2005 force field in *Maestro*. Compound alignment was carried out using the “Flexible Ligand Alignment” module in *Maestro* with compound **8** as a template. The aligned compounds were imported into the *Phase* program. A partial least-squares (PLS) method was used to correlate the activities of these compounds with the 3-D structural features calculated by *Phase* with default settings. The QSAR model was then validated by performing five leave-5-out training/test sets.

Crystallization and Structure Refinement. The crystallization of *MtDXR* was performed according to a published procedure.¹⁸ In brief, purified *MtDXR* (10 mg/mL) was mixed with an equal volume of the crystallization buffer containing 2 mM MnCl_2 , 3 mM NADPH, 0.2 M $(\text{NH}_4)_2\text{SO}_4$, 25% poly(ethylene glycol) 3350, and 0.1 M Bis-Tris, pH 5.7. The protein crystals appeared in ~ 1 week at room temperature, which were soaked for 1 h in the crystallization buffer containing an inhibitor (0.2 M). Data were collected to a resolution of 2.56 Å using a Rigaku FR-E+ SuperBright X-ray source at Baylor College of Medicine and were then processed using the program HKL2000.²¹ The initial structure was obtained by molecular replacement using the program *Phaser*²² with the coordinates of 2J CZ as a target. The refinement was carried out using the program CNS.²³ Since the inhibitor **4** was synthesized in a racemic form, both (*R*)- and (*S*)-**4** were built into the model with the occupancy of each enantiomer set to 0.5. The data collection and refinement statistics are provided in Table 4.

■ ASSOCIATED CONTENT

S Supporting Information. Figures S1–S5 showing Sequence alignment for *EcDXR* and *MtDXR*, superimposed structures of *EcDXR*:**1** and *MtDXR*:**1**, the overall structure of the *MtDXR*:**4** complex, the electron density of inhibitor **4**, and Ligplot interaction diagram for (*R*)-**4** in *MtDXR*. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Accession Codes

The coordinates for the *Mycobacterium tuberculosis* DXR:**4** complex have been deposited in Protein Data Bank as entry 3RAS.

AUTHOR INFORMATION

Corresponding Author

*Address: Department of Pharmacology, Baylor College of Medicine, 1 Baylor Plaza, Houston, TX 77030. Tel: 713-798-7415. E-mail: ysong@bcm.edu.

Author Contributions

[§]These authors contributed equally.

ACKNOWLEDGMENT

This work was supported by a grant (R21AI088123) from the National Institute of Allergy and Infectious Diseases (NIAID) to Y.S., grants AI049151 and AI065357 from NIAID to D.C.C., and a grant (Q1279) from Robert Welch Foundation to B.V.V.P. The recombinant *M. tuberculosis* DXR expression plasmid was kindly provided by Dr. T. Alwyn Jones (Uppsala University, Sweden). We thank the staff of the X-ray Crystallography Facility at Baylor College of Medicine for assistance in data collection.

ABBREVIATIONS

DXR, 1-deoxy-D-xylulose-5-phosphate reductoisomerase; EcDXR, DXR from *E. coli*; MtdDXR, DXR from *Mycobacterium tuberculosis*; DXP, 1-deoxy-D-xylulose-5-phosphate; IPP, isopentenyl diphosphate; DMAPP, dimethylallyl diphosphate; MEP, 2-C-methyl-D-erythritol-4-phosphate; SAR, structure activity relationship; QSAR, quantitative structure activity relationship.

REFERENCES

- (1) See a recent review and the references cited therein: Hunter, W. N. The non-mevalonate pathway of isoprenoid precursor biosynthesis. *J. Biol. Chem.* **2007**, *282*, 21573–21577.
- (2) (a) Singh, N.; Cheve, G.; Avery, M. A.; McCurdy, C. R. Targeting the methyl erythritol phosphate (MEP) pathway for novel antimalarial, antibacterial and herbicidal drug discovery: inhibition of 1-deoxy-D-xylulose-5-phosphate reductoisomerase (DXR) enzyme. *Curr. Pharm. Des.* **2007**, *13*, 1161–1177. (b) Rodriguez-Concepcion, M. The MEP pathway: a new target for the development of herbicides, antibiotics and antimalarial drugs. *Curr. Pharm. Des.* **2004**, *10*, 2391–2400. (c) Testa, C. A.; Brown, M. J. The methylerythritol phosphate pathway and its significance as a novel drug target. *Curr. Pharm. Biotechnol.* **2003**, *4*, 248–259. (d) Dhiman, R. K.; Schaeffer, M. L.; Bailey, A. M.; Testa, C. A.; Scherman, H.; Crick, D. C. 1-Deoxy-D-xylulose-5-phosphate reductoisomerase (IspC) from *Mycobacterium tuberculosis*: towards understanding mycobacterial resistance to fosmidomycin. *J. Bacteriol.* **2005**, *187*, 8395–8402.
- (3) Kuzuyama, T.; Shimizu, T.; Takahashi, S.; Seto, H. Fosmidomycin, a specific inhibitor of 1-deoxy-D-xylulose 5-phosphate reductoisomerase in the nonmevalonate pathway for terpenoid biosynthesis. *Tetrahedron Lett.* **1998**, *39*, 7913–7916.
- (4) Mine, Y.; Kamimura, T.; Nonoyama, S.; Nishida, M.; Goto, S.; Kuwahara, S. *In vitro* and *in vivo* antibacterial activities of FR-31564, a new phosphonic acid antibiotic. *J. Antibiot. (Tokyo)* **1980**, *33*, 36–43.
- (5) (a) Jomaa, H.; Wiesner, J.; Sanderbrand, S.; Altincicek, B.; Weidemeyer, C.; Hintz, M.; Turbachova, I.; Eberl, M.; Zeidler, J.; Lichtenthaler, H. K.; Soldati, D.; Beck, E. Inhibitors of the nonmevalonate pathway of isoprenoid biosynthesis as antimalarial drugs. *Science* **1999**, *285*, 1573–1576. (b) Missinou, M. A.; Borrmann, S.; Schindler, A.; Issifou, S.; Adegnika, A. A.; Matsiegui, P. B.; Binder, R.; Lell, B.; Wiesner, J.; Baranek, T.; Jomaa, H.; Kremsner, P. G. Fosmidomycin for malaria. *Lancet* **2002**, *360*, 1941–1942. (c) Borrmann, S.; Adegnika, A. A.; Matsiegui, P. B.; Issifou, S.; Schindler, A.; Mawili-Mboumba, D. P.; Baranek, T.; Wiesner, J.; Jomaa, H.; Kremsner, P. G. Fosmidomycin-clindamycin for *Plasmodium falciparum* infections in African children.

J. Infect. Dis. **2004**, *189*, 901–908. (d) Borrmann, S.; Lundgren, I.; Oyakhrome, S.; Impouma, B.; Matsiegui, P. B.; Adegnika, A. A.; Issifou, S.; Kun, J. F.; Hutchinson, D.; Wiesner, J.; Jomaa, H.; Kremsner, P. G. Fosmidomycin plus clindamycin for treatment of pediatric patients aged 1 to 14 years with *Plasmodium falciparum* malaria. *Antimicrob. Agents Chemother.* **2006**, *50*, 2713–2718. (e) Borrmann, S.; Adegnika, A. A.; Moussavou, F.; Oyakhrome, S.; Esser, G.; Matsiegui, P. B.; Ramharter, M.; Lundgren, I.; Kombila, M.; Issifou, S.; Hutchinson, D.; Wiesner, J.; Jomaa, H.; Kremsner, P. G. Short-course regimens of artesunate-fosmidomycin in treatment of uncomplicated *Plasmodium falciparum* malaria. *Antimicrob. Agents Chemother.* **2005**, *49*, 3749–3754. (f) Oyakhrome, S.; Issifou, S.; Pongratz, P.; Barondi, F.; Ramharter, M.; Kun, J. F.; Missinou, M. A.; Lell, B.; Kremsner, P. G. Randomized controlled trial of fosmidomycin-clindamycin versus sulfadoxine-pyrimethamine in the treatment of *Plasmodium falciparum* malaria. *Antimicrob. Agents Chemother.* **2007**, *51*, 1869–1871.

(6) Sakamoto, Y.; Furukawa, S.; Ogihara, H.; Yamasaki, M. Fosmidomycin resistance in adenylate cyclase deficient (*cya*) mutants of *Escherichia coli*. *Biosci. Biotechnol. Biochem.* **2003**, *67*, 2030–2033.

(7) Brown, A. C.; Parish, T. DXR is essential in *Mycobacterium tuberculosis* and fosmidomycin resistance is due to a lack of uptake. *BMC Microbiol.* **2008**, *8*, 78–86.

(8) For development based on fosmidomycin, see (a) Shtannikov, A. V.; Sergeeva, E. E.; Biketov, S. F.; Ostrovskii, D. N. Evaluation of *in vitro* antibacterial activity of fosmidomycin and its derivatives. *Antibiot. Khimioter* **2007**, *52*, 3–9. (b) Kuntz, L.; Tritsch, D.; Grosdemange-Billiard, C.; Hemmerlin, A.; Willem, A.; Bach, T. J.; Rohmer, M. Isoprenoid biosynthesis as a target for antibacterial and antiparasitic drugs: phosphonohydroxamic acids as inhibitors of deoxyxylulose phosphate reductoisomerase. *Biochem. J.* **2005**, *386*, 127–135. (c) Merckle, L.; de Andres-Gomez, A.; Dick, B.; Cox, R. J.; Godfrey, C. R. A fragment-based approach to understanding inhibition of 1-deoxy-D-xylulose-5-phosphate reductoisomerase. *ChemBioChem* **2005**, *6*, 1866–1874. (d) Munos, J. W.; Pu, X.; Liu, H. W. Synthesis and analysis of a fluorinated product analogue as an inhibitor for 1-deoxy-D-xylulose 5-phosphate reductoisomerase. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 3090–3094. (e) Ortmann, R.; Wiesner, J.; Silber, K.; Klebe, G.; Jomaa, H.; Schlitzer, M. Novel deoxyxylulosephosphate-reductoisomerase inhibitors: fosmidomycin derivatives with spacious acyl residues. *Arch. Pharm. (Weinheim)* **2007**, *340*, 483–490. (f) Silber, K.; Heidler, P.; Kurz, T.; Klebe, G. AFMoC enhances predictivity of 3D QSAR: a case study with DOXP-reductoisomerase. *J. Med. Chem.* **2005**, *48*, 3547–3563. (g) Woo, Y. H.; Fernandes, R. P.; Proteau, P. J. Evaluation of fosmidomycin analogs as inhibitors of the *Synechocystis* sp. PCC6803 1-deoxy-D-xylulose 5-phosphate reductoisomerase. *Bioorg. Med. Chem.* **2006**, *14*, 2375–2385. (h) Kurz, T.; Schlüter, K.; Kaula, U.; Bergmann, B.; Walter, R. D.; Geffken, D. Synthesis and antimalarial activity of chain substituted pivaloxyloxy-methyl ester analogues of Fosmidomycin and FR900098. *Bioorg. Med. Chem.* **2006**, *14*, 5121–5135. (i) Kurz, T.; Schlüter, K.; Pein, M.; Behrendt, C.; Bergmann, B.; Walter, R. D. Conformationally restrained aromatic analogues of fosmidomycin and FR900098. *Arch. Pharm. Chem. Life Sci.* **2007**, *340*, 339–344. (j) Haemers, T.; Wiesner, J.; Van Poecke, S.; Goeman, J.; Henschker, D.; Beck, E.; Jomaa, H.; Van Calenbergh, S. Synthesis of α -substituted fosmidomycin analogues as highly potent *Plasmodium falciparum* growth inhibitors. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 1888–1891. (k) Devreux, V.; Wiesner, J.; Jomaa, H.; Rozenski, J.; Van der Eycken, J.; Van Calenbergh, S. Divergent strategy for the synthesis of α -aryl-substituted fosmidomycin analogues. *J. Org. Chem.* **2007**, *72*, 3783–3789.

(9) For nonfosmidomycin-like inhibitors, see (a) Yajima, S.; Hara, K.; Sanders, J. M.; Yin, F.; Ohsawa, K.; Wiesner, J.; Jomaa, H.; Oldfield, E. Crystallographic structures of two bisphosphonate-1-deoxyxylulose-5-phosphate reductoisomerase complexes. *J. Am. Chem. Soc.* **2004**, *126*, 10824–10825. (b) Deng, L.; Sundriyal, S.; Rubio, V.; Shi, Z.; Song, Y. Coordination chemistry based approach to lipophilic inhibitors of 1-deoxy-D-xylulose-5-phosphate reductoisomerase. *J. Med. Chem.* **2009**, *52*, 6539–6542. (c) Deng, L.; Endo, K.; Kato, M.; Cheng, G.; Yajima, S.; Song, Y. Structures of 1-deoxy-D-xylulose-5-phosphate

reductoisomerase/lipophilic phosphonate complexes. *ACS Med. Chem. Lett.* **2011**, *2*, 165–170.

(10) For high-throughput screening, see Gottlin, E. B.; Benson, R. E.; Conary, S.; Antonio, B.; Duke, K.; Payne, E. S.; Ashraf, S. S.; Christensen, D. J. High-throughput screen for inhibitors of 1-deoxy-D-xylulose 5-phosphate reductoisomerase by surrogate ligand competition. *J. Biomol. Screen* **2003**, *8*, 332–339.

(11) (a) Yajima, S.; Nonaka, T.; Kuzuyama, T.; Seto, H.; Ohsawa, K. Crystal structure of 1-deoxy-D-xylulose 5-phosphate reductoisomerase complexed with cofactors: implications of a flexible loop movement upon substrate binding. *J. Biochem.* **2002**, *131*, 313–317. (b) Steinbacher, S.; Kaiser, J.; Eisenreich, W.; Huber, R.; Bacher, A.; Rohdich, F. Structural basis of fosmidomycin action revealed by the complex with 2-C-methyl-D-erythritol 4-phosphate synthase (IspC). Implications for the catalytic mechanism and anti-malaria drug development. *J. Biol. Chem.* **2003**, *278*, 18401–18407. (c) Mac Sweeney, A.; Lange, R.; Fernandes, R. P.; Schulz, H.; Dale, G. E.; Douangamath, A.; Proteau, P. J.; Oefner, C. The crystal structure of *E. coli* 1-deoxy-D-xylulose-5-phosphate reductoisomerase in a ternary complex with the antimalarial compound fosmidomycin and NADPH reveals a tight-binding closed enzyme conformation. *J. Mol. Biol.* **2005**, *345*, 115–127. (d) Yajima, S.; Hara, K.; Iino, D.; Sasaki, Y.; Kuzuyama, T.; Ohsawa, K.; Seto, H. Structure of 1-deoxy-D-xylulose 5-phosphate reductoisomerase in a quaternary complex with a magnesium ion, NADPH and the antimalarial drug fosmidomycin. *Acta Crystallogr., Sect. F Struct. Biol. Cryst. Commun.* **2007**, *63*, 466–470.

(12) *Glide*, version 5.5; Schrödinger, LLC: New York, 2010.

(13) *Schrödinger Suite*, version 2010; Schrödinger, LLC: New York, 2010.

(14) *Phase*, version 3.2; Schrödinger, LLC: New York, 2010.

(15) *Maestro*, version 9.1; Schrödinger, LLC: New York, 2010.

(16) (a) Altincicek, B.; Hintz, M.; Sanderbrand, S.; Wiesner, J.; Beck, E.; Jomaa, H. Tools for discovery of inhibitors of the 1-deoxy-D-xylulose 5-phosphate (DXP) synthase and DXP reductoisomerase: an approach with enzymes from the pathogenic bacterium *Pseudomonas aeruginosa*. *FEMS Microbiol. Lett.* **2000**, *190*, 329–333. (b) Woo, Y. H.; Fernandes, R. P.; Proteau, P. J. Evaluation of fosmidomycin analogs as inhibitors of the *Synechocystis* sp. PCC6803 1-deoxy-D-xylulose 5-phosphate reductoisomerase. *Bioorg. Med. Chem.* **2006**, *14*, 2375–2385. (c) Carretero-Paulet, L.; Ahumada, I.; Cunillera, N.; Rodriguez-Concepcion, M.; Ferrer, A.; Boronat, A.; Campos, N. Expression and molecular analysis of the *Arabidopsis* DXR gene encoding 1-deoxy-D-xylulose 5-phosphate reductoisomerase, the first committed enzyme of the 2-C-methyl-D-erythritol 4-phosphate pathway. *Plant Physiol.* **2002**, *129*, 1581–1591. (d) Grolle, S.; Bringer-Meyer, S.; Sahm, H. Isolation of the *dxr* gene of *Zymomonas mobilis* and characterization of the 1-deoxy-D-xylulose 5-phosphate reductoisomerase. *FEMS Microbiol. Lett.* **2000**, *191*, 131–137. Mueller, C.; Schwender, J.; Zeidler, J.; Lichtenthaler, H. K. Properties and inhibition of the first two enzymes of the non-mevalonate pathway of isoprenoid biosynthesis. *Biochem. Soc. Trans.* **2000**, *28*, 792–793. (e) Schwender, J.; Muller, C.; Zeidler, J.; Lichtenthaler, H. K. Cloning and heterologous expression of a cDNA encoding 1-deoxy-D-xylulose-5-phosphate reductoisomerase of *Arabidopsis thaliana*. *FEBS Lett.* **1999**, *455*, 140–144. (f) Rohdich, F.; Lauw, S.; Kaiser, J.; Feicht, R.; Kohler, P.; Bacher, A.; Eisenreich, W. Isoprenoid biosynthesis in plants: 2C-methyl-D-erythritol-4-phosphate synthase (IspC protein) of *Arabidopsis thaliana*. *FEBS J.* **2006**, *273*, 4446–4458.

(17) See a recent review and the references cited therein: Jassal, M.; Bishai, W. R. Extensively drug-resistant tuberculosis. *Lancet Infect. Dis.* **2009**, *9*, 19–30.

(18) Henriksson, L. M.; Unge, T.; Carlsson, J.; Aqvist, J.; Mowbray, S. L.; Jones, T. A. Structures of *Mycobacterium tuberculosis* 1-deoxy-D-xylulose-5-phosphate reductoisomerase provide new insights into catalysis. *J. Biol. Chem.* **2007**, *282*, 19905–19916.

(19) Koppisch, A. T.; Fox, D. T.; Blagg, B. S.; Poulter, C. D. *E. coli* MEP synthase: steady-state kinetic analysis and substrate binding. *Biochemistry* **2002**, *41*, 236–243.

(20) Kuzuyama, T.; Takahashi, S.; Takagi, M.; Seto, H. Characterization of 1-deoxy-D-xylulose 5-phosphate reductoisomerase, an enzyme

involved in isopentenyl diphosphate biosynthesis, and identification of its catalytic amino acid residues. *J. Biol. Chem.* **2000**, *275*, 19928–19932.

(21) Otwinowski, Z.; Minor, W. Processing of X-ray diffraction data collected in oscillation mode. In *Macromolecular Crystallography, Part A, Method Enzymology*, Carter, C. W. Jr., Sweet, R. M., Eds.; Academic Press: New York, 1997; Vol. 276, pp 307–326.

(22) McCoy, A. J.; Grosse-Kunstleve, R. W.; Adams, P. D.; Winn, M. D.; Storoni, L. C.; Read, R. J. *Phaser* crystallographic software. *J. Appl. Crystallogr.* **2007**, *40*, 658–674.

(23) Brunger, A. T.; Adams, P. D.; Clore, G. M.; DeLano, W. L.; Gros, P.; Grosse-Kunstleve, R. W.; Jiang, J. S.; Kuszewski, J.; Nilges, M.; Pannu, N. S.; Read, R. J.; Rice, L. M.; Simonson, T.; Warren, G. L. Crystallography & NMR system: A new software suite for macromolecular structure determination. *Acta Crystallogr., Sect. D* **1998**, *54*, 905–921.