

# A Nonheme, High-Spin {FeNO}<sup>8</sup> Complex that Spontaneously Generates $N_2O$

Alex M. Confer,<sup>†</sup> Alison C. McQuilken,<sup>†</sup> Hirotoshi Matsumura,<sup>‡</sup> Pierre Moënne-Loccoz,<sup>\*,‡</sup> and David P. Goldberg<sup>\*,†</sup>

<sup>†</sup>Department of Chemistry, Johns Hopkins University, Baltimore, Maryland 21218, United States

<sup>‡</sup>Institute of Environmental Health, Oregon Health & Science University, Portland, Oregon 97239, United States

**Supporting Information** 

**ABSTRACT:** One-electron reduction of  $[Fe(NO)-(N3PyS)]BF_4$  (1) leads to the production of the metastable nonheme  $\{FeNO\}^8$  complex, [Fe(NO)-(N3PyS)] (3). Complex 3 is a rare example of a high-spin (S = 1)  $\{FeNO\}^8$  and is the first example, to our knowledge, of a mononuclear nonheme  $\{FeNO\}^8$  species that generates N<sub>2</sub>O. A second, novel route to 3 involves addition of Piloty's acid, an HNO donor, to an Fe<sup>II</sup> precursor. This work provides possible new insights regarding the mechanism of nitric oxide reductases.

 $\mathbf{N}$  itric oxide (NO<sup>•</sup>) contributes to cellular stress through the secondary production of reactive oxygen and nitrogen species. To combat the toxicity associated with NO<sup>•</sup>, anaerobic microorganisms have evolved detoxification strategies based on the transition metal mediated reduction of NO<sup>•</sup> to N<sub>2</sub>O: 2 NO<sup>•</sup> + 2  $e^-$  + 2 H<sup>+</sup> → N<sub>2</sub>O + H<sub>2</sub>O. Denitrifying bacteria utilize a mixed heme–nonheme diiron motif to reductively activate NO<sup>•</sup> (cytochrome *c*-dependent NOR, cNOR), whereas nondenitrifying bacteria, archaea, and protozoans utilize a nonheme flavodiiron protein (FDP) to catalyze the same reaction (Figure 1).<sup>1</sup> Dinuclear Fe active sites



Figure 1. Fe-containing active sites involved in NO<sup>•</sup> reduction. Left: FDPs. Middle: heme-nonheme NORs. Right: P450nor.

can accommodate two Fe–NO units in close proximity, which may promote the key N–N bond formation required for N<sub>2</sub>O production and may also help stabilize NO<sup>•</sup> coupling intermediates (e.g., hyponitrite (ONNO<sup>2–</sup>)) or oxygencontaining coproducts (e.g., O<sup>2–</sup>, OH<sup>–</sup>). While both enzymatic<sup>1d,2</sup> and biomimetic<sup>3</sup> studies have suggested that preorientation of two Fe–NO monomers provides an efficient mechanism for N<sub>2</sub>O production, the example of fungal P450 NO<sup>•</sup> reductase (P450nor) demonstrates that NO<sup>•</sup> reduction to N<sub>2</sub>O can proceed at a mononuclear heme iron center (Figure 1). Alternative mechanisms have been proposed for the diiron active sites of cNOR and FDP that involve N–N bond formation and  $N_2O$  production occurring at a single nonheme iron center.  $\!\!\!^4$ 

Reduction of  $\{FeNO\}^{x}$  (x = 6, 7) species to the reactive  $\{FeNO\}^{8}$  or  $\{FeN(H)O\}^{8}$  states is often invoked as a prerequisite for N<sub>2</sub>O formation in enzymatic systems. There has been considerable interest in the preparation and characterization of  $\{FeNO\}^{8}/\{FeN(H)O\}^{8}$  model complexes,<sup>5</sup> but their synthesis is challenging because of their instability. A limited number of nonheme  $\{FeNO\}^{8}$  complexes have been characterized, <sup>5a-g</sup> and these complexes exhibit low-spin S = 0 ground states,<sup>5a-f</sup> with one exception of a high-spin S = 1 complex.<sup>5g</sup> However, there are no reports of a monouclear  $\{FeNO\}^{8}$  leading to N<sub>2</sub>O production in solution.<sup>6</sup> In the case of dinuclear Fe complexes, there are only two examples that produce N<sub>2</sub>O from ( $\{FeNO\}^{7}$ )<sub>2</sub> precursors.<sup>3a,b</sup> Biomimetic systems that not only stabilize the  $\{FeNO\}^{8}$  unit for spectroscopic interrogation but also direct reactivity toward N<sub>2</sub>O production would be valuable for study.

We previously described the synthesis of a mononuclear {FeNO}<sup>7</sup> complex as a model of the NO-bound form of the nonheme iron enzyme cysteine dioxygenase (CDO). This complex, [Fe(NO)(N3PyS)]BF<sub>4</sub> (1), prepared from addition of NO<sup>•</sup>(g) to the Fe<sup>II</sup> precursor  $[Fe^{II}(CH_3CN)(N3PyS)]BF_4$  $(2)^7$  was crystallographically characterized and showed excellent reversibility for both one-electron oxidation and reduction processes by cyclic voltammetry.<sup>8</sup> Herein we describe the one-electron reduction chemistry of 1, which leads to the generation of a metastable, nonheme {FeNO}<sup>8</sup> complex, [Fe(NO)(N3PyS)] (3). This complex spontaneously generates N2O upon standing in solution at 23 °C and is the first example, to our knowledge, of a mononuclear {FeNO}<sup>8</sup> species that reacts to give N2O. Spectroscopic and computational investigations also indicate that complex 3 exhibits a rare highspin S = 1 ground state. It is also shown that reaction of **2** with the HNO donor Piloty's acid (P.A.) leads to complex 3, suggesting that a nonheme Fe<sup>II</sup> center could serve as a target for nitroxyl (HNO/NO<sup>-</sup>) reactivity.

Previous examination of the {FeNO}<sup>7</sup> complex 1 by cyclic voltammetry revealed two reversible waves at  $E_{1/2} = 0.013$  and -1.18 V (vs Fc<sup>+</sup>/Fc in CH<sub>3</sub>CN).<sup>8a</sup> The well-behaved reversibility of the redox couple centered at  $E_{1/2} = -1.18$  V

Received: May 29, 2017

suggested that the one-electron-reduced {FeNO}<sup>8</sup> species may be stable enough for synthesis and characterization by chemical methods. The one-electron reductant decamethylcobaltocene (CoCp\*<sub>2</sub>) has a redox potential of  $E_{1/2} = -1.91$  V (vs Fc<sup>+</sup>/Fc in CH<sub>3</sub>CN),<sup>9</sup> indicating that it should be thermodynamically competent to reduce the {FeNO}<sup>7</sup> complex. Addition of CoCp\*<sub>2</sub> (1 equiv) to a dark brown solution of 1 in CH<sub>3</sub>CN at -40 °C results in the immediate formation of an air- and lightsensitive purple species with absorbance maxima at 520 and 720 nm ( $\varepsilon \approx 3400$  and 2200 M<sup>-1</sup> cm<sup>-1</sup>, respectively), corresponding to the {FeNO}<sup>8</sup> complex, [Fe(NO)(N3PyS)] (3) (Scheme 1).

Scheme 1. Reversible One-Electron Reduction of  $[Fe(NO)(N3PyS)]BF_4$  (1) at -40 °C



Spectral titrations at -40 °C show an isosbestic conversion of 1 to 3, reaching maximal formation of 3 upon addition of 1.0 equiv of reductant (Figure 2). These data confirm the 1:1



**Figure 2.** Left: UV–vis spectral titration showing the one-electron reduction of  $[Fe(NO)(N3PyS)]BF_4$  (1) (brown line) to [Fe(NO)-(N3PyS)] (3) (purple line) upon addition of  $CoCp^*_2$  (0.2–1.0 equiv) in CH<sub>3</sub>CN/toluene at -40 °C. Right: Plot of absorbance at 720 nm versus  $CoCp^*_2/1$ .

stoichiometry of the reduction reaction. Titration of 3 with the one-electron oxidant [FeCp<sub>2</sub>]<sup>+</sup>PF<sub>6</sub><sup>-</sup> results in loss of the spectrum for 3 and the quantitative recovery of the spectrum for the  $\{FeNO\}^7$  starting material 1 (Figure S2). The isosbestic point at 470 nm is retained during the reoxidation, although a second isosbestic point at 334 nm is likely obscured by absorbance from FeCp<sub>2</sub>. These findings demonstrate that the interconversion of the FeNO<sup>7</sup> and FeNO<sup>8</sup> species is both chemically reversible and high-yielding at  $-40 \circ C$ , and provide strong evidence that the Fe(NO)(N3PyS) formulation remains unchanged during reduction and oxidation cycles. The reduced 3 exhibits good stability over several hours under the spectral titration conditions (0.1 mM Fe complex) at -40 °C. However, efforts to concentrate and isolate 3 as a solid at low temperature have thus far been unsuccessful. Reduction of 1 at 23 °C also leads to complex 3 as seen by UV-vis, but the product has limited stability ( $t_{1/2} \approx 0.5$  h at 0.1 mM) at this temperature.

The {FeNO}<sup>8</sup> complex was further characterized by NMR and EPR spectroscopies. The <sup>1</sup>H NMR spectrum of **1** in CD<sub>3</sub>CN/toluene- $d_8$  (95/5 v/v) reveals well-separated, relatively sharp, paramagnetically shifted resonances between -2 and 35 ppm (Figure 3). Upon reduction with 1 equiv of CoCp\*<sub>2</sub>, the



Figure 3. Paramagnetic <sup>1</sup>H NMR spectra of 1 (red) and 3 (purple) in  $CD_3CN$ /toluene- $d_8$  (95/5 v/v) at 23 °C.

characteristic peaks for 1 disappear and are replaced by a new set of paramagnetic signals between -10 and 40 ppm. These new peaks uniformly diminish over 2 h at 23 °C, leaving only peaks in the diamagnetic region of the spectrum. These data are consistent with a paramagnetic, one-electron reduced species 3 decaying to diamagnetic products.

The magnetic moment for 3, prepared from freshly mixed 1 +  $CoCp_{2}^{*}$  in  $CD_{3}CN$ /toluene, was measured by the Evans method (see Supporting Information (SI)) and gave  $\mu_{\text{eff}} = 2.9$  $\mu_{\rm B}$ , which is in excellent agreement with the theoretical spinonly value for an S = 1 system (2.83  $\mu_{\rm B}$ ). In support of this assignment, DFT calculations yield a triplet (S = 1) ground state for 3, with a singlet (S = 0) state 11.8 kcal mol<sup>-1</sup> higher in energy (see SI for details). Mulliken population analysis of triplet 3 reveals spin densities of +3.156 on Fe and -1.154 on NO, suggestive of an electronic structure that is more in line with hs  $Fe^{I}$ -NO<sup>•</sup> than hs  $Fe^{II}$ -<sup>3</sup>NO<sup>-</sup>. The EPR spectrum for 1 at 15 K (g = [2.047, 2.007, 1.962]) arises from the S = 1/2 spin ground state of the {FeNO}<sup>7</sup> starting material and disappears upon one-electron reduction, leaving only a residual EPR signal near  $g \approx 2$  that can be assigned to unreacted CoCp<sub>2</sub>\* (Co<sup>II</sup>,  $d^7$ , S = 1/2) (Figure S5). These spectra indicate that the product 3 is EPR-silent, which, coupled with a paramagnetic <sup>1</sup>H NMR spectrum, is consistent with an S = 1 ground state. Monitoring the slow decay of 3 at 23 °C by EPR shows no new peaks, further indicative of diamagnetic decay products as observed by <sup>1</sup>H NMR.

The {FeNO}<sup>8</sup> complex **3** was characterized by lowtemperature resonance Raman (RR) spectroscopy and compared to the {FeNO}<sup>7</sup> starting material **1**. The RR spectra of **1** obtained with 458 nm excitation show a  $\nu$ (N–O) at 1641 cm<sup>-1</sup> and expected <sup>15</sup>N<sup>18</sup>O shifts that match frequencies previously measured by low-temperature FTIR (Figure 4). We



Figure 4. Resonance Raman spectra of 1 (top traces) and 3 (bottom traces) with  $^{14}\rm N^{16}O$  (black),  $^{15}\rm N^{18}O$  (red), and difference spectra (blue) at 110 K.

previously showed that 1 exhibits spin-crossover behavior, with a low-spin (S = 1/2) ground state and a thermally accessible high-spin (S = 3/2) excited state.<sup>8b</sup> The low-spin species exhibited a  $\nu(N-O)$  at 1641 cm<sup>-1</sup> at cryogenic temperatures, whereas the high-spin (S = 3/2)  $\nu$ (N–O) was observed at 1737  $cm^{-1}$  above 150 K (CD<sub>3</sub>CN). The RR spectrum of 3 (Figure 4)  $(\lambda_{\text{exc}} = 647 \text{ nm})$  reveals a prominent, broad band at 1588 cm<sup>-1</sup> that shifts to 1518 cm<sup>-1</sup> upon substitution with <sup>15</sup>N<sup>18</sup>O; this 70 cm<sup>-1</sup> downshift is in agreement with an N-O harmonic oscillator model (calculated  $\Delta \nu = -71$  cm<sup>-1</sup>), confirming the assignment of this band to the N-O stretch of the new {FeNO}<sup>8</sup> species. A vibration at 498 cm<sup>-1</sup>, which also shifts upon isotopic substitution with <sup>15</sup>N<sup>18</sup>O, is tentatively assigned as a  $\nu$ (Fe-NO) mode (Figure S7). The broadness of the  ${\rm FeNO}^{8}$  signals (half-widths of ~25 cm<sup>-1</sup>) is noteworthy and may reflect rotational conformers of the nitrosyl ligand. The  $\nu$ (N–O) frequency of 3 is ~150 cm<sup>-1</sup> lower than in the highspin  $\{FeNO\}^7$  complex 1, and can be compared with the 1618 cm<sup>-1</sup> frequency observed by Lehnert and co-workers for the four N-donor ligand (TMG3tren).5g For the few nonheme {FeNO}<sup>8</sup> complexes reported to date, a modest decrease in the  $\nu$ (N–O) frequency has been attributed to reduction of primarily Fe, rather than the nitrosyl ligand.<sup>5c,f,g</sup> In our case, unrestricted corresponding orbital (UCO) analysis also supports a largely metal-centered reduction of 1 to 3 (Figures S12, S13).

Solutions of **3** undergo a gradual color change from purple to red-orange at 23 °C, corresponding to slow decay of the UV– vis feature at 720 nm ( $t_{1/2} \approx 0.5$  h). This apparent self-decay reaction suggested the possibility of reductive N–N bond formation between two isolated {FeNO}<sup>8</sup> monomers, leading to N<sub>2</sub>O formation. Gas chromatography (GC) analysis of the headspace above the reaction of **1** (0.1 mM) with CoCp\*<sub>2</sub> (1 equiv) over 20 h showed the formation of N<sub>2</sub>O in ~54% yield based on an assumed 2:1 stoichiometry for {FeNO}<sup>8</sup> self-decay into N<sub>2</sub>O (Scheme 2). Control experiments established that the

Scheme 2. Production of  $N_2O$  from 3 and Conversion of 3 to 1 by Acid in CH<sub>3</sub>CN/toluene at 23 °C



 $\{FeNO\}^7$  starting material 1 does not produce any N<sub>2</sub>O under equivalent conditions, demonstrating the necessity of a reduced  $\{FeNO\}^8$  species for the critical N–N bond forming reaction. Although the final Fe decay products have not been identified, the diamagnetic NMR spectra of final reaction mixtures are consistent with magnetically coupled multi-iron clusters. Thus, complex 1 is a mononuclear nonheme iron complex that mediates N<sub>2</sub>O formation upon one-electron reduction.

It is well-known that HNO will rapidly dimerize to give  $N_2O$  and  $H_2O$  in aqueous solutions.<sup>10</sup> To examine the possibility of  $N_2O$  production from 3 via protonation from adventitious proton sources (e.g.,  $H_2O$ ) and subsequent release of HNO, we reacted freshly generated 3 with the weak acid  $[Me_3NH]^+[BPh_4]^-$  (1 equiv) at 23 °C. Rather than increasing the yield of  $N_2O$ , the formation of  $N_2O$  was greatly suppressed (~4%). This result argues against a mechanism involving

release of free HNO from 3 and subsequent N<sub>2</sub>O formation. Addition of a strong acid such as H<sup>+</sup>BArF<sup>-</sup> (1 equiv) also led to almost complete inhibition of N<sub>2</sub>O production. The reaction of 3 with acids instead led to the one-electron oxidation of 3 to give the {FeNO}<sup>7</sup> starting material 1 (confirmed by <sup>1</sup>H NMR and EPR spectroscopies). This reaction is precedented by the proton-induced oxidation of {FeNO}<sup>8</sup> complexes supported by sterically unencumbered porphyrins, which give the corresponding {FeNO}<sup>7</sup> species upon addition of acid.<sup>5h-l</sup>

Chemical reduction and bulk electrolysis of  $\{FeNO\}^7$  complexes are the known methods for the generation of  $\{FeNO\}^8$  complexes. However, to the best of our knowledge, there is no report of an  $\{FeNO\}^8$  complex resulting from the reaction of an Fe(II) precursor and a nitroxyl anion (NO<sup>-</sup>) donor. The HNO donor *N*-hydroxybenzenesulfonamide (Piloty's acid (P.A.)) can also furnish NO<sup>-</sup> under basic conditions.<sup>11</sup> We hypothesized that deprotonation of P.A. with 2 equiv of a strong base in the presence of  $[Fe^{II}(N_3PyS)-(CH_3CN)]BF_4$  (2) would provide access to the  $\{FeNO\}^8$  complex 3 through an Fe<sup>II</sup> + NO<sup>-</sup> trapping reaction (Figure 5).



Figure 5. UV–vis spectra showing addition of 'BuOK to a premixed solution of 2 and Piloty's acid in  $CH_3CN$  at -40 °C.

Mixing of 2 with P.A. (1 equiv) in  $CH_3CN$  at -40 °C shows no reaction by UV-vis; however, addition of <sup>t</sup>BuOK/18-crown-6 (1 equiv) triggers the slow decay of the  $Fe^{II}$  bands at 325, 418, and 493 nm over 1 h (Figure 5) to give a new spectrum with weaker bands at 418 and 501 nm. This new spectrum shows no further change for at least 6 h at -40 °C. However, addition of a second equiv of <sup>t</sup>BuOK/18-crown-6 results in immediate conversion to the characteristic spectrum of the  ${FeNO}^{8}$  complex 3 (Figure 5, purple trace). These results show that the overall reaction of 2 with P.A. and 2 equiv of base is a net transfer of  $NO^-$  to the  $Fe^{II}$  starting material to give 3. The observed intermediate formed after only 1 equiv of base (Figure 5, red trace) could be either an Fe-HNO adduct or a singly deprotonated Fe<sup>II</sup>-P.A. complex. Future work will focus on characterizing this species. It should be noted that the {FeNO}<sup>8</sup> product 3 formed at -40 °C under these conditions has reduced stability (full decay within  $\sim 5 \text{ min}$ ) as compared to one-electron reduction of 1. The lower stability in this case may be due to reaction with one or more of the byproducts of the deprotonation reaction (e.g., K<sup>+</sup>, <sup>t</sup>BuOH). Reaction of 2 with P.A. provides a novel approach for the preparation of {FeNO}<sup>8</sup> complexes that bypasses {FeNO}<sup>7</sup> reduction and suggests that nonheme Fe<sup>II</sup> sites in biology could be targets for NO<sup>-</sup> donors.

In summary, a nonheme {FeNO}<sup>8</sup> complex was prepared by chemical reduction of an {FeNO}<sup>7</sup> precursor and characterized by UV–vis, <sup>1</sup>H NMR, EPR, and RR spectroscopies. This complex exhibits a rare high-spin (S = 1) ground state and is the first example of a mononuclear nonheme {FeNO}<sup>8</sup> species that leads to production of N<sub>2</sub>O. These observations support the "super-reduced" mechanism for nonheme iron-dependent

## Journal of the American Chemical Society

NO reductases, which requires reduction of  $\{FeNO\}^7$  centers to  $\{FeNO\}^8$  prior to  $N_2O$  formation. Although this mechanism has been debated for a number of years, up to now there has been no direct characterization of a synthetic  $\{FeNO\}^8$  species that can generate  $N_2O$ . We have also demonstrated a novel method for forming an  $\{FeNO\}^8$  species from a readily available HNO donor and base. This work suggests that mononuclear nonheme iron centers in biology could be sites for NO reductive activation, as well as for the transport of HNO/NO<sup>-</sup>.

# ASSOCIATED CONTENT

# **S** Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.7b05549.

Syntheses, spectroscopy, GC traces, and DFT (PDF)

#### AUTHOR INFORMATION

#### Corresponding Authors

\*dpg@jhu.edu \*moennelo@ohsu.edu

# ORCID <sup>©</sup>

David P. Goldberg: 0000-0003-4645-1045

# Notes

The authors declare no competing financial interest.

### ACKNOWLEDGMENTS

The NSF (CHE1566007 to D.P.G.) and NIH (GM074785 to P.M.L.) are gratefully acknowledged for financial support. A.M.C. thanks the Harry and Cleio Greer Fellowship for support. We thank Prof. J. P. Toscano (JHU) and S. Nourian for use of instrumentation (headspace GC). The Kirin cluster at JHU KSAS is thanked for CPU time to A.M.C.

## REFERENCES

(1) (a) Kurtz, D. M., Jr. Dalton Trans. 2007, 4115–4121. (b) Wasser, I. M.; de Vries, S.; Moënne-Loccoz, P.; Schröder, I.; Karlin, K. D. Chem. Rev. 2002, 102, 1201. (c) Watmough, N. J.; Field, S. J.; Hughes, R. J.; Richardson, D. J. Biochem. Soc. Trans. 2009, 37, 392. (d) Sato, N.; Ishii, S.; Sugimoto, H.; Hino, T.; Fukumori, Y.; Sako, Y.; Shiro, Y.; Tosha, T. Proteins: Struct., Funct., Genet. 2014, 82, 1258.

(2) Silaghi-Dumitrescu, R.; Kurtz, D. M., Jr.; Ljungdahl, L. G.; Lanzilotta, W. N. *Biochemistry* **2005**, *44*, 6492–6501. (b) Hayashi, T.; Caranto, J. D.; Wampler, D. A.; Kurtz, D. M., Jr.; Moënne-Loccoz, P. *Biochemistry* **2010**, *49*, 7040–9.

(3) (a) Jiang, Y.; Hayashi, T.; Matsumura, H.; Do, L. H.; Majumdar, A.; Lippard, S. J.; Moënne-Loccoz, P. J. Am. Chem. Soc. 2014, 136, 12524. (b) Zheng, S.; Berto, T. C.; Dahl, E. W.; Hoffman, M. B.; Speelman, A. L.; Lehnert, N. J. Am. Chem. Soc. 2013, 135, 4902. (c) Kindermann, N.; Schober, A.; Demeshko, S.; Lehnert, N.; Meyer, F. Inorg. Chem. 2016, 55, 11538–11550.

(4) (a) Moënne-Loccoz, P. Nat. Prod. Rep. 2007, 24, 610.
(b) Timóteo, C. G.; Pereira, A. S.; Martins, C. E.; Naik, S. G.; Duarte, A. G.; Moura, J. J.; Tavares, P.; Huynh, B. H.; Moura, I. Biochemistry 2011, 50, 4251. (c) Duarte, A. G.; Cordas, C. M.; Moura, J. J.; Moura, I. Biochim. Biophys. Acta, Bioenerg. 2014, 1837, 375.
(d) Blomberg, L. M.; Blomberg, M. R.; Siegbahn, P. E. J. Biol. Inorg. Chem. 2006, 12, 79.

(5) (a) Serres, R. G.; Grapperhaus, C. A.; Bothe, E.; Bill, E.; Weyhermüller, T.; Neese, F.; Wieghardt, K. J. Am. Chem. Soc. 2004, 126, 5138. (b) Montenegro, A. C.; Amorebieta, V. T.; Slep, L. D.; Martín, D. F.; Roncaroli, F.; Murgida, D. H.; Bari, S. E.; Olabe, J. A. Angew. Chem., Int. Ed. 2009, 48, 4213. (c) Patra, A. K.; Dube, K. S.; Sanders, B. C.; Papaefthymiou, G. C.; Conradie, J.; Ghosh, A.; Harrop, T. C. Chem. Sci. 2012, 3, 364. (d) Sanders, B. C.; Patra, A. K.; Harrop, T. C. J. Inorg. Biochem. 2013, 118, 115. (e) Chalkley, M. J.; Peters, J. C. Angew. Chem., Int. Ed. 2016, 55, 11995. (f) Kupper, C.; Rees, J. A.; Dechert, S.; DeBeer, S.; Meyer, F. J. Am. Chem. Soc. 2016, 138, 7888. (g) Speelman, A. L.; Lehnert, N. Angew. Chem., Int. Ed. 2013, 52, 12283. (h) Choi, I. K.; Liu, Y.; Feng, D.; Paeng, K. J.; Ryan, M. D. Inorg. Chem. 1991, 30, 1832. (i) Pellegrino, J.; Bari, S. E.; Bikiel, D. E.; Doctorovich, F. J. Am. Chem. Soc. 2010, 132, 989. (j) Goodrich, L. E.; Roy, S.; Alp, E. E.; Zhao, J.; Hu, M. Y.; Lehnert, N. Inorg. Chem. 2013, 52, 7766. (k) Abucayon, E. G.; Khade, R. L.; Powell, D. R.; Zhang, Y.; Richter-Addo, G. B. J. Am. Chem. Soc. 2016, 138, 104. (l) Rahman, M. H.; Ryan, M. D. Inorg. Chem. 2017, 56, 3302. (m) Hu, B.; Li, J. Angew. Chem., Int. Ed. 2015, 54, 10579. (n) Kundakarla, N.; Lindeman, S.; Rahman, M. H.; Ryan, M. D. Inorg. Chem. 2016, 55, 2070.

(6) A site-isolated Fe center in a metal-organic framework (MOF) was shown to mediate reductive disproportionation of NO<sup>•</sup> to give  $N_2O$  and  $NO_2^{-}$ . See: Brozek, C. K.; Miller, J. T.; Stoian, S. A.; Dincă, M. J. Am. Chem. Soc. **2015**, 137, 7495.

(7) McQuilken, A. C.; Jiang, Y.; Siegler, M. A.; Goldberg, D. P. J. Am. Chem. Soc. 2012, 134, 8758.

(8) (a) McQuilken, A. C.; Ha, Y.; Sutherlin, K. D.; Siegler, M. A.; Hodgson, K. O.; Hedman, B.; Solomon, E. I.; Jameson, G. N. L.; Goldberg, D. P. J. Am. Chem. Soc. **2013**, 135, 14024. (b) McQuilken, A. C.; Matsumura, H.; Dürr, M.; Confer, A. M.; Sheckelton, J. P.; Siegler, M. A.; McQueen, T. M.; Ivanović-Burmazović, I.; Moënne-Loccoz, P.; Goldberg, D. P. J. Am. Chem. Soc. **2016**, 138, 3107.

(9) Connelly, N. G.; Geiger, W. E. Chem. Rev. 1996, 96, 877.

(10) Miranda, K. M. Coord. Chem. Rev. 2005, 249, 433.

(11) (a) Switzer, C. H.; Miller, T. W.; Farmer, P. J.; Fukuto, J. M. J. Inorg. Biochem. **2013**, 118, 128. (b) Chen, J.; Campos, J.; Mercado, B. Q.; Crabtree, R. H.; Balcells, D. Organometallics **2014**, 33, 4417.