This article was downloaded by: [University of Tennessee At Martin] On: 07 October 2014, At: 00:19 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Triphenylphosphine/2,3-Dichloro-5,6dicyanobenzoquinone (DDQ)/[n-Bu₄N]OCN as a Useful System for the Efficient Conversion of Tetrahydropyranyl (THP) Ethers to the Corresponding Alkyl Isocyanates

Batool Akhlaghinia ^{a b} & Sima Samiei ^b

^a Department of Chemistry, Faculty of Sciences , Ferdowsi University of Mashhad , Mashhad, Iran

^b Department of Chemistry, Damghan University of Basic Sciences, Damghan, Iran Published online: 21 Sep 2009.

To cite this article: Batool Akhlaghinia & Sima Samiei (2009) Triphenylphosphine/2,3-Dichloro-5,6-dicyanobenzoquinone (DDQ)/[n-Bu₄N]OCN as a Useful System for the Efficient Conversion of Tetrahydropyranyl (THP) Ethers to the Corresponding Alkyl Isocyanates, Phosphorus, Sulfur, and Silicon and the Related Elements, 184:10, 2525-2529, DOI: <u>10.1080/10426500802508212</u>

To link to this article: http://dx.doi.org/10.1080/10426500802508212

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



Triphenylphosphine/2,3-Dichloro-5,6dicyanobenzoquinone (DDQ)/[*n*-Bu₄N]OCN as a Useful System for the Efficient Conversion of Tetrahydropyranyl (THP) Ethers to the Corresponding Alkyl Isocyanates

Batool Akhlaghinia^{1,2} and Sima Samiei²

¹Department of Chemistry, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran ²Department of Chemistry, Damghan University of Basic Sciences, Damghan, Iran

Triphenylphosphine/2,3-dichloro-5,6-dicyanobenzoquinone/tetrabutylammonium cyanate was used as an efficient system for the conversion of tetrahydropyranyl ethers to the corresponding alkyl isocyanates.

 $\label{eq:keywords} \begin{array}{l} \mbox{Alkyl isocyanate; $2,3$-dichloro-5,6$-dicyanobenzoquinone (DDQ); tetrahydropyranyl ether (ROTHP); triphenylphosphine (TPP) \end{array}$

INTRODUCTION

Isocyanates are useful compounds capable of participating in a variety of reactions including nucleophilic addition reactions with alcohols and amines to produce carbamates and ureas, cycloaddition reactions to generate heterocycles, and polymerization reactions to produce commodities such as polyurethanes.¹ The high yields and the lack of byproducts associated with isocyanate chemistry have led to their widespread application in the pharmaceutical, agrochemical, and polymer industries.

Typically, aliphatic and aromatic isocyanates are generated from amines in the reaction with $phosgene^2$ or phosgene equivalents, such as diphosgene (trichloromethyl chloroformate)³ or triphosgene [bis(trichloromethyl) carbonate],⁴ or via thermal dissociation of

Received 5 September 2008; accepted 24 September 2008.

We gratefully acknowledge the partial support of this study by Damghan University Research Council.

Address correspondence to Batool Akhlaghinia, Department of Chemistry, Ferdowsi University of Mashhad, Mashhad, Iran. E-mail: akhlaghinia@um.ac.ir or b.akhlaghinia@dubs.ac.ir

carbamic acid derivatives using chloroformates,⁵ diphenylcarbonate,⁶ or N,N'-carbonyldiimidazole.⁷ Aryl isocyanates can also be generated from non-amine precursors via the rearrangements of acyl azides (Curtius rearrangement)⁸ and hydroxamic acids (Lossen rearrangement).⁹

Tetrahydropyranylation is one of the most practical and popular ways of protecting hydroxy groups of alcohols and phenols, especially in the synthesis of multifunctional organic molecules. THP ethers show remarkable stability towards some reagents such as organometallic reagents and hydrides, as well as reducing and oxidizing agents. Chemical transformation of protected functional groups to other functional groups is an important step in organic synthesis.¹⁰ Literature search reveals that only a few reports are available for the direct conversion of THP ethers to other functional groups. The known examples include conversion of THP ethers to bromides,^{11,12} iodides,¹³ sulfides,¹⁴ acetates,¹⁵ and esters.¹⁶

In this study, a novel and simple method for the efficient conversion of tetrahydropyranyl ethers to alkyl isocyanates is reported.

RESUTS AND DISCUSSION

Recently, we have reported the application of $Ph_3P/DDQ/R_4NX$ (X = Cl, Br, I, CN, N₃, NO₂, OCN) and 2,4,6-trichloro[1,3,5]triazine (TT)/R₄NX (X = CN, NO₂) systems for the conversion of alcohols, thiols, selenols, trimethylsilyl ethers, N₃, OCN, and tetrahydropyranyl ethers to the corresponding alkyl halides, cyanides, azides, nitrites, and isocyanates.¹⁷ Due to the wide applicability of tetrahydropyranyl ethers in multistep syntheses and in continuation of the study on the conversion of tetrahydropyranyl ethers to alkyl cyanides^{17d} and azides,^{17e} the transformation of THP ethers to alkyl isocyanates using the $Ph_3P/DDQ/[n-Bu_4N]OCN$ system under neutral and mild reaction condition was investigated (Scheme 1).

In order to optimize the reaction conditions, at first the effects of different ratios of ROTHP/PPh₃/DDQ/[*n*-Bu₄N]OCN, solvent, and temperature for the conversion of PhCH₂OTHP to PhCH₂NCO were

$$RO O \xrightarrow{Ph_3P/DDQ/n-Bu_4N[OCN]} RNCO$$

R = primary, secondary, and tertiary alkyl

SCHEME 1

ROTHP	Time (h)	Conversion $(\%)^a$	Isolated yield (%)
отнр	8	100	95
	11	100	90
	11.5	100	94
	30	100	89
OTHP	9	100	97
OTHP	8.5	100	96
ОТНР	7	100	98
OTHP	5	100	98
H ₃ CO THP	9	100	93
	11	100	90
OTHP	34	100	97

 TABLE I Conversion of Tetrahydropyranyl Ethers to Alkyl Isocyanates in Dry Acetonitrile Under Reflux Conditions

^aGC yield using internal standard.

examined. Employing the ratio of 1:2:2:2 in refluxing CH_3CN gave the best result and produced benzyl isocyanate after 7 h in 98% yield. These optimized conditions were applied for the conversion of structurally different tetrahydropyranyl ethers to the corresponding alkyl isocyanates. The results are shown in Table I.

As shown in Table I, this method is very suitable for the conversion of primary, secondary, and tertiary tetrahydropyranyl ethers to alkyl isocyanates in excellent yields.

A ¹³C NMR signal at 127 ppm is assigned to the quaternary carbon atom of the N=C=O group. A strong and sharp absorption band at 2280–2230 cm⁻¹ and a weak absorption band at 1450–1380 cm⁻¹ are typical for the isocyanate moiety.

In conclusion, the present investigation has demonstrated that the use of $Ph_3P/DDQ/[n-Bu_4N]OCN$ offers a simple, novel, and convenient method for the conversion of a wide variety of tetrahydropyranyl ethers to the corresponding alkyl isocyanates.

EXPERIMENTAL

Chemicals were obtained from Merck and Fluka. FT-IR spectra were recorded on a Perkin Elmer RXI spectrophotometer. NMR spectra were recorded with a Bruker Avance DPX 250 MHz instrument. The products were purified by column chromatography, and the purity of the products was determined by GC on a Shimadzu model GC 10-A instrument using *n*-octane as internal standard or by thin layer chromatography on silica gel polygram on SIL G/UV 254 plates.

Typical Procedure for Conversion of Tetrahydropyranyl Ethers to Alkyl Isocyanate

To a flask containing a mixture of DDQ (2 mmol, 0.454 g) and Ph₃P (2 mmol, 0.524 g) in dry CH₃CN (5 mL), while stirring Bu₄N[OCN] (2 mmol, 0.568 g) was added. Tetrahydropyranyl ether (1 mmol) was then added to this mixture. The reaction mixture was refluxed. GC analysis showed the completion of the reaction. The solvent was evaporated under reduced pressure. Column chromatography of the crude product on silica-gel using *n*-hexane as eluent gave alkyl isocyanates. The product was identified by comparison of its physical constants and spectral data with those of an authentic sample.

REFERENCES

- Books and reviews on isocyanates:(a) H. Ulrich, Chemistry and Technology of Isocyanates (Wiley, New York, 1996); (b) S. Ozaki, Chem. Rev., 72, 457 (1977); (c) B. A. Arbuzov and N. N. Zobova, Synthesis, 7, 433 (1982).
- (a) W. Hentschel, Ber. Dtsch. Chem. Ges., 17, 1284 (1884); (b) W. Siefken, Liebigs Ann. Chem., 562, 75 (1949); (c) R. J. Slocombe, E. E. Hardy, J. H. Saunder, and R. L. Jenkins, J. Am. Chem. Soc., 72, 1888 (1950).
- [3] (a) K. Kurita and Y. Iwakura, J. Org. Chem., 41, 2070 (1976); (b) K. Kurita and Y. Iwakura, Org. Synth. Coll., 6, 715 (1988).
- [4] (a) H. Eckert and B. Forster, Angew. Chem., Int. Ed. Engl., 26, 894 (1987); (b) L. Cotarca, P. Delogu, A. Nardelli, and C. V. Sunji, Synthesis, 553 (1996).
- [5] (a) H. R. Kricheldorf, Synthesis, 649 (1970); (b) H. R. Kricheldorf, Angew. Chem., 84, 107 (1972); (c) M. Sakamoto, M. Mochizuki, and M. Yoshioka, JP 10072428 (1998).
- [6] G. Greber and H. R. Kricheldorf, Angew. Chem., 80, 1028 (1968).
- [7] H. A. Staab and W. Benz, Angew. Chem., 73, 66 (1961).
- [8] (a) R. Bonjouklian and R. A. Ruden, J. Org. Chem., 42, 4095 (1977); (b) C. Kaiser and J. Weinstock, Org. Synth., 51, 48 (1971); (c) S. Heyden and G. Wilbert, Chem. Ind., 33, 1406 (1967).
- [9] (a) S. Bittner, S. Grinberg, and I. Kartoon, *Tetrahedron Lett.*, **15**, 1965 (1974); (b) F.
 A. Daniher, J. Org. Chem., **34**, 2908 (1969).
- [10] (a) P. J. Kocienski, Protecting Groups (Thieme, New York, 1994); (b) T. W. Green and P. G. M. Wuts, Protective Groups in Organic Synthesis, 2nd ed. Wiley, New

York, 1991); (c) A. J. Pearson and N. J. Roush, Handbook of Reagents for Organic Synthesis: Activating Agents and Protecting Groups, 1st ed. (Wiley, New York, 1999).

- [11] A. Wagner, M. Heitz, and C. Mioskowski, Tetrahedron Lett., 30, 557 (1989).
- [12] A. Tanaka and T. Oritani, Tetrahedron Lett., 38, 1955 (1997).
- [13] H. Firouzabadi, N. Iranpoor, and H. Hazarkhani, Tetrahedron Lett., 43, 713 (2002).
- [14] T. Sato, J. Otera, and H. Nozaki, J. Org. Chem., 55, 4770 (1990).
- [15] S. Chandrrasekhar, T. Ramachandar, M. V. Reddy, and M. Takhi, J. Org. Chem., 65, 4729 (2000).
- [16] S. Kim and W. J. Lee, Synth. Commun., 16, 659 (1986).
- [17] (a) N. Iranpoor, H. Firouzabadi, B. Akhlaghinia, and N. Nowrouzi, J. Org. Chem.,
 69, 2562 (2004); (b) N. Iranpoor, H. Firouzabadi, B. Akhlaghinia, and N. Nowrouzi, Tetrahedron Lett., 45, 3291 (2004); (c) B. Akhlaghinia and A. R. Pourali, Synthesis, 1747 (2004); (d) B. Akhlaghinia, Phosphorus, Sulfur, and Silicon, 179, 1783 (2004); (e) B. Akhlaghinia, Phosphorus, Sulfur, and Silicon, 1601 (2005); (f) B. Akhlaghinia, Synthesis, 1955 (2005); (g) B. Akhlaghinia and E. Roohi, Lett. Org. Chem., 2, 725 (2005); (h) B. Akhlaghinia and E. Roohi, Lett. Org. Chem., 3, 220 (2006); (i) B. Akhlaghinia and S. Samiei, J. Braz. Chem. Soc., 18, 1311 (2007); (j) B. Akhlaghinia and S. Samiei, Turk. J. Chem., 31, 35 (2007).