


A Chemoenzymatic Synthesis of Hept-6-ene-2,5-diol Stereomers: Application to Asymmetric Synthesis of Decarestrictine L, Pyrenophorol, and Stagonolide E

Sucheta Chatterjee,[†] Sneha Ghadigaonkar,[†] Payel Sur, Anubha Sharma,* and Subrata Chattopadhyay*

Bio-Organic Division, Bhabha Atomic Research Centre, Mumbai 400 085, India

S Supporting Information

ABSTRACT: The stereomers of hept-6-ene-2,5-diol derivatives were conceived as useful chiral intermediates and were synthesized starting from sulcatol using two lipase-catalyzed acylation reactions as the key steps. The versatility of the intermediates was demonstrated by converting them to the titled tetrahydropyran, macrolide, and macrodiolide compounds using standard synthetic protocols.



INTRODUCTION

Despite impressive progress, development of simple and efficient strategies for asymmetric syntheses of complex organic molecules remains a challenge.¹ Designing flexible and stereodivergent protocols, leading to several targets from a single starting molecule, has vital significance in organic synthesis. Small chiral intermediates with high functional density and multiple stereogenic centers are especially attractive to synthesize targets of structural and stereochemical diversities. The past two decades have witnessed extensive use of biocatalysts for various enantioselective organic transformations.² In particular, the lipases are most frequently used in organic synthesis since they do not require any cofactor, operate on a wide range of substrates, retain good catalytic activity in different media, display good stereoselectivity, and are easy to handle.³ Our group has extensively employed biocatalytic routes with the aim of developing new chiral intermediates and their subsequent transformations into several natural compounds.⁴ In our ongoing research on the development of potential immunomodulatory, anti-inflammatory, and antineoplastic agents,⁵ we identified several bioactive compounds 1–6 (Figure 1) that contain the hept-6-ene-2,5-diol derivatives 7 (different stereomers and Pgs) as the common structural motif.

This structural feature offered a remarkable opportunity to formulate a divergent strategy for the synthesis of a wide array of complex target molecules starting from the diol derivatives 7 as the ideal small-molecule chiral intermediates. Hence, in the present work, we have formulated a highly efficient biocatalytic route to all the four stereomers of 7 (Pg = TBDPS) as well as two stereomers of another derivative (Pg = PMB), and used




Figure 1. Some representative natural products bearing the hept-6-ene-2,5-diol structural motif.

some of these to devise asymmetric syntheses of (i) decarestrictine L (1), isolated from *Penicillium simplicissimum*;^{6a,b} (ii) pyrenophorol (2), isolated from *Byssoschlamys nivea*,^{6c} *Pyrenophora avenae*,^{6d} and *Stemphylium radicinum*,^{6e} cultures as well as from the imperfect fungus *Alternaria alternata*;^{6f} and (iii) stagonolide E (3), produced by *Stagonospora cirsi*, a fungal pathogen of *Cirsium arvense*.^{6g,h} The decarestrictines including 1 inhibit cholesterol biosynthesis in HEP-G2 liver cancer cells and in vivo.^{6a} On the other hand, the diolide 2 is specifically phytotoxic to the host *Avena sterilis* (wild oat), but not to other related plant species.^{7a} It also showed antimicrobial activity against *Microbotryum violaceum*, *Chlorella fusca*, *Escherichia coli*, and *Bacillus megaterium*.^{7b,c} In addition, it is a promising

Received: June 5, 2014

his group have obtained (*R*)-**10** in high yield and ee by combining the lipase-catalyzed acylation with in situ inversion or, alternatively, dynamic kinetic resolution using a Ru catalyst.^{14e} However, this protocol is unsuitable for the present work, because we wanted to synthesize all the stereoisomers of the key intermediate that warranted the availability of both (*R*)-**10** and (*S*)-**10** or the corresponding alcohols **9**.


The enantiomers of the alcohol **9** were individually silylated with *tert*-butyldiphenylsilyl chloride (TBDPSCI) in the presence of imidazole and 4-dimethylaminopyridine (DMAP) to obtain (*R*)-**11** and (*S*)-**11**. Next, the olefin function of (*R*)-**11** and (*S*)-**11** was subjected to reductive ozonolysis (O_3/Ph_3P) in CH_2Cl_2 to get the aldehydes (*R*)-**12** and (*S*)-**12**, respectively. Reaction of (*R*)-**12** with commercially available vinylmagnesium bromide furnished the allylic alcohol (3*R*,6*R*)-**13** as a 1:1 mixture of C-3 epimers. Its Novozym 435-catalyzed acetylation furnished (3*R*,6*R*)-**13** and (3*S*,6*R*)-**14** in >98% ees at 50% conversion (~6 h). The reaction was highly stereoselective and did not proceed further even under an extended period (Scheme 1.). A similar sequence of reaction with (*S*)-**12** also proceeded uneventfully and provided the target intermediate diastereomers (3*R*,6*S*)-**13** and (3*S*,6*S*)-**14** in enantiomerically pure forms. The absolute configurations of the alcohol and acetate were empirically assigned based on the fact that the Novozym 435-catalyzed acetylation of allylic alcohols has been found to furnish the (*S*)-acetate.^{12,17} This is also consistent with Kazlauskas' empirical rule.¹⁸ These results suggested that chirality at the distant C-6 center did not have any bearing on the diastereoselectivity of the chosen lipase-catalyzed resolution of **13**. Earlier, depending on the choice of hydrophobic supports, different immobilized CAL-B lipase preparations showed different enantioselectivities in the hydrolysis of certain racemic esters.¹⁹ Hence, we also carried out acetylation of (3*R*,6*R*)-**13** and (3*R*,6*S*)-**13** using another recombinant CAL-B lipase preparation (Sigma, L4777), expressed in *Aspergillus niger* and adsorbed on a macroporous acrylic resin. However, the reaction was too slow to be of any use in preparative chemistry.

Previously, the synthesis of the analogues of (3*R*,6*S*)-**13** and (3*S*,6*S*)-**14** containing the *tert*-butyldimethylsilyl protection at C-6 has been described, wherein the C-3 stereogenic center was installed using a lipase PS "Amano"-catalyzed acetylation with vinyl acetate in diisopropyl ether. However, the reaction required higher temperature (40 °C) and extended stirring (72 h). Moreover, this protocol is not amenable to the corresponding 6*R*-stereoisomers as it was derived from (*S*)-ethyl lactate.²⁰ To exclude any possible involvement of the protecting group at C-6 in the reaction, the alcohol (*S*)-**9** was converted to the *para*-methoxybenzyl (PMB) derivative (*S*)-**15** with *para*-methoxybenzyl chloride (PMBCl) in the presence of NaH. Oxidative cleavage (OsO_4/NMO ; $NaIO_4$) of its olefin function, followed by addition of vinylmagnesium bromide to the resultant aldehyde **16**, gave (3*R*,6*S*)-**17**, another derivative of the key intermediate. As expected, the Novozym 435-catalyzed acetylation of (3*R*,6*S*)-**17** also furnished (3*R*,6*S*)-**17** and (3*S*,6*S*)-**18** in excellent ees at ~50% conversion (~6 h). The enantiomeric ratios (*E*) of the lipase-catalyzed acetylation reactions were >195. Overall, the above simple synthetic strategy provided the target intermediate in its diastereomeric forms, and as differently protected derivatives. With all the stereoisomers of the intermediate in hand, we proceeded for the synthesis of the target compounds **1–3**. This also confirmed

the assigned configurations of the intermediates **13**, **14**, **17**, and **18** (all compound 7 equivalents).

Synthesis of Decastrictine L 1. For the synthesis, the acetate (3*S*,6*R*)-**14** was hydrolyzed to the diol derivative (3*S*,6*R*)-**13**, which was subjected to a cross-metathesis reaction²¹ with methyl vinyl ketone (MVK) in the presence of Grubbs' II catalyst by refluxing in dry CH_2Cl_2 to furnish compound **19**. To limit dimerization of **13**, the reaction was carried out using an excess of the less reactive olefin, MVK. Nevertheless, the homodimerized product of **13** was also obtained in ~7–8% yield. The reaction required use of freshly distilled MVK to get the product in an acceptable yield. The *E*-geometry of the olefin **19** as ascertained from its ¹H NMR spectrum was consistent with the proposed model for the cross-metathesis reaction.²² Desilylation of **19** with Bu_4NF directly afforded the target compound **1** via a concomitant Michael addition of the C-8 alcohol function to the conjugated ketone moiety (Scheme 2.). The physical and spectroscopic data of compound **1** were in accordance with the reported data of natural decastrictine L.^{8a}


Scheme 2^a



^a(i) $K_2CO_3/MeOH/25\text{ }^\circ C/6\text{ h}$, (ii) $CH_2=CHCOMe/Grubbs' II\text{ catalyst}/CH_2Cl_2/reflux/22\text{ h}$, (iii) $Bu_4NF/THF/0\text{ to }25\text{ }^\circ C/3\text{ h}$.

Synthesis of Pyrenophorol 2. For the synthesis, (3*S*,6*R*)-**13** was subjected to a cross-metathesis reaction with ethyl acrylate in the presence of Hoveyda Grubbs' II catalyst to furnish **20**. Its carbinol function was silylated with *tert*-butyldimethylsilyl chloride (TBSCl) in the presence of imidazole and DMAP to obtain **21**. This, on alkaline hydrolysis, furnished the acid **22**. In a parallel sequence, (3*S*,6*S*)-**13** was benzylated with BnBr in the presence of NaH as the base to furnish **23** as a mixture of rotamers, which, on desilylation with Bu_4NF , furnished the alcohol **24**. Esterification of **22** with **24** under the Mitsunobu conditions afforded the ester **25** (mixture of rotamers). Its desilylation with aqueous HF in CH_3CN afforded the diol **26**. The Novozym 435-catalyzed acrylation of **26** with ethyl acrylate furnished the desired acrylate **27** exclusively. The excellent regioselectivity of the reaction is noteworthy, given that Novozym 435 can acylate both 2-alkanols and allylic alcohols.¹² Moreover, the reaction avoided the use of hygroscopic, hazardous, and toxic acryloyl chloride, as well as vinyl acrylate that needs to be synthesized separately.²³ The reaction was slow, but proceeded without any side reaction, and the acrylate was conveniently isolated by filtering the reaction mixture, followed by solvent removal. The *R*-stereochemistry of the methylcarbinol moiety of **26** matched with the inherent enantioselectivity of the chosen lipase. Hence, this strategy may be useful in asymmetric syntheses of compounds possessing a chiral methylcarbinol moiety that is often encountered in various bioactive compounds.²⁴


The stage was set for the RCM reaction of **27** that was carried out with Grubbs' II catalyst to obtain **28** in modest yield. Control of the olefin geometry in the RCM-mediated macrocyclization is difficult due to the secondary olefin isomerization. This often leads to *E/Z* olefin mixtures with preponderance of the *E*-olefin that is governed by factors such

Scheme 3^a

^a(i) $\text{CH}_2=\text{CHCO}_2\text{Et}$ /Hoveyda Grubbs' II catalyst/ CH_2Cl_2 /25 °C/3 h, (ii) TBSCl /imidazole/ DMAP / CH_2Cl_2 /25 °C/7 h, (iii) aqueous 20% NaOH/MeOH /25 °C/2 h, (iv) $\text{K}_2\text{CO}_3/\text{MeOH}$ /25 °C/6 h, (v) $\text{NaH}/\text{BnBr}/\text{Bu}_4\text{NI}/\text{THF}/\text{reflux}/4$ h, (vi) $\text{Bu}_4\text{NF}/\text{THF}$ /0 to 25 °C/8 h, (vii) **22**/ $\text{Ph}_3\text{P}/\text{DIAD}/\text{THF}$ /0 to 25 °C/18 h, (viii) aqueous HF/MeCN /25 °C/16 h, (ix) $\text{CH}_2=\text{CHCO}_2\text{Et}/\text{Novozyme 435}/\text{diisopropyl ether}/25$ °C/30 h, (x) Grubbs' II catalyst/ CH_2Cl_2 /reflux/72 h, (xi) $\text{TiCl}_4/\text{CH}_2\text{Cl}_2$ /0 to 25 °C/0.5 h.

as solvent, catalyst, temperature, substrate structure, and product ring size. Nevertheless, we did not isolate any *Z*-isomer in the conversion of **27** to **28**, as revealed from the ^1H NMR spectrum. A TiCl_4 -catalyzed debenzoylation of **28** completed the synthesis of (5*S*,8*R*,13*S*,16*R*)-**2** (Scheme 3).²⁵

Synthesis of Stagonolide E 3. The synthesis of **3** commenced by the Hoveyda Grubbs' II catalyzed cross-metathesis of the intermediate (3*R*,6*S*)-**17** with acrolein to obtain the conjugated aldehyde **29**. After protecting its carbinol function with TBSCl /imidazole/ DMAP , the resultant compound **30** was subjected to a *Z*-selective Wittig–Horner reaction with ethyl diphenylphosphonoacetate²⁶ and NaH in THF to get the ester **31** in a 91/9 *Z/E* ratio (based on isolation after column chromatography). Oxidative removal of the PMB group with DDQ to the hydroxy ester **32**, followed by alkaline hydrolysis, furnished the seco acid **33**. This was converted to the target compound **3** by Mitsunobu lactonization to **34** and subsequent desilylation with aqueous HF (Scheme 4).

Scheme 4^a

^a(i) acrolein/Hoveyda Grubbs' II catalyst/ CH_2Cl_2 /25 °C/22 h, (ii) TBSCl /imidazole/ $\text{DMAP}/\text{CH}_2\text{Cl}_2$ /25 °C/7 h, (iii) $\text{NaH}/\text{THF}/(\text{PhO})_2\text{P}(\text{O})\text{CH}_2\text{CO}_2\text{Et}$ /0 °C/1 h; −78 °C/3 h, (iv) $\text{DDQ}/\text{aqueous CH}_2\text{Cl}_2$ /25 °C/3 h, (v) $\text{LiOH}/\text{THF}-\text{H}_2\text{O}$ /0 °C/6 h, (vi) $\text{Ph}_3\text{P}/\text{DIAD}/\text{toluene}/25$ °C/16 h, (vii) aqueous $\text{HF}/\text{MeCN}/25$ °C/48 h.

CONCLUSION

Overall, in consideration of hept-6-ene-2,5-diol as a common structural motif of several natural products, the stereoisomers and different derivatives (**13**, **14**, **17**, and **18**) of the diol were synthesized using two highly enantioselective lipase-catalyzed carbinol acetylation as the key steps. The versatility of the intermediates was established by converting some of these derivatives to the natural stereoisomers of decarestrictine **1**, pyrenophorol, and stagonolide **E**. Although these targets have been made before on numerous occasions, we elaborated a new general synthetic strategy, allowing control of both ring size and the configurations of stereogenic carbons into a range of significant materials using the key intermediate. Moreover, since our methodology gives access to all possible stereoisomers of the key intermediate, it would be possible to access all the stereoisomers of these natural products based on the described methodology. The yields of the target compound via the present stereodivergent strategy were comparable to those of the previously described target-specific syntheses. Given that the carbinol stereochemistry can be easily inverted, the synthetic strategy is also amenable for enantioconvergent routes that would provide the individual targets in much improved yields. Use of inexpensive reagents/materials and application of operationally simple reactions and environmentally benign biocatalysts were the other hallmarks of the flexible, efficient, and scalable syntheses. In addition to its use for traditional kinetic resolution applications, we also demonstrate that the Novozyme 435 lipase can be employed to effect regioselective acylation using a nontraditional acyl donor (ethyl acrylate). This transformation is particularly noteworthy and elevates the significance of the work.

EXPERIMENTAL SECTION

(±)-6-Methyl-5-hepten-2-ol 9. To a cooled (0 °C) and stirred suspension of LiAlH_4 (2.11 g, 55.56 mmol) in Et_2O (30 mL) was dropwise added **8** (10.00 g, 79.37 mmol) in Et_2O (70 mL). After

stirring for 2 h, the mixture was treated with aqueous saturated Na_2SO_4 and diluted with Et_2O , and the supernatant was filtered. The filtrate was carefully concentrated, and the residue was distilled to obtain pure **9** (9.2 g, 91%). colorless oil; bp: 90 °C/20 mm; IR: 3373, 1642 cm^{-1} ; ^1H NMR: δ 1.14 (d, J = 6.2 Hz, 3H), 1.48–1.59 (m, 2H), 1.65 (s, 3H), 1.71 (s, 3H), 2.00–2.15 (m, 3H), 3.74–3.90 (m, 1H), 5.11–5.19 (m, 1H); ^{13}C NMR: δ 17.5, 23.3, 24.4, 25.6, 39.1, 67.7, 124.0, 131.8.

(R)-6-Acetoxy-2-methyl-2-heptene 10. A mixture of (\pm)-**9** (7.50 g, 58.59 mmol), vinyl acetate (8.1 mL, 87.89 mmol), and Novozym 435 (0.75 g, ~12 mg/mmol) in hexane (30.0 mL) was agitated on an orbital shaker at 110 rpm for 50 min. The reaction mixture was filtered, and the solution was concentrated in vacuo to get a residue, which, on column chromatography (silica gel, 0–10% EtOAc /hexane), gave pure (**S**)-**9** (3.1 g, 41%) and (**R**)-**10** (4.4 g, 44%). (**S**)-**9**: colorless oil; $[\alpha]_{\text{D}}^{25} +10.9$ (c 1.30, CHCl_3) (lit.^{16b} $[\alpha]_{\text{D}}^{25} +10.5$ (c 0.4, CHCl_3)); IR: 3437 cm^{-1} ; ^1H NMR: δ 1.19 (d, J = 6.0 Hz, 3H), 1.41–1.57 (m, 2H), 1.62 (s, 3H), 1.69 (s, 3H), 1.93 (broad s, 1H), 2.00–2.14 (m, 2H), 3.80 (sextet, J = 6.0 Hz, 1H), 5.12–5.22 (m, 1H); ^{13}C NMR: δ 17.7, 23.4, 24.5, 25.7, 39.2, 67.9, 124.1, 132.0. Anal. Calcd for $\text{C}_8\text{H}_{16}\text{O}$: C, 74.94; H, 12.58%; Found: C, 74.77; H, 12.68%.

(**R**)-**10**: colorless oil; $[\alpha]_{\text{D}}^{24} -6.5$ (c 1.01, CHCl_3), $[\alpha]_{\text{D}}^{25} -6.8$ (c 1.23, EtOH) (lit.^{16c} $[\alpha]_{\text{D}}^{25} +7.7$ (c 0.03, EtOH)); IR: 1736, 1243 cm^{-1} ; ^1H NMR: δ 1.21 (d, J = 6.6 Hz, 3H), 1.43–1.57 (m, 2H), 1.59 (s, 3H), 1.68 (s, 3H), 1.97–2.04 (merged m and s at δ 2.03, 5H), 4.83–4.91 (m, 1H), 5.05–5.11 (m, 1H); ^{13}C NMR: δ 17.6, 21.4, 24.0, 25.7, 29.7, 36.0, 70.7, 123.5, 132.1, 170.8. Anal. Calcd for $\text{C}_{10}\text{H}_{18}\text{O}_2$: C, 70.55; H, 10.66%. Found: C, 70.38; H, 10.36%.

(R)-6-Methyl-5-hepten-2-ol (R)-9. Reduction of (**R**)-**10** (2.70 g, 15.88 mmol) with LiAlH_4 (0.480 g, 12.70 mmol) in Et_2O (60 mL), followed by work up as above, furnished pure (**R**)-**9** (1.9 g, 92%). colorless oil; $[\alpha]_{\text{D}}^{22} -11.7$ (c 1.06, CHCl_3); IR: 3050, 1642 cm^{-1} ; ^1H NMR: δ 1.16 (d, J = 6.0 Hz, 3H), 1.39–1.55 (m, 2H), 1.59 (s, 3H), 1.68 (s, 3H), 1.83 (broad s, 1H), 2.00–2.09 (m, 2H), 3.72–3.83 (m, 1H), 5.03–5.18 (m, 1H); ^{13}C NMR: δ 17.7, 23.5, 24.5, 25.8, 39.2, 67.9, 124.1, 132.0. Anal. Calcd for $\text{C}_8\text{H}_{16}\text{O}$: C, 74.94; H, 12.58%. Found: C, 74.77; H, 12.68%.

(R)-6-tert-Butyldiphenylsilyloxy-2-methyl-2-heptene (R)-11. To a stirred solution of (**R**)-**9** (2.50 g, 19.53 mmol), imidazole (1.73 g, 25.39 mmol), and DMAP (catalytic) in CH_2Cl_2 (20 mL) was dropwise added TBDPSCI (6.98 g, 25.39 mmol). After stirring the mixture for 7 h at room temperature, it was poured into ice-cold water (20 mL), the organic layer was separated, and the aqueous portion was extracted with CHCl_3 (3 \times 10 mL). The combined organic extracts were successively washed with H_2O (2 \times 10 mL) and brine (1 \times 5 mL) and dried. Removal of solvent in vacuo, followed by column chromatography (silica gel, 0–5% EtOAc /hexane) of the residue, afforded pure (**R**)-**11** (6.3 g, 88%). colorless oil; $[\alpha]_{\text{D}}^{24} +11.7$ (c 1.04, CHCl_3); IR: 3050, 3013 cm^{-1} ; ^1H NMR: δ 1.05 (merged s and d, J = 6.0 Hz, 12H), 1.18–1.26 (m, 2H), 1.53 (s, 3H), 1.63 (s, 3H), 1.88–2.01 (m, 2H), 3.79–3.87 (m, 1H), 4.95–5.00 (m, 1H), 7.32–7.44 (m, 6H), 7.65–7.70 (m, 4H); ^{13}C NMR: δ 17.6, 18.4, 19.3, 23.2, 24.0, 25.7, 26.8, 26.9, 39.6, 69.4, 124.5, 127.4, 127.5, 127.6, 129.4, 129.5, 131.2, 134.3, 134.6, 135.0, 135.6, 135.9. Anal. Calcd for $\text{C}_{24}\text{H}_{34}\text{OSi}$: C, 78.63; H, 9.35%. Found: C, 78.68; H, 9.51%.

(S)-6-tert-Butyldiphenylsilyloxy-2-methyl-2-heptene (S)-11. As above, silylation of (**S**)-**9** (2.20 g, 17.19 mmol) using TBDPSCI (6.12 g, 22.34 mmol), imidazole (1.52 g, 22.34 mmol), and DMAP (catalytic) in CH_2Cl_2 (20 mL) furnished (**S**)-**11** (5.7 g, 91%). colorless oil; $[\alpha]_{\text{D}}^{25} -11.8$ (c 1.03, CHCl_3); IR: 3071, 3049, 998 cm^{-1} ; ^1H NMR: δ 1.03 (merged s and d, J = 6.2 Hz, 12H), 1.40–1.45 (m, 1H), 1.50–1.55 (m containing a s at δ 1.54, 4H), 1.63 (s, 3H), 1.89–2.01 (m, 2H), 3.84–3.86 (m, 1H), 4.96–4.99 (m, 1H), 7.35–7.42 (m, 6H), 7.67–7.69 (m, 4H); ^{13}C NMR: δ 17.7, 18.5, 19.4, 23.2, 24.1, 25.8, 26.9, 27.1, 39.7, 69.4, 124.5, 127.5, 127.7, 129.5, 129.6, 131.3, 134.7, 135.0, 135.6, 136.0. Anal. Calcd for $\text{C}_{24}\text{H}_{34}\text{OSi}$: C, 78.63; H, 9.35%. Found: C, 78.31; H, 9.23%.

(R)-4-tert-Butyldiphenylsilyloxypentanal (R)-12. Ozone was bubbled through a cooled (-78 °C) solution of (**R**)-**11** (4.84 g, 13.22 mmol) in CH_2Cl_2 (20 mL) for 1 h. After 0.5 h, the excess O_3 was

removed by purging with N_2 , Ph_3P (5.20 g, 19.84 mmol) was added, and the mixture was stirred for 18 h at room temperature and concentrated in vacuo. The residue was taken in hexane (30 mL), concentrated in vacuo, and purified by column chromatography (silica gel, 0–10% Et_2O /hexane) to obtain pure (**R**)-**12** (3.6 g, 81%). colorless oil; $[\alpha]_{\text{D}}^{22} +2.2$ (c 1.01, CHCl_3); IR: 3070, 2717, 1726 cm^{-1} ; ^1H NMR: δ 1.06 (merged s and d, J = 6.3 Hz, 12H), 1.70–1.83 (m, 2H), 2.44–2.51 (m, 2H), 3.91–3.95 (m, 1H), 7.37–7.43 (m, 6H), 7.66–7.68 (m, 4H), 9.68 (s, 1H); ^{13}C NMR: δ 19.3, 23.0, 27.0, 27.1, 31.3, 39.6, 68.5, 127.5, 127.7, 129.6, 129.7, 134.1, 134.5, 135.8, 135.9, 202.4. Anal. Calcd for $\text{C}_{21}\text{H}_{28}\text{O}_2\text{Si}$: C, 74.07; H, 8.29%. Found: C, 74.18; H, 8.51%.

(S)-4-tert-Butyldiphenylsilyloxypentanal (S)-12. Reductive ozonolysis of (**S**)-**11** (5.01 g, 13.69 mmol) using O_3 and Ph_3P (5.38 g, 20.53 mmol) in CH_2Cl_2 (20 mL), followed by isolation as above, furnished pure (**S**)-**12** (3.8 g, 82%). colorless oil; $[\alpha]_{\text{D}}^{22} -1.9$ (c 1.19, CHCl_3); IR: 1727, 2717, 3050 cm^{-1} ; ^1H NMR: δ 1.07 (merged s and d, J = 6.1 Hz, 12H), 1.71–1.83 (m, 2H), 2.47–2.49 (m, 2H), 3.92–3.96 (m, 1H), 7.37–7.43 (m, 6H), 7.66–7.68 (m, 4H), 9.68 (s, 1H); ^{13}C NMR: δ 19.3, 23.0, 27.0, 27.1, 31.3, 39.6, 68.5, 127.4, 127.5, 127.6, 127.7, 129.6, 129.7, 134.1, 134.5, 135.8, 135.9, 136.0, 202.4. Anal. Calcd for $\text{C}_{21}\text{H}_{28}\text{O}_2\text{Si}$: C, 74.07; H, 8.29%. Found: C, 74.18; H, 8.51%.

(3R,6R)-6-tert-Butyldiphenylsilyloxyhept-1-en-3-ol (3R,6R)-13. To a cooled (-78 °C) and stirred solution of (**R**)-**12** (3.14 g, 9.24 mmol) in THF (30 mL) was added vinylmagnesium bromide (18.5 mL, 18.5 mmol, 1 M in THF). After stirring for 3 h, the mixture was treated with aqueous saturated NH_4Cl (10 mL), the organic was layer separated, and the aqueous portion was extracted with Et_2O (2 \times 20 mL). The combined organic extracts were washed with H_2O (2 \times 10 mL) and brine (1 \times 5 mL), dried, and concentrated in vacuo. The residue was purified by column chromatography (silica gel, 0–10% Et_2O /hexane) to afford pure (3R,6R)-**13** (2.8 g, 84%). colorless liquid; $[\alpha]_{\text{D}}^{23} +36.1$ (c 1.22, CHCl_3); IR: 3365, 3070, 3050, 1644 cm^{-1} ; ^1H NMR: δ 1.04 (merged s and d, J = 6.2 Hz, 12H), 1.45–1.69 (m, 5H), 3.85–3.87 (m, 1H), 3.98–4.00 (m, 1H), 5.01–5.20 (m, 2H), 5.69–5.86 (m, 1H), 7.30–7.40 (m, 6H), 7.63–7.68 (m, 4H); ^{13}C NMR: δ 19.3, 23.1, 27.1, 29.7, 32.3, 32.6, 34.7, 35.1, 69.3, 69.5, 73.0, 73.2, 114.5, 127.5, 127.6, 129.5, 129.6, 134.4, 134.7, 135.0, 136.4, 141.2. Anal. Calcd for $\text{C}_{23}\text{H}_{32}\text{O}_2\text{Si}$: C, 74.95; H, 8.75%. Found: C, 74.78; H, 8.84%.

(3R,6S)-6-tert-Butyldiphenylsilyloxyhept-1-en-3-ol (3R,6S)-13. As above, reaction of (**S**)-**12** (2.30 g, 6.76 mmol) with vinylmagnesium bromide (13.6 mL, 13.6 mmol, 1 M in THF) in THF (20 mL) and usual purification afforded pure (3R,6S)-**13** (2.1 g, 84%). colorless liquid; $[\alpha]_{\text{D}}^{24} -12.1$ (c 1.16, CHCl_3); IR: 3364, 3071, 997 cm^{-1} ; ^1H NMR: δ 1.04 (merged s and d, J = 6.2 Hz, 12H), 1.45–1.62 (m containing a s at δ 1.58, 5H), 3.85–3.87 (m, 1H), 3.96–4.02 (m, 1H), 5.01–5.20 (m, 2H), 5.69–5.86 (m, 1H), 7.33–7.40 (m, 6H), 7.63–7.67 (m, 4H); ^{13}C NMR: δ 19.2, 23.0, 27.0, 32.2, 32.5, 34.6, 35.0, 69.3, 69.4, 73.0, 73.2, 114.5, 127.4, 127.5, 129.4, 129.5, 135.9, 141.1. Anal. Calcd for $\text{C}_{23}\text{H}_{32}\text{O}_2\text{Si}$: C, 74.95; H, 8.75%. Found: C, 75.18; H, 8.51%.

(3S,6R)-3-Acetoxy-6-tert-butyldiphenylsilyloxyhept-1-ene (3S,6R)-14. Acetylation of (3R,6R)-**13** (4.16 g, 11.30 mmol) with vinyl acetate (5.0 mL) and Novozym 435 (0.50 g, ~50 mg/mmol) was carried out as described above to obtain pure (3R,6R)-**13** (2.0 g, 48%) and (3S,6R)-**14** (2.1 g, 46%). (3R,6R)-**13**: colorless oil; $[\alpha]_{\text{D}}^{26} +10.4$ (c 1.14, CHCl_3); ^1H NMR: δ 1.06 (merged s and d, J = 6.0 Hz, 12H), 1.47–1.55 (m, 5H), 3.88–3.91 (m, 1H), 3.98–4.00 (m, 1H), 5.01–5.18 (m, 2H), 5.76–5.82 (m, 1H), 7.35–7.43 (m, 6H), 7.67–7.69 (m, 4H); ^{13}C NMR: δ 19.2, 23.0, 27.0, 32.6, 35.1, 69.4, 73.2, 114.5, 127.4, 127.5, 129.5, 129.6, 134.3, 134.6, 135.9, 141.1. Anal. Calcd for $\text{C}_{23}\text{H}_{32}\text{O}_2\text{Si}$: C, 74.95; H, 8.75%. Found: C, 75.15; H, 8.92%.

(3S,6R)-**14**: colorless oil; $[\alpha]_{\text{D}}^{26} +13.1$ (c 1.03, CHCl_3); IR: 3070, 1739, 1647 cm^{-1} ; ^1H NMR: δ 1.04 (merged s and d, J = 7.2 Hz, 12H), 1.35–1.52 (m, 2H), 1.54–1.70 (m, 2H), 2.01 (s, 3H), 3.81–3.90 (m, 1H), 5.09–5.20 (m, 3H), 5.61–5.78 (m, 1H), 7.31–7.42 (m, 6H), 7.63–7.68 (m, 4H); ^{13}C NMR: δ 19.3, 21.2, 23.1, 27.0, 27.1, 29.6, 34.5, 69.0, 74.7, 116.6, 127.5, 129.5, 129.6, 134.4, 134.7, 135.8, 135.9,

136.0, 136.5, 170.3. Anal. Calcd for $C_{25}H_{34}O_3Si$: C, 73.13; H, 8.35%. Found: C, 73.15; H, 8.32%.

(3S,6S)-3-Acetoxy-6-tert-butylidiphenylsilyloxyhept-1-ene (3S,6S)-14. Following the same procedure (3R,6S)-13 (1.90 g, 5.16 mmol) was acetylated in vinyl acetate (5.0 mL) using Novozym 435 (0.25 g, ~50 mg/mmol) to obtain pure (3R,6S)-13 (0.860 g, 45%) and (3S,6S)-14 (1.0 g, 47%). (3R,6S)-13: colorless oil; $[\alpha]_D^{25}$ -17.7 (c 1.24, $CHCl_3$); IR: 3360, 3065, 998 cm^{-1} ; 1H NMR: δ 1.02 (merged s and d, $J = 6.0$ Hz, 12 H), 1.45–1.59 (m, 5H), 3.84–3.93 (m, 1H), 3.96–4.02 (m, 1H), 5.04–5.21 (m, 2H), 5.71–5.87 (m, 1H), 7.30–7.41 (m, 6H), 7.64–7.68 (m, 4H); ^{13}C NMR: δ 19.2, 23.0, 27.0, 32.3, 34.6, 69.3, 73.0, 114.6, 127.4, 127.5, 129.5, 134.4, 134.7, 135.9, 141.1. Anal. Calcd for $C_{23}H_{32}O_2Si$: C, 74.95; H, 8.75%. Found: C, 75.18; H, 8.51%.

(3S,6S)-14: colorless oil; $[\alpha]_D^{24}$ -13.5 (c 1.05, $CHCl_3$); IR: 1736, 1231 cm^{-1} ; 1H NMR: δ 1.04 (merged s and d, $J = 6.0$ Hz, 12H), 1.35–1.50 (m, 2H), 1.52–1.68 (m, 2H), 2.02 (s, 3H), 3.79–3.88 (m, 1H), 5.10–5.20 (m, 3H), 5.60–5.77 (m, 1H), 7.32–7.42 (m, 6H), 7.63–7.69 (m, 4H); ^{13}C NMR: δ 19.2, 21.2, 23.1, 27.0, 29.7, 34.5, 69.1, 74.8, 116.6, 127.4, 127.5, 129.4, 129.5, 134.3, 134.7, 135.8, 136.4, 170.3. Anal. Calcd for $C_{25}H_{34}O_3Si$: C, 73.13; H, 8.35%. Found: C, 73.32; H, 8.04%.

(2S)-6-para-Methoxybenzyloxy-2-methylhept-2-ene 15. To a cooled ($-5^\circ C$) and stirred suspension of hexane-washed NaH (4.52 g, 94.14 mmol, 50% suspension in oil) in DMF (10 mL) was added (S)-9 (4.82 g, 37.66 mmol) in DMF (10 mL). After 1 h, PMBCl (6.13 mL, 45.19 mmol) was dropwise added into the mixture, and stirring continued until completion of the reaction (cf. TLC). Water (20 mL) was added to the mixture, which was extracted with Et_2O (3×15 mL). The combined organic extracts were washed with H_2O (2×10 mL) and brine (1×5 mL), dried, and concentrated in vacuo to get a residue, which, on column chromatography (silica gel, 0–5% Et_2O /hexane), afforded pure (S)-15 (7.6 g, 82%). colorless liquid; $[\alpha]_D^{23}$ $+29.9$ (c 1.08, $CHCl_3$); IR: 3090, 1613 cm^{-1} ; 1H NMR: δ 1.18 (d, $J = 7.0$ Hz, 3H), 1.43–1.44 (m, 2H), 1.60 (s, 3H), 1.68 (s, 3H), 2.04–2.06 (m, 2H), 3.47–3.50 (m, 1H), 3.79 (s, 3H), 4.38 (d, $J = 14.0$ Hz, 1H), 4.49 (d, $J = 14.0$ Hz, 1H), 5.09 (t, $J = 2.4$ Hz, 1H), 6.86 (d, $J = 7.0$ Hz, 2H), 7.26 (d, $J = 7.0$ Hz, 2H); ^{13}C NMR: δ 17.6, 19.6, 24.1, 25.7, 36.7, 55.2, 69.9, 74.1, 113.7, 124.3, 129.1, 131.2, 131.5, 159.0. Anal. Calcd for $C_{16}H_{24}O_2$: C, 77.38; H, 9.74%. Found: C, 77.33; H, 9.61%.

(4S)-4-para-Methoxybenzyloxypentanal 16. To a stirred solution of (S)-15 (4.46 g, 17.98 mmol) in acetone:water (8:1, 30 mL) was added a mixture of NMO (4.20 g, 35.96 mmol) and OsO_4 (0.230 g, 0.89 mmol) in t -BuOH (2 mL). After stirring for 24 h, the mixture was treated with aqueous saturated Na_2SO_3 , stirred for 0.5 h, and subsequently extracted with $EtOAc$ (3×10 mL). The combined organic extracts were washed with H_2O (2×10 mL) and brine (1×10 mL), dried, and concentrated in vacuo to get a residue, which, on column chromatography (0–30% $EtOAc$ /hexane), furnished pure diol (4.8 g, 95%). colorless oil. $[\alpha]_D^{24}$ $+29.2$ (c 1.03, $CHCl_3$); IR: 3418, 1341, 1138 cm^{-1} ; 1H NMR: δ 1.12–1.26 (merged s and d, $J = 6.4$ Hz, 9H), 1.56–1.81 (m, 4H), 2.56 (broad s, 2H), 3.34 (t, $J = 7.4$ Hz, 1H), 3.45–3.61 (m, 1H), 3.78 (s, 3H), 4.37 (d, $J = 11.4$ Hz, 1H), 4.54 (d, $J = 11.4$ Hz, 1H), 6.87 (d, $J = 8.7$ Hz, 2H), 7.26 (d, $J = 8.7$ Hz, 2H); ^{13}C NMR: 19.4, 19.5, 23.3, 23.4, 26.4, 26.5, 27.5, 27.6, 33.7, 33.8, 55.3, 70.1, 72.9, 73.0, 74.5, 74.7, 78.4, 113.8, 129.4, 129.5, 130.5, 130.7, 159.2. Anal. Calcd for $C_{16}H_{26}O_4$: C, 68.06; H, 9.28%. Found: C, 67.88; H, 9.61%.

To a cooled ($10^\circ C$) and stirred solution of the above diol (4.08 g, 14.47 mmol) in aqueous 60% CH_3CN (20 mL) was added $NaIO_4$ (6.19 g, 28.92 mmol) in portions. After stirring for 2 h, the mixture was filtered and the filtrate was treated with aqueous 10% $NaHSO_3$ and extracted with $CHCl_3$ (3×15 mL). The organic layer was washed with H_2O (2×10 mL) and brine (1×5 mL), dried, and concentrated in vacuo to get a residue, which, on column chromatography (silica gel, 0–10% $EtOAc$ /hexane), furnished pure (S)-16 (3.2 g, quant.). colorless liquid; $[\alpha]_D^{24}$ $+45.5$ (c 1.03, $CHCl_3$); IR: 2724, 1723 cm^{-1} ; 1H NMR: δ 1.20 (d, $J = 6.0$ Hz, 3H), 1.79–1.86 (m, 2H), 2.46–2.54 (m, 2H), 3.50–3.56 (m, 1H), 3.80 (s, 3H), 4.36 (d, $J = 11.4$ Hz, 1H), 4.55 (d, $J = 11.4$ Hz, 1H), 6.88 (d, $J = 8.4$ Hz, 2H), 7.28 (d, $J = 8.4$ Hz,

2H), 9.74 (t, $J = 1.5$ Hz, 1H); ^{13}C NMR: δ 19.6, 29.3, 40.3, 55.4, 70.2, 73.5, 113.9, 129.4, 130.7, 159.2, 202.7. Anal. Calcd for $C_{13}H_{18}O_3$: C, 70.24; H, 8.16%. Found: C, 70.05; H, 8.08%.

(3R,6S)-6-para-Methoxybenzyloxyhept-1-en-3-ol 17. As described earlier, reaction of (S)-16 (2.28 g, 10.27 mmol) with vinylmagnesium bromide (15.40 mL, 15.40 mmol, 1 M in THF) (30 mL), followed by the usual isolation and purification, furnished (3R,6S)-17 (1.9 g, 73%). colorless liquid; $[\alpha]_D^{24}$ $+16.3$ (c 1.02, $CHCl_3$); IR: 3410 cm^{-1} ; 1H NMR: δ 1.15 (d, $J = 6.3$ Hz, 3H), 1.47–1.66 (m, 4H), 2.50 (broad s, 1H), 3.44–3.52 (m, 1H), 3.74 (s, 3H), 3.98–4.05 (m, 1H), 4.33 (d, $J = 11.4$ Hz, 1H), 4.47 (d, $J = 11.4$ Hz, 1H), 5.01–5.18 (m, 2H), 5.74–5.86 (m, 1H), 6.82 (d, $J = 8.4$ Hz, 2H), 7.22 (d, $J = 8.4$ Hz, 2H); ^{13}C NMR: δ 19.5, 32.4, 33.0, 55.3, 70.0, 72.9, 74.3, 74.5, 113.8, 114.4, 129.4, 130.8, 141.3, 159.1. Anal. Calcd for $C_{15}H_{22}O_3$: C, 71.97; H, 8.86%. Found: C, 72.21; H, 8.88%.

(3S,6S)-3-Acetoxy-6-para-methoxybenzyloxyhept-1-ene 18. As described earlier, acetylation of (3R,6S)-17 (1.80 g, 7.20 mmol) with vinyl acetate (1.0 mL, 10.8 mmol) in the presence of Novozym 435 (0.540 mg, ~75 mg/mmol) in diisopropyl ether (25 mL), followed by isolation, furnished (3R,6S)-17 (0.756 g, 42%) and 18 (0.946 g, 45%). (3R,6S)-17: colorless liquid; $[\alpha]_D^{23}$ $+20.7$ (c 1.00, $CHCl_3$); IR: 3410 cm^{-1} ; 1H NMR: δ 1.19 (d, $J = 6.3$ Hz, 3H), 1.54–1.72 (m, 4H), 2.43 (broad s, 1H), 3.49–3.59 (m, 1H), 3.79 (s, 3H), 4.03–4.11 (m, 1H), 4.38 (d, $J = 11.4$ Hz, 1H), 4.52 (d, $J = 11.4$ Hz, 1H), 5.07–5.26 (m, 2H), 5.80–5.91 (m, 1H), 6.87 (d, $J = 8.7$ Hz, 2H), 7.26 (d, $J = 8.7$ Hz, 2H); ^{13}C NMR: δ 19.5, 32.4, 33.0, 55.3, 70.1, 72.9, 74.5, 113.8, 114.5, 129.4, 130.8, 141.2, 159.1. Anal. Calcd for $C_{15}H_{22}O_3$: C, 71.97; H, 8.86%. Found: C, 72.15; H, 8.75%.

18: colorless liquid; $[\alpha]_D^{24}$ $+12.4$ (c 1.01, $CHCl_3$); IR: 1736, 1646, 990 cm^{-1} ; 1H NMR: δ 1.18 (d, $J = 6.3$ Hz, 3H), 1.39–1.80 (m, 4H), 2.03 (s, 3H), 3.46–3.56 (m, 1H), 3.80 (s, 3H), 4.36 (d, $J = 11.4$ Hz, 1H), 4.49 (d, $J = 11.4$ Hz, 1H), 5.13–5.25 (m, 3H), 5.70–5.81 (m, 1H), 6.87 (d, $J = 8.4$ Hz, 2H), 7.26 (d, $J = 8.4$ Hz, 2H); ^{13}C NMR: δ 19.6, 21.3, 30.1, 32.0, 55.3, 70.0, 73.9, 74.7, 113.8, 116.7, 129.2, 131.0, 136.5, 159.1, 170.4. Anal. Calcd for $C_{17}H_{24}O_4$: C, 69.84; H, 8.27%. Found: C, 69.88; H, 8.33%.

(3S,6R)-6-tert-Butyldiphenylsilyloxyhept-1-en-3-ol (3S,6R)-13. A mixture of (3S,6R)-14 (3.20 g, 7.80 mmol) and K_2CO_3 (1.30 g, 9.36 mmol) in MeOH (20 mL) was magnetically stirred until completion of the reaction (cf. TLC, ~6 h). After solvent removal in vacuo, water (25 mL) was added to the residue, followed by extraction with $EtOAc$ (3×20 mL). The combined organic extracts were washed with H_2O (2×10 mL) and brine (1×5 mL), dried, and concentrated in vacuo to get a residue, which, on column chromatography (silica gel, 0–15% $EtOAc$ /hexane), furnished pure (3S,6R)-13 (2.7 g, 92%). colorless liquid; $[\alpha]_D^{24}$ $+15.0$ (c 1.12, $CHCl_3$); IR: 3420, 3050, 997 cm^{-1} ; 1H NMR: δ 1.07 (merged s and d, $J = 6.3$ Hz, 12H), 1.42–1.60 (m, 5H), 3.85–3.92 (m, 1H), 3.96–4.02 (m, 1H), 5.07–5.19 (m, 2H), 5.78–5.83 (m, 1H), 7.35–7.43 (m, 6H), 7.67–7.69 (m, 4H); ^{13}C NMR: δ 19.3, 23.1, 27.0, 27.2, 32.3, 34.6, 69.3, 73.0, 114.6, 127.5, 127.6, 129.5, 129.6, 134.4, 134.8, 135.9, 141.2. Anal. Calcd for $C_{23}H_{32}O_3Si$: C, 74.95; H, 8.75%. Found: C, 74.77; H, 8.43%.

(3E,5S,8R)-8-tert-Butyldiphenylsilyloxy-5-hydroxynon-3-en-2-one 19. A mixture of (3S,6R)-13 (1.0 g, 2.72 mmol), MVK (1.33 g, 19.02 mmol), and Grubbs' II catalyst (5 mol %) in CH_2Cl_2 (5 mL) was refluxed for 22 h. After concentrating the mixture in vacuo, the residue was subjected to column chromatography (silica gel, 0–15% $EtOAc$ /hexane) to give pure 19 (0.936 g, 84%). colorless oil; $[\alpha]_D^{22}$ $+15.7$ (c 0.981, $CHCl_3$); IR: 3414, 1710 cm^{-1} ; 1H NMR: δ 1.07 (merged s and d, $J = 6.8$ Hz, 12H), 1.21–1.35 (m, 4H) 1.61 (broad s, 1H), 2.26 (s, 3H), 3.91–3.97 (m, 1H), 4.18–4.30 (m, 1H), 6.24 (d, $J = 16.0$ Hz, 1H), 6.70 (dd, $J = 5.0, 16.0$ Hz, 1H), 7.34–7.47 (m, 6H), 7.65–7.69 (m, 4H); ^{13}C NMR: δ 19.2, 22.7, 27.0, 27.4, 32.0, 34.9, 69.2, 71.2, 127.5, 127.6, 129.0, 129.6, 129.7, 134.0, 134.2, 135.8, 148.7, 198.5. Anal. Calcd for $C_{25}H_{34}O_3Si$: C, 73.13; H, 8.35%. Found: C, 73.39; H, 8.02%.

Decarestrictine L 1. To a cooled ($0^\circ C$) and stirred solution of 19 (0.700 g, 1.71 mmol) in THF (5 mL) was added Bu_4NF (1.7 mL, 1.71 mmol, 1 M in THF). The reaction mixture was brought to room temperature and stirred until the reaction was complete (cf. TLC, 3 h).

The mixture was poured into ice-cold H₂O (15 mL) and extracted with EtOAc (2 × 10 mL). The organic extract was washed with water (2 × 10 mL) and brine (1 × 5 mL) and dried. Removal of solvent, followed by column chromatography of the residue (silica gel, 0–15% EtOAc/hexane), furnished **1** (0.21 g, 71%). colorless oil; $[\alpha]_D^{25} + 28.4$ (c 1.14, CHCl₃), lit.^{8a} $[\alpha]_D^{25} + 28.8$ (c 0.49, CHCl₃); IR: 3441, 1710 cm⁻¹; ¹H NMR: δ 1.22 (d, *J* = 6.0 Hz, 3H), 1.55–1.58 (m, 1H), 1.69–1.75 (m, 2H), 1.83–1.91 (m, 2H), 2.21 (s, 3H), 2.73 (d, *J* = 6.6 Hz, 2H), 3.40–3.49 (m, 1H), 3.84–3.95 (m, 1H), 4.00 (q, *J* = 6.6 Hz, 1H); ¹³C NMR: δ 21.2, 29.6, 31.0, 32.8, 47.1, 70.3, 73.6, 78.3, 209.0. Anal. Calcd for C₉H₁₆O₃: C, 62.77; H, 9.36. Found: C, 62.59; H, 9.42.

Ethyl (2*E*,4*S*,7*R*)-4-Hydroxy-7-*tert*-butyldiphenylsilyloxyoct-2-enoate 20. Cross-metathesis between (3*S*,6*R*)-**13** (0.730 g, 1.98 mmol) and ethyl acrylate (1.99 g, 19.8 mmol) in the presence of Hoveyda Grubbs' II catalyst (5 mol) in CH₂Cl₂ (15 mL), followed by isolation, furnished **20** (0.870 g, quant.). colorless oil; $[\alpha]_D^{28} + 28.9$ (c 1.02, CHCl₃); IR: 3453, 1720, 1656, 984 cm⁻¹; ¹H NMR: δ 1.04 (merged s and d, *J* = 6.0 Hz, 12H), 1.29 (t, *J* = 7.2 Hz, 3H), 1.59–1.67 (m, 5H), 3.85–3.94 (m, 1H), 4.14–4.25 (merged m and q at δ 4.20, *J* = 7.2 Hz, 3H), 5.99 (dd, *J* = 1.6, 15.6 Hz, 1H), 6.88 (dd, *J* = 4.6, 15.6 Hz, 1H), 7.35–7.42 (m, 6H), 7.63–7.68 (m, 4H); ¹³C NMR: δ 14.2, 19.2, 22.9, 27.0, 31.6, 34.2, 60.4, 69.1, 70.8, 120.2, 127.4, 127.6, 129.5, 129.6, 134.0, 134.4, 135.8, 135.9, 150.1, 166.5. Anal. Calcd for C₂₆H₃₆O₄Si: C, 70.87; H, 8.23%. Found: C, 70.65; H, 8.36%.

Ethyl (2*E*,4*S*,7*R*)-4-*tert*-Butyldimethylsilyloxy-7-*tert*-butyldiphenylsilyloxyoct-2-enoate 21. As described for **11**, silylation of **20** (0.870 g, 1.98 mmol) with TBSCl (0.445 g, 2.97 mmol), imidazole (0.202 g, 2.97 mmol), and DMAP (catalytic) in CH₂Cl₂ (15 mL) furnished **21** (1.0 g, 92%) after isolation and purification. colorless oil; $[\alpha]_D^{22} + 25.8$ (c 1.04, CHCl₃); IR: 3071, 3048, 1720, 1656, 997 cm⁻¹; ¹H NMR: δ -0.04 (s, 6H), 0.85 (s, 9H), 1.01 (merged s and d, *J* = 6.0 Hz, 12H), 1.23–1.31 (m containing a t at δ 1.28, *J* = 7.0 Hz, 4H), 1.44–1.50 (m, 3H), 3.77–3.83 (m, 1H), 4.08–4.22 (merged m and q at δ 4.17, *J* = 7.0 Hz, 3H), 5.88 (dd, *J* = 1.8, 15.6 Hz, 1H), 6.82 (dd, *J* = 4.6, 15.6 Hz, 1H), 7.28–7.40 (m, 6H), 7.61–7.65 (m, 4H); ¹³C NMR: δ 14.3, 18.1, 19.2, 23.2, 25.8, 27.0, 32.8, 34.4, 60.3, 69.2, 71.4, 119.8, 127.4, 127.5, 129.4, 129.5, 134.4, 134.8, 135.8, 135.9, 150.9, 166.7. Anal. Calcd for C₃₂H₅₀O₄Si₂: C, 69.26; H, 9.08%. Found: C, 69.47; H, 9.23%.

(2*E*,4*S*,7*R*)-4-*tert*-Butyldimethylsilyloxy-7-*tert*-butyldiphenylsilyloxyoct-2-enoic acid 22. To a stirred solution of **21** (1.01 g, 1.82 mmol) in MeOH (20 mL) was added aqueous 20% NaOH (8 mL). After stirring for 2 h at room temperature, the mixture was concentrated in vacuo. The residue was acidified with aqueous 2 N HCl and extracted with Et₂O (3 × 30 mL). The organic layer was washed with H₂O (2 × 20 mL) and brine (1 × 10 mL). Removal of solvent in vacuo, followed by column chromatography of the residue (silica gel, 0–30% EtOAc/hexane), afforded pure **22** (0.948 g, 99%). colorless oil; $[\alpha]_D^{24} + 27.1$ (c 1.17, CHCl₃); IR: 3500–2500, 1698 cm⁻¹; ¹H NMR: δ 0.01 (s, 6H), 0.90 (s, 9H), 1.06 (merged s and d, *J* = 6.2 Hz, 12H), 1.27–1.30 (m, 2H), 1.51–1.63 (m, 3H), 3.82–3.88 (m, 1H), 4.11–4.19 (m, 1H), 5.96 (dd, *J* = 1.2, 15.6 Hz, 1H), 6.98 (dd, *J* = 4.4, 15.6 Hz, 1H), 7.34–7.42 (m, 6H), 7.66–7.70 (m, 4H); ¹³C NMR: δ 18.1, 19.2, 23.2, 25.8, 27.0, 29.7, 32.7, 34.3, 69.1, 71.2, 119.1, 127.4, 127.5, 129.4, 129.6, 134.3, 134.7, 135.8, 135.9, 153.7, 172.2. Anal. Calcd for C₃₀H₄₆O₄Si₂: C, 68.39; H, 8.80%. Found: C, 68.04; H, 9.19%.

(3*S*,6*S*)-6-*tert*-Butyldiphenylsilyloxyhept-1-en-3-ol (3*S*,6*S*)-13**.** Alkaline hydrolysis of (3*S*,6*S*)-**14** (1.20 g, 2.93 mmol) with K₂CO₃ (0.490 g, 3.55 mmol) in MeOH (15 mL), followed by the usual isolation as above, furnished pure (3*S*,6*S*)-**13** (0.981 g, 91%). colorless liquid; $[\alpha]_D^{24} - 10.1$ (c 1.06, CHCl₃); IR: 3420, 3050, 997 cm⁻¹; ¹H NMR: δ 1.04 (merged s and d, *J* = 5.6 Hz, 12H), 1.50–1.61 (m, 5H), 3.84–3.98 (m, 2H), 5.00–5.19 (m, 2H), 5.69–5.85 (m, 1H), 7.31–7.40 (m, 6H), 7.62–7.68 (m, 4H); ¹³C NMR: δ 19.0, 22.7, 26.8, 32.3, 34.8, 69.1, 73.0, 114.3, 127.2, 127.3, 129.2, 129.3, 134.4, 135.6, 140.9. Anal. Calcd for C₂₃H₃₂O₂Si: C, 74.95; H, 8.75%. Found: C, 74.71; H, 8.83%.

(3*S*,6*S*)-3-Benzoyloxy-6-*tert*-butyldiphenylsilyloxyhept-1-ene 23. To a stirred suspension of hexane-washed NaH (0.14 g, 5.64

mmol, 50% suspension in oil) in THF (10 mL) was added (3*S*,6*S*)-**13** (0.690 g, 1.88 mmol) in THF (5 mL). After refluxing the mixture for 1 h, it was brought to room temperature, BnBr (0.48 g, 2.82 mmol) and Bu₄NI (10 mol %) were added, and the mixture was refluxed until completion of the reaction (cf. TLC, 4 h). It was brought to room temperature and extracted with Et₂O (2 × 20 mL). The organic extract was washed with H₂O (2 × 10 mL) and brine (1 × 3 mL) and dried. Removal of solvent in vacuo, followed by column chromatography (silica gel, 0–5% EtOAc/hexane) of the residue, afforded pure **23** (0.835 g, 97%). colorless oil; $[\alpha]_D^{23} - 25.0$ (c 1.20, CHCl₃); IR: 3069, 1643 cm⁻¹; ¹H NMR: δ 1.03 (merged s and d, *J* = 6.0 Hz, 12H), 1.49–1.57 (m, 4H), 3.56–3.66 (m, 1H), 3.78–3.89 (m, 1H), 4.30 (d, *J* = 11.8 Hz, 1H), 4.54 (d, *J* = 11.8 Hz, 1H), 5.06–5.19 (m, 2H), 5.55–5.73 (m, 1H), 7.25–7.29 (m, 5H), 7.32–7.43 (m, 6H), 7.63–7.68 (m, 4H); ¹³C NMR: δ 19.2, 23.2, 27.0, 31.0, 35.0, 69.5, 69.9, 80.7, 117.1, 127.4, 127.6, 127.7, 128.3, 129.4, 134.4, 134.8, 135.9, 138.8, 138.9. Anal. Calcd for C₃₀H₃₈O₂Si: C, 78.55; H, 8.35%. Found: C, 78.37; H, 8.28%.

(2*S*,5*S*)-5-Benzoyloxyhept-6-en-2-ol 24. Desilylation of **23** (0.742 g, 1.62 mmol) with Bu₄NF (3.23 mL, 3.23 mmol, 1 M in THF) in THF (10 mL) at 0 °C, followed by isolation, as described earlier, furnished **24** (0.271 g, 76%); colorless oil; $[\alpha]_D^{25} - 20.6$ (c 1.06, CHCl₃); IR: 3400, 3065, 3030, 1643 cm⁻¹; ¹H NMR: δ 1.17 (d, *J* = 6.0 Hz, 3H), 1.64–1.66 (m, 5H), 3.72–3.83 (m, 2H), 4.34 (d, *J* = 11.8 Hz, 1H), 4.59 (d, *J* = 11.8 Hz, 1H), 5.17–5.26 (m, 2H), 5.67–5.84 (m, 1H), 7.25–7.32 (m, 5H); ¹³C NMR: δ 23.3, 31.6, 34.9, 67.6, 70.1, 80.5, 117.3, 127.4, 127.7, 128.3, 138.3, 138.6. Anal. Calcd for C₁₄H₂₀O₂: C, 76.33; H, 9.15%. Found: C, 76.37; H, 9.51%.

(9*E*,3*S*,6*R*,11*S*,14*R*)-3-Benzoyloxy-11-*tert*-butyldimethylsilyloxy-14-*tert*-butyldiphenylsilyloxy-6-methyl-7-oxa-8-oxopenta-1,9-diene 25. To a stirred solution of **24** (0.270 g, 1.23 mmol) in THF (10 mL) were added PPh₃ (0.48 g, 1.85 mmol) and **22** (0.775 g, 1.47 mmol). The reaction mixture was cooled to 0 °C, DIAD (0.36 g, 1.85 mmol) was added, and the mixture was stirred for 18 h. The mixture was extracted with EtOAc (2 × 15 mL), and the organic extract was washed with H₂O (2 × 10 mL) and brine (1 × 5 mL) and dried. After concentrating in vacuo, the residue was subjected to column chromatography (silica gel, 0–20% EtOAc/hexane) to give pure **25** (0.592 g, 66%). colorless oil; $[\alpha]_D^{23} + 7.7$ (c 1.06, CHCl₃); IR: 1715, 1657, 995 cm⁻¹; ¹H NMR: δ -0.03 (s, 6H), 0.86 (s, 9H), 1.03 (merged s and d, *J* = 6.0 Hz, 12H), 1.22 (d, *J* = 6.2 Hz, 3H), 1.46–1.57 (m, 8H), 3.72–3.85 (m, 2H), 4.12–4.14 (m, 1H), 4.33 (d, *J* = 11.8 Hz, 1H), 4.58 (d, *J* = 11.8 Hz, 1H), 4.93–4.97 (m, 1H), 5.16–5.25 (m, 2H), 5.63–5.77 (m, 1H), 5.86 (dd, *J* = 1.6, 15.6 Hz, 1H), 6.80 (dd, *J* = 4.6, 15.6 Hz, 1H), 7.31–7.37 (m, 11H), 7.63–7.67 (m, 4H); ¹³C NMR: δ 18.0, 19.1, 19.9, 23.2, 25.8, 27.0, 31.1, 31.5, 32.8, 34.4, 69.1, 70.0, 70.4, 71.3, 79.9, 117.3, 120.1, 127.3, 127.4, 127.6, 128.2, 129.3, 129.4, 134.2, 134.6, 135.7, 138.5, 138.7, 150.5, 166.1. Anal. Calcd for C₄₄H₆₄O₅Si₂: C, 72.48; H, 8.85%. Found: C, 72.52; H, 8.70%.

(6*E*,2*R*,5*S*,10*R*,13*S*)-13-Benzoyloxy-10-methyl-9-oxa-8-oxopenta-6,14-diene-2,5-diol 26. A mixture of **25** (0.583 g, 0.80 mmol) in CH₃CN (10 mL) and aqueous HF (0.4 mL) in a Teflon vessel was stirred at room temperature for 16 h. The mixture was concentrated in vacuo, and the residue was extracted with EtOAc (3 × 20 mL). The organic layer was washed with H₂O (2 × 10 mL) and brine (1 × 5 mL) and concentrated in vacuo, and the residue was column chromatographed (silica gel, 0–40% EtOAc/hexane) to afford pure **26** (0.210 g, 70%). colorless oil; $[\alpha]_D^{23} - 24.0$ (c 1.05, CHCl₃); IR: 3417, 3087, 3030, 1714, 990 cm⁻¹; ¹H NMR: δ 1.17 (d, *J* = 6.0 Hz, 3H), 1.20 (d, *J* = 6.0 Hz, 3H), 1.49–1.75 (m, 8H), 3.14 (broad s, 2H), 3.70–3.82 (m, 2H), 4.29–4.35 (merged m and d, *J* = 11.8 Hz, 2H), 4.57 (d, *J* = 11.8 Hz, 1H), 4.92–4.96 (m, 1H), 5.16–5.25 (m, 2H), 5.62–5.79 (m, 1H), 5.99 (dd, *J* = 1.4, 15.6 Hz, 1H), 6.89 (dd, *J* = 4.8, 15.6 Hz, 1H), 7.26–7.33 (m, 5H); ¹³C NMR: δ 19.9, 23.6, 31.1, 31.6, 33.2, 35.1, 68.1, 70.0, 70.8, 71.0, 80.0, 117.5, 120.4, 127.4, 127.7, 128.3, 138.5, 138.6, 150.0, 166.3. Anal. Calcd for C₂₂H₃₂O₅: C, 70.18; H, 8.57%. Found: C, 69.84; H, 8.53%.

(6*E*,2*R*,5*S*,10*R*,13*S*)-2-Acryloxy-13-benzoyloxy-10-methyl-9-oxa-8-oxopenta-6,14-dien-5-ol 27. A solution of **26** (0.210 g, 0.56

mmol), ethyl acrylate (0.450 g, 4.48 mmol), and Novozym 435 (0.50 g, ~890 mg/mmol) in diisopropyl ether (5 mL) was agitated on an orbital shaker at 120 rpm for 30 h. The reaction mixture was filtered, the filtrate was concentrated in vacuo, and the residue was column chromatographed (silica gel, 0–25% EtOAc/hexane) to furnish pure **27** (0.171 g, 71%). colorless oil; $[\alpha]_D^{25}$ –7.1 (c 1.13, CHCl₃); IR: 3479, 3065, 3030, 1728, 1714, 1657, 985 cm⁻¹; ¹H NMR: δ 1.22 (d, *J* = 6.2 Hz, 3H), 1.25 (d, *J* = 6.4 Hz, 3H), 1.54–1.80 (m, 8H), 3.51 (broad s, 1H), 3.68–3.79 (m, 1H), 4.29–4.35 (merged m and d, *J* = 11.8 Hz, 2H), 4.58 (d, *J* = 11.8 Hz, 1H), 4.88–5.06 (m, 2H), 5.16–5.25 (m, 2H), 5.66–5.83 (m, 2H), 5.95–6.15 (m, 2H), 6.38 (dd, *J* = 1.6, 17.2 Hz, 1H), 6.88 (dd, *J* = 5.0, 15.8 Hz, 1H), 7.29–7.33 (m, 5H); ¹³C NMR: δ 20.0, 31.1, 31.4, 31.6, 32.0, 70.0, 70.5, 70.7, 70.9, 80.0, 117.5, 120.8, 127.4, 127.7, 128.3, 128.7, 130.6, 138.5, 138.6, 149.5, 165.9, 166.0. Anal. Calcd for C₂₃H₃₄O₆: C, 69.74; H, 7.96%. Found: C, 69.44; H, 7.94%.

(5S,8R,13S,16R)-Pyrenophorol Monobenzyl Ether 28. A stirred solution of **27** (0.142 g, 0.33 mmol) and Grubbs' II catalyst (20 mol %) in degassed CH₂Cl₂ (10 mL) was refluxed for 72 h. The reaction mixture was concentrated in vacuo, and the residue was subjected to preparative TLC to obtain pure **28** (57 mg, 61% based on conversion) and recovered **27** (40 mg). colorless oil; $[\alpha]_D^{25}$ –39.0 (c 1.07, CHCl₃); IR: 3470, 1713, 1643 cm⁻¹; ¹H NMR: δ 1.24 (t, *J* = 6.2 Hz, 6H), 1.60–1.86 (m, 9H), 3.78–3.88 (m, 1H), 4.13–4.22 (m, 1H), 4.37 (d, *J* = 12.0 Hz, 1H), 4.55 (d, *J* = 12.0 Hz, 1H), 5.03–5.17 (m, 2H), 5.91 (d, *J* = 16.0 Hz, 2H), 6.74 (dd, *J* = 7.8, 16.0 Hz, 1H), 6.82 (dd, *J* = 6.8, 15.6 Hz, 1H), 7.32 (s, 5H); ¹³C NMR: δ 13.9, 18.4, 29.7, 30.9, 33.4, 43.7, 69.8, 70.7, 77.2, 79.2, 83.4, 124.5, 127.7, 128.6, 146.9, 178.1. Anal. Calcd for C₂₃H₃₀O₆: C, 68.64; H, 7.51%. Found: C, 68.56; H, 7.78%.

(5S,8R,13S,16R)-Pyrenophorol 2. A mixture of **28** (40 mg, 0.1 mmol) and TiCl₄ (11.0 μ L, 0.1 mmol) in CH₂Cl₂ (5 mL) was stirred at room temperature until the completion of the reaction (cf. TLC, 30 min). After concentration in vacuo, the residue was purified by preparative thin layer chromatography to get pure **(5S,8R,13S,16R)-2** (23 mg, 74%). sticky solid; $[\alpha]_D^{25}$ –2.9 (c 1.00, acetone), (lit.^{9f} $[\alpha]_D^{25}$ –3.2 (c 0.25, acetone)); IR: 3440, 1714, 1651, 985 cm⁻¹; ¹H NMR: δ 1.26 (d, *J* = 6.6 Hz, 6H), 1.64–1.93 (m, 8H), 2.30 (broad s, 2H), 4.26–4.32 (m, 2H), 5.10–5.17 (m, 2H), 5.98 (dd, *J* = 1.6, 15.6 Hz, 2H), 6.90 (dd, *J* = 5.2, 15.6 Hz, 2H); ¹³C NMR: δ 18.2, 28.9, 30.5, 69.8, 70.4, 122.2, 149.5, 165.0. Anal. Calcd for C₁₆H₂₄O₆: C, 61.52; H, 7.74%. Found: C, 61.45; H, 7.92%.

(2E,4R,7S)-4-Hydroxy-7-*para*-methoxybenzyloxyoct-2-enal 29. A mixture of (3R,6S)-**17** (0.690 g, 2.76 mmol), acrolein (0.92 mL, 13.80 mmol), and Hoveyda Grubbs' II catalyst (86 mg, 0.14 mmol) in CH₂Cl₂ (20 mL) was stirred for 22 h at room temperature. After concentrating the mixture in vacuo, the residue was subjected to column chromatography (silica gel, 0–25% EtOAc/hexane) to give pure **29** (0.682 g, 89%). colorless liquid; $[\alpha]_D^{26}$ +16.6 (c 1.01, CHCl₃); IR: 3426, 2729, 1688 cm⁻¹; ¹H NMR: δ 1.23 (d, *J* = 6.0 Hz, 3H), 1.66–1.83 (m, 5H), 3.44–3.61 (m, 2H), 3.82 (s, 3H), 4.36 (d, *J* = 11.2 Hz, 1H), 4.56 (d, *J* = 11.2 Hz, 1H), 6.25–6.37 (m, 1H), 6.80 (dd, *J* = 4.2, 15.6 Hz, 1H), 6.86–6.93 (m, 2H), 7.25–7.31 (m, 2H), 9.57 (d, *J* = 7.8 Hz, 1H); ¹³C NMR: δ 19.2, 19.7, 32.4, 55.3, 70.3, 70.7, 74.4, 113.8, 113.9, 129.3, 129.5, 130.1, 130.8, 159.4, 193.7. Anal. Calcd for C₁₆H₂₂O₄: C, 69.04; H, 7.97%. Found: C, 68.93; H, 7.89%.

(2E,4R,7S)-4-*tert*-Butyldimethylsilyloxy-7-*para*-methoxybenzyloxyoct-2-enal 30. Silylation of **29** (1.08 g, 3.88 mmol) with TBSCl (0.700 g, 4.66 mmol), imidazole (0.400 g, 5.83 mmol), and DMAP (catalytic) in CH₂Cl₂ (30 mL), followed by the usual isolation, afforded pure **30** (1.23 g, 81%). colorless liquid; $[\alpha]_D^{28}$ –12.7 (c 1.00, CHCl₃); IR: 2722, 1692 cm⁻¹; ¹H NMR: δ 0.06 (s, 3H), 0.10 (s, 3H), 0.95 (s, 9H), 1.22 (d, *J* = 6.2 Hz, 3H), 1.47–1.78 (m, 4H), 3.49–3.55 (m, 1H), 3.84 (s, 3H), 4.36–4.57 (m, 3H), 6.23–6.35 (m, 1H), 6.81 (dd, *J* = 4.4, 15.4 Hz, 1H), 6.87–6.95 (m, 2H), 7.26–7.31 (m, 2H), 9.60 (d, *J* = 7.8 Hz, 1H); ¹³C NMR: δ –4.9, –4.7, 18.1, 19.5, 25.7, 32.9, 33.2, 55.2, 69.9, 71.6, 74.2, 113.7, 129.1, 130.7, 130.9, 159.1, 160.0, 193.6. Anal. Calcd for C₂₂H₃₆O₄Si: C, 67.30; H, 9.24%. Found: C, 67.26; H, 9.41%.

Ethyl (2Z,4E,6R,9S)-6-*tert*-Butyldimethylsilyloxy-9-*para*-methoxybenzyloxydeca-2,4-dienoate 31. To a cooled (0 °C) and stirred suspension of hexane-washed NaH (0.142 mg, 2.95 mmol, 50% suspension in oil) in THF (5 mL) was added ethyl diphenylphosphonoacetate (0.750 g, 2.36 mmol) in THF (5 mL). After 1 h, the solution was cooled to –78 °C, **30** (0.77 g, 1.96 mmol) in THF (5 mL) was injected into it, and stirring continued at –78 °C for 3 h. The mixture was poured into ice–water and extracted with Et₂O (3 \times 10 mL). The ether layer was washed with H₂O (2 \times 10 mL) and brine (1 \times 5 mL), dried, and concentrated in vacuo to get a residue, which, on column chromatography (silica gel, 0–10% EtOAc/hexane), furnished pure **31** (0.620 g, 68%). colorless liquid; $[\alpha]_D^{24}$ –12.9 (c 1.00, CHCl₃); IR: 1716, 1641 cm⁻¹; ¹H NMR: δ 0.04 and 0.1 (two s, 6H), 0.90 (s, 9H), 1.16 (d, *J* = 6.0 Hz, 3H), 1.27 (t, *J* = 7.6 Hz, 3H), 1.48–1.74 (m, 4H), 3.47–3.53 (m, 1H), 3.80 (s, 3H), 4.15–4.22 (m, 3H), 4.39–4.50 (m, 2H), 5.64 (d, *J* = 11.4 Hz, 1H), 5.96 (dd, *J* = 6.3, 15.3 Hz, 1H), 6.54 (t, *J* = 11.4 Hz, 1H), 6.87 (d, *J* = 8.4 Hz, 2H), 7.26 (d, *J* = 8.4 Hz, 2H), 7.42 (dd, *J* = 11.4, 15.3 Hz, 1H); ¹³C NMR: δ –4.8, –4.3, 14.3, 18.2, 19.6, 22.7, 25.9, 29.7, 30.2, 31.7, 32.1, 33.7, 55.2, 59.9, 69.9, 72.7, 73.0, 74.1, 74.5, 113.7, 117.5, 125.6, 129.2, 131.1, 144.2, 146.8, 159.1, 166.3. Anal. Calcd for C₂₆H₄₂O₅Si: C, 67.49; H, 9.15%. Found: C, 67.74; H, 9.21%.

Ethyl (2Z,4E,6R,9S)-6-*tert*-Butyldimethylsilyloxy-9-hydroxydeca-2,4-dienoate 32. To a solution of **31** (0.620 g, 1.34 mmol) in CH₂Cl₂–H₂O was added DDQ (0.394 g, 1.74 mmol). After stirring for 3 h at room temperature, the reaction mixture was filtered through a pad of Celite. The filtrate was dried and concentrated in vacuo to get a residue, which, on column chromatography (silica gel, 0–10% EtOAc/hexane), furnished pure **32** (0.353 g, 77%). colorless liquid; $[\alpha]_D^{24}$ –7.2 (c 0.82, CHCl₃); IR: 3415, 1715, 1641 cm⁻¹; ¹H NMR: δ 0.07 (s, 6H), 0.89 (s, 9H), 1.18 (d, *J* = 6.3 Hz, 3H), 1.22 (t, *J* = 6.3 Hz, 3H), 1.31–1.69 (m, 4H), 2.04 (broad s, 1H), 3.78–3.81 (m, 1H), 4.19 (q, *J* = 7.0 Hz, 2H), 4.29–4.35 (m, 1H), 5.65 (d, *J* = 11.9 Hz, 1H), 5.98 (dd, *J* = 15.4, 6.3 Hz, 1H), 6.56 (t, *J* = 11.9 Hz, 1H), 7.39–7.46 (m, 1H); ¹³C NMR: δ –4.8, –4.3, 14.3, 18.2, 22.7, 23.4, 25.9, 29.7, 31.9, 33.8, 34.4, 59.9, 67.8, 72.8, 114.0, 117.6, 125.8, 139.2, 144.0, 146.3, 166.3. Anal. Calcd for C₁₈H₃₄O₄Si: C, 63.11; H, 10.00%. Found: C, 63.18; H, 9.84%.

(2Z,4E,6R,9S)-6-*tert*-Butyldimethylsilyloxy-9-hydroxydeca-2,4-dienoic acid 33. A solution of **32** (0.080 g, 0.234 mmol) and aqueous 1 M LiOH (0.351 mL, 0.351 mmol) in THF:H₂O (1:1, 10 mL) was stirred at 0 °C for 6 h. Removal of solvent in vacuo gave a residue, which was purified by preparative thin layer chromatography (silica gel, 40% EtOAc/hexane) to afford **33** (60 mg, 82%). viscous liquid; $[\alpha]_D^{24}$ –6.98 (c 1.39, CHCl₃); IR: 3500–2500, 1693, 1639 cm⁻¹; ¹H NMR: δ 0.04 (s, 3H), 0.06 (s, 3H), 0.90 (s, 9H), 1.18 (d, *J* = 6.2 Hz, 3H), 1.48–1.68 (m, 4H), 2.34 (broad s, 1H), 3.75–3.85 (m, 1H), 4.28–4.36 (m, 1H), 5.66 (d, *J* = 11.0 Hz, 1H), 6.04 (dd, *J* = 15.4, 6.0 Hz, 1H), 6.65 (t, *J* = 11.4 Hz, 1H), 7.45 (dd, *J* = 15.4, 11.4 Hz, 1H); ¹³C NMR: δ –4.7, –4.2, 18.1, 23.1, 25.9, 29.7, 33.5, 33.9, 66.9, 72.5, 126.0, 126.4, 137.9, 142.4, 175.0. Anal. Calcd for C₁₆H₃₀O₄Si: C, 61.11; H, 9.61%. Found: C, 61.04; H, 9.76%.

(3Z,5E,7R,10R)-7-*tert*-Butyldimethylsilyloxy-10-methyl-7,8,9,10-tetrahydrooxecin-2-one 34. To a cooled (0 °C) and stirred solution of **33** (0.210 g, 0.67 mmol) and PPh₃ (0.352 g, 1.34 mmol) in anhydrous toluene (96 mL) was slowly added DIAD (0.26 mL, 1.34 mmol). The reaction mixture was stirred at room temperature for 16 h and concentrated in vacuo, and the residue was purified by preparative thin layer column chromatography (silica gel, 20% EtOAc/hexane) to furnish **34** (130 mg, 73% based on conversion) along with unreacted **33** (20 mg). colorless liquid; $[\alpha]_D^{24}$ –51.8 (c 0.4, CHCl₃); IR: 1702, 1643, 1360 cm⁻¹; ¹H NMR: δ 0.04 (s, 3H), 0.08 (s, 3H), 0.89 (s, 9H), 1.32 (d, *J* = 6.2 Hz, 3H), 1.46–1.59 (m, 2H), 1.78–2.01 (m, 2H), 4.02–4.15 (m, 1H), 4.75–4.93 (m, 1H), 5.58 (d, *J* = 11.6 Hz, 1H), 5.90 (dd, *J* = 8.2, 15.4 Hz, 1H), 6.49 (t, *J* = 11.6 Hz, 1H), 7.09 (dd, *J* = 15.4, 11.6 Hz, 1H); ¹³C NMR: δ –4.8, –4.0, 18.2, 20.9, 25.8, 29.7, 33.7, 35.1, 72.1, 75.3, 119.6, 126.2, 141.9, 146.1, 166.2. Anal. Calcd for C₁₆H₂₈O₃Si: C, 64.82; H, 9.52%. Found: C, 64.47; H, 9.22%.

(3Z,5E,7R,10R)-7-Hydroxy-10-methyl-7,8,9,10-tetrahydro-oxecin-2-one (Stagonolide E) **3**. Desilylation of **34** (40 mg, 0.14 mmol) with aqueous HF (0.4 mL) in CH₃CN (10 mL), followed by isolation, as described above, and purification by preparative thin layer chromatography (silica gel, 15% EtOAc/hexane) furnished pure **3** (19 mg, 77%). colorless oil; [α]_D²⁵ –178.0 (c 0.342, CHCl₃), (lit.^{11a} [α]_D²⁵ –181 (c 0.2, CHCl₃); IR: 3478, 1697, 1642 cm^{–1}; ¹H NMR: δ 1.26 (d, *J* = 6.2 Hz, 3H), 1.50–1.76 (m, 4H), 1.90–2.01 (m, 1H), 4.25–4.29 (m, 1H), 4.94–5.12 (m, 1H), 5.68 (d, *J* = 11.4 Hz, 1H), 6.06 (dd, *J* = 15.4, 5.0 Hz, 1H), 6.56 (t, *J* = 11.4 Hz, 1H), 6.77 (dd, *J* = 15.4, 11.2 Hz, 1H); ¹³C NMR: δ 19.8, 31.8, 32.8, 71.5, 72.6, 118.5, 125.2, 143.0, 145.9, 166.7. Anal. Calcd for C₁₀H₁₄O₃: C, 65.91; H, 7.74%. Found: C, 65.79; H, 7.42%.

■ ASSOCIATED CONTENT

Supporting Information

¹H and ¹³C NMR spectra for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Authors

*Fax: +91 22 25505151. E-mail: schatt@barc.gov.in (S.C.).

*E-mail: anubhas@barc.gov.in (A.S.).

Author Contributions

[†]Equal contributions.

Notes

The authors declare no competing financial interest.

■ REFERENCES

- (a) Nicolaou, K. C.; Vourloumis, D.; Winssinger, N.; Baran, P. S. *Angew. Chem., Int. Ed.* **2000**, *39*, 44–122. (b) Corey, E. J.; Guzman-Perez, A. *Angew. Chem., Int. Ed.* **1998**, *37*, 388–340. (c) Noyori, R. *Asymmetric Catalysis in Organic Synthesis*; Wiley: New York, 1994.
- (a) Michels, P. C.; Khmelnitsky, Y. L.; Dordick, J. S.; Clark, D. S. *Trends Biotechnol.* **1998**, *16*, 210–215. (b) Fessner, W.-D.; Helaine, V. *Curr. Opin. Biotechnol.* **2001**, *12*, 574–586. (c) Clouthier, C. M.; Pelletier, J. N. *Chem. Soc. Rev.* **2012**, *41*, 1585–1605. (d) Reetz, M. T. *J. Am. Chem. Soc.* **2013**, *135*, 12480–12496. (e) Nestl, B. M.; Hammer, S. C.; Nebel, B. A.; Hauer, B. *Angew. Chem., Int. Ed.* **2014**, *53*, 3070–3095.
- (a) Reetz, M. T. *Curr. Opin. Chem. Biol.* **2002**, *6*, 145–150. (b) Faber, K. *Biotransformations in Organic Chemistry*, 5th ed.; Springer: Heidelberg, Germany, 2004. (c) Bornscheuer, U. T.; Kazlauskas, R. J. *Hydrolases in Organic Synthesis*, 2nd ed.; Wiley-VCH: Weinheim, Germany, 2005. (d) Gotor-Fernández, V.; Busto, E.; Gotor, V. *Adv. Synth. Catal.* **2006**, *348*, 797–812. (e) Haraldsson, G. G. *The Application of Lipases in Organic Synthesis* [Online]; John Wiley & Sons, Ltd.: Hoboken, NJ, 2010. DOI: 10.1002/9780470772508.ch24. (f) Sharma, S.; Kanwar, S. S. *Sci. World J.* **2014**, *2014*, No. 625258.
- (a) Sharma, A.; Chattopadhyay, S. *J. Org. Chem.* **1998**, *63*, 6128–6131. (b) Sharma, A.; Chattopadhyay, S. *J. Org. Chem.* **1999**, *64*, 8059–8062. (c) Kulkarni, B. A.; Sharma, A.; Gamre, S.; Chattopadhyay, S. *Synthesis* **2003**, 595–599. (d) Kumar, A. N.; Bhatt, S.; Chattopadhyay, S. *Tetrahedron: Asymmetry* **2009**, *20*, 205–209. (e) Sharma, A.; Gamre, S.; Chattopadhyay, S. *Tetrahedron: Asymmetry* **2009**, *20*, 1164–1167. (f) Ghadigaonkar, S.; Koli, M. R.; Gamre, S. S.; Choudhary, M. K.; Chattopadhyay, S.; Sharma, A. *Tetrahedron: Asymmetry* **2012**, *23*, 1093–1099.
- (a) Maity, B.; Chattopadhyay, S. *Curr. Bioact. Compd.* **2008**, *4*, 225–244. (b) Guha, P.; Dey, A.; Sarkar, B.; Dhyani, M. V.; Chattopadhyay, S.; Bandyopadhyay, S. K. *J. Pharmacol. Exp. Ther.* **2009**, *328*, 1–10. (c) Guha, P.; Dey, A.; Chatterjee, A.; Chattopadhyay, S.; Bandyopadhyay, S. K. *Br. J. Pharmacol.* **2010**, *159*, 726–734. (d) Patro, B. S.; Maity, B.; Chattopadhyay, S. *Antioxid. Redox Signaling* **2010**, *12*, 945–960. (e) Patro, B. S.; Tyagi, M.; Saha, J.; Chattopadhyay, S. *Bioorg. Med. Chem.* **2010**, *18*, 7043–7051. (f) Yadav, S. K.; Adhikary, B.; Chand, S.; Maity, B.; Bandyopadhyay, S. K.; Chattopadhyay, S. *Free Radical Biol. Med.* **2012**, *52*, 1175–1187. (g) Maity, B.; Yadav, S. K.; Patro, B. S.; Bandyopadhyay, S. K.; Chattopadhyay, S. *Free Radical Biol. Med.* **2012**, *52*, 1680–1691.
- (a) Grabley, S.; Hammann, P.; Hutter, K.; Kirsch, R.; Kluge, H.; Thiericke, R.; Mayer, M.; Zeeck, A. *J. Antibiot.* **1992**, *45*, 1176–1182. (b) Grabley, S.; Granzer, E.; Hutter, K.; Ludwig, D.; Mayer, M.; Thiericke, R.; Till, G.; Wink, J.; Philipps, S.; Zeeck, A. *J. Antibiot.* **1992**, *45*, 56–65. (c) Kis, Z.; Furger, P.; Sigg, H. P. *Experientia* **1969**, *25*, 123–124. (d) Nozoe, S.; Hirai, K.; Tsuda, K.; Ishibashi, K.; Shirasaka, M. *Tetrahedron Lett.* **1965**, 4675–4677. (e) Grove, J. F. *J. Chem. Soc.* **1971**, *12*, 2261–2263. (f) Kind, R.; Zeeck, A.; Grabley, S.; Thiericke, R.; Zerlin, M. *J. Nat. Prod.* **1996**, *59*, 539–540. (g) Evidente, A.; Capasso, R.; Andolfi, A.; Vurro, M.; Chiara Zonno, M. *Nat. Toxins* **1999**, *6*, 183. (h) Evidente, A.; Cimmino, A.; Berestetskiy, A.; Mitina, G.; Andolfi, A.; Motta, A. *J. Nat. Prod.* **2008**, *71*, 31–34.
- (a) Kastanias, M. A.; Chrysai-Tokousbalides, M. *Pest Manage. Sci.* **2000**, *56*, 227–232. (b) Karsten, K.; Umar, F.; Ulrich, F.; Barbara, S.; Siegfried, D.; Gennaro, P.; Piero, S.; Sándor, A.; Tibor, K. *Eur. J. Org. Chem.* **2007**, 3206–3211. (c) Zhang, W.; Krohn, K.; Egdol, H.; Draeger, S.; Schulz, B. *Eur. J. Org. Chem.* **2008**, 4320–4328. (d) Kind, R.; Zeeck, A.; Grabley, S.; Thiericke, R.; Zerlin, M. *J. Nat. Prod.* **1996**, *59*, 539–540. (e) Christner, C.; Kullertz, G.; Fischer, G.; Zerlin, M.; Grabley, S.; Thiericke, R.; Taddei, A.; Zeeck, A. *J. Antibiot.* **1998**, *51*, 368–371.
- (a) Machinaga, N.; Kibayashi, C. *Tetrahedron Lett.* **1993**, *34*, 5739–5742. (b) Clark, J. S.; Whitlock, G. A. *Tetrahedron Lett.* **1994**, *35*, 6381–6382. (c) Nokami, J.; Taniguchi, T.; Ogawa, Y. *Chem. Lett.* **1995**, *24*, 43–44. (d) Solladie, G.; Arce, E.; Bauder, C.; Carreno, M. C. *J. Org. Chem.* **1998**, *63*, 2332–2337. (e) Esumi, T.; Kimura, R.; Mori, M.; Iwabuchi, Y.; Irie, H.; Hatakeyama, S. *Heterocycles* **2000**, *52*, 525–528. (f) Lukesh, J. M.; Donaldson, W. A. *Tetrahedron: Asymmetry* **2003**, *14*, 757–762. (g) Clark, J. S.; Fessard, T. C.; Whitlock, G. A. *Tetrahedron* **2006**, *62*, 73–78. (h) Garcia, I.; Gomez, G.; Teixeira, M.; Teran, C.; Fall, Y. *Tetrahedron Lett.* **2006**, *47*, 1333–1335. (i) Riatto, V. B.; Pilli, R. A.; Victor, M. M. *Tetrahedron* **2008**, *10*, 2279–2300. (j) Rawat, V.; Chouthaiwale, P. V.; Suryavanshi, G.; Sudalai, A. *Tetrahedron: Asymmetry* **2009**, *20*, 2173–2177. (k) Zúñiga, A.; Pérez, M.; Pazos, G.; Gómez, G.; Fall, Y. *Synthesis* **2010**, 2446–2450.
- (a) Dommerholdt, E. J.; Thijs, L.; Zwanenburg, B. *Tetrahedron Lett.* **1991**, *32*, 1499–1502. (b) Machinaga, N.; Kibayashi, C. *Tetrahedron Lett.* **1993**, *34*, 841–844. (c) Amigoni, S.; Le Floch, Y. *Tetrahedron: Asymmetry* **1997**, *8*, 2827–2831. (d) Yadav, J. S.; Reddy, U. V. S.; Reddy, B. V. S. *Tetrahedron Lett.* **2009**, *50*, 5984–5986. (e) Oh, H.-S.; Kang, H.-Y. *Bull. Korean Chem. Soc.* **2011**, *32*, 2869–2870. (f) Yadav, J. S.; Reddy, G.; Rao, T. S.; Reddy, B. V. S.; Ghamdi, A. A. K. *Synthesis* **2012**, 783–787.
- Trost, B. M.; Quintard, A. *Angew. Chem., Int. Ed.* **2012**, *51*, 6704–6708.
- (a) Sabitha, G.; Padmaja, P.; Reddy, P. N.; Jadav, S. S.; Yadav, J. S. *Tetrahedron Lett.* **2010**, *51*, 6166–6168. (b) Das, T.; Nanda, S. *Tetrahedron Lett.* **2012**, *53*, 256–258. (c) Schmidt, B.; Kunz, O. *Beilstein J. Org. Chem.* **2013**, *9*, 2544–2555.
- (a) Sharma, A.; Gamre, S.; Chattopadhyay, S. *Tetrahedron: Asymmetry* **2009**, *20*, 1164–1167. (b) Sharma, A.; Mahato, S.; Chattopadhyay, S. *Tetrahedron Lett.* **2009**, *50*, 4986–4988. (c) Biswas, S.; Chattopadhyay, S.; Sharma, A. *Tetrahedron: Asymmetry* **2010**, *21*, 27–32.
- (a) Ohtani, T.; Nakatsukasa, H.; Kamezawa, M.; Tachibana, H.; Naoshima, Y. *J. Mol. Catal. B: Enzym.* **1998**, *4*, 53–60. (b) Fuji, M.; Fukumura, M.; Hory, Y.; Hirai, Y.; Akita, H.; Nakamura, K.; Torizuka, K.; Ida, Y. *Tetrahedron: Asymmetry* **2006**, *17*, 2292–2298. (c) Felluga, F.; Forzato, C.; Ghelfi, F.; Nitti, P.; Pitacco, G.; Pagnoni, U. M.; Roncaglia, F. *Tetrahedron: Asymmetry* **2007**, *18*, 527–536.
- (a) Mori, K. *Tetrahedron* **1981**, *37*, 1341–1342. (b) Nakamura, K.; Kinoshita, M.; Ohno, A. *Tetrahedron* **1995**, *51*, 8799–8808. (c) Kinoshita, M.; Ohno, A. *Tetrahedron* **1996**, *52*, 5397–5406. (d) Steinreiber, A.; Stadler, A.; Mayer, S. F.; Faber, K.; Kappe, C. O. *Tetrahedron Lett.* **2001**, *42*, 6283–6286. (e) Steinreiber, A.; Edegger, K.; Mayer, S. F.; Faber, K. *Tetrahedron: Asymmetry* **2001**, *12*, 2067–

2071. (f) Fantin, G.; Fogagnolo, M.; Medici, A.; Pedrini, P.; Fontana, S. *Tetrahedron: Asymmetry* **2000**, *11*, 2367–2373. (g) Belan, A.; Bolte, J.; Fauve, A.; Gourcy, J. G.; Veschambre, H. *J. Org. Chem.* **1987**, *52*, 256–260. (h) Fantin, G.; Fogagnolo, M.; Giovannini, P. P.; Medici, A.; Pedrini, P. *Tetrahedron: Asymmetry* **1995**, *6*, 3047–3053. (i) Clair, N. S.; Wang, Y.-F.; Margolin, A. L. *Angew. Chem.* **2000**, *112*, 388–391. (j) Chen, S. L.; Hu, Q. Y.; Loh, T. P. *Org. Lett.* **2004**, *6*, 3365–3367. (15) Dale, J. A.; Mosher, S. H. *J. Am. Chem. Soc.* **1973**, *95*, 512–519. (16) (a) Patil, P. N.; Chattopadhyay, A.; Udupa, S. R.; Banerji, A. *Biotechnol. Lett.* **1993**, *15*, 367–372. (b) Ferreira, H. V.; Rocha, L. C.; Severino, R. P.; Viana, R. B.; da Silva, A. B. F.; Porto, A. L. M. *J. Iran. Chem. Soc.* **2010**, *7*, 883–889. (c) Burgess, K.; Henderson, I. *Tetrahedron: Asymmetry* **1990**, *1*, 57–60. (17) Jana, N.; Mahapatra, T.; Nanda, S. *Tetrahedron: Asymmetry* **2009**, *20*, 2622–2628. (18) Kazlauskas, R. J.; Weissfloch, A. N. E.; Rappaport, A. T.; Cuccia, L. A. *J. Org. Chem.* **1991**, *56*, 2656–2665. (19) Cabrera, Z.; Fernandez-Lorente, G.; Fernandez-Lafuente, R.; Palomo, J. M.; Guisan, J. M. *J. Mol. Catal. B: Enzym.* **2009**, *57*, 171–176. (20) Enders, D.; Nguyen, D. *Synthesis* **2000**, 2092–2098. (21) Reviews: (a) Armstrong, S. K. *J. Chem. Soc., Perkin Trans. 1* **1998**, 371–388. (b) Schrock, R. R. *Tetrahedron* **1999**, *55*, 8141–8153. (c) Blackwell, H. E.; O'Leary, D. J.; Chatterjee, A. K.; Washenfelder, R. A.; Bussmann, D. A.; Grubbs, R. H. *J. Am. Chem. Soc.* **2000**, *122*, 58–71. (d) Fürstner, A. *Angew. Chem., Int. Ed.* **2000**, *39*, 3012–3043. (e) Trnka, T. M.; Grubbs, R. H. *Acc. Chem. Res.* **2001**, *34*, 18–29. (f) Cannon, S. J.; Blechert, S. *Angew. Chem., Int. Ed.* **2003**, *42*, 1900–1923. (g) Nicolaou, K. C.; Bulger, P. G.; Sarlah, D. *Angew. Chem., Int. Ed.* **2005**, *44*, 4490–4527. (h) Lin, Y. A.; Davis, B. G. *Beilstein J. Org. Chem.* **2010**, *6*, 1219–1228. For some leading references, see: (i) Galan, B. R.; Kalbarczyk, K. P.; Szczepankiewicz, S.; Keister, J. B.; Diver, S. T. *Org. Lett.* **2007**, *9*, 1203–1206. (j) Lipshutz, B. H.; Aquinaldo, G. T.; Ghorai, S.; Voigtritter, K. *Org. Lett.* **2008**, *10*, 1325–1328. (k) Ferrié, L.; Bouzbouz, S.; Cossy, J. *Org. Lett.* **2009**, *11*, 5446–5448. (l) Voigtritter, K.; Ghorai, S.; Lipshutz, B. H. *J. Org. Chem.* **2011**, *76*, 4697–4702. (m) Meek, S. J.; O'Brien, R. V.; Llaveria, J.; Schrock, R. R.; Hoveyda, A. H. *Nature* **2011**, *471* (7339), 461–466. (22) Chatterjee, A. K.; Choi, T.-L.; Sanders, D. P.; Grubbs, R. H. *J. Am. Chem. Soc.* **2003**, *125*, 11360–11370. (23) Sundby, E.; Perk, L.; Anthonsen, T.; Aasen, A. J.; Hansen, T. V. *Tetrahedron* **2004**, *60*, 521–524. (24) (a) Simpson, T. J. *Nat. Prod. Rep.* **1987**, *4*, 339–376. (b) Mori, K. *Tetrahedron* **1989**, *45*, 3233–3298. (25) Gurjar, M. K.; Nagaprasad, R.; Ramana, C. V. *Tetrahedron Lett.* **2003**, *44*, 2873–2875. (26) Ando, K. *J. Org. Chem.* **1997**, *62*, 1934–1939.