

A short synthesis of (+)-hygrine

Enzo B. Arévalo-García^{a,b,*}, Juan Carlos Q. Colmenares^c

^a Department of Chemistry, Wright State University, Dayton, OH 45435, USA

^b Department of Chemistry, C.W. Post. L.I.U., Long Island, NY 11548, USA

^c Loker Hydrocarbon Research Institute, USC, Los Angeles, CA 90089, USA

Received 18 March 2008; revised 12 April 2008; accepted 15 April 2008


Available online 20 April 2008

Abstract

The synthesis of the alkaloid hygrine in a series of six steps starting from readily available proline *N*-methyl derivative (**5**) is described. © 2008 Elsevier Ltd. All rights reserved.

Keywords: Hygrine; Alkaloid; Dess–Martin periodinane

Hygrine (**1**) is an alkaloid present in coca leaves as well as in a variety of other plants. It has been the subject of several biological and pharmacological studies,¹ especially since it is known that hygrine is the precursor of hyoscyamine and scopolamine, both compounds are used in the preparation of a vast number of pharmaceutical products; consequently, there is an increased interest in the development of synthetic routes to this alkaloid in order to investigate structure–activity relationship. Hygrine (**1**) has been prepared previously² by rather




lengthy and tedious synthetic routes. We now present an exceptionally simple synthesis route which provides **1** in good overall yield (Scheme 1); As a result, our approach presents an advantage over a recently published synthesis of hygrine.³

Thus, treatment of the readily available **5**⁴ with DIBAL-H in CH₂Cl₂ at –30 °C afforded aldehyde **4**. This product, without further purification, was then reacted with (methoxymethyl)triphenylphosphonium bromide and with KO-*t*-Bu/THF to generate a methyl vinyl ether. The subsequent acid hydrolysis of the vinyl ether provided the homologated aldehyde **3**; treatment of this compound with methyl magnesium bromide afforded **2** in a mixture 2.5:1 (hygroline:pseudohygroline). Finally, the transformation of **2** to hygrine **1** was carried out by the Dess–Martin⁵ periodinane procedure; the final product was then purified with flash column chromatography (hexane–AcOEt = 6:4) and its identification was accomplished by comparison of the obtained data with those previously described in the literature.^{2,3} All the analytical data are in agreement with the values reported for the optically pure compound, included: bp 87–88 °C (23 mm); (lit.^{2c,5} bp 83–84 °C (21 mm)); hygrine·HCl⁶ [α]_D +34.0 (*c* = 0.5, H₂O); (lit.³ [α]_D +34.5 (*c* = 0.5, H₂O)).

Thus, an efficient 6-step synthesis of hygrine was accomplished in 35% overall yield.

* Corresponding author. Tel.: +1 516 4245959; fax: +1 516 2993944.
E-mail address: barevalo@excite.com (E. B. Arévalo-García).



Scheme 1.

Acknowledgements

The authors are grateful to Wright State University, C.W. Post Long Island University Chemistry Departments, and Loker Hydrocarbon Research Institute for partial support of this work.

References and notes

1. Some examples: (a) Berkov, S.; Zayed, R.; Doncheva, T. *Fitoterapia* **2006**, 77, 179; (b) Jordan, M.; Humam, M.; Bieri, S.; Christen, C.; Poblete, E.; Munoz, O. *Phytochemistry* **2006**, 67, 570; (c) Doncheva, T.; Berkov, S.; Philipov, S. *Biochem. Syst. Ecol.* **2006**, 34, 478; (d) Berkov, S.; Pavlov, A.; Georgiev, V.; Stanimirova, P.; Kovacheva, P.; Philipov, S. *Annuaire Univ. Sofia Fac. Biol.* **2005**, 96, 95; (e) Vitale, A.; Acher, A.; Pomilio, A. *J. Ethnopharmacol.* **1995**, 49, 81; (f) Humam, M.; Munoz, O.; Christen, P.; Hostettmann, K. *Nat. Prod. Commun.* **2007**, 2, 743.
2. Some examples: (a) Nagasaka, T.; Yamamoto, H.; Hayashi, I.; Watanabe, M.; Hamaguchi, F. *Heterocycles* **1989**, 29, 155; (b) Langenskiöld, T.; Lounasmaa, M. *Heterocycles* **1983**, 20, 671; (c) Shono, T.; Matsumura, Y.; Tsubata, K. *J. Am. Chem. Soc.* **1981**, 103, 1172; (d) Hill, R. K.; Loeffler, L. J. *J. Org. Chem.* **1960**, 25, 2028.
3. Lee, J.-H.; Jeong, B.-S.; Ku, J.-M.; Jew, S.-s.; Park, H.-g. *J. Org. Chem.* **2006**, 71, 6690.
4. Commercially available. It is also easily synthesized from proline by treatment with $\text{SOCl}_2/\text{MeOH}$, and subsequent addition of MeI and NaH.
5. Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, 48, 4155.
6. Hygrine was readily transformed into its hydrochloride salt for comparison with the literature data of the same compound (see Ref. 3).