

Diastereoselective hydrophosphonylation of imines using (*R,R*)-TADDOL phosphite. Asymmetric synthesis of α -aminophosphonic acid derivatives†

Francisco Palacios,* Tomasz K. Olszewski and Javier Vicario

Received 17th February 2010, Accepted 8th July 2010

DOI: 10.1039/c003004j

Efficient synthesis of α -aminophosphonic acid derivatives is achieved, the key step being a diastereoselective hydrophosphonylation of *N*-diphenylphosphinyl imines using a readily available chiral cyclic (*R,R*)-TADDOL-phosphite derived from inexpensive natural tartaric acid.

Introduction

α -Aminophosphonic acids¹ are structural analogous to α -amino acids, obtained by isosteric substitution of the carboxylic acid by a phosphonate moiety. As expected from this analogy the single α -aminophosphonic acid molecules or their phosphonate esters as well as phosphapeptides containing α -aminophosphonic units show an assorted biological activity² as happens of catalytic antibodies,^{3a} peptide mimetics,^{3b} enzyme inhibitors,^{3c-e} and antibacterial agents.^{3f} Moreover they are key compounds in agrochemistry and have found several applications as fungicides^{4a} or herbicides.^{4b}

The Pudovik reaction⁵ is one of the most straight methods for the synthesis of α -aminophosphonic acid derivatives, which involves a variant of Strecker reaction based in the nucleophilic addition of phosphites to imines. Because the biological activity of α -aminophosphonic acid derivatives depends very often on the absolute configuration of their α carbon, a strong effort has been made in the study of stereoselective Pudovik reactions,⁶ mainly focused in the catalytic hydrophosphonylation of imines.⁷ However, there are only few examples of diastereoselective Pudovik reactions with acyclic imines using chiral phosphites.⁶ Addition of (–)-bornyl and (–)-menthyl phosphites **I** and **II** (Fig. 1) to *N*-benzylimine derived from benzaldehyde has been reported to afford α -aminophosphonates with moderate to good diastereoselectivity.⁸

Last years we have been involved in the synthesis of α -⁹ and β -aminophosphonate¹⁰ derivatives and we even used TADDOL phosphoimide derivatives as catalyst for the asymmetric synthesis of α -dehydroaminoester derivatives.¹¹ In this context, we thought^{12,13} that TADDOL derived phosphite¹⁴ **IV** (Fig. 1) would be a suitable candidate as phosphorus nucleophile for Pudovik reactions. (*R,R*)-TADDOL auxiliary is easily prepared in a simple two step procedure from cheap natural tartaric acid¹⁵ and, unlike natural alcohols, only one unit of chiral auxiliary relative to phosphorus is needed for the preparation of the hydrogenphosphonate ester. Moreover, TADDOL phosphite derivative shows a C-2 symmetry, implying the presence of a non chiral phosphorus(v)





Fig. 1 (–)-Borneol **I**, (–)-Menthol **II**, (*R*)-(+)-BINOL **III** and TADDOL **IV** derived phosphites.

which avoids the presence of mixtures of diastereoisomers and their separation. Furthermore, the endocyclic P–O bonds in the 7-membered ring, now including four sp^3 carbons, would be more stable than in the case of BINOL phosphite **III** as this is a less strained structure. Now, as a continuation of our work in the field of aminophosphorus compounds we report here a diastereoselective hydrophosphonylation of imines using a TADDOL derived phosphite and applied to the stereoselective synthesis of α -aminophosphonic acids.

Results and discussion

(*R,R*)-TADDOL derived phosphite **3** is easily synthesized by generation of chlorophosphite **2** from (*R,R*)-TADDOL **1** and PCl_3 and subsequent hydrolysis ‘*in situ*’ with triethylamine in aqueous media (Scheme 1, 95%).




Scheme 1 Synthesis of (*R,R*)-TADDOL phosphite **3**.


First we tested the hydrophosphonylation reaction of several aldimines derived from benzaldehyde in THF or CH_2Cl_2 (Table 1). In terms of chiral induction, it was established that the use of diphenylphosphinylimines is preferred if compared to tosyl or benzylamines. The larger size of diphenylphosphinyl substituent at the nitrogen might be the reason for this behaviour. The benefits of diphenylphosphinyl substituent on the electronic activation

Departamento de Química Orgánica I, Facultad de Farmacia, Universidad del País Vasco, Apartado 450, 01080 Vitoria, Spain. E-mail: francisco.palacios@ehu.es; Fax: (+34)-945-013049; Tel: (+34) 945 183103

† Electronic supplementary information (ESI) available: Preparation and full characterization of compounds **3**, **4f–h** and **5** and copies of 1H , ^{13}C of α -aminophosphonates **4**. See DOI: 10.1039/c003004j

Table 1 Hydrophosphonylation of aldimines derived from benzaldehyde with (*R,R*)-TADDOL phosphite **3**


Entry	<i>R</i>	Base	Solvent	<i>T</i> /°C	%Conv ^a	dr ^a
1	Bz	LDA (1 eq.)	THF	−80	97	56 : 44
2	Bz	ZnEt ₂ /TMEDA (1 eq.)	THF	−80	48	65 : 35
3	Ts	LDA (1 eq.)	THF	−80	88	75 : 25
4	Ts	ZnEt ₂ /TMEDA (1 eq.)	THF	−80	79	62 : 38
5	POPh ₂	Et ₃ N (1 eq.)	CH ₂ Cl ₂	r.t.	90	45 : 55
6	POPh ₂	No	CH ₂ Cl ₂	r.t.	0	—
7	POPh ₂	No	CH ₂ Cl ₂	40	0	—
8	POPh ₂	LDA	THF	−80	75	74 : 26
9	POPh ₂	ZnEt ₂ (1 eq.)	THF	−80	70	70 : 30
10	POPh ₂	ZnEt ₂ /TMEDA (1 eq.)	THF	−80	82	92 : 8
11	POPh ₂	ZnEt ₂ /TMEDA (1.2 eq.)	THF	−80	90	>95 : 5
12	POPh ₂	ZnEt ₂ /TMEDA (1.5 eq.)	THF	−80	95	>95 : 5
13	POPh ₂	ZnEt ₂ /TMEDA (1.8 eq.)	THF	−80	97	>95 : 5
14	POPh ₂	ZnEt ₂ /TMEDA (2.1 eq.)	THF	−80	97	>95 : 5

^a Determined by ³¹P NMR.**Table 2** Hydrophosphonylation of *N*-diphenylphosphinyl aldimines with (*R,R*)-TADDOL phosphite **3**


Entry	Comp.	<i>R</i>	%Yield ^a	³¹ P NMR (major)	³¹ P NMR (minor)	dr ^b
1	4a	Ph	80	18.2, 25.0	17.1, 24.8	>95 : 5
2	4b	1-Naphthyl	75	18.6, 25.1	16.5, 25.0	>95 : 5
3	4c	2-Naphthyl	73	18.0, 25.1	16.9, 25.5	>95 : 5
4	4d	<i>p</i> -MeO-C ₆ H ₄	82	18.3, 24.5	17.2, 24.5	>95 : 5
5	4e	<i>p</i> -CF ₃ -C ₆ H ₄	75	17.2, 25.3	16.2, 25.1	>95 : 5
6	4f	2-Furyl	80	14.5, 25.4	14.5, 25.3	91 : 9
7	4g	2,4,6-Me ₃ C ₆ H ₂	59	18.6, 24.7	20.4, 23.8	85 : 15
8	4h	<i>i</i> -Pr	65	19.2, 23.0	18.9, 22.7	87 : 13

^a Isolated yield. ^b Determined by ³¹P NMR.

of aldimines as well as its effects in stereoselective nucleophilic addition reactions is well documented.¹⁶

Moreover, although very good conversions were obtained in general at low temperature when the reaction was performed using one equivalent of LDA as a base (Table 1, Entries 1, 3, 8), much better diastereoselectivities with still a good yield are obtained when ZnEt₂ is used instead (Table 1, Entries 2, 4, 10). The presence of one equivalent of tetramethylethylenediamine (TMEDA) is crucial in order to favour the solubility of ZnEt₂ in THF at low temperature by a coordination effect between the Zn and N atoms, since very low yields were obtained when TMEDA was not present (Table 1, Entry 9). Switching the solvent to CH₂Cl₂ did not improve the diastereoselectivity when Et₃N

was used as a base (Table 1, Entry 5) and, in the same solvent, the hydrophosphonylation reaction did not proceed in the absence of a base (Table 1, Entries 6–7). The conversions in THF at low temperature were finally improved to almost quantitative when the amount of ZnEt₂/TMEDA was increased (Table 1, Entries 10–14) without dropping in the diastereoselectivities.


With these results in hands we extended the hydrophosphonylation reaction to several aromatic diphenylphosphinyl aldimines (Table 2). Treatment of *N*-diphenylphosphinyl aldimines in THF at −80 °C with TADDOL phosphonate **3** in the presence of 1.2 eq. of ZnEt₂ and TMEDA during 12 h afforded in good yields the corresponding aminophosphonates **4** with very good diastereoselectivities.

The reaction is applicable to *N*-diphenylphosphinyl aryl aldimines ($R = \text{Ph}$, 1-naphthyl, 2-naphthyl), bearing electron rich ($R = p\text{-MeO-C}_6\text{H}_4$, Table 2, Entry 4) and electron poor ($R = p\text{-CF}_3\text{-C}_6\text{H}_4$, Table 2, Entry 5) substituents as well as heteroaromatic groups ($R = 2\text{-furyl}$, Table 2, Entry 6) at the α position. A substantial drop in the yield and diastereoselectivity was observed when an ortho substituted arylimine was used which may be due to the steric crowding in the electrophilic imine carbon ($R = 2,4,6\text{-Me}_3\text{C}_6\text{H}_2$, Table 2, Entry 7). Previous reports have described the generation '*in situ*' of *N*-phosphinylimines from phosphorylated aminosulfones using ZnEt_2 as base.¹⁷ In our case, an important fall in the diastereoselectivity and conversion was observed when aromatic *N*-phosphinylimines were generated '*in situ*' with ZnEt_2 and subsequently treated with TADDOL phosphite **3** in the presence of TMEDA (60% yield, d.r. = 65 : 45).

The diastereoselective addition of TADDOL phosphite **3** was also extended to imines derived from aliphatic aldehydes. Although required aliphatic imines are normally only available '*in situ*', isolation of *N*-phosphinylimine derived from isobutyraldehyde, from the corresponding aminosulfone, by β -elimination of sulfonic acid using NaHCO_3 as base, followed by filtration and evaporation is feasible. Treatment of unpurified *N*-phosphinyl isobutyraldimine in the optimized conditions afforded the target aminophosphonate **4h** in moderate yield and diastereoselectivity (Table 2, Entry 8).

It should be emphasized that pure samples of the major diastereoisomer can be obtained in all cases by simple purification of the crude mixtures by chromatography. This strategy represents an improvement not only in conversions but also in the diastereoselectivities respect to the other two reported examples of addition of chiral phosphites to imines.

The synthesis of enantiopure aminophosphonic acids **5** is achieved by simultaneous hydrolysis of (*R,R*)-TADDOL phosphite and diphenylphosphinyl groups of the major diastereoisomer of α -aryl and α -alkyl α -aminophosphonates **4a,h** in refluxing aqueous HCl 4 N (Scheme 2, 77–82%). The monitoring of selective deprotection of amino group in milder conditions, at room temperature and increasing the concentration of hydrochloric acid from 0.1 M to 2 M, showed mixtures of mono-deprotected phosphonate and double-deprotected phosphonic acid. Comparison of the optical rotation with literature values showed an (*R*) absolute configuration for the aminophosphonic analog of (*R*)-phenylglycine **5a** ($R = \text{Ph}$) and (*R*)-valine **5b** ($R = i\text{-Pr}$).



Scheme 2 Simultaneous deprotection of amino and phosphonate groups.

According to the configuration of the stereogenic center we propose a model for the addition of phosphite¹⁸ where the seven membered ring adopts a more stable boat conformation fixed by the *trans* configuration of the five-membered ring, where the two heteroatoms adopt the most stable equatorial orientation with the two hydrogens in axial conformation (see Fig. 2). The alkoxy group




Fig. 2 Model for the hydrophosphonylation of *N*-diphenylphosphinoyl imines with (*R,R*)-TADDOL phosphite **3**.

in the metalated phosphite tautomer is expected to adopt an axial orientation due to the anomeric effect and the Zn counteranion ionically bonded to the oxygen would be coordinated to the lone pair of the iminic nitrogen. The diphenylphosphinoyl group pointing opposite to the Zn makes the imine approach to the phosphite lone pair with the appropriate orientation for the nucleophilic attack of the phosphorus to the *Re* face.

Conclusions

In summary we report an easy synthesis of enantiopure α -aminophosphonic acid derivatives through diastereoselective hydrophosphonylation of *N*-diphenylphosphinylimines using an efficient (*R,R*)-TADDOL derived phosphorus nucleophile. Easily available (*R,R*)-TADDOL phosphite, prepared from very cheap commercially available natural tartaric acid, affords (*R*)- α -aminophosphonic acids. This is not trivial, since it is well known that in most of the occasions (*R*) forms of α -aminophosphonic acids^{2,6} are more active than their (*S*) isomers. As far as we know the results reported here represent the best conversions and/or diastereoselectivities reported so far in the addition of chiral phosphites to imines.

Experimental section

Representative example for the diastereoselective hydrophosphonylation of imines using TADDOL phosphite **3**.

Synthesis of 1,3-dioxolo[4,5-e][1,3,2]dioxaphosphepin, tetrahydro-2,2-dimethyl-6-[diphenylphosphinoylamino-1-phenylmethyl]-4,4,8,8 tetraphenyl-, 6-oxide, (*3aR,8aR*) (**4a**)

TADDOL phosphite **3** (1.0 g, 1.95 mmol) was dissolved in dry THF (15 mL) and the solution was cooled down to -80°C . A 1.1 M solution of Et_2Zn in toluene (2.14 mL) and TMEDA (0.3 mL, 1.95 mmol) were injected and the mixture was stirred for 15 min at -80°C . After that time the solution of *N*-benzylidene-*P,P*-diphenylphosphinic amide (0.60 g, 1.95 mmol) in dry THF (15 mL) was added dropwise. The stirring was continued for 12 h at -80°C before quenching with sat. aq. NH_4Cl (15 mL). The layers were separated and the aqueous layer was extracted with CH_2Cl_2 (3×30 mL). The combined organic extracts were dried over anhydrous MgSO_4 and concentrated under reduced pressure affording crude product that was further purified by column chromatography (SiO_2 , EtOAc/Hexane) affording 1.28 g (80%) of **4a** as a white solid. Mp. $165\text{--}167^\circ\text{C}$. $[\alpha]_D^{20} -111.4$ (c 1.0, CH_2Cl_2). R_f ($\text{EtOAc/Pentane } 2 : 1$): 0.40. $^1\text{H NMR}$ (300 MHz, CDCl_3): δ 0.43 (s, 3H, CH_3), 0.88 (s, 3H, CH_3), 3.75–3.80 (m, 1H, NH), 4.68–4.83 (m, 1H, CHP), 4.96 (d, $^3J_{\text{HH}} = 8.0$ Hz, 1H, CHO), 5.54 (d,

$^3J_{\text{HH}} = 8.0$ Hz, 1H, CHO), 6.96–7.90 (m, 35H, CH_{ar}). ^{13}C NMR (75 MHz, CDCl₃): δ 26.4 (CH₃), 27.2 (CH₃), 53.4 (d, $^1J_{\text{PC}} = 161.7$ Hz, CHP), 79.1 (d, $^3J_{\text{PC}} = 2.0$ Hz, CHO), 80.2 (d, $^3J_{\text{PC}} = 2.0$ Hz, CHO), 87.1 (d, $^2J_{\text{PC}} = 8.4$ Hz, CPh₂), 91.0 (d, $^2J_{\text{PC}} = 13.4$ Hz, CPh₂), 113.8 (C(CH₃)₂), 126.6 (2 $\times$ CH_{ar}), 127.2 (2 $\times$ CH_{ar}), 127.3 (2 $\times$ CH_{ar}), 127.4 (2 $\times$ CH_{ar}), 127.6 (CH_{ar}), 127.8 (CH_{ar}), 127.9 (CH_{ar}), 128.0 (2 $\times$ CH_{ar}), 128.1 (d, $^3J_{\text{PC}} = 5.4$ Hz, 2 $\times$ CH_{ar}), 128.3 (2 $\times$ CH_{ar}), 128.4 (4 $\times$ CH_{ar}), 128.5 (CH_{ar}), 128.6 (d, $^3J_{\text{PC}} = 7.8$ Hz, 2 $\times$ CH_{ar}), 128.9 (2 $\times$ CH_{ar}), 129.8 (2 $\times$ CH_{ar}), 131.2 (d, $^1J_{\text{PC}} = 102.3$ Hz, 2 $\times$ Cq), 131.9 (CH_{ar}), 132.0 (CH_{ar}), 132.1 (2 $\times$ CH_{ar}), 132.2 (CH_{ar}), 132.3 (2 $\times$ CH_{ar}), 132.6 (d, $^3J_{\text{PC}} = 9.1$ Hz, 2 $\times$ CH_{ar}), 136.4 (C_q), 139.4 (C_q), 139.6 (C_q), 143.7 (C_q), 144.3 (d, $^2J_{\text{PC}} = 7.6$ Hz, C_q). ^{31}P NMR (120 MHz, CDCl₃): δ 18.2 (d, $^3J_{\text{PP}} = 39.7$ Hz, PO(OR)₂), 25.2 (d, $^3J_{\text{PP}} = 39.7$ Hz, POPh₂). FTIR (neat) $\nu_{\text{max}}/\text{cm}^{-1}$: 3432 (N–H st), 1225 (P=O st). CIMS m/z (amu): 818 ([M+H], 100). HRMS (amu) Calcd for C₅₀H₄₆NO₆P₂ (M+H)⁺ 818.2800; Found, 818.2772.

Acknowledgements

The present work has been supported by the Dirección General de Investigación del Ministerio de Ciencia e Innovación (MICINN, Madrid DGI, CTQ2009-12156) and by the Departamento de Educación, Universidades e Investigación del Gobierno Vasco and Universidad del País Vasco (GV, IT 422-10; UPV, GIU-09/57). J. V. thanks the Ministerio de Ciencia e Innovación for a “Ramón y Cajal contract”. T.K.O. acknowledges the contract from Universidad del País Vasco, programme Ayudas de Especialización para Investigadores Doctores 2008. UPV/EHU-SGIker technical support (MICINN, GV/EJ, European Social Fund) is gratefully acknowledged.

Notes and references

- 1 For an excellent book see: V. P. Kukhar, H. R. Hudson, ed. in *Aminophosphonic and Aminophosphinic Acids. Chemistry and Biological Activity*. J Wiley, Chichester, 2000.
- 2 (a) For reviews see: P. Kafarski and B. Lejczak, *Curr. Med. Chem.: Anti-Cancer Agents*, 2001, **1**, 301–312; (b) J. Grembecka and P. Kafarski, *Mini-Rev. Med. Chem.*, 2001, **1**, 133–144; (c) P. Kafarski and B. Lejczak, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1991, **63**, 193–215.
- 3 (a) R. Hirschmann, A. B. Smith III, C. M. Taylor, P. A. Benkovic, S. D. Taylor, K. M. Yager, P. A. Sprengeler and S. J. Venkovic, *Science*, 1994, **265**, 234–237; (b) B. Lejczak, P. Kafarski, H. Sztajer and P. Mastarlerz, *J. Med. Chem.*, 1986, **29**, 2212–2217; (c) J. Oleksyszyn and J. C. Powers, *Biochemistry*, 1991, **30**, 485–493; (d) P. A. Bartlett, J. E. Hanson and P. G. Giannousis, *J. Org. Chem.*, 1990, **55**, 6268–6274; (e) M. C. Allen, W. Fuhrer, B. Tuck, R. Wade and J. M. Wood, *J. Med. Chem.*, 1989, **32**, 1652–1661; (f) F. R. Atherton, C. H. Hassall and R. W. Lambert, *J. Med. Chem.*, 1986, **29**, 29–40.
- 4 (a) W. W. Smith and P. A. Bartlett, *J. Am. Chem. Soc.*, 1998, **120**, 4622–4628; (b) D. Bonarska, H. Kleszczyńska and J. Sarapuk, *Cell Mol. Biol. Lett.*, 2002, **7**, 929–935.
- 5 (a) V. A. Alfonso, *Phosphorus, Sulfur Silicon Relat. Elem.*, 2008, **183**, 2637–2644; (b) A. N. Pudovik, *Dokl. Akad. Nauk SSSR*, 1952, **83**, 865–868.
- 6 (a) P. Merino, E. Marques-Lopez and R. P. Herrera, *Adv. Synth. Catal.*, 2008, **350**, 1195–1208; (b) T. Vilaivan, W. Bhanthumnavin and Y. Sritana-Anant, *Curr. Org. Chem.*, 2005, **9**, 1315–1392; (c) H. Gröger and B. Hammer, *Chem.-Eur. J.*, 2000, **6**, 943–948.
- 7 (a) Recent contributions: F. Shi and B. Song, *Org. Biomol. Chem.*, 2009, **7**, 1292–1298; (b) S. Nakamura, H. Nakashima, A. Yamamura, N. Shibata and T. Toru, *Adv. Synth. Catal.*, 2008, **350**, 1209–1212; (c) J. P. Abell and H. Yamamoto, *J. Am. Chem. Soc.*, 2008, **130**, 10521–10523; (d) G. D. Joly and E. N. Jacobsen, *J. Am. Chem. Soc.*, 2004, **126**, 4102–4103.
- 8 O. I. Kolodiazny, E. V. Grishkun, S. Sheiko, O. Demchuk, H. Thoennessen, P. G. Jones and R. Schmutzler, *Tetrahedron: Asymmetry*, 1998, **9**, 1645–1649.
- 9 (a) For a recent review see: F. Palacios, C. Alonso and J. M. de los Santos, *Chem. Rev.*, 2005, **105**, 899–931; (b) F. Palacios, A. M. Ochoa de Retana, J. Oyarzabal, S. Pascual and G. Fernandez de Troconiz, *J. Org. Chem.*, 2008, **73**, 4568–4574.
- 10 (a) J. Vicario, D. Aparicio and F. Palacios, *J. Org. Chem.*, 2009, **74**, 452–455; (b) J. M. De los Santos, R. Ignacio, D. Aparicio and F. Palacios, *J. Org. Chem.*, 2007, **72**, 5202–5206; (c) F. Palacios, J. Vicario, A. Maliszewska and D. Aparicio, *J. Org. Chem.*, 2007, **72**, 2682–2685; (d) F. Palacios, A. M. Ochoa de Retana and J. M. Alonso, *J. Org. Chem.*, 2005, **70**, 8895–8901; (e) F. Palacios, D. Aparicio, Y. López and J. M. de los Santos, *Tetrahedron Lett.*, 2004, **45**, 4345–4348; (f) F. Palacios, D. Aparicio, A. M. Ochoa de Retana, J. M. de los Santos, J. I. Gil and R. Lopez de Munain, *Tetrahedron: Asymmetry*, 2003, **14**, 689–700; (g) F. Palacios, D. Aparicio, A. M. Ochoa de Retana, J. M. de los Santos, J. I. Gil and J. M. Alonso, *J. Org. Chem.*, 2002, **67**, 7283–7288.
- 11 (a) J. Vicario and F. Palacios, *Org. Lett.*, 2006, **8**, 5405–5408; (b) J. Vicario and F. Palacios, *Synthesis*, 2007, 3923–3925.
- 12 During our preliminary analysis of several chiral phosphorus nucleophiles we found that cleavage of endocyclic P–O bonds in BINOL phosphite (III) is very easy due to the twisted structure of BINOL auxiliary and the low stability of the 7-membered ring including four sp² carbons. For instance, *trans* esterification is fast if BINOL phosphites are dissolved in alcoholic solvents. That might be the reason of the low yields achieved in the addition of BINOL phosphite (III) towards BF₃-activated 3-thiazolines. (See ref. 17).
- 13 I. Schlemminger, A. Lützen, A. Willecke, W. Maison, R. Koch, W. Saak and J. Martens, *Tetrahedron Lett.*, 2000, **41**, 7285–7288.
- 14 (a) For recent contributions with TADDOL phosphite IV see: M. R. Garret, J. C. Tarr and J. S. Johnson, *J. Am. Chem. Soc.*, 2007, **129**, 12944–12945; (b) D. Enders, L. Tedeschi and D. Förster, *Synthesis*, 2006, 1447–1469; (c) L. Ackermann, C. J. Gschrei, A. Althammer and M. Riederer, *Chem. Commun.*, 2006, 1419–1421; (d) M. R. Nahm, J. R. Potnick, P. S. White and J. S. Johnson, *J. Am. Chem. Soc.*, 2006, **128**, 2751–2756.
- 15 F. Toda and K. Tanaka, *Tetrahedron Lett.*, 1988, **29**, 551–554.
- 16 (a) Some selected examples of diastereoselective nucleophilic additions to diphenylphosphinyl aldimines: R. B. Lettan II, C. C. Woodward and K. A. Scheidt, *Angew. Chem., Int. Ed.*, 2008, **47**, 2294–2297; (b) H. Morimoto, S. H. Wiedemann, A. Yamaguchi, S. Harada, Z. Chen, S. Matsunaga and M. Shibasaki, *Angew. Chem., Int. Ed.*, 2006, **45**, 3146–3150; (c) M. Sugita, A. Yamaguchi, N. Yamagiwa, S. Handa, S. Matsunaga and M. Shibasaki, *Org. Lett.*, 2005, **7**, 5339–5342; (d) T. Yoshida, H. Morimoto, N. Kumagai, S. Matsunaga and M. Shibasaki, *Angew. Chem., Int. Ed.*, 2005, **44**, 3470–3474; (e) G. Grach, J. Sopkova-de Oliveira Santos, J. Lohier, L. Mojovic, N. Plé, A. Turck, V. Reboul and P. Metzner, *J. Org. Chem.*, 2006, **71**, 9572–9579; (f) S. Matsunaga, T. Yoshida, H. Morimoto, N. Kumagai and M. Shibasaki, *J. Am. Chem. Soc.*, 2004, **126**, 8777–8785.
- 17 (a) A. Coté, A. A. Boezio and A. B. Charette, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 5405–5410; (b) J.-N. Desrosiers, A. Coté and A. B. Charette, *Tetrahedron*, 2005, **61**, 6186–6192.
- 18 J. L. Pirat, J. Monbrun, D. Virieux, J.-N. Volle, M. Tillard and H. J. Cristau, *J. Org. Chem.*, 2005, **70**, 7035–7041.