

Resolution of alkyl pyridyl sulfoxides by complexation with a chiral host compound derived from tartaric acid

Jin Zhu,^a Yong Qin,^a Ze He,^a Fang-min Fu,^a Zhong-yuan Zhou, Jin-gen Deng^{*},^{a,*} Yao-zhong Jiang^{* a} and Tay-yuan Chau^b

^a Union Laboratory of Asymmetric Synthesis, Chengdu Institute of Organic Chemistry, Chinese Academy of Sciences, Chengdu 610041, China

^b Department of Chemistry, National Chung-Hsing University, Taichung 40227, Taiwan

Abstract: Alkyl pyridyl sulfoxides 2 were prepared in high enantiomeric excess by resolution through inclusion complexation with a chiral host compound, (2R,3R) or (2S,3S)-(-)-trans-4,5-bis(hydroxydiphenylmethyl)-2,2-dimethyl-1,3-dioxolane 1. The X-ray crystallographic structure analysis of complex (R,R)-(-)-1 and (R)-(-)-2b is reported. © 1997 Published by Elsevier Science Ltd

Chiral sulfoxides are useful synthons for asymmetric carbon-carbon bonds formation.¹ The preparation of non-racemic chiral sulfoxides has been developed.²⁻¹¹ However, reports about the preparation of alkyl pyridyl sulfoxides with high enantiomeric excess are few,^{8,10} despite their great potential for the development of new ways for asymmetric carbon-carbon bonds formation.¹²⁻¹⁴

Inclusion crystallization has been used since the early 1980s, to selectively and reversibly include chiral guest molecules in host lattices of chiral molecules.¹⁵ Recently, we found this method is effective for the resolution of alkyl pyridyl sulfoxides **2**, which form inclusion complexes with chiral host compounds **1**, and optically active alkyl pyridyl sulfoxides with high enantiomeric excess are obtained (Table 1).



A solution of (R,R)-(-)-1 (1.174 g, 2.52 mmol) and rac-2b (0.776 g, 5 mmol) in toluene (3 ml) was kept at room temperature for 12 h, a 1:1 inclusion complex of (R,R)-(-)-1 and (R)-(-)-2b was obtained as colourless crystals (1.265 g, 81% yield). Two recrystallizations of the crystals from toluene gave pure inclusion crystals (1.060 g, 68% yield, mp 122-123°C), and (R)-(-)-2b of 100% e.e. [0.240 g, 62% yield, $[\alpha]_D^{16}$ -168 (c=0.78, MeOH)] was obtained by chromatography on silica gel. A crude (S)-(+)-2b (0.39 g, 15% e.e.) was recovered from the filtrate by column chormatography on silica gel, and then the crude (S)-(+)-2b (0.39 g, 2.5 mmol) and (S,S)-(+)-1 (1.166 g, 2.5 mmol) were dissolved in toluene (2 ml) and kept at room temperature for 12 h. The crude inclusion complex of (S,S)-(+)-1 and (S)-(+)-2b was recrystallized twice from toluene and subsequently separated by column chromatography on silica gel to give (S)-(+)-2b of 98% e.e. [0.210 g, 54% yield, $[\alpha]_D^{16}$ +167 (c=0.79, MeOH)]. The recovered compounds 1 could be recycled and affect on the resolution was observed.

X-Ray structure analytical data showed that inclusion complexes of compounds 1 and 2 are produced by the formation of hydrogen bonds between the two components. The X-ray ORTEP drawing of

^{*} Corresponding author. Email: locmaq@ntr.cdb.ac.cn

sulfoxide	chiral host compound	mp of	product	yield (%)	e.e (%)°
		inclusion complex (°C)*			
2 a	(R,R)-(-)-1	119-123	(R)-(+)-2a	61	95
	(S,S)-(+)-1	132-133	(S)-(-)-2a	42	96
2b	(R,R)-(-)-1	122-123	(R)-(-)-2b	62	100
	(S,S)-(+)-1	122-124	(S)-(+)-2b	54	98

Table 1. Resolution of 2 through inclusion complexation with I by crystallization from toluene

a. Inclusion complex was purified by two recrystallizations from toluene.

b. The absolute configuration was confirmed by X-ray analysis.

c. The enantiomeric excess of 2 was determined by HPLC useing Chiralcel OB as the chiral solid phase and n-hexane /2-propanol (85 : 15) as ehute.



Figure 1. X-Ray ORTEP drawing of inclusion complex of (R,R)-(-)-1 and (-)-2b.

inclusion complex of (R,R)-(-)-1 and (-)-2b is shown in Figure 1.¹⁶ It is first report of an X-ray structure of a sulfoxide inclusion complex: a 1:1 complex of (-)-1 and (-)-2b was formed. The sulfonyl oxygen of (-)-2b is included by forming a hydrogen bond with the hydroxyl group of (-)-1, and the distance between O(1) and O(5) is 2.650 Å. The other remaining hydroxyl group O(2) of host 1 also makes an intramolecular hydrogen bond with the hydroxyl group of O(1), and the distance between O(1) and O(2) is 2.650 Å. The crystal of (-)-1–(-)-2b was found to be monoclinic and the space group to be p2₁. The absolute configuration of (-)-2b was confirmed to be (R), which was assigned by the (R,R) absolute configuration of the host 1 as an internal reference. The absolute configuration of (+)-2a was also confirmed to be (R) by X-ray and HPLC analysis.¹⁷

NMR spectra showed that the molar ratio of compound 1 and 2 in the inclusion complexes is one to one.¹⁸ IR spectra of the inclusion complexes of 1 and 2 also showed that there is hydrogen bonding between the host 1 and guest 2 in a crystalline state.¹⁹ The IR spectrum of the inclusion complex (-)-1-(+)-2a exhibits the sulfonyl absorption band at 1028 cm⁻¹, while that of (+)-2a shows the corresponding band at 1036 cm⁻¹, the hydrogen bonding between the sulfonyl oxygen of (+)-2a and the hydroxyl group of (-)-1 thus causes a lower wave number shift of 8 cm⁻¹. Inclusion complex (-)-1-(+)-2a exhibits a hydrogen bond around 3280 cm⁻¹, while chiral host (-)-1 exhibits intramolecular hydrogen bond at 3434 cm⁻¹ and intermolecular hydrogen bonding 3206 cm⁻¹. The IR spectrum of the inclusion complex (-)-1-(-)-2b shows the same case.

Acknowledgements

We are grateful for the financial support of the National Natural Science Foundation of China.

References

- (a) A. J. Walker, *Tetrahedron: Asymmetry*, **1992**, 3, 961; (b) G. H. Posner, Acc. Chem. Res., **1987**, 20, 72; (c) G. Solladie, Pure & Appl. Chem., **1988**, 60, 1699; (d) O. De Lucci and L. Pasquato, *Tetrahedron*, **1988**, 44, 6755.
- 2. K. K. Andersen, J. Org. Chem., 1964, 29, 1953.
- 3. J. K. Whitesell and M.-S. Wong, J. Org. Chem., 1991, 56, 4552.
- 4. C. Mioskowski and G. Solladie, Tetrahedron, 1980, 36, 227.
- 5. J. M. Llera, I. Fernandez and F. Alcudia, Tetrahedron Lett., 1991, 32, 7299.
- 6. F. Rebiere, O. Samuel, L. Ricard and H. B. Kagan, J. Org. Chem., 1991, 56, 5991.
- 7. (a) S. C. Benson and J. K. Sanyder, *Tetrahedron Lett.*, 1991, 32, 5885; (b) F. Wudl and T. B. K. Lee, J. Am. Chem. Soc., 1973, 95, 6349.
- (a) P. Pitchen, E. Dunach, M. N. Deshmukh and H. B. Kagan, J. Am. Chem. Soc., 1984, 106, 8188; (b) S. H. Zhao, O. Samuel and H. B. Kagan, Tetrahedron, 1987, 43, 5135; (c) F. Di Furia, G. Modena and R. Seraglia, Synthesis, 1984, 325.
- 9. F. A. Davis, R. T. Reddy, M. C. Weismiller and J. P. McCauley, J. Am. Chem. Soc., 1989, 111, 5964.
- 10. S. Colonna, N. Gaggero, L. Casella, G. Carrer and P. Pasta, Tetrahedron: Asymmetry, 1992, 3, 95.
- (a) H. Ohta, Y. Kato and G. Tsuchihashi, *Chem. Lett.*, **1986**, 217; (b) P. J. Cox, A. Persad and N. S. Simpkins, *Synlett*, **1992**, 194.
- 12. T. Y. Chau, H. C. Lin and W. Hu, Synthetic Commun., 1996, 26, 1867.
- 13. H. Takayama, A. Lyobe and T. Koizumi, J. Chem. Soc., Chem. Commun., 1986, 771.
- 14. G. Demailly, C. Greck and G. Solladie, Tetrahedron Lett., 1984, 25, 4113.
- 15. (a) F. Toda, Top. Curr. Chem., 1987, 140, 43; (b) F. Toda, K. Tanaka and T. Okada, J. Chem. Soc., Chem. Commun., 1995, 639; (c) G. Kaupp, Angew. Chem. Int. Ed. Engl., 1994, 3, 728.
- 16. Crystal data for 1:1 inclusion complex of (R,R)-(-)-1 and (-)-2b: $C_{38}H_{39}NO_5S$; Mr=621.76; colorless prism (0.38×0.28×0.2 mm); monoclinic; space group P2₁, a=9.701(2), b=9.953(2), c=17.392(2) Å; V=1678.1(5) Å³; Z=2; Dc=1.230 g/cm³; F(000)=660; μ (Mo-K α)=0.140 cm⁻¹; SIEMENS P4 diffractometer; MoK α radiation (0.71073 Å): graphite crystal monochromator; T=294 K; scan type, ω ; scan speed, 6° min⁻¹; scan ring, $\Delta \omega=1.4$; scan limits, 2°<20<48°; standard reflection, 3 per 97 reflections; crystal stability, standard reflection collected, 2335 (Rint=0.012). The skeletal structure was solved by the direct method and successive Fourier syntheses. All hydrogens were found by the difference Fourier syntheses. Full matrix least-squares refinement of positional and thermal parameters, including anomalous scattering factor of oxygen and carbon atoms, lead to the final convergence with R=0.0351 and WR₂=0.0772. The absolute configuration was assigned as shown by using the (2R,3R) absolute configuration of the host 1 as an internal reference.

- 17. J. Zhu, J. G. Deng, A. Q. Mi, Y. Z. Jiang, unpublished results. We obtained a crystal from the filtrate of the suspension in hexane of (-)-1 and 2a (1:1 mol/mol) and the X-ray structure analysis shows 2:1:1 inclusion complex of (2R,3R)-(-)-1 and (-)-2a and water. The absolute configuration of (-)-2a was confirmed to be (S) by the (2R,3R) absolute configuration of host 1 as an internal reference and retention time of (-)-2a is 11.3' on HPLC. However, that of (+)-2a resolved by the formation of inclusion complex of (2R,3R)-(-)-1 is 17.1' on HPLC and thus the absolute configuration of (+)-2a was confirmed to be (R). This configuration of (+)-2a was also verified according to the order of elute on Chiralcel OB column.¹⁰
- 18. ¹H-NMR (300 MHz, CDCl₃ and D₂O). Complex (-)-1-(+)-2a: δ (ppm) 0.97 (6H, s), 2.74 (3H, s), 4.48 (2H, s), 7.16-7.48 (21H, m), 7.87-7.91 (2H, m), 8.53-8.54 (1H, m). Complex (-)-1-(-)-2b: δ (ppm) 0.96 (6H, s), 1.08 (3H, J=7 Hz, t) 2.80 (2H, J=7 Hz, q), 4.48 (2H, s), 7.12-7.47 (21H, m), 7.81-7.83 (2H, m), 8.50-8.51 (1H, m).
- 19. IR spectrum (firm). Complex (-)-1-(+)-2a (v_{max}): 3280, 3086, 3058, 3032, 2957, 2854, 1576, 1562, 1494, 1448, 1425, 1410, 1377, 1368, 1333, 1240, 1225, 1171, 1102, 1083, 1052, 1028, 940, 887, 794, 771, 757, 743, 723, 706 cm⁻¹. Complex (-)-1-(-)-2b (v_{max}): 3282, 2958, 2872, 2854, 1598, 1579, 1561, 1492, 1423, 1378, 1271, 1236, 1202, 1171, 1082, 1051, 1024, 1011, 962, 887, 783, 760, 743, 722, 697 cm⁻¹.

(Received in Japan 20 May 1997)