

# 1,2-FDTA: A Ferrocene-Based Redox-Active EDTA Analogue with a High $\text{Ca}^{2+}/\text{Mg}^{2+}$ and $\text{Ca}^{2+}/\text{Sr}^{2+}$ Selectivity in Aqueous Solution

Herbert Plenio\*, Dirk Burth, and Raphael Vogler

Institut für Anorganische und Analytische Chemie,  
Albertstraße 21, D-79104 Freiburg, Germany  
E-mail: plenio@ruf.uni-freiburg.de

Received May 20, 1997

**Keywords:** Ferrocenes / EDTA / Cyclic voltammetry / Alkaline earth metals / Potentiometry

1,2-Ferrocenemethylenediyldiamino tetraacetate (1,2-FDTA) is a redox-active relative of EDTA and was synthesized in the reaction of 1,2-bis(chloromethyl)ferrocene with iminodiacetate methyl ester followed by cleavage of the ester with methanolic KOH. Potentiometric determination of the stability constants of the complexes of 1,2-FDTA with the alkaline earth metal ions showed this ligand to display a higher  $\text{Ca}^{2+}/\text{Mg}^{2+}$  and  $\text{Ca}^{2+}/\text{Sr}^{2+}$  selectivity ( $\text{Ca}^{2+} \log K = 6.56$ ,  $\text{Mg}^{2+} \log$

$K = 4.65$ ,  $\text{Sr}^{2+} \log K = 3.32$ ) than EDTA itself. The redox potential of 1,2-FDTA- $\text{Ca}^{2+}$  determined in aqueous solution by cyclic voltammetry was found to be anodically shifted by +50 mV with respect to that of the calcium free complex. Consequently the redox-inactive calcium can be determined electrochemically in aqueous solution with a high selectivity with respect to the other alkaline earth metal ions.

EDTA (ethylenediamino tetraacetate) has found widespread use in analytical chemistry for the determination of metals, since it forms very stable complexes with a large number of metal ions.<sup>[1]</sup> It is well suited for the complexation of alkaline earth metal ions and thus very useful for the quantitative determination of calcium, because it displays a remarkable  $\text{Ca}^{2+}/\text{Mg}^{2+}$  selectivity.<sup>[2]</sup> Ferrocene continues to be among the most prominent organometallic compounds and displays numerous interesting properties among which its ability to undergo reversible one-electron transfer processes is one of the most important.<sup>[3]</sup> The combination of ferrocenes with macrocyclic polyethers<sup>[4]</sup> or chelating ligands<sup>[5]</sup> has resulted in a number of attractive compounds, which have been used as redox-receptors for cations and anions,<sup>[6]</sup> redox-switched crown ethers<sup>[7]</sup> or as hydrogenase models<sup>[8]</sup> since the redox potential of the metallocene is influenced reversibly by coordination of metal cations by the appended macrocyclic or chelating ligand.

With a view to the development of amperometric sensor devices for alkaline earth metal ions<sup>[9]</sup> it therefore seemed very attractive to synthesize an electrochemically active ligand, which has the same coordination characteristics as EDTA.

We have recently described the synthesis of a 1,1'-substituted ferrocene which carries one iminodiacetate group on each cyclopentadienyl ring and determined the stability constants of the complexes with alkaline earth metal ions.<sup>[10]</sup> The complexes with  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  are reasonably strong in water [ $\log K(\text{Mg}^{2+}) = 4.63$ ,  $\log K(\text{Ca}^{2+}) = 4.97$ ] but a selectivity for any of the group II metal ions was not observed. We therefore anticipated that attaching the functional groups of EDTA at the 1,2-positions of one cyclopentadienyl ring of ferrocene would render the resulting

complexon much more similar to EDTA than the highly flexible 1,1'-substituted ferrocene species.

In this paper the synthesis of 1,2-ferrocenemethylenediyldiamino tetraacetate (1,2-FDTA), the electrochemical behavior and the potentiometric determinations of the stability constants of the complexes of alkaline earth metal ions with this ligand will be described and it will be shown that the complexation characteristics of 1,2-FDTA are closely related to those of EDTA.

## Results and Discussion

### Syntheses of the Ferrocenylimino Diacetates

The reaction of chloromethylferrocene with  $\text{HN}(\text{CH}_2\text{COOMe})_2$  in the presence of a base results in the formation of methyl ferrocenylimino diacetate in a yield of 64%. The methyl ester was hydrolyzed to the ferrocenylimino diacetate by refluxing a methanolic KOH solution for 4 h. A yellow precipitate consisting of the dipotassium salt of ferrocenylimino diacetate (FIA) could be isolated and was used for all following investigations (Scheme 1).

For the synthesis of 1,2-FDTA, 1,2-di(hydroxymethyl)ferrocene was treated with  $\text{PCl}_3$  resulting in the formation of the previously unknown 1,2-bis(chloromethyl)ferrocene. This highly reactive compound was treated with two equivalents of  $\text{HN}(\text{CH}_2\text{COOMe})_2$  to yield 1,2- $[\text{CH}_2\text{N}(\text{CH}_2\text{COOMe})_2]_2$ -ferrocene in a 44% yield. The ester cleavage was again performed in methanolic KOH to generate the respective tetraacetate [1,2-FDTA-(4  $\text{K}^+$ )] in 71% yield, which was used for all further investigations (Scheme 1), since treatment of this salt with HCl did not lead to the free ferrocenyl diiminotetraacetic acid but only resulted in decomposition of the ferrocene instead.<sup>[11]</sup> The tetraacetate 1,1'-FDTA was available from previous investigations (Scheme 1).<sup>[10]</sup>

Scheme 1. Synthesis and ferrocene-based EDTA relatives and proposed structure of 1,2-FDTCa; reaction and conditions: a) iminodiacetic acid methyl ester,  $\text{Et}_3\text{N}$  in  $\text{Et}_2\text{O}$ ; b)  $\text{KOH}/\text{MeOH}$ , reflux; c)  $\text{PCl}_3$  in  $\text{THF}$ ; d) iminodiacetic acid methyl ester,  $\text{Et}_3\text{N}$  in  $\text{Et}_2\text{O}$ ; e)  $\text{KOH}/\text{MeOH}$ , reflux




Table 1.  $\text{p}K_3$  and  $\text{p}K_4$  values of  $(\text{HOOCCH}_2)_2\text{N}-(\text{CH}_2)_n-\text{N}(\text{CH}_2\text{COOH})_2$  and 1,1'-FDTCa and 1,2-FDTCa ( $n$  is the number of  $\text{CH}_2$  groups separating the iminodiacetate groups)

$n =$	2	3	4	5	1,1'-FDTCa	1,2-FDTCa
$\text{p}K_3$	6.16	7.90	9.07	9.50	9.18	7.73
$\text{p}K_4$	10.26	10.27	10.45	10.58	10.27	8.90

FDTCa- $\text{Ca}^{2+}$ ). Prior to looking at stability constants derived from these data it can be easily seen in these schemes upon looking at the concentrations of free  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  that the former metal forms much more stable complexes than  $\text{Mg}^{2+}$  with 1,2-FDTCa. The calculated overall stability constants  $\beta$  of the  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ , and  $\text{Sr}^{2+}$  complexes of 1,2-FDTCa are listed in Table 2, while a listing of the stabilities of the different 1:1 complexes of these metal ions with 1,2-FDTCa and related ligands is found in Table 3. Upon comparing the 1:1-magnesium complexes of 1,1'-FDTCa- $\text{Mg}^{2+}$  and 1,2-FDTCa- $\text{Mg}^{2+}$  virtually the same stabilities ( $\log \beta_1 = 4.63$  vs. 4.65) were found. Very much in contrast the 1:1-calcium complex of 1,2-FDTCa is much more stable than that of 1,1'-FDTCa with the same metal ion. More significantly so 1,2-FDTCa- $\text{Ca}^{2+}$  is by almost two orders of magnitudes more stable than the complex of  $\text{Mg}^{2+}$  with the same ligand, hence documenting a very pronounced selectivity of  $\text{Ca}^{2+}$  over  $\text{Mg}^{2+}$  complexation. The difference between the two  $\log K$  values amounts to  $\Delta \log K = 1.91$  which is even slightly larger than the  $\text{Ca}^{2+}/\text{Mg}^{2+}$ -selectivity displayed by EDTA itself ( $\Delta \log K = 1.75$ ). The strontium complex of 1,2-FDTCa is much less stable than those of calcium and magnesium resulting in a  $\text{Ca}^{2+}/\text{Sr}^{2+}$  selectivity of  $\Delta \log K = 3.24$ , which is much better than the  $\text{Ca}^{2+}/\text{Sr}^{2+}$  selectivity displayed by EDTA ( $\Delta \log K = 2.16$ ). However, in general the stabilities of the complexes of 1,2-FDTCa are much lower than those of EDTA, but nonetheless the calcium complex of 1,2-FDTCa is almost as stable as that of the tetraacetate in which three  $\text{CH}_2$  units separate the two iminodiacetate groups (Table 3).

#### Determination of the Stability Constants

The stability constants of the complexes of the alkaline earth metal ions  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ , and  $\text{Sr}^{2+}$  with FIA and 1,2-FDTCa were determined by potentiometric titrations and the evaluation of the data was conducted with the program TITFIT.<sup>[12]</sup>

The  $\text{p}K_2$  value of FIA thus calculated is 9.58 and is comparable to that of  $(\text{HOOCCH}_2)_2\text{N}-(\text{CH}_2)_5-\text{N}(\text{CH}_2\text{COOH})_2$  in which the two nitrogen centers are virtually independent from each other (Table 1). The complexes of  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ , and  $\text{Sr}^{2+}$  with FIA are weak and the stability constants are in the same range as observed for the complexes of iminodiacetate or *N*-benzyliminodiacetate (Table 3). For calcium and strontium the 1:1 complexes are preferentially formed, whereas in the magnesium complex of FIA a  $\text{MgL}_2$  species is important ( $\log \beta_2 = 7.16$ ). In conclusion the complexes of FIA are equally weak for the three metal ions investigated and are not suitable as redox-active equivalents of EDTA.


In the case of 1,2-FDTCa the  $\text{p}K_3$  and  $\text{p}K_4$  values were determined to be 7.73 and 8.90. While the  $\text{p}K_3$  value is in the range found for other EDTA derivatives with closely spaced nitrogen atoms (Table 1), the  $\text{p}K_4$  value is surprisingly low, since in all other EDTA relatives the first protonation occurs at close to  $\text{p}K_4$  ca. 10.4. An explanation for this low  $\text{p}K_4$  value is not close at hand. In this respect one should, however, remember that due to the 1,2-substitution pattern at the cyclopentadienyl ring two orientations of the nitrogen atoms with respect to the ferrocene can be distinguished, which lead to an unusual stereochemistry in this molecule.

The stabilities of the complexes of 1,2-FDTCa with  $\text{Mg}^{2+}$ ,  $\text{Ca}^{2+}$ , and  $\text{Sr}^{2+}$  were determined by potentiometric titrations and the distribution curves for the concentrations of the different species, depending on the pH value, are depicted in Scheme 2 (1,2-FDTCa- $\text{Mg}^{2+}$ ) and Scheme 3 (1,2-

#### Cyclic Voltammetry

The fundamental difference between the ferrocene based ligands described here and EDTA is the ability of the for-

Scheme 2. Distribution curves of the 1,2-FDTCa- $\text{Mg}^{2+}$  complexes in a solution containing 1,2-FDTCa ( $0.904 \cdot 10^{-3}$  mol/l) and  $\text{Mg}^{2+}$  ( $1.000 \cdot 10^{-3}$  mol/l)



Scheme 3. Distribution curves of the 1,2-FDTA- $\text{Ca}^{2+}$  complexes in a solution containing 1,2-FDTA ( $0.904 \cdot 10^{-3}$  mol/l) and  $\text{Ca}^{2+}$  ( $1.018 \cdot 10^{-3}$  mol/l)




Table 2. Stability constants for the complexes of 1,2-FDTA with  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , and  $\text{Sr}^{2+}$

species	$\log \beta(\text{Mg}^{2+})$	$\log \beta(\text{Ca}^{2+})$	$\log \beta(\text{Sr}^{2+})$
ML	4.65	6.56	3.32
M(LH)	12.92	12.63	12.36
M(L)(LH)	19.36	20.06	17.59

Table 3. Stability constants of the 1:1 complexes of 1,2-FDTA, 1,1'-FDTA, EDTA, and related complexons with alkaline earth metal cations

ligand	$\text{Mg}^{2+}$	$\text{Ca}^{2+}$	$\text{Sr}^{2+}$	$\text{Ba}^{2+}$	reference
iminodiacetate	2.94	2.59	2.23	1.67	[18]
N-benzyliminodiacetate	2.63	3.13	—	—	[19]
FIA	3.17	3.55	2.94	—	this work
EDTA	8.69	10.44	8.28	8.0	[18]
$[\text{CH}_2]_3[\text{N}(\text{CH}_2\text{COOH})_2]_2$	6.02	7.12	5.18	4.24	[20]
1,1'-FDTA	4.63	4.97	3.91	3.06	[10]
1,2-FDTA	4.65	6.56	3.32	—	this work

mer to undergo (reversible) one-electron redox processes, which adds a new dimension to this class of ligands, especially when the coordination chemistry of otherwise redox-inactive metal ions is investigated. Even though the possible redox potential window in water is fairly small due to oxidative or reductive cleavage of the solvent, the redox potentials of the ferrocenes studied here are easily accommodated in the allowed potential range, thereby presenting no problems for the electrochemical studies.

The redox potentials of the potassium salts of FIA ( $E_{1/2} = +0.20$  V), 1,2-FDTA ( $E_{1/2} = +0.26$  V) and 1,1'-FDTA ( $E_{1/2} = +0.24$  V) and those of their respective metal complexes were determined in water solution and are given in Table 4. Complexation of metal cations by these ligands leads to anodic shifts  $\Delta E_{1/2}$  of the ferrocene redox potential, whose amount depends on the degree of electronic communication between the two metal centers.<sup>[13]</sup>

The redox potential shifts  $\Delta E_{1/2}$  upon complexation of the dicationic metal ions are listed relative to the potentials of the respective potassium complexes. It is obvious from these data that the redox shifts  $\Delta E_{1/2}$  of FIA group-II metal

Table 4. Redox potentials  $E_{1/2}$  of FIA, 1,2-FDTA, and 1,1'-FDTA and their alkaline earth metal complexes in aqueous solution at pH = 10;  $\Delta E_{1/2}$  is the anodic shift of the redox potential upon addition of alkaline earth metal salts relative to the  $E_{1/2}$  of the potassium complex of the respective ferrocene

ferrocene	$E_{1/2}$ [V]	$\Delta E_{1/2}$ [mV]
FIA- $\text{K}^+$	+0.20	—
FIA- $\text{Mg}^{2+}$	+0.22	+20
FIA- $\text{Ca}^{2+}$	+0.25	+50
FIA- $\text{Sr}^{2+}$	+0.23	+30
FIA- $\text{Ba}^{2+}$	+0.22	+20
1,2-FDTA- $\text{K}^+$	+0.26	—
1,2-FDTA- $\text{Mg}^{2+}$	+0.36	+100
1,2-FDTA- $\text{Ca}^{2+}$	+0.31	+50
1,2-FDTA- $\text{Sr}^{2+}$	+0.28	+20
1,1'-FDTA- $\text{K}^+$	+0.24	—
1,1'-FDTA- $\text{Mg}^{2+}$	+0.31	+70
1,1'-FDTA- $\text{Ca}^{2+}$	+0.32	+80

complexes are small regardless of the metal salt added. In the complexes of 1,2-FDTA and 1,1'-FDTA larger  $\Delta E_{1/2}$  values of up to +100 mV result from the complexation of metal ions, which can be easily monitored despite the typically much broader redox waves displayed in water solution. Noteworthy is the calcium induced +50 mV redox potential shift of 1,2-FDTA, which is smaller than the  $\Delta E_{1/2} = +100$  mV of  $\text{Mg}^{2+}$  but the strong preference of  $\text{Ca}^{2+}$  vs.  $\text{Mg}^{2+}$  complexation allows the easy detection of the heavier metal ion even in the presence of excess magnesium.

## Conclusions

We have demonstrated that it is possible to synthesize an organometallic ferrocene-based relative of EDTA. The complexon (1,2-Ferrocenemethylenediyl)-Diamino Tetra-Acetate (1,2-FDTA) displays important characteristics of EDTA, which are a high selectivity for the complexation of  $\text{Ca}^{2+}$  vs.  $\text{Mg}^{2+}$  as well as a fairly high stability constant of the  $\text{Ca}^{2+}$  complex ( $\log K = 6.56$ ). In addition the appended redox-active ferrocene unit allows the highly selective electrochemical detection of an otherwise redox-inactive metal such as calcium even in the presence of excess magnesium or strontium, thus making this compound a potential amperometric sensor component.

This work was supported by the *Deutsche Forschungsgemeinschaft* (Graduiertenkolleg "Ungepaarte Elektronen in Chemie und Biologie"). We wish to thank cand.-chem. R. Kiefer for experimental assistance, Prof. Dr. H. Vahrenkamp for his support and Prof. Dr. A. D. Zuberbühler for providing us with the program TITFIT.

## Experimental Section

Commercially available solvents and reagents were purified according to literature procedures. — NMR spectra were recorded at 300 K with a Bruker AC200 F ( $^1\text{H}$  NMR: 200 MHz;  $^{13}\text{C}$  NMR: 50 MHz).  $^1\text{H}$  NMR was referenced to residual hydrogen in the deuterated solvents and  $^{13}\text{C}$  NMR to the signals of the solvent:  $\text{CDCl}_3$  ( $\delta = 7.26, 77.0$ ) or in  $\text{D}_2\text{O}$  with respect to TSP standard ( $\delta = 0.00$ ). — Elemental analyses were performed at the Mikroanalytisches Laboratorium der Chemischen Laboratorien, Universität Freiburg. — Infrared spectra of KBr tablets were recorded with a Bruker IFS 25. — Cyclic voltammetry: The standard electrochemi-

cal instrumentation consisted of an Amel System 5000 potentiostat/galvanostat. Cyclic voltammograms were recorded in demineralized water at ambient temperature using Amel software for a PC. A three-electrode configuration was employed. The working electrode was a Pt disk (diameter 1 mm) sealed in soft glass, the counter electrode was a Pt wire, the potentials were referenced with an Ag/AgCl reference electrode. KCF<sub>3</sub>SO<sub>3</sub> (0.1 M) was used as a supporting electrolyte. – Starting materials were commercially available or synthesized according to literature procedures: 1,2-bis(hydroxymethyl)ferrocene,<sup>[14]</sup> chloromethylferrocene,<sup>[15]</sup> HN(CH<sub>2</sub>COOMe)<sub>2</sub>.<sup>[16]</sup>

**Potentiometry:** Experimental conditions were as described previously.<sup>[17]</sup> Overall stability constant  $\beta_{mlh}$  ( $m$  = number of metal ions,  $l$  = number of ligand molecules,  $h$  = number of protons):  $\beta_{110}$ :  $M + L = ML$ ;  $\beta_{111}$ :  $M + L + H = [M(LH)]$ ;  $\beta_{121}$ :  $M + 2L + H = M(L)(LH)$ . Stepwise stability constants  $K_{mlh}$ :  $K_{111}$ :  $M + LH = (MLH)$ ,  $\log K_{111} = \log \beta_{111} - \log \beta_{110}$ .

$$\sigma_N = \sqrt{\frac{\sum_{n=1}^N (m_{l_{exp}} - m_{l_{calc}})^2}{(n_m - n_{var})}}$$

$\sigma(ml)$  determines the quality of the fit of the proposed species distribution for TITFIT,  $m_{l_{exp}}$  and  $m_{l_{calc}}$  = added or calculated volume of NaOH solution,  $n_m$  = number of titrant additions,  $n_{var}$  = number of variables in the model. For a summary of potentiometric data see Table 5. Even though we did not observe any clouding of the solution due to the precipitation of Mg(OH)<sub>2</sub> this could be possible at pH  $\geq 10$ .

Table 5. Summary of selected potentiometric data

Measurement	Species	log $\beta_{MLH}$	$\sigma$ (ML)	$\frac{C_{ligand} \cdot C_{metal}}{[10^{-3} \text{ mol/l}]}$
FIA	011	9.58(4)	0.0028	1.005
FIA + Ca <sup>2+</sup>	CaL	3.55(9)	0.0033	1.000
FIA + Mg <sup>2+</sup>	MgL	3.17(14)	0.0034	1.019
	MgL <sub>2</sub>	7.16(8)		
FIA + Sr <sup>2+</sup>	SrL	2.94(14)	0.0030	1.000
1,2-FDTA	011	7.73(7)	0.0015	0.904
	012	8.90(2)		
1,2-FDTA + Ca <sup>2+</sup>	CaL	6.56(8)	0.0025	1.018
	Ca(LH)	12.63(9)		
	Ca(L)(LH)	20.06(15)		
1,2-FDTA + Mg <sup>2+</sup>	MgL	4.65(9)	0.0019	1.000
	Mg(LH)	12.92(9)		
	Mg(L)(LH)	19.36(15)		
1,2-FDTA + Sr <sup>2+</sup>	SrL	3.32(14)1	0.0028	1.235
	Sr(LH)	2.36(10)1		
	Sr(L)(LH)	7.59(15)		

**Fc[CH<sub>2</sub>N(CH<sub>2</sub>COOMe)<sub>2</sub>]:** To a solution of chloromethylferrocene (1.94 g, 8.2 mmol) in diethyl ether (30 ml) were added Et<sub>3</sub>N (2 ml) and HN(CH<sub>2</sub>COOMe)<sub>2</sub> (3.0 g, 18.6 mmol). After stirring for 12 h, the reaction mixture was filtered and the ethereal solution washed with aq. Na<sub>2</sub>CO<sub>3</sub>. The organic layer was separated, dried with MgSO<sub>4</sub> and the solvent evaporated. The remaining pale yellow oil was purified by chromatography (silica gel, cyclohexane/ethyl acetate = 10:1). Yield: 1.9 g (64%). – <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>):  $\delta$  = 3.28 (s, 6 H, OCH<sub>3</sub>), 3.54 (s, 4 H, NCH<sub>2</sub>), 3.80 (s, 2 H, FcCH<sub>2</sub>), 3.92–4.14 (m, 9 H, FcH). – <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>):  $\delta$  = 50.78, 53.59, 53.76, 68.39, 70.26, 83.86, 171.23. – IR [ $\nu$ (CO)]:  $\tilde{\nu}$  = 1752, 1738 cm<sup>−1</sup>.

**Fc[CH<sub>2</sub>N(CH<sub>2</sub>COOK)<sub>2</sub>] (FIA-2K<sup>+</sup>):** Fc[CH<sub>2</sub>N(CH<sub>2</sub>COOMe)<sub>2</sub>] (0.95 g, 2.6 mmol) was dissolved in methanolic 2 N KOH (20 ml) and heated under reflux for 4 h, whereupon a yellow precipitate formed. This solid was centrifuged off, the liquid decanted and the precipitate washed with methanol three times. Yield: 0.63 g (61%). – <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta$  = 3.10 (s, 2 H, FcCH<sub>2</sub>), 3.68 (s, 4 H, CH<sub>2</sub>COO<sup>−</sup>), 4.23 (m, 7 H, FcH), 4.28 (“t”,  $J$  = 1.7 Hz, 2 H, FcH). – <sup>13</sup>C NMR (D<sub>2</sub>O):  $\delta$  = 54.81, 60.04, 71.05, 71.58, 84.14, 182.06. – IR [ $\nu$ (CO)]:  $\tilde{\nu}$  = 1613, 1590 cm<sup>−1</sup>.

**1,2-(CH<sub>2</sub>Cl)<sub>2</sub>-Fc:** To a solution of 1,2-(CH<sub>2</sub>OH)<sub>2</sub>-Fc (1.5 g, 6.1 mmol) and pyridine (2 ml) in THF (40 ml) was added freshly distilled PCl<sub>3</sub> (0.5 ml, 5.4 mmol), dissolved in THF (10 ml). After stirring for 3 h, the solution was filtered and the volatiles were removed from the filtrate. The remaining solid was recrystallized from petroleum ether. The yellow product decomposes at room temperature and should be stored at −20 °C. Yield: 1.6 g (90%). – <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>):  $\delta$  = 3.75 (s, 5 H, FcH), 3.78 (t,  $J$  = 3.8 Hz, 1 H, FcH), 3.94 (d,  $J$  = 3.7 Hz, 2 H, FcH), 3.97 (d,  $J$  = 12 Hz, CH<sub>2</sub>Cl), 4.07 (d,  $J$  = 12 Hz, CH<sub>2</sub>Cl). – <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>):  $\delta$  = 41.25, 68.29, 69.66, 70.28, 83.52.

**1,2-[CH<sub>2</sub>N(CH<sub>2</sub>COOMe)<sub>2</sub>]<sub>2</sub>-Fc:** To a solution of 1,2-(CH<sub>2</sub>Cl)<sub>2</sub>-Fc (1.6 g, 5.7 mmol) and Et<sub>3</sub>N (3.0 ml) in ether (50 ml) was added HN(CH<sub>2</sub>COOMe)<sub>2</sub> (2.0 g, 12.2 mmol). After stirring for 12 h, the reaction mixture was washed with aqueous Na<sub>2</sub>CO<sub>3</sub>. The ethereal layer was separated, dried with MgSO<sub>4</sub>, filtered, and the solvent evaporated. The remaining brown oil was purified by chromatography (cyclohexane/ethyl acetate = 1:1). Yield: 1.3 g (44%). – <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>):  $\delta$  = 3.32 (s, 12 H, OCH<sub>3</sub>), 3.68 (s, 8 H, NCH<sub>2</sub>), 3.84 (m, 6 H, FcH), 3.86 (d,  $J$  = 12.6 Hz, 2 H, FcCH<sub>2</sub>), 3.96 (d,  $J$  = 12.8 Hz, 2 H, FcCH<sub>2</sub>), 4.16 (d,  $J$  = 2.4 Hz, 2 H, FcH). – <sup>13</sup>C NMR (C<sub>6</sub>D<sub>6</sub>):  $\delta$  = 50.75, 52.10, 54.07, 67.11, 69.51, 71.08, 84.27, 171.59.

**1,2-[CH<sub>2</sub>N(CH<sub>2</sub>COOK)<sub>2</sub>]<sub>2</sub>-Fc (1,2-FDTA-4K<sup>+</sup>):** 1,2-[CH<sub>2</sub>N(CH<sub>2</sub>COOMe)<sub>2</sub>]<sub>2</sub>-Fc (1.3 g, 2.4 mmol) was dissolved in methanolic 2 N KOH (20 ml) and heated under reflux for 4 h, whereupon a brown precipitate formed. This solid was centrifuged off, the liquid decanted and the precipitate washed with methanol three times. The residue was dissolved in water and freeze-dried. Yield: 1.2 g (71%). – <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta$  = 3.05–3.30 (m, 8 H, CH<sub>2</sub>COO<sup>−</sup>), 3.37 (d,  $J$  = 12.2 Hz, 2 H, FcCH<sub>2</sub>), 3.84 (d,  $J$  = 12.3 Hz, 2 H, FcCH<sub>2</sub>), 4.16 (s, 2 H, FcH), 4.29 (d,  $J$  = 2.0 Hz, 2 H, FcH). – <sup>13</sup>C NMR (D<sub>2</sub>O):  $\delta$  = 55.94, 61.4 (br.), 70.04, 72.17, 75.34, 85.37, 181.85 (br.). – C<sub>20</sub>H<sub>20</sub>FeK<sub>4</sub>N<sub>2</sub>O<sub>8</sub> (628.63): C 38.21, H 3.21, N 4.46; found: C 38.67, H 3.76.

- [1] R. Pribil, *Analytical Applications of EDTA and Related Compounds*, Pergamon, Oxford, 1972.
- [2] G. Schwarzenbach, H. Flaschka, *Complexometric Titrations*, Methuen, London, 1969.
- [3] A. Togni, T. Hayashi (Eds.), *Ferrocenes*, VCH, Weinheim, 1995.
- [4] J. C. Medina, T. T. Goodnow, M. T. Rojas, J. L. Atwood, B. C. Lynn, A. E. Kaifer, G. W. Gokel, *J. Am. Chem. Soc.* **1992**, 114, 10583. – [4b] H. Plenio, D. Burth, *Organometallics* **1996**, 15, 1151. – [4c] H. Plenio, R. Diodone, *Inorg. Chem.* **1995**, 35, 3964. – [4d] H. Plenio, R. Diodone, *J. Organomet. Chem.* **1995**, 492, 73. – [4e] A. J. Pilgrim, P. D. Beer, *J. Chem. Soc., Faraday Trans.* **1995**, 91, 4331.
- [5] [5a] H. Plenio, D. Burth, *Organometallics* **1996**, 15, 4054. – [5b] H. Plenio, D. Burth, *Angew. Chem.* **1995**, 107, 881; *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 800. – [5c] E. T. Singewald, C. A. Mirkin, C. L. Stern, *Angew. Chemie* **1995**, 107, 1725; *Angew. Chem. Int. Ed.* **1995**, 34, 1624. – [5d] I. M. Lorkovic, M. S. Wrighton, W. M. Davis, *J. Am. Chem. Soc.* **1994**, 116, 6220. – [5e] M. J. L. Tenders, A. Benito, R. Martinez-Manez, J. Soto, J. Paya, A. J. Edwards, P. R. Raithby, *J. Chem. Soc., Dalton Trans.* **1996**, 343.
- [6] [6a] P. D. Beer, Z. Chen, M. G. B. Drew, J. Kingdom, M. Odgen,

- P. Spencer, *J. Chem. Soc., Chem. Commun.* **1993**, 1046. — [6b] P. D. Beer, C. Hazelwood, D. Hesek, J. Hodacova, S. E. Stokes, *J. Chem. Soc., Dalton Trans.* **1993**, 1327. — [6c] P. D. Beer, M. G. B. Drew, A. R. Graydon, *J. Chem. Soc., Dalton Trans.* **1996**, 4129. — [6d] P. D. Beer, K. Y. Wild, *Polyhedron* **1996**, *15*, 775.
- [7] [7a] P. D. Beer, *Chem. Soc. Rev.* **1989**, *18*, 409. — [7b] F. C. J. M. vanVeggel, W. Verboom, D. N. Reinhoudt, *Chem. Rev.* **1994**, *94*, 279.
- [8] [8a] R. T. Hembre, J. S. McQueen, V. W. Day, *J. Am. Chem. Soc.* **1996**, *118*, 798. — [8b] R. T. Hembre, J. S. McQueen, *Angew. Chem.* **1997**, *109*, 79; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 65.
- [9] P. D. Beer, *Adv. Inorg. Chem.* **1992**, *39*, 79.
- [10] H. Plenio, D. Burth, P. Gockel, *Chem. Ber.* **1993**, *126*, 2585.
- [11] The related 1,2-substituted benzene compound has been synthesized, but the stability constants of metal complexes were not described: T. Ando, K. Ueno, *Bull. Chem. Soc. Jpn.* **1966**, *39*, 2053.
- [12] A. D. Zuberbühler, T. A. Kaden, *Talanta* **1982**, *29*, 201.
- [13] H. Plenio, J. Yang, R. Diodone, J. Heinze, *Inorg. Chem.* **1994**, *33*, 4098.
- [14] S. I. Goldberg, W. D. Bailey, *J. Am. Chem. Soc.* **1974**, *96*, 6381.
- [15] S. B. Wilkes, I. R. Butler, A. E. Underhill, M. B. Hursthouse, D. E. Hibbs, K. M. A. Malik, *J. Chem. Soc., Dalton Trans.* **1995**, 897.
- [16] H. W. Rinehart, *J. Am. Chem. Soc.* **1926**, *48*, 2798.
- [17] P. Gockel, H. Vahrenkamp, A. D. Zuberbühler, *Helv. Chim. Acta* **1993**, *76*, 511.
- [18] G. Anderegg, *Helv. Chim. Acta* **1964**, *47*, 1801.
- [19] T. Ando, *Bull. Chem. Soc. Jpn.* **1962**, *35*, 1395.
- [20] R. Smith, J. L. Bullock, F. C. Beworth, A. E. Martell, *J. Org. Chem.* **1949**, *14*, 355.

[97119]