

# Assembly of Substituted 3-Methyleneisoindolin-1-ones via a CuI/L-Proline-Catalyzed Domino Reaction Process of 2-Bromobenzamides and Terminal Alkynes


Li Li,<sup>†</sup> Miao Wang,<sup>†</sup> Xiaojing Zhang,<sup>†</sup> Yongwen Jiang,<sup>\*,‡</sup> and Dawei Ma<sup>\*,‡</sup>

Shenyang Pharmaceutical University, 103 Wenhua Lu, Shenyang 110016, China, and  
State Key Laboratory of Bioorganic and Natural Products Chemistry, Shanghai  
Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu,  
Shanghai 200032, China

madw@mail.sioc.ac.cn

Received January 16, 2009

## ABSTRACT



CuI/L-proline catalyzed coupling of 2-bromobenzamides and terminal alkynes in *i*-PrOH (or DMF and DMSO) at 85–110 °C and subsequent additive cyclization produces substituted 3-methyleneisoindolin-1-ones. Variation of N-substituents, aromatic ring, and methylene part is possible by using suitable starting materials.

The 3-methyleneisoindolin-1-one is a core structure of numerous natural products or designed pharmaceutical molecules.<sup>1</sup> These compounds include fumaridine (Figure 1) that was extracted from vegetable sources,<sup>1a</sup> compound **1** that has local anesthetic activity superior to that of procaine,<sup>1b</sup> compound **2** that exhibited sedative activity,<sup>1h</sup> and muscaric

receptor ligand **3**.<sup>1f</sup> In addition, 3-methyleneisoindolin-1-ones have been found to be useful intermediates for the total synthesis of alkaloids such as lennoxamine.<sup>2</sup>

The classical methods to 3-methyleneisoindolin-1-ones are based on using phthalimides as starting materials, which undergo Wittig reaction<sup>3</sup> or addition of organometallic reagents and subsequent dehydration<sup>2a,4</sup> to give the target molecules. This approach suffers from poor regioselectivity in the cases of unsymmetrical substrates. To avoid this problem, a new method based on condensation of phosphorylated 4-alkoxy-

<sup>†</sup> Shenyang Pharmaceutical University.


<sup>‡</sup> Shanghai Institute of Organic Chemistry.

(1) (a) Blaskó, G.; Gula, D. J.; Shamma, M. *J. Nat. Prod.* **1982**, *45*, 105. (b) Laboratori Baldacci S. p. A. JP 59046268, *Chem. Abstr.* **1984**, *101*, 54922. (c) Kato, Y.; Ebiiike, H.; Achiwa, K.; Ashizawa, N.; Kurihara, T.; Kobayashi, F. *Chem. Pharm. Bull.* **1990**, *38*, 2060. (d) Kato, Y.; Takemoto, M.; Achiwa, K. *Chem. Pharm. Bull.* **1999**, *47*, 529. (e) Chia, Y.-C.; Chang, F.-R.; Teng, C.-M.; Wu, Y.-C. *J. Nat. Prod.* **2000**, *63*, 1160. (f) Botero Cid, H. M.; Tränkle, C.; Baumann, K.; Pick, R.; Mies-Klomfass, E.; Kostenis, E.; Mohr, K.; Holzgrabe, U. *J. Med. Chem.* **2000**, *43*, 2155. (g) Couture, A.; Deniau, E.; Grandclaudon, P.; Rybalko-Rosen, H.; Léonce, S.; Pfeiffer, B.; Renard, P. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3557. (h) Hulin, B.; Parker, J. C.; Piotrowski, D. W. US2005234065. (i) Toyooka, K.; Kanamitsu, N.; Yoshimura, M.; Kuriyama, H.; Tamura, T. WO2004048332.

(2) (a) Pigeon, P.; Decroix, B. *Tetrahedron Lett.* **1996**, *37*, 7707. (b) Moreau, A.; Couture, A.; Deniau, E.; Grandclaudon, P. *J. Org. Chem.* **2004**, *69*, 4527. (c) Rys, V.; Couture, A.; Deniau, E.; Lebrun, S.; Grandclaudon, P. *Tetrahedron* **2005**, *61*, 665. (d) Couty, S.; Meyer, C.; Cossy, J. *Tetrahedron Lett.* **2006**, *47*, 767. (e) Lamblin, M.; Couture, A.; Deniau, E.; Grandclaudon, P. *Org. Biomol. Chem.* **2007**, *5*, 1466.

(3) Flitsch, W.; Peters, H. *Tetrahedron Lett.* **1969**, *10*, 1161.

(4) (a) Griesbeck, A. G.; Warzecha, K.-D.; Neudörfl, J.-M.; Görner, H. *Synlett* **2004**, *13*, 2347. (b) Bousquet, T.; Fleury, J.-F.; Daïch, A.; Netchitaïlo, P. *Tetrahedron* **2006**, *62*, 706.



**Figure 1.** Structures of some biologically important 3-methyleneisindolin-1-ones and their derivatives.

isindolinones and aldehydes was developed. However, this method required additional steps to prepare phosphorylated reagents.<sup>2b,5</sup> The lack of an efficient approach to 3-methyleneisindolin-1-ones has stimulated intensive studies for assembly of these heterocycles via metal-catalyzed reactions.<sup>6</sup> The successful examples include a Sonogashira reaction of 2-iodobenzamides with terminal alkynes followed by NaOEt-mediated cyclization,<sup>6c</sup> Heck–Suzuki–Miyaura domino reactions involving ynamides,<sup>6d</sup> and a Sonogashira reaction–carbonylation–hydroamination of 2-bromiodobenzene.<sup>6f</sup> Among them, Kundu's method is quite attractive because both 2-halobenzamides and terminal alkynes are conveniently available.<sup>6c</sup> In Kundu's report,  $(\text{Ph}_3\text{P})_2\text{PdCl}_2$  and CuI were utilized for promoting the cross-coupling reaction, leading to a mixture of open-chain condensation products and heteroannulation products in most cases.<sup>6c</sup> The pure 3-methyleneisindolin-1-ones had to be obtained by treatment of the mixture with NaOEt. In addition, when aliphatic alkynes were employed a mixture of 3-methyleneisindolin-1-ones and isoquinolinones was formed.<sup>6c</sup> These shortcomings limited its application in organic synthesis.

In connection with our project to develop new heterocycle synthesis via amino acid-promoted Ullmann-type reactions,<sup>7–9</sup> we became interested in using CuI/amino acid to catalyze the coupling reaction of 2-bromobenzamides and

terminal alkynes.<sup>8e,10</sup> We were pleased to find after the reaction that substituted 3-methyleneisindolin-1-ones were isolated exclusively. Both aromatic and aliphatic alkynes are compatible with this process, thereby giving an inexpensive, diverse, and convenient approach to assemble these heterocycles. Herein, we disclose our results.

As indicated in Table 1, we initiated our studies by conducting a coupling reaction of *N*-benzyl-2-bromobenzamide

**Table 1.** Coupling of *N*-Benzyl-2-bromobenzamide with Phenylacetylene under Different Conditions<sup>a</sup>

| entry | ligand <sup>b</sup> | base                            | solvent        | yield <sup>c</sup> (%) |
|-------|---------------------|---------------------------------|----------------|------------------------|
| 1     | A                   | K <sub>2</sub> CO <sub>3</sub>  | DMF            | 85                     |
| 2     | A                   | K <sub>2</sub> CO <sub>3</sub>  | <i>i</i> -PrOH | 86                     |
| 3     | A                   | K <sub>2</sub> CO <sub>3</sub>  | DMSO           | 83                     |
| 4     | A                   | K <sub>2</sub> CO <sub>3</sub>  | dioxane        | 50                     |
| 5     | A                   | K <sub>2</sub> CO <sub>3</sub>  | toluene        | <10                    |
| 6     | A                   | K <sub>3</sub> PO <sub>4</sub>  | <i>i</i> -PrOH | 82                     |
| 7     | A                   | Cs <sub>2</sub> CO <sub>3</sub> | <i>i</i> -PrOH | 82                     |
| 8     | B                   | K <sub>2</sub> CO <sub>3</sub>  | <i>i</i> -PrOH | 37                     |
| 9     | C                   | K <sub>2</sub> CO <sub>3</sub>  | <i>i</i> -PrOH | 70                     |
| 10    | D                   | K <sub>2</sub> CO <sub>3</sub>  | <i>i</i> -PrOH | 47                     |
| 11    | E                   | K <sub>2</sub> CO <sub>3</sub>  | <i>i</i> -PrOH | 17                     |
| 12    | F                   | K <sub>2</sub> CO <sub>3</sub>  | <i>i</i> -PrOH | 30                     |
| 13    | no                  | K <sub>2</sub> CO <sub>3</sub>  | <i>i</i> -PrOH | 0                      |

<sup>a</sup> Reaction conditions: **4a** (1 mmol), **5a** (1.5 mmol), CuI (0.1 mmol), ligand (0.3 mmol), solvent (2 mL), 85 °C, 24 h. <sup>b</sup> Key: (A) L-proline; (B) *N,N*-dimethylglycine; (C) pipecolic acid; (D) 2-picolinic acid; (E) ethyl 2-cyclohexanecarboxylate; (F) *N,N*-dimethylethylenediamine. <sup>c</sup> Isolated yield.

with phenylacetylene. It was found that under the catalysis of CuI/L-proline and heating at 85 °C for 24 h in DMF, 3-methyleneisindolin-1-one **6a** was isolated as a single product in 85% yield (entry 1). Similar results were observed when the solvent was changed to *i*-PrOH and DMSO (entries 2 and 3). However, poor yields were obtained if the reaction was carried out in dioxane or toluene (entries 4 and 5). Switching base to K<sub>3</sub>PO<sub>4</sub> and Cs<sub>2</sub>CO<sub>3</sub> had little influence to this reaction (entries 6 and 7). Further investigation revealed that ligand is essential for this process, as evident from the unsatisfactory yields that were observed when *N,N*-dimethylglycine, pipecolic acid, 2-picolinic acid, ethyl 2-cyclohexanecarboxylate, and *N,N*-dimethylethylenediamine were used (entries 8–12) and the fact that no


(5) Couture, A.; Deniau, E.; Grandclaudeon, P.; Lebrun, S. *Synlett* **1997**, 1475.

(6) (a) Uozumi, Y.; Kawasaki, N.; Mori, E.; Mori, M.; Shibasaki, M. *J. Am. Chem. Soc.* **1989**, *111*, 3725. (b) Wu, M.-J.; Chang, L.-J.; Wei, L.-M.; Lin, C.-F. *Tetrahedron* **1999**, *55*, 13193. (c) Kundu, N. G.; Khan, M. W. *Tetrahedron* **2000**, *56*, 4777. (d) Couty, S.; Liegault, B.; Meyer, C.; Cossy, J. *Org. Lett.* **2004**, *6*, 2511. (e) Yao, T.; Larock, R. C. *J. Org. Chem.* **2005**, *70*, 1432. (f) Cao, H.; McNamee, L.; Alper, H. *Org. Lett.* **2008**, *10*, 5281. (g) Sun, C.; Xu, B. *J. Org. Chem.* **2008**, *73*, 7361.

(7) Ma, D.; Cai, Q. *Acc. Chem. Res.* **2008**, *41*, 1450.

(8) (a) Lu, B.; Ma, D. *Org. Lett.* **2006**, *8*, 6115. (b) Lu, B.; Wang, B.; Zhang, Y.; Ma, D. *J. Org. Chem.* **2007**, *72*, 5337. (c) Zou, B.; Yuan, Q.; Ma, D. *Angew. Chem., Int. Ed.* **2007**, *46*, 2598. (d) Zou, B.; Yuan, Q.; Ma, D. *Org. Lett.* **2007**, *9*, 4291. (e) Liu, F.; Ma, D. *J. Org. Chem.* **2007**, *72*, 4844. (f) Chen, Y.; Xie, X.; Ma, D. *J. Org. Chem.* **2007**, *72*, 9329. (g) Chen, Y.; Wang, Y.; Sun, Z.; Ma, D. *Org. Lett.* **2008**, *10*, 625. (h) Wang, B.; Lu, B.; Jiang, Y.; Zhang, Y.; Ma, D. *Org. Lett.* **2008**, *10*, 2761. (i) Yuan, Q.; Ma, D. *J. Org. Chem.* **2008**, *73*, 5159. (j) Guo, L.; Li, B.; Huang, W.; Pei, G.; Ma, D. *Synlett* **2008**, 1833.

**Table 2.** Synthesis of Substituted 3-Methyleneisindolin-1-ones via a Domino Coupling/Additive Cyclization Process from 2-Bromobenzamides and Terminal Alkynes<sup>a</sup>



| entry | X  | Y               | R  | R'                                       | time (h) | product   | yield <sup>b</sup> (%) |
|-------|----|-----------------|--|--|----------|-----------|------------------------|
| 1     | H  | H               | Bn   | 4-MeC <sub>6</sub> H <sub>4</sub>        | 24       | <b>6b</b> | 78                     |
| 2     | H  | H               | Bn   | 4-MeOC <sub>6</sub> H <sub>4</sub>       | 36       | <b>6c</b> | 81                     |
| 3     | H  | H               | Bn   | 3-FC <sub>6</sub> H <sub>4</sub>         | 24       | <b>6d</b> | 80                     |
| 4     | H  | H               | Bn   | BnOCH <sub>2</sub>                       | 36       | <b>6e</b> | 74                     |
| 5     | H  | H               | Bn   | <i>n</i> -C <sub>5</sub> H <sub>11</sub> | 24       | <b>6f</b> | 55 <sup>c</sup>        |
| 6     | H  | NO <sub>2</sub> | Bn   | Ph                                       | 48       | <b>6g</b> | 57 <sup>d</sup>        |
| 7     | H  | OMe             | Bn   | Ph                                       | 24       | <b>6h</b> | 78                     |
| 8     | Cl | H               | Bn   | Ph                                       | 36       | <b>6i</b> | 70                     |
| 9     | H  | H               | 4-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>  | Ph                                       | 48       | <b>6j</b> | 90                     |
| 10    | H  | H               | 4-MeOC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> | Ph                                       | 48       | <b>6k</b> | 88                     |
| 11    | H  | H               | allyl  | Ph                                       | 48       | <b>6l</b> | 88                     |
| 12    | H  | NO <sub>2</sub> | allyl  | Ph                                       | 36       | <b>6m</b> | 61 <sup>d,e</sup>      |
| 13    | H  | OMe             | allyl  | Ph                                       | 36       | <b>6n</b> | 93                     |
| 14    | H  | H               | C <sub>6</sub> H <sub>5</sub>                      | Ph                                       | 36       | <b>6o</b> | 92                     |
| 15    | H  | H               | H  | Ph                                       | 36       | <b>6p</b> | 70                     |
| 16    | H  | H               | <i>n</i> -C <sub>4</sub> H <sub>9</sub>            | Ph                                       | 48       | <b>6q</b> | 83                     |
| 17    | H  | H               | TBSO(CH <sub>2</sub> ) <sub>2</sub>                | Ph                                       | 48       | <b>6r</b> | 82                     |
| 18    | H  | H               | Et <sub>2</sub> N(CH <sub>2</sub> ) <sub>2</sub>   | 4-HOC <sub>6</sub> H <sub>4</sub>        | 48       | <b>6s</b> | 58 <sup>f,g,h</sup>    |

<sup>a</sup> Reaction conditions: aryl bromide (0.5 mmol), 1-alkyne (0.75 mmol), CuI (0.05 mmol), L-proline (0.15 mmol), *i*-PrOH (1 mL), 85 °C. <sup>b</sup> Isolated yield. <sup>c</sup> The reaction was carried out in DMSO at 100 °C. <sup>d</sup> A mixture of *E*- and *Z*-isomers was determined in a ratio of 5:1. <sup>e</sup> The reaction was carried out in DMSO at 85 °C. <sup>f</sup> The reaction was carried out in DMF at 100 °C. <sup>g</sup> 4-Ethynylphenyl acetate was used as a coupling partner. <sup>h</sup> A mixture of *E*- and *Z*-isomers was determined in a ratio of 14:1.

desired product was isolated in the absence of a ligand (entry 13). Based on these results, we chose L-proline as the ligand, K<sub>2</sub>CO<sub>3</sub> as the base, and *i*-PrOH as the solvent in the subsequent studies.

The reaction scope and limitations were then examined under the optimized conditions, and the results are summarized in Table 2. Some functionalized arylacetylenes also worked well, giving the corresponding products in good yields (entries 1–3). Gratifyingly, when aliphatic alkynes were employed, only 3-methyleneisindolin-1-ones **6e** and **6f** were isolated (entries 4 and 5). This result indicated that

the CuI-mediated additive cyclization of the amide moiety to the triple bond took place in a 5-*exo* manner exclusively, which is different with base or Lewis acid-mediated additive cyclization (both 5-*exo* and 6-*endo* attacks were observed).<sup>6c</sup> In cases of 1-heptyne as a substrates, the reaction was sluggish at 85 °C, and good conversion was obtained by increasing the reaction temperature and using DMSO as a solvent (entry 5). This result illustrated that aliphatic alkynes are less reactive substrates for this transformation.

We next explored the possibility of changing the substituents at the aromatic ring of aryl bromides and were pleased to find that both electron-rich and electron-deficient aryl bromides were compatible with these conditions (entries 6–8). It seemed that electron-rich aryl bromides are more reactive than electron-deficient ones, as longer reaction times were required for nitro- and chloro-substituted aryl bromides.

Further investigations demonstrated that a wide range of N-substituents were tolerated in this process, which include functionalized benzyl groups, allyl, phenyl, proton, and simple and functionalized alkyl groups (entries 9–18). These additional functional groups would allow further conversion to more useful heterocycles. For example, **6r** could be transformed into a known tetracyclic intermediate for alkaloid

(9) For related studies from other groups, see: (a) Cuny, G.; Bois-Choussy, M.; Zhu, J. *J. Am. Chem. Soc.* **2004**, *126*, 14475. (b) Klapars, A.; Parris, S.; Andersom, K. W.; Buchwald, S. L. *J. Am. Chem. Soc.* **2004**, *126*, 3529. (c) Altenhoff, G.; Glorius, F. *Adv. Synth. Catal.* **2004**, *346*, 1661. (d) Yang, T.; Lin, C.; Fu, H.; Jiang, Y.; Zhao, Y. *Org. Lett.* **2005**, *7*, 4781. (e) Hu, H.; Li, C. *Org. Lett.* **2006**, *8*, 5365. (f) Martin, R.; Rodríguez, R.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2006**, *45*, 7079. (g) Rivero, M. R.; Buchwald, S. L. *Org. Lett.* **2007**, *9*, 973. (h) Zheng, N.; Buchwald, S. L. *Org. Lett.* **2007**, *9*, 4749. (i) Jiang, B.; Tian, H.; Huang, Z.-G.; Xu, M. *Org. Lett.* **2008**, *10*, 2737. (j) Bao, W.; Liu, Y.; Lv, X.; Qian, W. *Org. Lett.* **2008**, *10*, 3899. (k) Coste, A.; Toumi, M.; Wright, K.; Razafimahaleo, V.; Couty, F.; Marrot, J.; Evano, G. *Org. Lett.* **2008**, *10*, 3841. (l) Yao, P.-Y.; Zhang, Y.; Hsung, R. P.; Zhao, K. *Org. Lett.* **2008**, *10*, 4275. (m) Chen, L.; Shi, M.; Li, C. *Org. Lett.* **2008**, *10*, 5285. (n) Hasegawa, K.; Kimura, N.; Arai, S.; Nishida, A. *J. Org. Chem.* **2008**, *73*, 6363. (o) Viire, R. D.; Evindar, G.; Batey, R. A. *J. Org. Chem.* **2008**, *73*, 3452. (p) Li, Z.; Sun, H.; Jiang, H.; Liu, H. *Org. Lett.* **2008**, *10*, 3263. (q) Ohta, Y.; Chiba, H.; Oishi, S.; Fujii, N.; Ohno, H. *Org. Lett.* **2008**, *10*, 3535. (r) Liu, X.; Fu, H.; Jiang, Y.; Zhao, Y. *Angew. Chem., Int. Ed.* **2009**, *48*, 348. (s) Martins, A.; Lemouzy, S.; Lautens, M. *Org. Lett.* **2009**, *11*, 181.


(10) Ma, D.; Liu, F. *Chem. Commun.* **2004**, 1934.

synthesis,<sup>4b</sup> while acylation of **6s** would provide compound **1** that has local anesthetic activity superior to that of procaine.<sup>1b</sup>

For stereochemistry of the present transformation, it was found that in most cases only *Z*-isomers were determined. However, in cases of two nitro-substituted aryl bromides (entries 6 and 12) and *N,N*-diethylethylene-substituted aryl bromide (entry 18) as substrates, *E*-isomers were isolated as a major product. Their geometry was established via NOESY studies and comparison of the analytical data with the known compounds. The reason for this difference is not clear yet, but this result indicated that some substituents have strong influence to the transition state in additive cyclization step.

For reaction process, we envisaged that coupling of aryl bromides with 1-alkynes should occur first as indicated in Scheme 1. The coupling products **A** might undergo depro-


Scheme 1



tonation of the amide moiety and subsequent addition to C—C triple bond under the action of the copper complex (as intermediates **B**) to afford substituted 3-methyleneisoindolin-1-ones. Indeed, in some cases the simple Sonogashira reaction products **A** could be isolated if the reaction time was shortened.

To further illustrate the synthetic usage of the present method, we developed a formal synthesis of lennoxamine as depicted in Scheme 2. Aryl bromide **7** was prepared from 2,3-dimethoxybenzoic acid in two steps based on known procedure, which was coupled with 1-alkyne **8** under the catalysis of CuI/L-proline in DMF at 110 °C to afford heteroannulation product **9** in 57% yield as a mixture of *Z*- and *E*-isomers (about 5.5:1 determined by <sup>1</sup>H NMR). Upon hydrogenation, this mixture was transformed into isoindoli-

Scheme 2



none **10** in quantitative yield. This intermediate has been converted into lennoxamine in two steps by Cossy and co-workers.<sup>2d</sup>

In conclusion, we have developed a CuI/L-proline-catalyzed coupling/additive cyclization domino reaction process for assembling substituted 3-methyleneisoindolin-1-ones from 2-bromobenzamides and terminal alkynes. Variation of *N*-substituents, aromatic ring, and methylene part was proven possible. Thus, this process gives a straightforward access to substituted 3-methyleneisoindolin-1-ones and may find applications in the synthesis of complex natural products or designed bioactive compounds.

**Acknowledgment.** We are grateful to the Chinese Academy of Sciences, National Natural Science Foundation of China (Grant Nos. 20621062 and 20572119) and Ministry of Science & Technology (Grant No. 2009CB940900) for their financial support.

**Supporting Information Available:** Experimental procedures and copies of <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra for all new products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL9000922