


ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: D. Gryko, M. Giedyk, K. Goliszewska and K. óProinsias, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC07363D.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Journal Name

COMMUNICATION

Cobalt(I)-catalysed CH-alkylation of terminal olefins, and beyond

Maciej Giedyk,^a Katarzyna Goliszewska,^a Keith ó Proinsias^a and Dorota Gryko^{a*}Received 00th January 20xx,
Accepted 00th January 20xx


DOI: 10.1039/x0xx00000x

www.rsc.org/

Cobalester, a natural nontoxic vitamin B₁₂ derivative was found to catalyse unusual olefinic sp² C-H alkylation with diazo reagents as a carbene source instead of the expected cyclopropanation.

Diazocarbonyl compounds are important and useful reagents in organic chemistry due to their ease of integration into numerous chemical transformations under mild conditions.¹ Particularly, they are utilised in cyclopropanation, direct insertion into sp³ C-H, aromatic sp² C-H, and sp C-H bonds as well as heteroatom-H bonds.²⁻⁷ Until recently, C-H functionalisation was mainly achieved with Rh and Cu catalysts,^{1d} as well as other group 11 metals.⁸ Nowadays, methodologies avoiding the use of harmful and toxic reagents have attracted a lot of attention with Fe,⁹ Co,¹⁰ and Cr¹¹ being recognised as valuable catalysts.


'supernucleophilicity' of the Co(I) species (B₁₂S) and radical character of the Co(II) form (B₁₂r) have been investigated in numerous chemical reactions, such as dehalogenation, additions to activated alkenes, functional group migration, dimerisation, cyclopropanation and cyclopropane ring opening.¹³ In 2004 Chen and Zhang reported that the reaction of styrenes with ethyl diazoacetate (EDA) in the presence of (HO)Cbl led to cyclopropane derivatives (Scheme 1).¹⁴



Scheme 1 Functionalisation of olefins with EDA.

Their proposed mechanism suggests that the reaction involves the formation of Co(II) species that when reacted with EDA generates C-centred alkylcobalamin radical. On the other hand, carbene complexes, formed via reaction of diazo compounds with metal salts, including cobalt, are known to trap nucleophiles and insert into C-H bonds.^{1a} On this basis we questioned whether and how diazo reagents, those of electrophilic character, would react with 'supernucleophilic' B₁₂S. We presumed that the reaction should provide alkylcobalamins that upon heating or light irradiation would generate radicals that could either disproportionate, abstract H', or self-couple.

Our recent investigations into vitamin B₁₂-catalysed dimerisation of benzyl bromides led us to use newly developed cobalester (**2**, Cble(III)) as a catalyst for the designed reaction.¹⁵ Firstly, EDA was reacted with cobalester **2** in the presence of a reducing system (Zn/NH₄Cl) to prove the formation of hypothesised alkylcobalamin. Indeed, mass spectrometry analysis (ESI MS) of the reaction mixture showed a peak at 1506.8 corresponding to the desired alkylcobalester (see ESI). Following this observation, EDA was reacted with 1,1-diphenylethene (**3**) in the presence of Cble (**2**) under light irradiation (Scheme 3, Table 1, entry 3). A mixture of two compounds formed; product **4a** and ester **5a** with the reduced

Fig. 1 Structure of vitamin B₁₂ and cobalester.

In this regard, vitamin B₁₂ (**1**, Cbl, cobalamin, Fig. 1), being a natural, environmentally benign Co-complex bears advantages over other transition metal compounds. Its catalytic properties are inherently connected with the ability to form alkyl derivatives via Co(II) and Co(I) reduced forms.¹² To date,


^a Institute of Organic Chemistry Polish Academy of Science, Kasprzaka 44/52, 01-224 Warsaw, (Poland), e-mail: dorota.gryko@icho.edu.pl, homepage: http://ww2.icho.edu.pl/gryko_group.

[†] Electronic Supplementary Information (ESI) available: details of optimization and mechanistic studies, and spectroscopic data for all new compounds. See DOI: 10.1039/x0xx00000x

COMMUNICATION

Journal Name

double bond. Surprisingly, ^1H NMR analysis revealed no evidence of cyclopropanation occurring.



Scheme 2 Cble-catalysed reaction of 1,1-diphenylethene (**3a**) with EDA: 1,1-diphenylethene (**3a**, 0.5 mmol), EDA (3 equiv.), Cble (**2**, 1 mol%), Zn (3 equiv.), NH_4Cl (3.4 equiv.), MeCN (2.5 mL), LED light, 18 h.

So far, the only examples representing functionalisation of double bonds are reactions of enones with diazocompounds reported by Ryu et al.,¹⁶ Noels,¹⁷ and Maruoka,¹⁸ and that of α,β -unsaturated oximes leading to pyridine *N*-oxides.^{6e} Our reaction represents the first example of alkylation of electron-rich double bonds with diazo reagents affording Heck-type products.

Subsequently, we systematically preformed optimisation of the reaction conditions with regard to solvent, catalyst loading, Zn, NH_4Cl , and time to favourably form the Heck-product. Among solvents allowing for the reduction of catalyst **2**, MeCN was found to be the solvent of choice (see ESI). Control experiments revealed that in the absence of light, at room temperature the extrusion of nitrogen from the reaction slowed and products **4a** and **5a** (1.4 : 1) formed in only 34% combined yield (Table 1, entry 1). An increase in the reaction's temperature to 42 °C (equivalent to photoreactor) improved the yield but decreased the selectivity (entry 4). Generally, photocatalytic reactions assured better selectivity (entries 2, 3), hence reactions were performed under light irradiation.

Table 1 Optimisation: the influence of light and temperature.^a

Entry	Light	T [°C]	Conversion [%]	Yield [%]	4a : 5a ratio
1	-	RT	63	34	1.4 : 1
2	LED	15	78	28	2.6 : 1
3	LED	-	94	74	2.7 : 1
4	-	42	94	65	1.3 : 1
5	-	60	86	68	1.5 : 1

^aReaction conditions: 1,1-diphenylethene (0.5 mmol), EDA (3 equiv.), Zn (3 equiv.), NH_4Cl (3.4 equiv.), Cble **2** (1 mol%), MeCN (2.5 mL), 18 h.

The product ratio strongly depended on the amount of Zn and NH_4Cl used (Table 2). Almost in all cases yields remained high while the selectivity increased with an increase in Zn amount and with a decrease in the amount of NH_4Cl reaching a **4a** : **5a** ratio of 13 : 1 (Table 2, compare entries 1 to 3 and 3 to 4).

Table 2 Optimisation: the influence of zinc and ammonium chloride.^a

Entry	Cble 2 [mol%]	Zn [equiv.]	NH_4Cl [equiv.]	Yield [%]	4a : 5a ratio
1	1	3	3.4	74	2.7 : 1
2	5	3	3.4	91	6 : 1
3	2	6	3.4	91	5 : 1
4	2	6	1.1	79	13 : 1

^aReaction conditions: 1,1-diphenylethene (0.5 mmol), EDA (3 equiv.), Zn, NH_4Cl , Cble **2** (2 mol%), MeCN (5 mL), 18 h light: 2x300 Lm LED warm light

Having established effective reaction conditions (Table entry 3) we next focused on examining the scope of various compounds with terminal double bonds (Table 3).

Table 3 Substrate scope of C-H functionalisation of olefins.^a

Entry	R ₁	R ₂	R ₃	t [h]	5	Yield [%]
1	Ph	Ph	Et	18	a	91
2	Ph	Ph	<i>t</i> -Bu	18	b	67
3	Ph	Ph	(CH ₂) ₃ Ph	18	c	60
4 ^b	Ph	Ph	CH ₂ Ph	18	d	71 ^b
5	Ph	CH ₂ C(CH ₃) ₂ Ph	Et	18	e	57
6	Ph	H	Et	6	f	44
7	4-FC ₆ H ₄	H	Et	6	g	1
8	4-ClC ₆ H ₄	H	Et	6	h	22 ^c
9	4-BrC ₆ H ₄	H	Et	6	i	2
10	4-MeOC ₆ H ₄	H	Et	6	j	38
11	3-MeOC ₆ H ₄	H	Et	6	k	23
12	2-MeOC ₆ H ₄	H	Et	6	l	18
13	4-CO ₂ MeC ₆ H ₄	H	Et	6	m	0
14	-CH ₂ CH ₂ CH ₂ CH ₂ -		Et	18	n	73
15 ^d	C ₆ H ₄ -2-Me	C ₆ H ₄ -2-Me	Et	18	o	68 ^d


^aReaction conditions: alkene (**3**, 0.5 mmol), EDA (3 equiv.), Cble (**2**) (2 mol%), Zn (6 equiv.), NH_4Cl (3.4 equiv.) MeCN (5 mL), light. ^bBenzyl moiety was cleaved during hydrogenation step, giving 4,4-diphenylbutanoic acid **5d** as a product.

^cProduct **5f** was isolated instead of expected halogenated compounds.

^dSubsequent hydrogenation was not performed as only unsaturated product was formed in the reaction


As the scope of the study focused on overall yield, the subsequent hydrogenation was performed exclusively furnishing the saturated product **5**. In styrene series **3f-l** (entries 6-13) only electron rich substrates **3f**, **3j-l** were reactive enough to furnish 1,2-disubstituted olefins as a mixture of *trans*- and *cis*-diastereoisomers. These results are in contrast to Hwang and Ryu work reporting sp^2 C-H bond functionalisation of only electron-deficient olefins - cycloenones.¹⁶ In the case of halogenated styrene derivatives, except the fluorinated one underwent dehalogenation giving compound **5f** in low yield (entry 8-9). It is worth to note that in all cases cyclopropanation was not observed. Furthermore an efficient reaction took place for methylenecyclopentane (**3n**, entry 10) and 1-methyl-2-[1-(2-methylphenyl)vinyl]benzene (**3o**, entry 15), with no saturation being detected. Presumably, once product **5o** is formed, the sterically hindered double bond has a limited access to the cobalt centre. As far as diazocompounds are concerned only monosubstituted derivatives furnished desired products. Furthermore, we envisaged that this approach demonstrated the potential for targeting electron rich double bonds such as those present in enol ethers, enamides, or vinyl sulfides. Enol ethers were examined by Davies and Ren, however only functionalisation at the allylic position with aryldiazoacetates was reported, while EDA predominantly afforded cyclopropanation products (Scheme 3).¹⁹ Later, they were found to react with

aryldiazoacetates in the presence $\text{Rh}(\text{OAc})_4$ to afford 1,4-dicarbonyl compounds via cyclopropanation and subsequent ring opening.²⁰



Scheme 3 Reactivity of enol ethers towards diazo reagents.

Our Cble-catalysed reaction allowed for functionalisation of not only these challenging substrates but also enamides **13** and vinyl sulfides **15** (Scheme 4). Although the list of substrates is not exhaustive it shows the potential of the reaction and some of its limitations.



Scheme 4 Scope of Cble-catalysed functionalisation of sp^2 CH bonds with EDA. ^a After treatment with TFA. ^b After hydrogenation.

In order to establish a mechanism, several experiments were conducted. Firstly, the formation of cyclopropane followed by ring opening was considered as an alternative pathway regardless of the fact that cyclopropane was not detected in our reaction but it could still act as a possible intermediate.^{13,14} Hence, ethyl 2,2-diphenylcyclopropane-carboxylate²¹ was subjected to the reaction conditions, but it was found to remain intact even after 24 h. On this basis the mechanism involving a cyclic intermediate was ruled out. Secondly, a series of Co(II) and Co(III) complexes were tested in the absence of a reducing agent (Table 4, entries 3, 4, 5) and in all cases compounds **4a**, **5a** were not observed contrary to reactions with the addition of a reductant (entry 6 and 7), lending support to our Co(I) theory. Thirdly, the ESI MS spectra of the crude reaction mixture (after 1 h) showed peaks at 1419.9 and 1718.9 suggesting species **17** and proving alkylcobalester **20** (see ESI; in MeOH, ligand exchange occurred) thus indicating the efficient formation of radical **A** and its fast addition to the double bond. In the absence of olefin **3a**, dimerization and reduction of diazoacetate occurred. The concentration of free radicals in the reaction mixture was too low to detect directly by EPR spectroscopy. Thus, the spin

trapping experiment was performed by adding a spin trap (phenyl-*N*-*t*-butyl-nitrone) to the reaction mixture (see ESI).

Table 4 C-H-functionalisation of terminal olefins – mechanistic studies.^a

Entry	Catalyst [mol%]	Light ^b	Reductant ^c	Yield [%] ^d	4a : 5a ratio
1	Cble/1	LED	+	74	2.7 : 1
2	-	LED	-	-	-
3	Co(acac) ₃ /2	LED	-	-	-
4 ^e	Cby/2	LED	-	-	-
5	Cble/2	LED	-	-	-
6	Co(acac) ₃ /2	LED	+	25	1 : 2.1
7	CoCl(Ph ₃ P) ₃ /2	LED	+	4	1 : 1
8	Co(TPP)	LED	+	23	15 : 1

^a Conditions: 1,1-diphenylethene (0.5 mmol), EDA (1.5 mmol, 3 equiv.), catalysts MeCN, light, 18 h. ^b LED – lamps. ^c Reductant: Zn, NH₄Cl. ^d Isolated yields. ^e Cby(II).


The formation of 'spin adducts' was ascertained supporting the radical mechanism. Moreover, addition of TEMPO as a radical scavenger, to the reaction mixture rendered it inactive.

The proposed mechanism for the cobalester-catalysed C-H insertion is shown in Scheme 5. Based on the experimental evidences, it is assumed to proceed via carbene transfer mechanism involving Co(I) species, contrary to previous reports. Firstly, cobalester (**2**) is reduced to the catalytically active Co(I) form **17** (I) which reacts with EDA affording alkylcobalester(III) **18** (II) (see ESI). In this reaction, addition of a proton present in complex **18** originates from NH₄Cl, which was clearly demonstrated by deuteration experiment (see ESI). Upon heating or light irradiation homolytic bond cleavage furnishes radical **A** (III). The subsequent, key C-C bond formation step involves the radical addition to the electron rich olefin (**IV**). Newly formed radical **B** can either recombine with Cble(II) species **19** possessing a radical character (**V**), abstract proton or disproportionate leading to by-product **5** (**VI**). Intermediate **20**, which presence was detected using ESI MS (see ESI) undergoes dehydrocobaltation generating desired product **4a** and hydridocobalester (Co(III)-H) **21** (VII) that upon reduction regenerates the catalyst (VIII) or reacts with olefin **4a** to produce unwanted by-product **5a** (XI).

Conclusions

In conclusion, we have demonstrated the use of vitamin B₁₂ derivative **2** as a catalyst for the reaction of electron rich double bonds with EDA leading to unexpected insertion into olefinic sp^2 C-H bond instead of the usual cyclopropanation. The reaction is likely to proceed via a radical pathway involving the Co(I) species, allowing the formation of alkylcobalester intermediate, thus Cble **2** can be considered a 'reversible carrier' for an alkyl.

Our findings complement known reactivity of diazo compounds which now can be inserted not only into sp^3 and sp^2 C-H bonds but also into sp^2 olefinic bonds. The developed methodology is not limited to electron-rich olefins but allows functionalisation of enol ethers, enamides, and vinyl sulfides. Moreover, this work gives solid foundation for future endeavors into diazo/vitamin B₁₂ reactions, releasing its stronghold on cyclopropanation reactions. Further studies



Scheme 5 Proposed mechanism.

on expanding this methodology are currently being undertaken and will be reported in due course.

The authors acknowledge the generous support from the National Science Centre: grant OPUS no. 2012/07/B/ST5/02016 and ETIUDA 2014/12/T/ST5/00113.

Notes and references

- 1 (a) T. Ye, M. A. McKervey, *Chem. Rev.* 1994, **94**, 1091. (b) Z. Zhang, J. Wang, *Tetrahedron*, 2008, **64**, 6577. (c) H. M. L. Davies, R. E. J. Beckwith, *Chem. Rev.* 2003, **103**, 2861. (d) M. P. Doyle, R. Duffy, M. Ratnikov, L. Zhou, *Chem. Rev.* 2010, **110**, 704. (e) Å. Kaupang, T. Bonge-Hansen, J. Beilstein, *Org. Chem.* 2013, **9**, 1407. (f) F. Hu, Y. Xia, Ch. Ma, Y. L. Zhang, J. Wang, *Chem. Commun.* 2015, **51**, 7986. (g) H. M. L. Davies, D. Morton, *Chem. Soc. Rev.* 2011, **40**, 1857. (h) A. C. B. Burtoloso, R. M. P. Dias, B. Bernardim, *Acc. Chem. Res.* 2015, **48**, 921. (i) T. Hashimoto, K. Maruoka, *Bull. Chem. Soc. Jpn.* 2013, **86**, 1217.
- 2 Cyclopropanation reactions: (a) M. Bordeaux, V. Tyagi, R. Fasan, *Angew. Chem. Int. Ed.* 2015, **54**, 1744. (b) A. Pons, H. Beucher, P. Ivashkin, G. Lemonnier, T. Poisson, A. B. Charette, P. Jubault, X. Pannecoucke, *Org. Lett.* 2015, **17**, 1790. (c) X. Xu, S. Zhu, X. Cui, L. Wojtas, X. P. Zhang, *Angew. Chem. Int. Ed.* 2013, **52**, 11857. (d) M. Otte, P. F. Kuijpers, O. Troeppner, I. Ivanović-Burmazović, J. H. Reek, B. de Bruin, *Chem. Eur. J.* 2014, **20**, 4880. (e) H. Lebel, J. -F. Marcoux, C. Molinaro, A. B. Charette, *Chem. Rev.* 2003, **103**, 977.
- 3 sp functionalisation: (a) F. Ye, X. Ma, Q. Xiao, H. Li, Y. Zhang, J. J. Wang, *Am. Chem. Soc.* 2012, **134**, 5742. (b) X. Zhao, Y. Hang, J. Wang, *Chem. Commun.* 2012, **48**, 10162. (c) A. Suarez, G. C. Fu, *Angew. Chem. Int. Ed.* 2004, **43**, 3580.
- 4 sp³ functionalisation: (a) A. R. Reddy, C. -Y. Zhou, Z. Guo, J. Wei, Ch. -M. Che, *Angew. Chem. Int. Ed.* 2014, **53**, 1. (b) H. Davies, J. Mannig, *Nature* 2008, **451**, 417. (c) X. Cui, X. Xu, L.-M. Jin, L. Wojtas, X. P. Zhang, *Chem. Sci.* 2015, **6**, 1219. (d) Ch. J. Flynn, C. J. Elcoate, S. E. Lawrence, A. R. Maguire, *J. Am. Chem. Soc.* 2010, **132**, 1184.
- 5 heteroatom-H functionalisation: (a) G. Sreenilayam, R. Fasan, *Chem. Commun.* 2015, **51**, 1532. (b) V. Tyagi, R. B. Bonn, R. Fasan, *Chem. Sci.* 2015, **6**, 2488. (c) D. Chen, X. Zhang, W.-Y. Qi, B. Xu, M.-H. Xu, *J. Am. Chem. Soc.* 2015, **137**, 5268. (d) H. M. Mbuvi, E. R. Klobukowski, G. M. Roberts, L. K. Woo, *J. Porphyrins Phthalocyanines* 2010, **14**, 284.
- 6 aromatic sp² functionalisation: (a) E. Tayama, M. Ishikawa, H. Iwamoto, E. Hasegawa, *Tetrahedron Lett.* 2012, **53**, 5159. (b)

- M. Delgado-Rebollo, A. Prieto, P. J. Pérez, *Chem. Cat. Chem.* 2014, **6**, 2047. (c) E. Tayama, T. Yanaki, H. Iwamoto, H. Hasegawa, *Eur. J. Org. Chem.* 2010, 6719. (d) H. M. L. Davies, S. J. Hedley, *Chem. Soc. Rev.* 2007, **36**, 1109. (e) Z. Shi, I. Koester, M. Boulfadakis-Arapinis, F. Glorius, *J. Am. Chem. Soc.* 2013, **135**, 12204. (f) H. M. Mbuvi, L. K. Woo, *J. Porphyrins Phtalocyanines* 2009, **13**, 137.
- 7 M. P. Doyle, M. A. McKervey, T. Ye, in *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds: From Cyclopropanes to Ylides*, Wiley, NY, 1998.
- 8 (a) M. M. Díaz-Requejo, P. J. Pérez, *J. Organomet. Chem.* 2005, **690**, 5441. (b) A. Caballero, P. J. Pérez, *J. Organomet. Chem.* 2015, **793**, 108. (c) N. Komine, J. A. Flores, K. Pal, K. C. Caulton, D. Mindiola, *J. Organometallics* 2013, **32**, 3185. (d) R. Gava, A. Olmos, B. Noverges, T. Varea, E. Álvarez, T. K. Belderrain, A. Caballero, G. Asensio, P. J. Pérez, *ACS Catal.* 2015, **5**, 3726. (e) J. A. Flores, N. Komine, K. Pal, B. Pinter, M. Pink, Ch. -H. Chen, K. G. Caulton, D. J. Mindiola, *ACS Catal.* 2012, **2**, 2066.
- 9 example: Y. Ma, D. Zhang, Z. Yan, M. Wag, C. Bian, X. Gao, E. E. Bunel, A. Lei, *Org. Lett.* 2015, **17**, 2174.
- 10 example: (a) D. Zhao, J. H. Kim, L. Stegemann, C. A. Strasser, F. Glorius, *Angew. Chem. Int. Ed.* 2015, **54**, 4508. (b) M. Moselage, N. Sauermann, S. C. Richter, L. Ackermann, *Angew. Chem. Int. Ed.* 2015, **54**, 6354. (c) Y. Chen, K. B. Fields, X. P. Zhang, *J. Am. Chem. Soc.* 2004, **126**, 14718.
- 11 example: N. D. Hahn, M. Nieger, K. H. Dötz, *Eur. J. Org. Chem.* 2004, 1049.
- 12 K. L. Brown, *Chem. Rev.* 2005, **105**, 2075.
- 13 (a) M. Giedyk, K. Goliszewska, D. Gryko, *Chem. Soc. Rev.* 2015, **44**, 3391. (b) Y. Hisaeda, T. Nishioka, Y. Inoue, K. Asada, T. Hayashi, *Coord. Chem. Rev.* 2000, **198**, 21.
- 14 Y. Chen, X. P. Zhang, *J. Org. Chem.* 2004, **69**, 2431.
- 15 M. Giedyk, S. N. Fedosov, D. Gryko, *Chem. Commun.* 2011, **50**, 4674.
- 16 S. I. Lee, B. Ch. Kang, G.-S. Hwang, D. H. Ryu, *Org. Lett.* 2011, **15**, 1428.
- 17 A. F. Noels, J. N. Braham, A. J. Hubert, Ph. Teyssie, *J. Org. Chem.* 1977, **42**, 1527.
- 18 T. Kano, T. Hashimoto, K. Maruoka, *J. Am. Chem. Soc.* 2006, **128**, 2174.
- 19 H. M. L. Davies, P. Ren, *J. Am. Chem. Soc.* 2001, **123**, 2070.
- 20 S. Muthusamy, P. Srinivasan, *Tetrahedron Lett.* 2006, **47**, 6297.
- 21 P. Müller, C. Baud, D. Ené, S. Motallebi, M. P. Doyle, B. P. Brandes, A. B. Dyatkin, M. M. See, *Helv. Chim. Acta* 1995, **78**, 459.
- 22 Cby(II) was used as a cobalester(II) substitute because there is no suitable method for its preparation.