

Discovery of *N*-{*N*-[(3-Cyanophenyl)sulfonyl]-4(*R*)-cyclobutylamino-(*L*)-prolyl}-4-[(3',5'-dichloroisonicotinoyl)amino]-(*L*)-phenylalanine (MK-0668), an Extremely Potent and Orally Active Antagonist of Very Late Antigen-4

Linus S. Lin,^{*,†} Thomas Lanza, Jr.,[†] James P. Jewell,[†] Ping Liu,[†] Carrie Jones,[†] Gerard R. Kieczkowski,[†] Kelly Treonze,[‡] Qian Si,[‡] Salony Manior,[‡] Gloria Koo,[‡] Xinchun Tong,[†] Junying Wang,[†] Anne Schuelke,[†] James Pivnichny,[†] Regina Wang,[§] Conrad Raab,[§] Stella Vincent,[§] Philip Davies,[‡] Malcolm MacCoss,[†] Richard A. Mumford,[‡] and William K. Hagmann[†]

Departments of Medicinal Chemistry, Immunology and Rheumatology, and Drug Metabolism, Merck Research Laboratories, Rahway, New Jersey 07065

Received March 1, 2009

Abstract: Extremely potent very late antigen-4 (VLA-4) antagonists with picomolar, whole blood activity and slow dissociation rates were discovered by incorporating an amino substituent on the proline fragment of the initial lead structure. This level of potency against the unactivated form of VLA-4 was shown to be sufficient to overcome the poor pharmacokinetic profiles typical of this class of VLA-4 antagonists, and sustained activity as measured by receptor occupancy was achieved in preclinical species after oral dosing.

VLA-4^a ($\alpha 4\beta 1$, CD49d/CD29) is a member of the integrin family that is expressed on all lymphocytes.¹ VLA-4 normally exists at a resting state or unactivated state. Although the precise mechanism of activation *in vivo* is not clear, VLA-4 can be activated *in vitro* by divalent cation Mn^{2+} or activating antibodies.² Once activated, VLA-4 binds strongly to ligands including (1) vascular cell adhesion molecule-1 (VCAM-1), which is expressed on activated endothelial cells at sites of inflammation and is produced in response to inflammatory cytokines, and (2) the CS-1 domain of fibronectin (FN) which arises by alternative splicing within a region of FN. The binding interactions between VLA-4 and VCAM-1/FN are proposed to be key mediators of cell–cell and cell–matrix adhesion. As such, these cell adhesion interactions may be essential for the activation, migration, proliferation, and differentiation of lymphocytes during normal and/or pathophysiological processes. Thus, inhibition of VLA-4 may result in a reduction in the migration and/or activation of cell types important to sustaining a prolonged inflammatory response.³ Anti- $\alpha 4$ antibodies and blocking peptides have been reported to be efficacious in a number of animal models of inflammatory diseases, including antigen-induced bronchial hyperresponsiveness,⁴ experimental allergic encephalomyelitis,⁵ adjuvant arthritis,⁶ and spontaneous chronic colitis.⁷




Figure 1. Structure of a clinical VLA-4 antagonist.




Figure 2. Lead structures of tight binding VLA-4 antagonists.

Clinical proof of concept was first achieved with an anti- $\alpha 4$ neutralizing antibody, natalizumab, which has been shown to be effective in human clinical trials of multiple sclerosis (MS) and Crohn's disease (CD) and has been approved by FDA for both indications.^{8,9} At a monthly iv dose of 3 and 6 mg, approximately 80% and 90% saturation, respectively, of $\alpha 4\beta 1$ receptors (receptor occupancy or RO) on peripheral blood lymphocytes was reported.⁸ Subsequently, a small molecule VLA-4 antagonist, valategrast (R411) (Figure 1), was shown to be beneficial for the treatment of asthma in a 12-week phase II clinical trial.^{10–12} The active metabolite of valategrast (RO0270608, Figure 1) was a dual $\alpha 4\beta 1$ – $\alpha 4\beta 7$ antagonist with high affinity for the activated states of both $\alpha 4\beta 1$ and $\alpha 4\beta 7$ integrins, but the affinity was lower and the dissociation was rapid when the receptors were not activated.¹¹

Previously, we identified a tight binding VLA-4 antagonist **1a** (Figure 2) with picomolar affinity on the unactivated form of VLA-4 along with a slow dissociation rate or off-rate.¹³ However, it suffered from poor bioavailability and a short half-life in preclinical species. After iv dosing (0.5 mpk) in rats, only 15% receptor occupancy was observed at 8 h postdosing. On the basis of the reported experience with natalizumab, we hypothesized that a small molecule antagonist would need to achieve $\geq 80\%$ RO at 12–24 h to be effective in the treatment of MS and CD. Therefore, much improvement in whole blood activity and/or pharmacokinetics over **1a** would be necessary to achieve this level of RO. Because of the peptidic nature and molecular weight of this lead, improving pharmacokinetic properties was a significant challenge. However, if a compound possessed very potent whole blood activity with a fast on-rate and also had a very slow off-rate once bound, sustained receptor occupancy might be achievable even after the drug was rapidly eliminated from circulation.

To test this hypothesis, we sought to improve the whole blood activity and the off-rate of lead structure **1a**. Compound **1a** is a carboxylic acid and is highly bound to human plasma proteins

* To whom correspondence should be addressed. Phone: 732-594-8391. Fax: 732-594-5966. E-mail: linus_lin@merck.com.

[†] Department of Medicinal Chemistry.

[‡] Department of Immunology and Rheumatology.


[§] Department of Drug Metabolism.

^a Abbreviations: VLA-4, very late antigen-4; VCAM-1, vascular cell adhesion molecule-1; FN, fibronectin; MS, multiple sclerosis; CD, Crohn's disease; RO, receptor occupancy; LOQ, limit of quantitation; TDI, time dependent inhibition.

Table 2. Receptor Occupancy and Pharmacokinetic Parameters of VLA-4 Antagonists¹⁶

compd	species	dose (mg/kg)	route	RO at X h (%) ^a	X (h) ^b	Cl _p ((mL/min)/kg) ^c	t _{1/2} (h) ^c	t _{min} (h) ^d
9a	rat	0.5	iv	64 ± 2	8	65	1.0	2
9a	dog	0.75	iv	72 ± 6	8	29	0.5	1
9a	monkey	0.75	iv	87 ± 6	8	33	1.1	6
9a	rat	1	po	46 ± 12	8	ND	ND	ND
9a-Et	rat	1	po	62 ± 16	8	ND	ND	ND
9e-Et	rat	1	po	51 ± 5	8	ND	ND	ND
9f	rat	5	po	67 ± 11	12	65	1.0	0
9f	dog	5	po	83 ± 8	12	29	0.5	6
9f	monkey	5	po	88 ± 1.8	8	33	1.1	0
9f-Et	rat	5	po	77 ± 13	12	ND	ND	0
9f-Et	dog	5	po	97 ± 7	12	ND	ND	12
9f-Et	monkey	5	po	95 ± 4	8	ND	ND	0

^a % receptor occupancy at X h. ^b Time when receptor occupancy was measured. ^c Cl_p and plasma half-life were determined after iv dose. ^d Latest time point when **9a** or **9f** could be detected (LOQ ≈ 0.1 nM).

Table 3. Cytochrome P450 Inhibition and clogP of VLA-4 Antagonists¹⁷


compd	X, Y	CYP3A4 IC ₅₀ (μM)	TDI rate (min ⁻¹)	clogP ^a
9e-Et	Cl, Cl	<0.4	0.05	5.5
9f-Et	CN, H	1.7, 2.4	0.04	3.5

^a Calculated by ACD8.1 software.

Compound **9a** and its ethyl ester **9a-Et** were then studied via oral dosing in rats. Receptor occupancies of 46 ± 12% and 62 ± 16%, respectively, were observed at 8 h postdosing (1 mg/kg). The ethyl ester of **9e** (**9e-Et**) was also studied by oral dosing (1 mg/kg) in rats and afforded comparable RO (51 ± 5% at 8 h postdosing). Although still moderate in comparison with clinical observations of natalizumab, the level of oral activity of **9a**/**9a-Et** and **9e**/**9e-Et** as assessed by receptor occupancy represented a breakthrough in the search for an orally active small molecule antagonist of the unactivated form of VLA-4.

Because the azetidine ring of **9a**/**9a-Et** was found to be acid labile, **9e**/**9e-Et** was selected for additional characterization toward the identification of a development candidate. Metabolism studies revealed that **9e-Et** is a potent and time-dependent inhibitor (TDI) of CYP3A4 with a TDI rate of 0.05 min⁻¹ (Table 3).¹⁷ The parent acid **9e** also displayed a propensity for bioactivation upon incubation with liver microsomes, and covalent adducts were formed with microsomal proteins after incubation for 1 h (rat, 108 pmol/mg; human, 150 pmol/mg).¹⁸ Because CYP inhibition and covalent adduct formation pose significant risks of drug–drug interactions and idiosyncratic toxicities, respectively,^{19,20} alternative structures were sought. It is well-known that CYP inhibition is often associated with high lipophilicity.²¹ Replacing the 3,5-dichlorophenyl group of **9e-Et** with a 3-cyanophenyl group (**9f-Et**) resulted in significant reduction in CYP3A4 inhibition, while the clogP value decreased from 5.5 to 3.5. Furthermore, **9f** also demonstrated decreased levels of covalent adduct formation when incubated with microsomal preparations (rat, 39 ± 2 pmol/mg; human, 56 ± 2 pmol/mg).

The amino acid **9f** (MK-0668) was a potent antagonist of the unactivated state of VLA-4 (IC₉₀ in whole blood: human

0.13 ± 0.03 nM, dog 0.19 ± 0.01 nM, rhesus 0.21 nM, rat 0.4 ± 0.1 nM; off-rate at 37 °C on human VLA-4, 64 ± 10% bound at 3 h). When evaluated in vivo (5 mg/kg po), **9f-Et** demonstrated sustained RO of 77–97% at 8–12 h postdosing across three species (Table 2). As a comparison, the acid **9f** was also studied and afforded somewhat lower levels of RO (67–88% at 8–12 h postdosing). Analogous to other compounds in the series, **9f**/**9f-Et** displayed poor pharmacokinetic properties in the preclinical species (*F* < 1%, *t*_{1/2} ≤ 1 h). Nevertheless, RO was maintained even after no compound was detectable in the plasma, consistent with slow dissociation rate of **9f** from VLA-4 receptors. If similar PK/PD relationship holds in humans, with a twice-daily oral dosing schedule, **9f** or its prodrug **9f-Et** would be expected to achieve RO coverage comparable to that of intravenously dosed natalizumab.

In summary, extremely potent VLA-4 antagonists with picomolar, whole blood activity and slow off-rates were discovered by incorporating an amino substituent on the proline fragment of the initial lead structure. This level of potency was shown to be sufficient to overcome the poor pharmacokinetic profiles typical of this class of VLA-4 antagonists, and sustained activity as measured by receptor occupancy was achieved after oral dosing.²²

Acknowledgment. We acknowledge the contributions of many members of the Labeled Synthesis Group and the Laboratory of Animal Resources of Merck Research Laboratories.

Note Added after ASAP Publication. The paper was published on May 14, 2009 with typographical errors. The revised version was published on May 20, 2009.

Supporting Information Available: Synthetic procedures and analytical data of VLA-4 antagonists and conditions for all the biological assays. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Hynes, R. O. Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* **1992**, *69*, 11–25.
- (2) Masumoto, A.; Hemler, M. E. Multiple activation states of VLA-4. Mechanistic differences between adhesion to CS1/fibronectin and to vascular cell adhesion molecule-1. *J. Biol. Chem.* **1993**, *268*, 228–234.
- (3) Postigo, A. A.; Teixido, J.; Sanchez-Madrid, F. The alpha 4 beta 1/VCAM-1 adhesion pathway in physiology and disease. *Res. Immunol.* **1993**, *144*, 723–735.
- (4) Abraham, W. M.; Gill, A.; Ahmed, A.; Sielczak, M. W.; Laureda, I. T.; Botinnikova, Y.; Lin, K. C.; Pepinsky, R. B.; Leone, D. R.; Lobb, R. R. A small-molecule, tight-binding inhibitor of the integrin α4β1 blocks antigen-induced airway responses and inflammation in experimental asthma in sheep. *Am. J. Respir. Crit. Care Med.* **2000**, *162*, 603–611.

- (5) Yednock, T.; Cannon, C.; Fritz, L. C.; Sanchez-Madrid, F.; Steinman, L.; Karlin, N. Prevention of experimental autoimmune encephalomyelitis by antibodies against $\alpha 4 \beta 1$ integrin. *Nature* **1993**, *356*, 63–66.
- (6) Seiffge, D. Protective effects of monoclonal antibody to VLA-4 on leukocyte adhesion and course of disease in adjuvant arthritis in rats. *J. Rheumatol.* **1996**, *23*, 2086–2091.
- (7) Podolsky, D. K.; Fiocchi, C. Cytokines, Chemokines, Growth Factors, Eicosanoids and Other Bioactive Molecules in Inflammatory Bowel Disease. In *Inflammatory Bowel Disease*; Kirsner, J. B., Ed.; W. B. Saunders: Chicago, IL, 2000; pp 191–207.
- (8) Miller, D. H.; Khan, O. A.; Sheremata, W. A.; Blumhardt, L. D.; Rice, G. P. A.; Libonati, M. A.; Willmer-Hulme, A. J.; Dalton, C. M.; Miszkiele, K. A.; O'Connor, P. W. For the International Natalizumab Multiple Sclerosis Trial Group. A controlled trial of natalizumab for relapsing multiple sclerosis. *N. Engl. J. Med.* **2003**, *348*, 15–23.
- (9) Ghosh, S.; Goldin, E.; Gordon, F. H.; Malchow, H. A.; Rask-Madsen, J.; Rutgeerts, P.; Vyhnaček, P.; Zádorová, Z.; Palmer, T.; Donoghue, S. For the Natalizumab Pan-European Study Group. Natalizumab for active Crohn's disease. *N. Engl. J. Med.* **2003**, *348*, 24–32.
- (10) Tilley, J. W.; Sidduri, A.; Lou, J. P.; Rossman, P.; Tare, N.; Cavallo, G.; Railkar, A.; Gerber, L.; Frank, K.; Renzetti, L. Identification of *N*-acyl-4-(3-pyridinyl)phenylalanine derivatives and their orally active prodrug esters as dual acting $\alpha 4$ - $\beta 1$ and $\alpha 4$ - $\beta 7$ receptor antagonists. Presented at the 234th National Meeting of the American Chemical Society; Boston, MA, Aug 19–23, 2007; MEDI-426.
- (11) Tilley, J. W. VLA-4 antagonists. *Expert Opin. Ther. Pat.* **2008**, *18*, 841–859.
- (12) Hijazi, Y.; Welker, H.; Dorr, A. E.; Tang, J.-P.; Blain, R.; Renzetti, L. M.; Abbas, R. Pharmacokinetics, safety, and tolerability of R411, a dual $\alpha 4 \beta 1$ – $\alpha 4 \beta 7$ integrin antagonist after oral administration at single and multiple once-daily ascending doses in healthy volunteers. *J. Clin. Pharmacol.* **2004**, *44*, 1368–1378.
- (13) Doherty, G. A.; Yang, G. X.; Borges, E.; Tong, S.; McCauley, E. D.; Treon, K. M.; Van Riper, G.; Pacholok, S.; Si, Q.; Koo, G. C.; Shah, K.; Mumford, R. A.; Hagmann, W. K. *N*-Isonicotinoyl-(L)-4-aminophenylalanine derivatives as tight binding VLA-4 antagonists. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 1891–1895.
- (14) For assay conditions, see Supporting Information.
- (15) For another study demonstrating high level of RO after iv or sc dosing, see the following: Leone, D. R.; Giza, K.; Gill, A.; Dolinski, B. M.; Yang, W.; Perper, S.; Scott, D. M.; Lee, W.-C.; Cornebise, M.; Wortham, K.; Nickerson-Nutter, C.; Chen, L. L.; LePage, D.; Spell, J. C.; Whalley, E. T.; Petter, R. C.; Adams, S. P.; Lobb, R. R.; Pepinsky, R. B. An assessment of the mechanistic differences between two integrin $\alpha 4 \beta 1$ inhibitors, the monoclonal antibody TA-2 and the small molecule BIO5192, in rat experimental autoimmune encephalomyelitis. *J. Pharmacol. Exp. Ther.* **2003**, *305*, 1150–1162.
- (16) For assay conditions, see Supporting Information.
- (17) For assay conditions, see Supporting Information.
- (18) For assay conditions and SAR to reduce covalent binding, see the following: (a) Samuel, K.; Yin, W.; Stearns, R. A.; Tang, Y. S.; Chaudhary, A. G.; Jewell, J. P.; Lanza, T., Jr.; Lin, L. S.; Hagmann, W. K.; Evans, D. C.; Kumar, S. Addressing the metabolic activation potential of new leads in drug discovery: a case study using ion trap mass spectrometry and tritium labeling techniques. *J. Mass Spectrom.* **2003**, *38*, 211–221. (b) Lin, L. S.; Lanza, T. J., Jr.; Jewell, J. P.; Liu, P.; Shah, S. K.; Qi, H.; Tong, X.; Wang, J.; Xu, S.; Fong, T. M.; Shen, C.-P.; Lao, J.; Chen, J.; Shearman, L. P.; Stribling, D. S.; Rosko, K.; Strack, A.; Marsh, D. J.; Feng, Y.; Kumar, S.; Samuel, K.; Yin, W.; Van der Ploeg, L.; Goulet, M. T.; Hagmann, W. K. Discovery of *N*-[(1*S*,2*S*)-3-(4-chlorophenyl)-2-(3-cyanophenyl)-1-methylpropyl]-2-methyl-2-[[5-(trifluoromethyl)pyridin-2-yl]oxy]propanamide (MK-0364), a novel, acyclic cannabinoid-1 receptor inverse agonist for the treatment of obesity. *J. Med. Chem.* **2006**, *49*, 7584–7587.
- (19) The MRL target threshold value for covalent binding was set at 50 pmol·equiv/mg protein. For more detailed discussions, see the following: Evans, D. C.; Watt, A. P.; Deborah, A.; Nicoll-Griffith, D. A.; Baillie, T. A. Drug–protein adducts: an industry perspective on minimizing the potential for drug bioactivation in drug discovery and development. *Chem. Res. Toxicol.* **2004**, *17*, 3–16.
- (20) Riley, R. J.; Grime, K.; Weaver, R. Time-dependent CYP inhibition. *Expert Opin. Drug Metab. Toxicol.* **2007**, *3*, 51–66.
- (21) Gleeson, M. P. Generation of a set of simple, interpretable ADMET rules of thumb. *J. Med. Chem.* **2008**, *51*, 817–834.
- (22) Additional profiling of **9f/9f**-Et will be reported by R. A. Mumford in a separate manuscript.

JM900257B