

Communication

**Pd(II)-Catalyzed Enantioselective #-C(sp)-H  
Functionalizations of Free Cyclopropylmethyamines**

Zhe Zhuang, and Jin-Quan Yu

*J. Am. Chem. Soc.*, **Just Accepted Manuscript** • DOI: 10.1021/jacs.0c04801 • Publication Date (Web): 30 Jun 2020

Downloaded from pubs.acs.org on July 1, 2020

**Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

# Pd(II)-Catalyzed Enantioselective $\gamma$ -C(sp<sup>3</sup>)-H Functionalizations of Free Cyclopropylmethylamines

Zhe Zhuang,<sup>†</sup> and Jin-Quan Yu<sup>\*,†</sup>

<sup>†</sup>Department of Chemistry, The Scripps Research Institute, 10550 N. Torrey Pines Road, La Jolla, California 92037, United States




*Supporting Information Placeholder*

**ABSTRACT:** Prized for their ability to reliably forge stereocenters with precise regiocontrol from simple and abundant starting materials, substrate-directable enantioselective reactions are widely used in modern organic synthesis. As such, enantioselective C(sp<sup>3</sup>)-H functionalization reactions directed by innate functional groups could provide new routes to introduce molecular complexity within the inert hydrocarbon moiety, but so far this approach has been met with little success. While free primary aliphatic amines are common, versatile intermediates in synthesis, they are traditionally unreactive in C(sp<sup>3</sup>)-H activation reactions. Herein we report the Pd-catalyzed enantioselective C(sp<sup>3</sup>)-H functionalization of free aliphatic amines (cyclopropylmethylamines) enabled by a chiral bidentate thioether ligand. This ligand's privileged bidentate coordination mode and thioether motif favor the generation of the requisite mono(amine)-Pd(II) intermediate, thus enabling the enantioselective C-H activation of free amines. The resulting C-Pd(II) species could engage in either Pd(II)/Pd(IV) or Pd(II)/Pd(o) catalytic cycles, enabling access to a diverse range of products through (hetero)arylation, carbonylation and olefination reactions. Consequently, this versatile reactivity offers medicinal chemists a general strategy to rapidly prepare and functionalize biologically relevant amines.

Ligand-controlled enantioselective reactions directed by innate functional groups (e.g. carboxylic acids, amines, and alcohols) comprise a classic toolkit to control the regio- and stereochemical outcomes of organic reactions<sup>1</sup>, as exemplified by the Sharpless's oxidation<sup>2</sup>, Noyori and Knowles's hydrogenation<sup>3,4</sup> reactions of alkenes. While such strategies highlight significant developments in the functionalization of unsaturated moieties, only recently has serious attention been directed toward complementary strategies for the functionalization of saturated hydrocarbons<sup>5</sup>. Despite advances in C-H functionalization directed by weakly coordinating free carboxylic acids<sup>6</sup>, aliphatic amines bearing free NH<sub>2</sub> groups – ubiquitous structural motifs among compounds of pharmaceutical, agrochemical, and societal importance – are less amenable to Pd-catalyzed C-H functionalization reactions<sup>7</sup>. This is due to the formation of stable but unreactive bis(amine)-Pd(II) species, stemming


from the strong metal binding properties of the amine<sup>8</sup>. Additionally, free amines are susceptible to oxidative degradation through  $\beta$ -hydride elimination and *N*-substitution reactions towards electrophiles under the conditions of C-H activation. In this context, extensive studies have been focused on the design of preinstalled<sup>9</sup> or transient<sup>10</sup> bidentate directing groups (DGs) on account of their ability to accelerate cyclometalation and suppress  $\beta$ -hydride elimination (Scheme 1A). Despite the elegance of DG or TDG strategies, these reactions are difficult to render enantioselective due to the persistent racemic background reactions driven by strong coordination, as well as the lack of available coordination sites for a chiral ligand. Bidentate coordination from the substrate also confines the reactivity to a Pd(II)/Pd(IV) catalytic cycle, owing to the lack of a vacant coordination site to engage the coupling partner in a Pd(II)/Pd(o) cycle. Most importantly, the requirement for substrate DGs precludes the use of native amino groups to direct C-H activation, which affords superior atom and step economies in the context of protecting-group-free synthesis. In fact, reported free aliphatic amine-directed C-H functionalization reactions are largely limited to bulky amines; a bulky group at the  $\alpha$  position of the free amine is required to favor dissociation of bis(amine)-Pd(II) complexes through steric repulsion, hamper oxidative degradation by blocking the  $\beta$  position, as well as weaken the nucleophilicity of amino group (Scheme 1A)<sup>11</sup>.

**Scheme 1. C(sp<sup>3</sup>)-H Functionalization of Aliphatic Amines**

**A Strategies towards C(sp<sup>3</sup>)-H activation of aliphatic amines****B Native amine strategy via ligand acceleration****C Ligand-enabled enantioselective C(sp<sup>3</sup>)-H functionalization of free amines**

In light of our approach using the combination of monodentate practical DGs and bidentate ligands to enable enantioselective C-H activation reactions<sup>12</sup>, we were particularly interested in applying this concept to the otherwise challenging chemical feedstocks such as aliphatic amines (Scheme 1B). We reasoned that due to the bidentate chelation mode of the ligand, unreactive bis(amine)-Pd(II) species would readily dissociate as a result of strong coordination from the ligand, generating the desired reactive mono(amine)-Pd(II) intermediate for cyclopalladation. The bidentate ligand also creates a rigid framework that is ideal for the transfer of chiral information to the substrate during C-H activation. Moreover, the privileged amidate group (*N*-acyl) on the ligand can not only participate as an internal base to facilitate C(sp<sup>3</sup>)-H activation over  $\beta$ -hydride elimination but is also labile enough to generate a vacant coordination site for coupling partners used in Pd(II)/Pd(o) chemistry. Herein we report the design of a thioether-based chiral bidentate ligand that enables Pd(II)-catalyzed enantioselective C-H functionalization reactions of free cyclopropylmethamines (Scheme 1C). Arylation via a Pd(II)/Pd(IV) catalytic cycle afforded chiral free aliphatic amines bearing aryl groups, including aza-heterocycles, at the  $\gamma$ -position. Preliminary results also showed that this catalyst enables the olefination and carbonylation reactions of primary and secondary free amines through a Pd(II)/Pd(o) catalytic cycle, providing valuable  $\gamma$ -olefinated free amines and  $\gamma$ -lactams respectively.

**Table 1. Ligand Investigation for  $\gamma$ -C(sp<sup>3</sup>)-H Arylation of Free Cyclopropylmethamine<sup>a,b</sup>**




<sup>a</sup>Conditions: **1a** (0.1 mmol), Pd(TFA)<sub>2</sub> (10 mol%), ligand (**L**) (10 mol%), 4-iodotoluene **2a** (2.0 equiv.), Ag<sub>2</sub>O (2.0 equiv.), HFIP (0.2 mL), 90 °C, 12 h. <sup>b</sup>The yields were determined by <sup>1</sup>H NMR analysis of the crude product using CH<sub>2</sub>Br<sub>2</sub> as the internal standard. The er values of the corresponding Boc-protected amine **3a** were determined on the SFC system using commercially available chiral columns.

Our previous research has disclosed two enantioselective C(sp<sup>3</sup>)-H functionalization reactions using a monodentate triflamide DG: the  $\gamma$ -C(sp<sup>3</sup>)-H arylation of cyclopropylmethamine enabled by a mono-*N*-protected amino acid (MPAA) ligand<sup>13</sup> and the  $\gamma$ -C(sp<sup>3</sup>)-H cross-coupling of 3-aminopentane enabled by acetyl-protected aminomethyl oxazoline ligands (APAO) ligand<sup>14</sup>. Bidentate picolinamide-directed benzylic C(sp<sup>3</sup>)-H arylation has also been reported using a BINOL phosphoric acid ligand with limited scope<sup>15</sup>. Due to the widespread presence of cyclopropyl moieties among drug and agrochemical molecules<sup>16</sup>, we selected free cyclopropylmethamine **1a** as a model substrate for our initial investigations. Upon extensive studies of the reaction conditions in the absence of ligand, a combination of Pd(TFA)<sub>2</sub>, 4-iodotoluene **2a**, Ag<sub>2</sub>O, and 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP) solvent produced a 6% <sup>1</sup>H NMR yield of the desired *cis*- $\gamma$ -arylation product **3a** with a considerable amount of the degradation product cyclopropane aldehyde. To test our hypothesis, chiral bidentate ligands that previously promoted enantioselective C(sp<sup>3</sup>)-H activation, including MPAA **L1**<sup>5c,13</sup>, *O*-methylhydroxamic acid **L2**<sup>12c</sup>, dimethylamine **L3**<sup>6b</sup>, oxazoline **L4**<sup>12b,14</sup>, and quinoline **L5**<sup>12a</sup>, were investigated under the standard reaction conditions (Table 1). To our delight, these ligands generally promoted the arylation reaction, delivering the arylation product **3a** in 30% to 40% yield with MPAA ligand **L1** being the superior ligand (40% yield, 85:15 er). However, further modifications of the ligand structure and optimization of the reaction conditions failed to improve the yield and er. Therefore, we turned our attention to using thioethers as soft  $\sigma$ -donors, believing that thioethers might facilitate dissociation of bis(amine)-Pd(II) complexes, as well as stabilize the Pd catalysts through their strong coordination. The racemic ligand **L7** developed for the olefination of free carboxylic acids<sup>17</sup> significantly improved the reaction to 57% yield (see Supporting Information Table S5). Through systematic modifications of the backbone of the thioether ligand we found that introduction of an isopropyl group gave optimal reactivity with 97.5:2.5 er in 68% yield (see Supporting Information Table S5). Notably, this optimal

thioether ligand **L6** could be easily synthesized in two steps with a single recrystallization in 92% yield from the commercially available Evans' oxazolidinone chiral auxiliary (see Supporting Information preparation of ligand section).

**Table 2. Free Cyclopropylmethylamine Scope for  $\gamma$ -C(sp<sup>3</sup>)-H arylation<sup>a,b</sup>**




<sup>a</sup>Conditions: **1** (0.1 mmol), Pd(TFA)<sub>2</sub> (10 mol%), **L6** (10 mol%), 4-iodotoluene **2a** (3.0 equiv.), Ag<sub>2</sub>O (2.0 equiv.), HFIP (0.2 mL), 90 °C, 12 h. <sup>b</sup>Isolated yields of the corresponding Bz or Boc-protected amines. The er values were determined on the SFC system using commercially available chiral columns.

With the optimal ligand and conditions in hand, we next explored the scope of aliphatic amines (Table 2). The  $\gamma$ -arylated products were isolated in the form of *tert*-butoxycarbonyl (Boc) or benzoyl (Bz)-protected amines for their ease of purification and analysis. Cyclopropylmethylamines (**1b**–**1g**) bearing various aliphatic chains at the  $\beta$  position, including cyclobutanes (**1e**), cyclopentanes (**1f**), and cyclohexanes (**1g**), were all compatible with the optimal protocol, affording the arylation products (**3b**–**3g**) in yields ranging from modest to good (54% to 70%) with excellent ers (greater than 97.5:2.5). A range of functionalities such as fluoro (**3i**), trifluoromethyl (**3j**),

tetrahydropyran (**3h**), benzyl (Bn)-protected hydroxyl (**3k**), methoxy (**3l**), and phenolic ether (**3m**) were all well tolerated, and products were obtained in moderate yields (45% to 66%) and high ers (greater than 96:4). Phenyl groups (**3a** and **3n**–**3s**) were compatible under the current conditions, and remained intact despite the potentially reactive aryl C(sp<sup>2</sup>)-H or benzylic C(sp<sup>3</sup>)-H bonds. A range of substituents on the aryl ring from electron-withdrawing (fluoro and chloro) (**3n**, **3o**, and **3p**) to electron-donating (methoxy) (**3q**) groups were also well tolerated, providing arylation products in good yields (up to 73%) and excellent ers (up to 99:1 er). The less reactive unsubstituted cyclopropylmethylamine **1t** also reacted in a synthetically useful yield (42%) and correspondingly high er (96.5:3.5). Extension of this desymmetrization protocol to isobutylamine and 3-pentanamine also afforded desired arylation products in good yields and moderate ers (see Supporting Information Table S9). However, cyclobutylmethylamine substrates were not reactive under these conditions.

Next, we surveyed the scope of aryl and heteroaryl iodides using **1a** as the model substrate (Table 3).  $\gamma$ -Arylation of **1a** with simple iodobenzene **2b** resulted in good yield (71%) and excellent er (97.5:2.5). A wide range of *para* substituents on the aryl iodides from electron-donating (Me and isopropyl) to electron-withdrawing (CF<sub>3</sub>, ester, and halide) groups were well tolerated, delivering  $\gamma$ -arylated products (**3a** and **4c**–**4g**) with high ers (greater than 96:4 er) in moderate to good yields (46% to 72%). *Meta*-substituted aryl iodides bearing methyl or coordinating OCF<sub>3</sub> and NO<sub>2</sub> groups consistently provided useful yields (51% to 68%) and ers greater than 94.5:5.5 (**4h**–**4j**). Indeed, the above products featured additional reactive groups such as esters (in **4e**) and halogens (chloro in **4g**) that could serve as useful synthetic handles for subsequent derivatizations. Given the ubiquity of heteroaromatics, especially pyridine-containing heterocycles, in small molecule drug discovery<sup>18</sup>, we examined the reactivity of aza-heteroaryl iodides under the aforementioned conditions. Although heteroaryl iodides were incompatible with the previously reported MPAA-enabled arylation of triflamide<sup>13</sup>, we reasoned that when switching to a strongly coordinating thioether ligand, catalyst poisoning by coordination of these aza-heterocycles might be dramatically suppressed. While unsubstituted 4-iodopyridine **2t** failed to give any desired arylation product under the standard conditions, we were pleased to find that a broad range of 2-substituted iodopyridines (**2k**–**2r**) proceeded smoothly with excellent ers (up to 97.5:2.5). A variety of functional groups including fluoro (**4k**), chloro (**4l** and **4m**), bromo (**4n**), methoxy (**4o**), and CF<sub>3</sub> (**4p**–**4r**) were compatible with the optimal conditions, yielding valuable *cis*- $\gamma$ -heteroaryl cyclopropylmethylamines. Electron-rich 5-iodoindole **2s** was also a suitable coupling partner, affording product **4s** in moderate yield (51%) with excellent er (96.5:3.5).

**Table 3. Aryl and Heteroaryl Iodide Scope for  $\gamma$ -C(sp<sup>3</sup>)-H Arylation<sup>a,b</sup>**




<sup>a</sup>Conditions: **1a** (0.1 mmol),  $\text{Pd}(\text{TFA})_2$  (10 mol% or 15 mol% for heteroaryl iodides), **L6** (10 mol% or 15 mol% for heteroaryl iodides), aryl or heteroaryl iodide **2** (3.0 equiv.),  $\text{Ag}_2\text{O}$  (2.0 equiv.), HFIP (0.2 mL), 90 °C, 12 h. <sup>b</sup>Isolated yields of the corresponding Boc-protected amines. The er values were determined on the SFC system using commercially available chiral columns.

To test the feasibility of extending this Pd(II)/thioether catalyst system to a Pd(II)/Pd(0) catalytic cycle, we embarked on the enantioselective carbonylation and olefination of free primary and secondary aliphatic amines (Table 4). Reported works on C(sp<sup>3</sup>)-H carbonylation<sup>19</sup> and olefination<sup>20</sup> are largely limited to DG strategies, and no asymmetric version has been achieved to date. Using  $\text{Mo}(\text{CO})_6$  as a nonhazardous and air-stable solid source of CO, carbonylation of primary amine **1a** delivered bicyclic  $\gamma$ -lactam **5a** in a synthetically useful yield (45%) and moderate er (89.5:10.5). A range of secondary amines (**1u**–**1x**) were also reactive using the optimal protocol, affording the valuable cyclopropane fused pyrrolidones (**5b**–**5e**) in moderate yields (up to 63%) and good ers (up to 97:3 er). Olefination of primary amines (**1a** and **1i**)

and secondary amines (**1u**–**1w**) with pentafluorostyrene proceeded smoothly, delivering the  $\gamma$ -olefinated aliphatic amines (**6a**–**6e**) with good to excellent ers (94:6 to 98.5:1.5). Under current conditions, other olefins such as acrylate and styrene failed to deliver desired olefination product.

**Table 4. Free Primary and Secondary Cyclopropylmethylamine Scope for  $\gamma$ -C(sp<sup>3</sup>)-H Carbonylation and Olefination<sup>a,b</sup>**



<sup>a</sup>Conditions for carbonylation: **1** (0.1 mmol),  $\text{Pd}(\text{OAc})_2$  (15 mol%), **L6** (15 mol%),  $\text{Mo}(\text{CO})_6$  (0.3 equiv.),  $\text{Ag}_2\text{O}$  (2.0 equiv.), NaOAc (1.0 equiv.), HFIP (0.1 mL), 90 °C, 12 h. Conditions for olefination: **1** (0.1 mmol),  $\text{Pd}(\text{OAc})_2$  (15 mol%), **L6** (15 mol%), pentafluorostyrene (3.0 equiv.),  $\text{Ag}_2\text{O}$  (2.0 equiv.), HFIP (0.1 mL), 90 °C, 6 h. <sup>b</sup>Isolated yields for secondary amines or isolated yields of the corresponding Boc-protected amines for primary amines. The er values were determined on the SFC system using commercially available chiral columns.

In summary, we have realized effective and general Pd-catalyzed enantioselective C(sp<sup>3</sup>)-H functionalizations of free aliphatic amines enabled by a single Pd(II) catalyst bearing a bidentate chiral thioether ligand. Both Pd(II)/Pd(IV) and Pd(II)/Pd(0) catalytic cycles are compatible with this chiral ligand, as demonstrated by (hetero)arylation, carbonylation and olefination reactions respectively. A broad range of chiral  $\gamma$ -(hetero)aryl,  $\gamma$ -olefinated free amines and  $\gamma$ -lactams could be easily accessed without installing exogenous directing groups.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Full experimental details and characterization of new compounds (PDF)

## AUTHOR INFORMATION



**Corresponding Author**

\*yuzoo@scripps.edu

**ORCID**

Zhe Zhuang: 0000-0001-6679-0496

Jin-Quan Yu: 0000-0003-3560-5774

**Notes**

The authors declare no competing financial interests.

**ACKNOWLEDGMENT**

We gratefully acknowledge The Scripps Research Institute and NIH (NIGMS, R01GM084019) for financial support. Z. Z. thanks Alastair N. Herron for proofreading.

**REFERENCES**

- (1) Hoveyda, A. H.; Evans, D. A.; Fu, G. C. Substrate-directable chemical reactions. *Chem. Rev.* **1993**, *93*, 1307–1370.
- (2) Katsuki, T.; Sharpless, K. B. The first practical method for asymmetric epoxidation. *J. Am. Chem. Soc.* **1980**, *102*, 5974–5976.
- (3) Noyori, R. Asymmetric catalysis: science and opportunities (Nobel lecture). *Angew. Chem., Int. Ed.* **2002**, *41*, 2008–2022.
- (4) Knowles, W. S. Asymmetric hydrogenations (Nobel lecture). *Angew. Chem., Int. Ed.* **2002**, *41*, 1998–2007.
- (5) For reviews, see: (a) Giri, R.; Shi, B.-F.; Engle, K. M.; Maugel, N.; Yu, J.-Q. Transition metal-catalyzed C–H activation reactions: diastereoselectivity and enantioselectivity. *Chem. Soc. Rev.* **2009**, *38*, 3242–3272. (b) Newton, C. G.; Wang, S.-G.; Oliveira, C. C.; Cramer, N. Catalytic enantioselective transformations involving C–H bond cleavage by transition-metal complexes. *Chem. Rev.* **2017**, *117*, 8908–8976. (c) Shao, Q.; Wu, K.; Zhuang, Z.; Qian, S.; Yu, J.-Q. From Pd(OAc)<sub>2</sub> to chiral catalysts: the discovery and development of bifunctional mono-N-protected amino acid ligands for diverse C–H activation reactions. *Acc. Chem. Res.* **2020**, *53*, 833–851.
- (6) (a) Shi, B.-F.; Zhang, Y.-H.; Lam, J. K.; Wang, D.-H.; Yu, J.-Q. Pd(II)-catalyzed enantioselective C–H olefination of diphenylacetic acids. *J. Am. Chem. Soc.* **2010**, *132*, 460–461. (b) Shen, P.-X.; Hu, L.; Shao, Q.; Hong, K.; Yu, J.-Q. Pd(II)-catalyzed enantioselective C(sp<sup>3</sup>)-H arylation of free carboxylic acids. *J. Am. Chem. Soc.* **2018**, *140*, 6545–6549. (c) Giri, R.; Maugel, N.; Li, J.-J.; Wang, D.-H.; Breazzano, S. P.; Saunderson, L. B.; Yu, J.-Q. Palladium-catalyzed methylation and arylation of sp<sup>3</sup> and sp<sup>3</sup> C–H bonds in simple carboxylic acids. *J. Am. Chem. Soc.* **2007**, *129*, 3510–3511. (d) Zhuang, Z.; Yu, J.-Q. Lactonization as a general route to β-C(sp<sup>3</sup>)-H functionalization. *Nature* **2020**, *577*, 656–659.
- (7) For reviews, see: (a) Daugulis, O.; Roane, J.; Tran, L. D. Bidentate, monoanionic auxiliary-directed functionalization of carbon–hydrogen bonds. *Acc. Chem. Res.* **2015**, *48*, 1053–1064. (b) He, G.; Wang, B.; Nack, W. A.; Chen, G. Syntheses and transformations of α-amino acids via palladium-catalyzed auxiliary directed sp<sup>3</sup> C–H functionalization. *Acc. Chem. Res.* **2016**, *49*, 635–645.
- (8) Ryabov, A. D. Mechanisms of intramolecular activation of C–H bonds in transition-metal complexes. *Chem. Rev.* **1990**, *90*, 403–424.
- (9) (a) Zaitsev, V. G.; Shabashov, D.; Daugulis, O. Highly regioselective arylation of sp<sup>3</sup> C–H bonds catalyzed by palladium acetate. *J. Am. Chem. Soc.* **2005**, *127*, 13154–13155. (b) He, G.; Chen, G. A practical strategy for the structural diversification of aliphatic scaffolds through the palladium-catalyzed picolinamide-directed remote functionalization of unactivated C(sp<sup>3</sup>)-H Bonds. *Angew. Chem. Int. Ed.* **2011**, *50*, 5192–5196. (c) Chan, K. S. L.; Wasa, M.; Chu, L.; Laforteza, B. N.; Miura, M.; Yu, J.-Q. Ligand-enabled cross-coupling of C(sp<sup>3</sup>)-H bonds with arylboron reagents via Pd(II)/Pd(0) catalysis. *Nat. Chem.* **2014**, *6*, 146–150.
- (10) (a) Liu, Y.; Ge, H. Site-selective C–H arylation of primary aliphatic amines enabled by a catalytic transient directing group. *Nat.*

*Chem.* **2017**, *9*, 26–32. (b) Wu, Y.; Chen, Y.-Q.; Liu, T.; Martin, D. E.; Yu, J.-Q. Pd-catalyzed γ-C(sp<sup>3</sup>)-H arylation of free amines using a transient directing group. *J. Am. Chem. Soc.* **2018**, *140*, 10363–10367. (c) Li, B.; Seth, K.; Niu, B.; Pan, L.; Yang, H.; Ge, H. Transient-ligand-enabled *ortho*-arylation of five-membered heterocycles: facile access to mechanochromic materials. *Angew. Chem., Int. Ed.* **2018**, *57*, 3401–3405. (d) Li, B.; Lawrence, B.; Li, G.; Ge, H. Ligand-controlled direct γ-C–H arylation of aldehydes. *Angew. Chem., Int. Ed.* **2020**, *59*, 3078–3082.

(11) (a) Lazareva, A.; Daugulis, O. Direct palladium-catalyzed *ortho*-arylation of benzylamines. *Org. Lett.* **2006**, *8*, 5211–5213. (b) McNally, A.; Haffemayer, B.; Collins, B. S. L.; Gaunt, M. J. Palladium-catalysed C–H activation of aliphatic amines to give strained nitrogen heterocycles. *Nature* **2014**, *510*, 129–133. (c) Calleja, J.; Pla, D.; Gorman, T. W.; Domingo, V.; Haffemayer, B.; Gaunt, M. J. A steric tethering approach enables palladium-catalysed C–H activation of primary amino alcohols. *Nat. Chem.* **2015**, *7*, 1009–1016. (d) Chen, K.; Wang, D.; Li, Z.-W.; Liu, Z.; Pan, F.; Zhang, Y.-F.; Shi, Z.-J. Palladium catalyzed C(sp<sup>3</sup>)-H acetoxylation of aliphatic primary amines to γ-amino alcohol derivatives. *Org. Chem. Front.* **2017**, *4*, 2097–2101.

(12) (a) Chen, G.; Gong, W.; Zhuang, Z.; Andrá, M. S.; Chen, Y.-Q.; Hong, X.; Yang, Y.-F.; Liu, T.; Houk, K. N.; Yu, J.-Q. Ligand-accelerated enantioselective methylene C(sp<sup>3</sup>)-H bond activation. *Science* **2016**, *353*, 1023–1027. (b) Wu, Q.-F.; Shen, P.-X.; He, J.; Wang, X.-B.; Zhang, F.; Shao, Q.; Zhu, R.-Y.; Mapelli, C.; Qiao, J. X.; Poss, M. A.; Yu, J.-Q. Formation of α-chiral centers by asymmetric β-C(sp<sup>3</sup>)-H arylation, alkenylation, and alkynylation. *Science* **2017**, *355*, 499–503. (c) Xiao, K.-J.; Lin, D. W.; Miura, M.; Zhu, R.-Y.; Gong, W.; Wasa, M.; Yu, J.-Q. Palladium(II)-catalyzed enantioselective C(sp<sup>3</sup>)-H activation using a chiral hydroxamic acid ligand. *J. Am. Chem. Soc.* **2014**, *136*, 8138–8142.

(13) Chan, K. S. L.; Fu, H.-Y.; Yu, J.-Q. Palladium(II)-catalyzed highly enantioselective C–H arylation of cyclopropylmethylamines. *J. Am. Chem. Soc.* **2015**, *137*, 2042–2046.

(14) Shao, Q.; Wu, Q.-F.; He, J.; Yu, J.-Q. Enantioselective γ-C(sp<sup>3</sup>)-H activation of alkyl amines via Pd(II)/Pd(0) catalysis. *J. Am. Chem. Soc.* **2018**, *140*, 5322–5325.

(15) Wang, H.; Tong, H. R.; He, G.; Chen, G. An enantioselective bidentate auxiliary directed palladium-catalyzed benzylic C–H arylation of amines using a BINOL phosphate ligand. *Angew. Chem., Int. Ed.* **2016**, *55*, 15387–15391.

(16) Rubin, M.; Rubina, M.; Gevorgyan, V. Transition metal chemistry of cyclopropenes and cyclopropanes. *Chem. Rev.* **2007**, *107*, 3117–3179.

(17) Zhuang, Z.; Yu, C.-B.; Chen, G.; Wu, Q.-F.; Hsiao, Y.; Joe, C. L.; Qiao, J. X.; Poss, M. A.; Yu, J.-Q. Ligand-enabled β-C(sp<sup>3</sup>)-H olefination of free carboxylic acids. *J. Am. Chem. Soc.* **2018**, *140*, 10363–10367.

(18) Chen, Y.-Q.; Wang, Z.; Wu, Y.; Wisniewski, S. R.; Qiao, J. X.; Ewing, W. R.; Eastgate, M. D.; Yu, J.-Q. Overcoming the limitations of γ- and δ-C–H arylation of amines through ligand development. *J. Am. Chem. Soc.* **2018**, *140*, 17884–17894.

(19) (a) Wang, C.; Zhang, L.; Chen, C.; Han, J.; Yao, Y.; Zhao, Y. Oxalyl amide assisted palladium-catalyzed synthesis of pyrrolidones via carbonylation of γ-C(sp<sup>3</sup>)-H bonds of aliphatic amine substrates. *Chem. Sci.* **2015**, *6*, 4610–4614. (b) Wang, P.-L.; Li, Y.; Wu, Y.; Li, C.; Lan, Q.; Wang, X.-S. Pd-catalyzed C(sp<sup>3</sup>)-H carbonylation of alkylamines: a powerful route to γ-lactams and γ-amino acids. *Org. Lett.* **2015**, *17*, 3698–3701. (c) Hernandez, E.; Villalva, J.; Martínez, Á. M.; Alonso, I.; Rodríguez, N.; Arrayás, R. G.; Carretero, J. C. Palladium-catalyzed carbonylative cyclization of amines via γ-C(sp<sup>3</sup>)-H activation: late-stage diversification of amino acids and peptides. *ACS Catal.* **2016**, *6*, 6868–6882.

(20) Jiang, H.; He, J.; Liu, T.; Yu, J.-Q. Ligand-enabled γ-C(sp<sup>3</sup>)-H olefination of amines: en route to pyrrolidines. *J. Am. Chem. Soc.* **2016**, *138*, 2055–2059.

