

Trisubstituted (*E*)-Alkene Dipeptide Isosteres as β -Turn Promoters in the Gramicidin S Cyclodecapeptide Scaffold


Jingbo Xiao,[†] Bernard Weisblum,[‡] and Peter Wipf^{*,†}

Department of Chemistry and Center for Chemical Methodologies & Library Development, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, and Pharmacology Department, University of Wisconsin Medical School, 1300 University Avenue, Madison, Wisconsin 53706

pwipf+@pitt.edu

Received July 18, 2006

ABSTRACT



A concise synthesis of a gramicidin S analogue with trisubstituted (*E*)-alkene dipeptide isostere (TEADI) replacements at both D-Phe-Pro positions was realized. Conformational analysis demonstrated that TEADIs can serve as type II β -turn promoters in a cyclic scaffold and successfully mimic a proline residue.

Peptides demonstrate a wide range of diverse physiological properties as hormones, enzyme inhibitors, growth promoters, signaling pathway modulators, antimicrobial agents, etc.¹ To overcome the limited bioavailability of peptides,² we are studying the synthesis and pharmacological evaluation of bioisosteric replacements of the amide bond.³ The relatively rigid trisubstituted (*E*)-alkene dipeptide isosteres ($\psi[(E)-C(R)=CH]$, $R \neq H$, TEADIs) maintain ω -angle planarity and represent useful structural surrogates of hydrolytically labile amide bonds.⁴ In addition, TEADIs were found to have potential as β -turn promoters in acyclic sequences in our previous studies.⁵ In a continuation of this work, we have now started to introduce these building blocks into biologically active cyclic peptide sequences as surrogates of the

powerful turn-inducing D-Phe-Pro sequence⁶ and evaluate their potential as β -turn promoters and modulators of biological and metabolic properties.⁷

Since its discovery in 1942,⁸ the cyclodecapeptide antibiotic gramicidin S (GS, *cyclo*-[(Val-Orn-Leu-D-Phe-Pro)₂])

(3) (a) Wipf, P.; Fritch, P. C. *J. Org. Chem.* **1994**, 59, 4875. (b) Henninger, T.; Wipf, P. In *Methods in Molecular Biology: Peptidomimetics*; Walker, J. M., Kazmierski, W. M., Eds.; Humana Press: Totowa, 1999; Vol. 23, pp 125–136. (c) Mu, Y.; Stephenson, C. R. J.; Kendall, C.; Saini, S. P. S.; Toma, D.; Ren, S.; Cai, H.; Strom, S. C.; Day, B. W.; Wipf, P.; Xie, W. *Mol. Pharmacol.* **2005**, 68, 403. (d) Wipf, P.; Werner, S.; Woo, G. H. C.; Stephenson, C. R. J.; Walczak, M. A. A.; Coleman, C. M.; Twining, L. A. *Tetrahedron* **2005**, 61, 11488. (e) Levinson, N.; Hinman, R.; Patil, A.; Stephenson, C. R. J.; Werner, S.; Woo, G. H. C.; Xiao, J.; Wipf, P.; Lynch, K. W. *RNA* **2006**, 12, 925. (f) Wipf, P.; Xiao, J.; Jiang, J.; Belikova, N. A.; Tyurin, V. A.; Fink, M. P.; Kagan, V. E. *J. Am. Chem. Soc.* **2005**, 127, 12460.

(4) (a) Tamamura, H.; Koh, Y.; Ueda, S.; Sasaki, Y.; Yamasaki, T.; Aoki, M.; Maeda, K.; Watai, Y.; Arikuni, H.; Otaka, A.; Mitsuya, H.; Fujii, N. *J. Med. Chem.* **2003**, 46, 1764. (b) Yang, H.; Sheng, X. C.; Harrington, E. M.; Ackermann, K.; Garcia, A. M.; Lewis, M. D. *J. Org. Chem.* **1999**, 64, 242. (c) Oishi, S.; Miyamoto, K.; Niida, A.; Yamamoto, M.; Ajito, K.; Tamamura, H.; Otaka, A.; Kuroda, Y.; Asai, A.; Fujii, N. *Tetrahedron* **2006**, 62, 1416.

(5) (a) Wipf, P.; Henninger, T. C.; Geib, S. J. *J. Org. Chem.* **1998**, 63, 6088. (b) Wipf, P.; Henninger, T. C. *J. Org. Chem.* **1997**, 62, 1586.

[†] University of Pittsburgh.

[‡] University of Wisconsin Medical School.

(1) (a) Kazmierski, W. M.; Kenakin, T. P.; Gudmundsson, K. S. *Chem. Biol. Drug Des.* **2006**, 67, 13. (b) Clark, R. J.; Fischer, H.; Dempster, L.; Daly, N. L.; Rosengren, K. J.; Nevin, S. T.; Meunier, F. A.; Adams, D. J.; Craik, D. J. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, 102, 13767. (c) Ladner, R. C.; Sato, A. K.; Gorzelany, J.; De Souza, M. *Drug Discuss. Today* **2004**, 9, 525.

(2) Conradi, R. A.; Hilgers, A. R.; Ho, N. F. H.; Burton, P. S. *Pharm. Res.* **1992**, 9, 435.

has served as an inspiration for the design of antibacterial agents and antimicrobial peptides and as a model system for conformational mimicry.⁹ GS is therefore a particularly significant target for the evaluation of alkene peptide isosteres as surrogates for the type II' β -turn inducing sequence.

The D-Phe-Pro reverse turn is a critical feature of the rigid amphipathic antiparallel β -pleated sheet conformation of GS.^{10,11} In earlier work,⁷ we were able to replace the Leu-D-Phe peptide bond in GS with a $\psi[(E)\text{-C}(\text{CF}_3)=\text{CH}]$ isostere with minimal perturbation of the secondary structure and biological activity, whereas the corresponding $\psi[(E)\text{-C}(\text{CH}_3)=\text{CH}]$ isostere failed to maintain the β -pleated sheet conformation according to CD and NMR analyses. A different situation presents itself when the D-Phe-Pro peptide bond is replaced. Because the D-Phe carbonyl group is not involved in intramolecular H-bonding or dipolar interactions, a $\psi[(E)\text{-C}(\text{CH}_3)=\text{CH}]$ surrogate should be as effective as a trifluoromethylated congener in conformationally preorganizing the chain. The restricted backbone rotation imposed by the A^{1,3}-strain across the trisubstituted alkene and, to a lesser extent, by the A^{1,2}-strain experienced by substituents attached to the alkene should be sufficient for both methyl and trifluoromethyl groups to impose the reverse turn. Furthermore, neither alkene isostere provides an NH hydrogen bond donor group that can lead to the stabilization of γ -turns or other competitive backbone folding patterns that would interfere with the desired β -turn motif. These properties lead, theoretically, to a close match of isostere and D-Phe-Pro features (Figure 1).^{5b} We now report an experimental confirmation of this hypothesis.




Figure 1. GS and its analogue with $\psi[(E)\text{-C}(\text{CH}_3)=\text{CH}]$ peptide bond surrogates.

Because of the lability of phenylacetaldimines, the sulfinyl adduct **3**¹² was employed in the organometallic allylation


(6) Wadhvani, P.; Afonin, S.; Ieronimo, M.; Buerck, J.; Ulrich, A. S. *J. Org. Chem.* **2006**, *71*, 55.

(7) Xiao, J.; Weisblum, B.; Wipf, P. *J. Am. Chem. Soc.* **2005**, *127*, 5742.

(8) Gauze, G. F.; Brazhnikova, M. G. *Am. Rev. Soviet Med.* **1944**, *2*, 134.

(9) See: Grotenbreg, G. M.; Buizert, A. E. M.; Llamas-Saiz, A. L.; Spalburg, E.; Van Hoof, P. A. V.; De Neeling, A. J.; Noort, D.; Van Raaij, M. J.; Van Der Marel, G. A.; Overkleeft, H. S.; Overhand, M. *J. Am. Chem. Soc.* **2006**, *128*, 7559, and references therein.

Scheme 1. Synthesis of Boc-D-Phe- $\psi[(E)\text{-C}(\text{CH}_3)=\text{CH}]$ -Pro-Val-OMe



reaction (Scheme 1).¹³ Using our hydrozirconation/ $\text{Zr} \Rightarrow \text{Zn}$ transmetalation methodology,^{14,15} the alkenylzinc species derived from the chiral internal alkyne **2**^{13b} was added to **3**, affording the allylic amide **4** in 64% yield as a ~1:1 mixture of diastereomers. Deprotection of the TBDPS group with TBAF provided the primary alcohol **5**. The two diastereomers could not be separated after conversion of **5** to the corresponding acetates;^{3f,7} however, a two-step oxidation with Dess–Martin periodinane,^{16,17} followed by coupling with valine methyl ester, provided pseudotripeptides **6** and **7** which were separated by preparative C₁₈ RP HPLC.¹⁸

Saponification of pseudotripeptide **7** followed by fragment coupling with dipeptide H-Orn(Cbz)-Leu-OMe in the presence of EDC as a coupling reagent afforded the pseudopentapeptide **8** in 96% yield over two steps. We initially envisioned a one-pot dimerization–cyclization of **8**; however, this approach resulted exclusively in the formation of cyclized pseudopentapeptide. In contrast, the stepwise coupling proceeded smoothly to give the pseudodecapeptide **9** in excellent yield. Saponification of **9** and stepwise removal of the Boc protecting group followed by macrolactamization afforded the desired bis-Cbz-protected GS analogue **10** in 50% yield after preparative C₁₈ RP HPLC purification (Scheme 2).

The chemical shifts of all amide protons in **10** were assigned using a combination of COSY, NOESY, HMQC,

(10) (a) Yamada, K.; Unno, M.; Kobayashi, K.; Oku, H.; Yamamura, H.; Araki, S.; Matsumoto, H.; Katakai, R.; Kawai, M. *J. Am. Chem. Soc.* **2002**, *124*, 12684. (b) Doi, M.; Fujita, S.; Katsuya, Y.; Sasaki, M.; Taniguchi, T.; Hasegawa, H. *Arch. Biochem. Biophys.* **2001**, *395*, 85.

(11) (a) Kondejewski, L. H.; Farmer, S. W.; Wishart, D. S.; Hancock, R. E. W.; Hodges, R. S. *Int. J. Pept. Protein Res.* **1996**, *47*, 460. (b) Kondejewski, L. H.; Farmer, S. W.; Wishart, D.; Kay, C. M.; Hancock, R. E. W.; Hodges, R. S. *J. Biol. Chem.* **1996**, *271*, 25261.

(12) Côté, A.; Boezio, A. A.; Charette, A. B. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5405.

(13) (a) Wipf, P.; Xiao, J.; Geib, S. J. *Adv. Synth. Catal.* **2005**, *347*, 1605. (b) Wipf, P.; Xiao, J. *Org. Lett.* **2005**, *7*, 103.


(14) (a) Wipf, P.; Jahn, H. *Tetrahedron* **1996**, *52*, 12853. (b) Wipf, P.; Kendall, C. *Top. Organomet. Chem.* **2004**, *8*, 1.

(15) (a) Wipf, P.; Kendall, C.; Stephenson, C. R. J. *J. Am. Chem. Soc.* **2001**, *123*, 5122. (b) Wipf, P.; Kendall, C.; Stephenson, C. R. J. *J. Am. Chem. Soc.* **2003**, *125*, 761.

(16) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155.

(17) Wipf, P.; Kim, Y.; Goldstein, D. M. *J. Am. Chem. Soc.* **1995**, *117*, 11106.

Scheme 2. Fragment Condensation and Synthesis of Pseudo-cyclodecapeptide **10**




and HMBC data sets collected in DMSO-*d*₆ at 338 K because some amide ¹H NMR signals were obscured at 298 K. Variable-temperature NMR was applied to probe the conformation of **10** in solution and determine the level of intramolecular hydrogen bonding (Table 1). Temperature

Table 1. Temperature Shift Coefficients ($\Delta\delta/\Delta T$) of Amide Protons in DMSO-*d*₆ (ppb/K) for **10** and **Cbz₂GS**

	Orn-NH	Leu-NH	Orn- δ -NH	Val-NH	D-Phe-NH
10	-4.8	-1.9	-6.3	-1.8	-5.9
Cbz₂GS	-6.7	-3.8	-8.6	-2.2	-11.8

shift coefficients for **10** were in close agreement with the values for bis-Cbz-protected GS (**Cbz₂GS**).⁷ The NH shifts of Leu and Val residues in **10** showed small temperature

(18) The structural assignment of **6** and **7** was based on the ozonolysis product **12**, which was identical to the methylation product obtained from Boc-D-Phe-OH (**11**) by chiral HPLC coinjection (Pace, R. D.; Kabalka, G. W. *J. Org. Chem.* **1995**, *60*, 4838).



coefficients of -1.9 and -1.8 ppb/K, respectively, whereas Orn-NH and D-Phe-NH were solvent exposed, thus indicating a hydrogen bonding array typical for an antiparallel β -pleated sheet conformation.¹⁹ Furthermore, NOESY spectra showed transannular Leu-NH-Val-NH and Val-NH-D-Phe-H α contacts for **10** in agreement with that found for **Cbz₂GS** (Figure 2). The resulting ten-membered intramolecular H-bonding




Figure 2. Observed NOEs for **10** and **Cbz₂GS**.

interaction between the valine amide NH and the carbonyl group of the leucine residue is typical for a type II' β -turn.²⁰

Further confirmation of the close match between the secondary structures of **10** and **Cbz₂GS** was provided by circular dichroism (CD) spectra in EtOH (Figure 3). The




Figure 3. CD spectra of **10** and **Cbz₂GS** in EtOH.

strong negative band at ~205–225 nm and a shoulder in the region of 225–235 nm for both **10** and **Cbz₂GS** are consistent with a combination of a type II' β -turn and a

β -sheet conformation in these compounds.²¹ This result, along with the data from NOESY and variable-temperature NMR experiments, further confirmed that the methyl (*E*)-alkene dipeptide isostere replacements at the former D-Phe-Pro positions strongly promoted the archetypical architecture of the parent peptide, gramicidin S. An MMFF-minimized structure for **1** that is in agreement with all experimental data is shown in Figure 4.²²





Figure 4. Stereoview of the minimized structure of **1** derived from an MMFF conformational search algorithm.

Our previous biological studies showed that free amine functions on the ornithine side chains were necessary to retain the antibacterial, antifungal, and hemolytic activities of GS.⁷ The Cbz protecting groups in **10** were successfully removed by hydrogenolysis in the presence of 10% Pd/C in a 0.02 M HCl/MeOH solution, without concomitant reduction of the trisubstituted (*E*)-alkene moieties (Scheme 3). As expected, the hydrochloride salt of **1** also exhibited functional mimicry of the natural product, with an MIC of $\sim 20 \mu\text{g/mL}$ against *Bacillus subtilis*, and thus was equipotent with GS hydrochloride (MIC $\sim 15 \mu\text{g/mL}$ in the same assay).

The development of proline mimics is of considerable current interest due to the helix-breaking and unique con-

Scheme 3. Synthesis of the Hydrochloride Salt of $[\text{CH}_3]_2\text{GS}$ (**1**·2HCl)



formational properties of this amino acid residue.²³ We have now demonstrated that a trisubstituted (*E*)-alkene dipeptide isostere can serve as a bioisosteric replacement for the D-Phe-Pro type II' β -turn in the cyclopeptide antibiotic gramicidin S. The solution conformational analysis and the biological assay of analogues **10** and **1**, respectively, provide strong validation of our design principles. Furthermore, the hydrazirone/Zn $\rightleftharpoons$ Zn transmetalation/imine addition methodology was key to a rapid synthetic access to the target compounds.

Acknowledgment. This work was supported by the NIH/NIGMS P50 program (GM067082). J.X. gratefully acknowledges an Andrew W. Mellon Predoctoral Fellowship for support and Dr. Corey R. J. Stephenson (ETHZ) for helpful discussions.

Supporting Information Available: Experimental procedures and spectral data, including copies of ^1H and ^{13}C NMR spectra for all new compounds, and 2D NMR spectra of **10** and **Cbz**₂GS. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0617704

(19) (a) Wipf, P.; Fritch, P. C.; Geib, S. J.; Seffler, A. M. *J. Am. Chem. Soc.* **1998**, *120*, 4105. (b) Imperiali, B.; Fisher, S. L.; Moats, R. A.; Prins, T. J. *J. Am. Chem. Soc.* **1992**, *114*, 3182. (c) Kessler, H. *Angew. Chem., Int. Ed.* **1982**, *21*, 512.

(20) Ball, J. B.; Hughes, R. A.; Alewood, P. F.; Andrews, P. R. *Tetrahedron* **1993**, *49*, 3467.

(21) (a) Tamaki, M.; Akabori, S.; Muramatsu, I. *Bull. Chem. Soc. Jpn.* **1993**, *66*, 3113. (b) Woody, R. W. In *Circular Dichroism*; Nakanishi, K., Berova, N., Woody, R. W., Eds.; VCH: New York, 1994; Chapter 17, pp 473–496.

(22) A 4300-step conformational equilibrium search was performed with: *Spartan 04*; Wavefunction, Inc.: Irvine, CA.

(23) (a) Jenkins, C. L.; Vasbinder, M. M.; Miller, S. J.; Raines, R. T. *Org. Lett.* **2005**, *7*, 2619. (b) Cordero, F. M.; Pisaneschi, F.; Batista, K. M.; Valenza, S.; Machetti, F.; Brandi, A. *J. Org. Chem.* **2005**, *70*, 856. (c) Wang, X. J.; Hart, S. A.; Xu, B.; Mason, M. D.; Goodell, J. R.; Etzkorn, F. A. *J. Org. Chem.* **2003**, *68*, 2343. (d) Halab, L.; Lubell, W. D. *J. Am. Chem. Soc.* **2002**, *124*, 2474. (e) Otaka, A.; Katagiri, F.; Kinoshita, T.; Odagaki, Y.; Oishi, S.; Tamamura, H.; Hamanaka, N.; Fujii, N. *J. Org. Chem.* **2002**, *67*, 6152.