


## Proline Derived Spirobarbiturates as Highly Effective $\beta$ -Turn Mimetics Incorporating Polar and Functionalizable Constraint Elements

Luelak Lomlim,<sup>†</sup> Juergen Einsiedel,<sup>†</sup> Frank W. Heinemann,<sup>‡</sup> Karsten Meyer,<sup>‡</sup> and Peter Gmeiner<sup>\*,†</sup>

Department of Chemistry and Pharmacy, Emil Fischer Center, Chair of Medicinal Chemistry, Friedrich Alexander University, Schuhstrasse 19, D-91052 Erlangen, Germany, and Department of Chemistry and Pharmacy, Chair of Inorganic Chemistry, Friedrich Alexander University, Egerlandstrasse 1, D-91058 Erlangen, Germany

gmeiner@pharmazie.uni-erlangen.de

Received November 30, 2007



A practical and efficient synthesis of spirobarbiturates of type **III** is reported when NH acidity of the imide function of the hydrophilic linker element allowed the introduction of different substituents. Structural characterization, which was based on both X-ray crystallography and spectroscopic investigations, indicated type II  $\beta$ -turn formation. Introduction of the molecular scaffold into solid phase peptide synthesis gave rise to spirocyclic neuropeptide analogs.

Since reverse-turn motifs play a crucial role in molecular recognition and signal transduction,<sup>1</sup> the development of privileged scaffolds nucleating turn structures has attracted remarkable interest.<sup>2</sup> Upon introduction of constraint elements into biologically active peptides, the number of degrees of




FIGURE 1. Modification of constraint elements.

conformational freedom and, thus, binding-associated loss of entropy can be decreased.<sup>3</sup> The most powerful  $\beta$ -turn mimetics have been derived from the functionally important amino acid proline<sup>4</sup> when the introduction of suitable constraint elements led to lactam-bridged molecular scaffolds.<sup>5</sup> Employing both saturated and unsaturated constraint elements leading to Freidinger-type (**I**)<sup>6</sup> and dehydro-Freidinger-type (**II**) spirocycles,<sup>7</sup> respectively, we were able to establish a molecular building-kit that allows adjustment of a wide range of dihedral angles (Figure 1).

To complement these investigations, we aimed to construct spirocyclic analogues incorporating polar, “backbone-like” constraint elements. Interestingly, such a concept was described for conformationally constrained nucleosides when a barbiturate acid moiety was incorporated into a spirocyclic system.<sup>8</sup>

Following the concept of privileged structures, we planned to synthesize and to conformationally evaluate spirobarbiturates of type **III**. Such molecular scaffolds should be available via cyclocondensation reactions as described for the preparation of barbiturate-type drugs.<sup>9</sup> NH acidity of the hydrophilic linker element should allow the introduction of different substituents. These could serve as molecular probes exploring binding pockets of complementary target proteins. We herein present a practical synthesis of model peptide surrogates of type **III**, solid-phase supported application toward two artificial neuropeptide mi-

<sup>†</sup> Chair of Medicinal Chemistry.

<sup>‡</sup> Chair of Inorganic Chemistry.

(1) (a) Ohage, E. C.; Graml, W.; Walter, M. M.; Steinbacher, S.; Steipe, B. *Protein Sci.* **1997**, *6*, 233–241. (b) Zhou, H. X.; Hoess, R. H.; deGrado, W. F. *Nat. Struct. Biol.* **1996**, *3*, 446–451. (c) Venkatachalam, C. M. *Biopolymers* **1968**, *6*, 1425–1436. (d) Tyndall, J. D. A.; Pfeiffer, B.; Abbenante, G.; Fairlie, D. P. *Chem. Rev.* **2005**, *105*, 793–826.

(2) For examples, see: (a) Weber, K.; Ohnmacht, U.; Gmeiner, P. *J. Org. Chem.* **2000**, *65*, 7406–7416. (b) Thomas, C.; Ohnmacht, U.; Nigier, M.; Gmeiner, P. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 2885–2890. (c) Haubner, R.; Schmitt, W.; Holzemann, G.; Goodman, S. L.; Jonczyk, A.; Kessler, H. *J. Am. Chem. Soc.* **1996**, *118*, 7881–7891. (d) Liu, R.; Dong, D. L.-Y.; Sherlock, R.; Nestler, H. P.; Gennari, C.; Mielgo, A.; Scolastico, C. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 847–852.

(3) For recent reviews and articles, see: (a) Cluzeau, J.; Lubell, W. D. *Biopolymers (Pept. Sci.)* **2005**, *80*, 98–150. (b) Maison, W.; Prenzel, A. H. G. P. *Biopolymers* **2005**, *7*, 1031–1048. (c) Hanessian, S.; McNaughton-Smith, G.; Lombart, H.-G.; Lubell, W. D. *Tetrahedron* **1997**, *53*, 12789–12854. (d) Freidinger, R. M. *J. Med. Chem.* **2003**, *46*, 5553–5566. (e) Halab, L.; Gosselin, F.; Lubell, W. D. *Biopolymers (Pept. Sci.)* **2000**, *55*, 101–122. (f) Hoffmann, T.; Waibel, R.; Gmeiner, P. *J. Org. Chem.* **2003**, *68*, 62–69.

(4) Paul, A.; Bittermann, H.; Gmeiner, P. *Tetrahedron* **2006**, *62*, 8919–8927, and references cited therein.


(5) (a) Belvisi, L.; Colombo, L.; Manzoni, L.; Potenza, D.; Scolastico, C. *Synlett* **2004**, *9*, 1449–1471. (b) Fernández, M. M.; Diez, A.; Rubiralta, M.; Montenegro, E.; Casamitjana, N. *J. Org. Chem.* **2002**, *67*, 7587–7599. (c) Jeannotte, G.; Lubell, W. D. *J. Am. Chem. Soc.* **2004**, *126*, 14334–14335. (d) Wang, W.; Xiong, C.; Hruby, V. J. *Tetrahedron Lett.* **2001**, *42*, 3159–3561. (e) Mueller, R.; Revesz, L. *Tetrahedron Lett.* **2001**, *35*, 4091–4092. (f) Ward, P.; Ewan, G. B.; Jordan, C. C.; Ireland, S. J.; Hagan, R. M.; Brown, J. R. *J. Med. Chem.* **1990**, *33*, 1848–1851. (g) Hoffmann, T.; Lanig, H.; Waibel, R.; Gmeiner, P. *Angew. Chem.* **2001**, *113*, 3465–3468. (h) Hoffmann, T.; Lanig, H.; Waibel, R.; Gmeiner, P. *Angew. Chem., Int. Ed.* **2001**, *40*, 3361–3364. (i) Einsiedel, J.; Lanig, H.; Waibel, R.; Gmeiner, P. *J. Org. Chem.* **2007**, *72*, 9102–9113.

(6) (a) Müller, G.; Hessler, G.; Decorez, H. Y. *Angew. Chem., Int. Ed.* **2000**, *39*, 894–896. *Angew. Chem.* **2000**, *112*, 926–928. (b) Hinds, M. G.; Welsh, J. H.; Brennand, D. M.; Fisher, J.; Glennie, M. J.; Richards, N. G. J.; Turner, D. L.; Robinson, J. A. *J. Med. Chem.* **1991**, *34*, 1777–1789. (c) Ward, P.; Ewan, G. B.; Jordan, C. C.; Ireland, S. J.; Hagan, R. M.; Brown, J. R. *J. Med. Chem.* **1990**, *33*, 1848–1851. (d) Bittermann, H.; Einsiedel, J.; Hübner, H.; Gmeiner, P. *J. Med. Chem.* **2004**, *47*, 5587–5590. (e) Bittermann, H.; Gmeiner, P. *J. Org. Chem.* **2006**, *71*, 97–102.

(7) Bittermann, H.; Boeckler, F.; Einsiedel, J.; Gmeiner, P. *Chem. Eur. J.* **2006**, *24*, 6315–6322.

(8) Renard, A.; Lhomme, J.; Kotera, M. *J. Org. Chem.* **2002**, *67*, 1302–1307.

(9) Fischer, E.; Diltz, A. *Liebigs Ann. Chem.* **1904**, *335*, 334–268.

SCHEME 1<sup>a</sup>

<sup>a</sup> Key: (a) *t*-BuOK, DMSO, rt, 16 h; (b) CH<sub>3</sub>NH<sub>2</sub>·HCl, HATU, DBU, NMP, rt, 15 min–3 h; (d) prep HPLC, silica gel, diisopropyl ether/acetonitrile; Dcb: 2,6-dichlorobenzyl.

metics and structural characterization which was based on both X-ray crystallography and spectroscopic investigations.

The construction of the novel spirocyclic scaffolds was envisioned by a base-promoted cytocondensation reaction of *N*-carbamoyl-substituted amino acids with a suitably *N*-protected α-carboxyproline diester (Scheme 1). Starting from *N*-Boc-protected diethylaminomalonate and 1,3-dibromopropane in the presence of base, the prochiral building block **1**<sup>10</sup> was prepared in 90% yield when the addition of KI significantly increased the efficiency of the cyclization. Our initial attempts to the synthesis of the spirocyclic scaffold including the treatment of the dicarboxylate **1** with *N*-carbamoyl glycine benzylester and EtONa or *t*-BuOK failed. This is putatively due to the high energetic demand for the generation of a 1,2-dianion from the initially formed ester enolate. To circumvent this problem, we planned to react diester **1** with free *N*-carbamoylamino acids in presence of an excess of base, because formation of a carboxylate anion should prevent from deprotonation at C<sub>α</sub> and facilitate optical integrity in case of chiral amino acid derivatives. To approach to representative building blocks, the urea derivatives **2a–c** were synthesized by *N*-carbamoylation of glycine, (*S*)-phenylalanine and *O*-(2,6-dichlorobenzyl)-(*S*)-tyrosine with KOCN, according to previously reported protocols.<sup>11</sup> In fact, deprotonation of the carbamoyl amino acids **2a–c** by 3 equiv of *t*-BuOK in DMSO and subsequent addition of the pyrrolidine

dicarboxylic ester **1** resulted in a 55–71% formation of the spirobarbiturate **3a–c** (Scheme 1). Employing HATU as an activating agent, carboxamide formation could be performed giving access to the *N*-protected Pro-GlyNHMe surrogate **4a** in racemic form. Furthermore, the conformationally constrained Pro-Phe-NHMe and Pro-Tyr(ODcb)-NHMe analogues **4b** and **4c**, both as 1:1 mixtures of diastereomers, were obtained. Finally, preparative HPLC gave rise to the isomerically pure model peptide surrogates **4b1**, **4b2**, **4c1**, and **4c2**.

X-ray diffraction analysis of the model peptide surrogate **4b1** was performed to provide helpful information on both the conformational properties in solid state and the absolute stereochemistry of the spirocyclic center which clearly revealed (*R*)-configuration (Figure 2). Due to reverse CIP priority-based assignments, the disposition of the backbone-forming carboxamide function of **4b1** is identical to that of natural (*S*)-proline amide. Based on the X-ray structure of the reference scaffold **4b1**, overall similarities of <sup>1</sup>H NMR data and analogous elution profiles allowed determination of the spirocyclic stereogenic centers of **4c1** and **4c2** to be (*R*) and (*S*), respectively.




FIGURE 2. Thermal ellipsoid plot of the molecular structure of **4b1** (50% probability ellipsoids).

Taking advantage of the *NH* acidity of our linker elements, we intended to introduce different substituents which can be exploited as biomolecular probes. To facilitate a chemoselective *N*-alkylation, orthogonal protection of the carboxylic acid function should be first conducted. TCE-esters are known to be resistant toward acidic and basic conditions and can be readily removed by reductive cleavage with zinc dust.<sup>12</sup> In detail, EDC/DMAP promoted esterification of **3a–c** with 2,2,2-trichloroethanol afforded the 2,2,2-trichloroethyl (TCE) esters **5a–c**. Upon deprotonation of the imides **5a–c** by NaH and treatment with methyl iodide, benzyl bromide or (DcbO)-benzyl bromide,<sup>13</sup> *N*-alkylation was observed giving access to the functionalized scaffolds **6a–e** in 62–85% yield. Zn-mediated TCE deprotection of the representatives **6b–e** could be accomplished when we obtained the free carboxylic acid derivatives **7a–d**. To approach to the model peptide surrogates of type **8**, HATU-promoted peptide coupling and subsequent aminolysis were performed.


Finally, separation of the diastereomers **8a1/8a2** and **8b1/8b2** was done by HPLC using normal-phase conditions, followed by an NMR-based configurational assignment using **4b1** as a reference structure.

(10) (a) Galons, H.; Damianos, N.; Miocque, M.; Bram, G. In *Second Forum on Peptides*; Aubry, A., Marraud, M., Vitoux, B., Eds.; Colloque INSERM/John Libbey Eurotext Ltd.: Paris, 1989; Vol. 174, pp 281–288. (b) Aboujaoude, E. E.; Collignon, N.; Savignac, P.; Bensoam, J. *Phosphorus Sulfur* **1987**, 34, 93–104.

(11) Preparation of the urea derivatives according to: (a) Moschel, R. C.; Hudgins, R. W.; Dipple, A. J. *Org. Chem.* **1986**, 51, 4180–4185. (b) Taillades, J.; Boiteau, L.; Beuzelin, I.; Lagrille, O.; Biron, J.-P.; Vayaboury, W.; Vandennebeele-Trambouze, O.; Giani, O.; Commeyras, A. J. *Chem. Soc., Perkin Trans. 2* **2001**, 1247, 1254. (c) **2b**: [α]<sub>D</sub><sup>20</sup> +37.8 (c 1.0, MeOH); [α]<sub>D</sub><sup>20</sup> +39.2 (c 1.0, MeOH) is reported in: Sarges, R.; Howard, H. R.; Donahue, K. M.; Welch, W. M.; Dominy, B. W.; Weissman, A.; Koe, B. K.; Bordners, J. J. *Med. Chem* **1986**, 29, 8–19. (d) Casimir, J. R.; Tourwé, D.; Iterbeke, K.; Guichard, G. J. *Org. Chem.* **2000**, 65, 6487–6492.


(12) Jou, G.; Gonzales, F.; Albericio, F.; Lloyd-Williams, P.; Giralt, E. J. *Org. Chem.* **1997**, 62, 354–366.

(13) Warrellow, G. J.; Porter, J. R.; Archibald, S. C.; Head, J. C. *Chem. Abstr.* **2000**, 132, 265095 WO 2000020396, 2000.

SCHEME 2<sup>a</sup>

<sup>a</sup> Key: (a) 2,2,2-trichloroethanol, EDC, DMAP, CH<sub>2</sub>Cl<sub>2</sub>, rt, 16 h; (b) CH<sub>3</sub>I or 2,6-Cl<sub>2</sub>Bn-OBnBr, NaH, DMF, rt; (c) Zn dust, 1 M NH<sub>4</sub>OAc, THF, rt, 24 h; (d) CH<sub>3</sub>NH<sub>2</sub>·HCl, HATU, DBU, NMP, rt, 15 min–3 h; (e) HPLC; Dcb: 2,6-dichlorobenzyl.

To apply our methodology to the synthesis of biologically relevant peptide mimetics, we intended to incorporate our molecular scaffolds of type **III** into NT(8–13) (H-Arg-Arg-Pro-Tyr-Ile-Leu-OH), the active portion of the neuropeptide neurotensin.<sup>14</sup> Since the Pro-Tyr fragment is known to be of special importance for receptor recognition, introduction of the structural congeners **3c** and **7a** should give access to valuable molecular probes of type **9** and **10**, respectively (Scheme 3). Thus, Fmoc-isoleucine was coupled using PYBOP as an activating reagent to *N*-deprotected Leu immobilized on Wang resin. Microwave acceleration proved to be advantageous for both Fmoc-deprotection of the resin and acylation. Subsequently,  $\mu$ -wave assisted coupling of the building blocks **3c** and **7a** was performed when a second acylation with HATU was necessary to complete the ligation of **3c**. Cleavage of the Boc function was done according to a previously published protocol employing 10% sulfuric acid in dioxane at 8 °C, thus preventing liberation of the peptide from the solid support.<sup>15</sup> Since the pyrrolidine nitrogen of the spirobarbiturate system was not susceptible to acylation reactions with common peptide coupling reagents, we employed the recently introduced BTC (bis-trichloromethylcarbonate),<sup>16</sup> which proved successful for the coupling of Fmoc(Pbf)-arginine to the sterically hindered amine function. After a further deprotection

SCHEME 3<sup>a</sup>

<sup>a</sup> Key: (a) Fmoc-Leu-Wang resin, piperidine/DMF (1:4),  $\mu$ : 5 × 5 s, 100W, 5 × cooling to –10 °C; (b) Fmoc-Ile-OH, PyBOP, DIPEA, HOBt, DMF,  $\mu$ : 15 × 10 s, 50 W, 15 × cooling to –10 °C; (c) (1) Fmoc deprotection (see a), (2) **3c** or **7a**, conditions see (b) for **3c**: reagents for a 2nd  $\mu$ -assisted coupling: HATU, DIPEA, DMF; (d) H<sub>2</sub>SO<sub>4</sub>/dioxane (1:9), 8 °C, 2 h; (e) Fmoc-Arg(Pbf)-OH, BTC, 2,6-lutidine, rt, 1 h; (f) (1) Fmoc deprotection (see a), (2) coupling of Fmoc-Arg(Pbf)-OH (see b), (3) Fmoc deprotection (see a); (g) TFA, phenol, H<sub>2</sub>O, triisopropylsilane 88:5:5:2, 2 h; (h) RP-HPLC; (i) H<sub>2</sub>, 20% Pd(OH)<sub>2</sub>/C, MeOH; (j) RP-HPLC.

and coupling with Fmoc(Pbf)-arginine, TFA-promoted cleavage, preparative HPLC and palladium-catalyzed hydrogenolysis of the 2,6-dichlorobenzyl group, the target peptides **9** and **10** were isolated in isomerically pure form.

Employing X-ray crystallographic data, we clearly identified a type II  $\beta$ -turn structure for the model peptide scaffold **4b1** (Figure 3) when an intramolecular hydrogen bond with a distance of 2.21 Å<sup>17</sup> could be unambiguously deduced between the methyl amide *N*-H<sub>*i*+3</sub> and the C=O function in position *i*. The backbone dihedral angles of proline ( $\phi_{i+1}$ : –61.1°/ $\psi_{i+1}$ : 137.7°) and phenylalanine ( $\phi_{i+2}$ : 65.9°/ $\psi_{i+2}$ : 17.4°) were deviating by less than 30°<sup>18</sup> from the canonical angles of –60°, 120°, 80°, and 0°, respectively, thus correlating well with the ideal form of a type II  $\beta$ -turn.<sup>19</sup> The distance from the *tert*-butoxy oxygen and the methylamide carbon (*d* O<sub>*i*</sub>–C<sub>*i*+3</sub> representing the distance between C $\alpha_i$  and C $\alpha_{i+3}$ , respectively) was found substantially shorter than 7 Å (5.07 Å), indicating that a U-turn is adopted. Additionally, the low pseudodihedral angle  $\beta$  (C<sub>*i*</sub>–C $\alpha_{i+1}$ –C $\alpha_{i+2}$ –N<sub>*i*+3</sub>) of 9.3° revealed an almost coplanar spatial arrangement and, thus, antiparallel pleated  $\beta$ -sheet nucleating properties.




FIGURE 3. Geometric parameters of **4b1**.

To evaluate the structural behavior of the model peptide mimetic **4b1** in solution, we performed conformational studies based on FT-IR and <sup>1</sup>H NMR spectroscopy. To exclude intermolecular interactions, spectra were recorded at 2 mM concentrations.<sup>20</sup> <sup>1</sup>H NMR spectra of **4b1** showed two peaks for

(14) Hong, F.; Cusack, B.; Fauq, A.; Richelson, E. *Curr. Med. Chem.* **1997**, 4, 412–434.

(15) Trivedi, H. S.; Anson, M.; Steel, P. G.; Worley, J. *Synlett* **2001**, 12, 1932–1934.

(16) Falb, E.; Yechezkel, T.; Salitra, Y.; Gilon, C. *J. Peptide Res.* **1999**, 53, 507–517.

(17) Belvisi, L.; Gennari, G.; Mielgo, A.; Potenza, D.; Scolastico, C. *Eur. J. Org. Chem.* **1999**, 38, 9–400.

(18) Lewis, P. N.; Momany, F. A.; Scheraga, H. A. *Biochim. Biophys. Acta* **1973**, 303, 249–255.

(19) Ball, J. B.; Hughes, R. A.; Alewood, P. F.; Andrews, P. R. *Tetrahedron* **1993**, 49, 3467–3478.

(20) Boussard, G.; Marraud, M. *J. Am. Chem. Soc.* **1985**, 107, 1825–1828.



amide *NH*, a major peak at 6.52 ppm (95%) and a minor peak at 5.87 ppm (5%) indicating the presence of two conformers. Temperature dependent chemical shifts as a measure for the stability of secondary structures were studied between 273 and 313 K in CDCl<sub>3</sub> recorded in 10 K steps, revealing a low  $\Delta\delta\Delta T$  value of  $-3.1$  ppb/K for the *NH* of the major conformer. IR spectra of **4b1** (CHCl<sub>3</sub>) displayed *NH* stretching vibrations at 3381 cm<sup>-1</sup> and a very weak shoulder above 3450 cm<sup>-1</sup>. Thus, we concluded, that an intramolecular hydrogen bond is strongly favored. The above-mentioned diagnostic values for the spirobarbiturates **4a,b2,c1/2** and **8a1/2,b1/2** ( $\delta(\text{NH})$ : 6.41–6.56 ppm;  $\nu(\text{NH})$ : 3380–3402 cm<sup>-1</sup>;  $\Delta\delta\Delta T$ :  $(-4.8)$ – $(-2.4)$  ppb/K) were very similar to the data measured for reference peptide surrogate **4b1** clearly indicating the preferred formation of type II  $\beta$ -turn secondary structures for all compounds investigated.

In conclusion, an efficient methodology was established that gives access to a novel type of  $\beta$ -turn nucleating scaffolds incorporating polar, “backbone-like” constraint elements.

## Experimental Section

**General Method for the Synthesis of the Barbiturate Scaffolds 3a–c.** To a solution of *t*-BuOK in DMSO (0.5 mmol/mL) was added the respective urea (**2a–c**) dissolved in DMSO (0.15 mmol/mL) at 10 °C. After 10 min of stirring, a solution of pyrrolidine-1,2,2-tricarboxylic acid 1-*tert*-butyl ester 2,2-diethyl ester (**1**) in DMSO (0.3 mmol/mL) was added. The reaction mixture stirred at rt for 20 h, and then 5% aq citric acid was added and extraction with EtOAc was performed. After the organic layer was washed with H<sub>2</sub>O and brine and dried with MgSO<sub>4</sub>, the solvent was evaporated and the residue was purified by flash column chromatography.

**General Method for the Synthesis of the Methyl Amides 4a–c and 8a,b.** A solution of the carboxylic acid, methylamine hydrochloride, and HATU in NMP (2 mL) was stirred for 10 min at rt, and then a solution of DBU in NMP (2 mL) was added dropwise. After 15 min–3 h, 5% citric acid was added, and extraction with EtOAc was performed. The organic layer was washed with H<sub>2</sub>O and brine, dried over MgSO<sub>4</sub>, and evaporated. The product was purified by flash column chromatography.

**General Method for the Synthesis of 2,2,2-Trichloroethyl Ester Derivatives 5a–c.** To a stirred solution of the respective carboxylic acid, 2,2,2-trichloroethanol, and DMAP in CH<sub>2</sub>Cl<sub>2</sub> was added EDC at 0 °C. After 1 h, the solution was allowed to stir at rt for 18 h. Thereafter, the solvent was removed in vacuo, H<sub>2</sub>O

was added, and a pH  $\approx 7$  was adjusted by the addition of 5% citric acid. After extraction with EtOAc, the combined organic layer was washed with H<sub>2</sub>O and brine and dried over MgSO<sub>4</sub>. The solvent was removed in vacuo, and the product was purified by flash column chromatography.

**General Method for the *N*-Alkylation of the Barbiturate Scaffolds to Give 6a–e.** An oily suspension of NaH (60%) was washed 2 $\times$  with dry hexane, and then DMF (4 mL) was added and the mixture cooled to 0 °C. A solution of the corresponding barbiturate derivative in DMF (3–10 mL) was added dropwise. After the development of H<sub>2</sub>, a solution of the respective alkyl halide in DMF (3 mL) was added, and the reaction mixture was allowed to stir at rt for 16 h. Thereafter, 5% citric acid was added, and extraction with ethylacetate (3  $\times$  ) was performed. The organic layer was washed with H<sub>2</sub>O and brine, dried over MgSO<sub>4</sub>, and evaporated, and the residue was purified by flash column chromatography.

**General Method for the Reductive Ester Cleavage To Give 7a–d.** To a solution of the respective 2,2,2-trichloroethyl ester in THF was added zinc dust and then an aqueous 1 M NH<sub>4</sub>OAc solution. After vigorous stirring for 24 h at rt, the mixture was filtered through Celite and washed with THF. Subsequently, the solvent was removed in vacuo, and the residue was partitioned between 5% citric acid and EtOAc. The organic layer was washed with H<sub>2</sub>O and brine and dried over MgSO<sub>4</sub>, and the solvent was removed in vacuo. The residue was purified by flash column chromatography.

**Acknowledgment.** We thank I. Torres-Berger, M. Kettler, and Dr. R. Waibel for technical assistance and helpful discussions, respectively. This work was supported by the DFG (Gm 13/7).

**Supporting Information Available:** Experimental and analytical data (for compounds **1**, **2c**, and **3–10** including epimerization study of **3b,c**, HPLC data and conformational analysis data for **4a,b1/b2,c** and **8a1/2,b1/2** as well as details of the X-ray crystal structure determination of **4b1** in CIF format. This material is available free of charge via the Internet at <http://pubs.acs.org>. CCDC-668906 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

JO702573Z