



Contents lists available at ScienceDirect

## Bioorganic &amp; Medicinal Chemistry

journal homepage: [www.elsevier.com/locate/bmc](http://www.elsevier.com/locate/bmc)

# Synthesis, biological evaluation and 3D-QSAR studies of 3-keto salicylic acid chalcones and related amides as novel HIV-1 integrase inhibitors

Horrick Sharma<sup>a</sup>, Shivaputra Patil<sup>a</sup>, Tino W. Sanchez<sup>b</sup>, Nouri Neamati<sup>b</sup>, Raymond F. Schinazi<sup>c</sup>, John K. Buolamwini<sup>a,\*</sup>

<sup>a</sup> Department of Pharmaceutical Sciences, College of Pharmacy, University of Tennessee Health Science Center, Memphis, 847 Monroe Avenue, Suite 327, TN 38163, USA

<sup>b</sup> Department of Pharmacology and Pharmaceutical Sciences, School of Pharmacy, University of Southern California, 1985 Zonal Avenue, Los Angeles, CA 90089, USA

<sup>c</sup> Emory University School of Medicine/VA Medical Center, Medical Research 151H, 1670 Clairmont Road, Decatur, GA 30033, USA

## ARTICLE INFO

## Article history:

Received 24 November 2010

Revised 16 January 2011

Accepted 24 January 2011

Available online 1 March 2011

## Keywords:

HIV-1

HIV integrase

Integrase inhibitors

3-Keto salicylic acids

Antiviral activity

Strand transfer inhibitors

Chalcones

Synthesis

3D-QSAR

CoMFA

CoMSIA

## ABSTRACT

HIV-1 integrase is one of the three most important enzymes required for viral replication and is therefore an attractive target for anti retroviral therapy. We herein report the design and synthesis of 3-keto salicylic acid chalcone derivatives as novel HIV-1 integrase inhibitors. The most active compound, 5-bromo-2-hydroxy-3-[3-(2,3,6-trichlorophenyl)acryloyl]benzoic acid (**25**) was selectively active against integrase strand transfer, with an  $IC_{50}$  of 3.7  $\mu$ M. While most of the compounds exhibited strand transfer selectivity, a few were nonselective, such as 5-bromo-3-[3-(4-bromophenyl)acryloyl]-2-hydroxybenzoic acid (**15**), which was active against both 3'-processing and strand transfer with  $IC_{50}$  values of  $11 \pm 4$  and  $5 \pm 2$   $\mu$ M, respectively. The compounds also inhibited HIV replication with potencies comparable with their integrase inhibitory potencies. Thus, 5-bromo-2-hydroxy-3-[3-(2,3,6-trichlorophenyl)acryloyl]benzoic acid (**25**) and 5-bromo-3-[3-(4-bromophenyl)acryloyl]-2-hydroxybenzoic acid (**15**) inhibited HIV-1 replication with  $EC_{50}$  values of 7.3 and 8.7  $\mu$ M, respectively. A PHASE pharmacophore hypothesis was developed and validated by 3D-QSAR, which gave a predictive  $r^2$  of 0.57 for an external test set of ten compounds. Pharmacophore derived molecular alignments were used for CoMFA and CoMSIA 3D-QSAR modeling. CoMSIA afforded the best model with  $q^2$  and  $r^2$  values of 0.54 and 0.94, respectively. This model predicted all the ten compounds of the test set within 0.56 log units of the actual  $pIC_{50}$  values; and can be used to guide the rational design of more potent novel 3-keto salicylic acid integrase inhibitors

© 2011 Elsevier Ltd. All rights reserved.

## 1. Introduction

Acquired immunodeficiency syndrome (AIDS) has been a devastating pandemic with around 33 million people infected with it worldwide. The human immunodeficiency virus (HIV) is implicated in the etiology of AIDS. The *pol* gene of HIV-1 encodes three enzymes that are required for viral replication viz., reverse transcriptase (RT), integrase (IN) and protease (PR).<sup>1</sup> Currently FDA approved antiretroviral drugs target mainly RT<sup>2</sup> and PR<sup>3</sup> and are widely used in combination as highly active antiretroviral therapy (HAART). Entry inhibitors, for example, Maraviroc<sup>4</sup> and fusion inhibitors such as Enfuvirtide<sup>5</sup> have also recently been approved

**Abbreviations:** HIV, human immunodeficiency virus; ST, strand transfer; FDA, food and drug administration; 3D-QSAR, three dimensional quantitative structure activity relationship; CoMFA, comparative molecular field analysis; CoMSIA, comparative molecular similarity analysis; VdW, Van der Waals; NMR, nuclear magnetic resonance; HPLC, high performance liquid chromatography.

\* Corresponding author. Tel.: +1 901 448 7533; fax: +1 901 448 6828.

E-mail address: [jbuolamwini@uthsc.edu](mailto:jbuolamwini@uthsc.edu) (J.K. Buolamwini).

as anti-HIV agents. Though, HAART has been effective in reducing morbidity and mortality, it does not eliminate the virus from patients.<sup>6</sup> Furthermore, the emergence of multidrug resistant viral strains along with severe side effect<sup>7,8</sup> associated with existing drugs and the life-long chronic nature of the infection, demand the development of novel drugs that target other crucial steps of viral replication and survival.

The integration of viral cDNA into host chromosomes is an essential step in the viral replication process<sup>9</sup> and is responsible for stable infection. Moreover, IN has no known structural analogs in human cells.<sup>10–13</sup> Inhibition of this enzyme therefore provides an attractive strategy for antiretroviral drug development.

The function of IN is to integrate viral cDNA into the host genome. This process occurs in two steps: 3'-end processing (cleavage; 3'-P) and joining reaction (strand transfer, ST). In the processing reaction which occurs in the cytoplasm of an infected cell, a site specific endonucleolytic activity removes the highly conserved 3' terminal CA dinucleotide from the 3'-end of the of viral DNA to generate a reactive nucleophilic 3'-hydroxyl group.<sup>14,15</sup> In the strand transfer reaction, the processed 3'-hydroxyl end of the viral

DNA does a nucleophilic attack at the phosphodiester bond of the host DNA to effect integration.

$\beta$ -Diketo acid derivatives as represented by S-1360<sup>16</sup> was the first integrase inhibitor class to have entered clinical trials. Naphthyridine carboxamides such as L-870,810<sup>17</sup> have also advanced to clinical trials as have novel mono keto quinolone carboxylic acids<sup>18,19</sup> such as GS-9137 which is currently undergoing Phase III clinical trials. In spite of over two decades of extensive research leading to promising IN inhibitors,<sup>20–25</sup> hitherto only one IN inhibitor, raltegravir,<sup>26</sup> a pyrimidinone carboxamide, has been approved for clinical use. Moreover, emergence of resistance against raltegravir due to viral mutations mostly usually T66I, N155H, Q148H/R and E92Q,<sup>27</sup> demands exploration of novel scaffolds for the treatment of HIV infection.

We have previously reported novel DKA IN inhibitors with aryl moiety modifications that were derived from a phenanthridinone scaffold.<sup>28</sup> In continuing our efforts, we report herein the synthesis and biological evaluation of chalcone derivatives having 3-keto salicylic acid moieties as a new class of IN inhibitors. We have undertaken substantial analog synthesis and applied the PHASE pharmacophore analysis program in conjunction with three dimensional structure–activity relationship (3D-QSAR) studies to gain insights into their binding at the IN active site. CoMFA and CoMSIA 3D-QSAR models were derived to study the influence of stereoelectronic, hydrophobic and hydrogen bonding interactions on activity; and to establish predictive tools for design of novel inhibitors.

## 2. Results and discussion

### 2.1. Chemistry

In the present study, a novel series of IN inhibitors was discovered based on modification of the diketo acid pharmacophoric group as represented by structure **III** (Fig. 1). The 2-position OH and 3-position keto groups together mimic the keto-enol tautomer of the  $\beta$ -diketo group. Attachment of the carboxylic acid at the 1-position gave the 3-keto salicylic acid moiety of **III**, intended to interact with IN similar to the  $\beta$ -diketo acids. A styryl group was attached to the 3-position keto group to create chalcones (**IV**) (Fig. 1), the type of which have not yet been investigated as HIV IN inhibitors to our knowledge. The chalcone moiety was also reduced to give an ethylene linker as shown in compound **60**. Furthermore, replacement of the enone group of the chalcones with an amide group was also investigated.

The synthesis of target chalcones was performed according to reaction Scheme 1. The 5-substituted salicylic acid (X = COOH) starting material **4** was first acetylated with acetic anhydride and

concentrated sulfuric acid. The acetylated intermediate **5** was then subjected to Fries rearrangement reaction using anhydrous aluminum chloride and heating at 160 °C to transfer the acetyl group from the 2-position oxygen to the 3-position to yield 3-acetyl salicylic acid intermediate **6**.<sup>29</sup> The next step involved the Claisen–Schmidt condensation reaction of intermediate **6** with various aryl aldehydes at room temperature using NaOH to obtain the target chalcones (**10–57**).<sup>30</sup> To determine the importance of the  $\alpha,\beta$ -unsaturated bond (chalcone scaffold) to IN inhibitory activity, the synthesis of compound **60** was undertaken by reduction of compound **36** by Pd/C-catalyzed hydrogenation (Scheme 2). The amide **61** was synthesized according to reaction Scheme 3 such that, commercially available 5-bromo salicylic acid (**2**) was first formylated with chloroform at the 2-position (ortho) by Reimer–Tiemann reaction to yield the aldehyde intermediate **7**. However, the low yield of the above reaction, prompted to employ an alternative synthetic route. Thus, 3-formyl salicylaldehyde **1** was brominated by NBS in acetonitrile to afford intermediate **7** in higher yields. In the next step, compound **7** was condensed with 4-fluorobenzyl amine to afford intermediate **8** which was subsequently subjected to oxidation with KMnO<sub>4</sub> to yield **61**. For SAR purposes, compounds **62** and **63** were synthesized by coupling with *p*-fluorobenzyl amine following step c of reaction Scheme 3, from the commercially available starting materials **3a** and **3b**.

### 2.2. HIV-1 integrase inhibition structure–activity relationship (SAR)

After synthesis, the target compounds were tested for integrase inhibitory activity to assess their potency and study the SAR. The integrase 3'-cleavage and strand transfer (ST) inhibitory activity data are presented in Table 1 along with compound structures. In general, the compounds behaved as typical diketo acid IN inhibitors, being selective against the strand transfer (ST) step compared to the 3'-end processing (3'-P).

#### 2.2.1. Substitution on aromatic ring A

Compound **9**, 2-hydroxy-3-(3-phenyl-propyl)-benzoic acid, in which salicylic acid is connected to an unsubstituted phenyl ring B by a propylene linker exhibited only weak inhibitory activity with IC<sub>50</sub> of >100  $\mu$ M for ST and 91  $\mu$ M for 3'-end processing. Modification to a 3-keto salicylic acid along with substitution with Br at R<sub>1</sub> and an  $\alpha,\beta$ -unsaturated linker (chalcone) in compound **10** (used as reference) demonstrated significant increase in both activity and selectivity (IC<sub>50</sub> 25  $\mu$ M for ST vs IC<sub>50</sub> 75  $\mu$ M for 3'-end processing). With the presence of a bromine substituent at the R<sub>1</sub> on aromatic ring A being shown to be important for activity, chlorine and fluorine substituents were also investigated. Compounds with a Br substituent were the most active followed by F which did a little better than the corresponding Cl substituted analogs. To determine the effect of hydrophobic interaction, a methyl group was substituted for the halogens at R<sub>1</sub> (see compound **57**). Compound **57** was found to be 13 times less active than the corresponding Br-substituted compound **25**. Since Br and a methyl group are similar in steric terms, this implies that it is not the hydrophobic effect of the halogen, but possibly the electronegativity and/or polarizability that might be dominant. The 3-keto salicylic acid pharmacophore was designed to form coordinate bonds with Mg<sup>2+</sup>. To test this hypothesis, the carboxylic acid was replaced with NO<sub>2</sub> functionality (**58** and **59**) and this modification diminished HIV inhibitory activity. Replacement of the carboxyl group with another polar group but with weaker coordination ability, CN (compound **56**) also resulted in reduced inhibitory activity. This probably indicates that the formal negative charge of the free ionizable acidic moiety, that would chelate better with the metal ion,






Figure 1. Design of 3-keto salicylic acid pharmacophore of synthesized chalcones.



**Scheme 1.** Reagents and conditions: (a)  $\text{Ac}_2\text{O}$ ,  $\text{H}_2\text{SO}_4$ ; (b)  $\text{AlCl}_3$ ,  $160^\circ\text{C}$ ; (c)  $\text{ArCHO}$ ,  $\text{NaOH}$ ,  $\text{EtOH}$ .



**Scheme 2.** Reagents and conditions: (a)  $\text{H}_2$ ,  $\text{Pd/C}$  (10 mol %), methanol.

is required for potent activity, supporting the hypothesis. Of course a chelation study will have to be done to confirm this.


### 2.2.2. Substitution on aromatic ring B

Various groups were substituted on aromatic ring B to further explore the SAR. The presence of halogens on phenyl ring B produced noticeable improvements in potency based on the type, number and position on the aromatic ring. Fluorine substitution in the *ortho*- and *meta*-positions (compounds **12** and **55**) diminished activity, as did iodine substitution at the *para*-position (compound **16**). However, substitution with Cl enhanced activity specifically for ST in all the three positions (compounds **11**, **13** and **14**). Interestingly, compounds **54** and **15** with *m*-Br and *p*-Br substituent, respectively, were potent against both catalytic steps (compound **15** showed  $\text{IC}_{50}$  of  $5\text{ }\mu\text{M}$  against ST activity and  $11\text{ }\mu\text{M}$  against 3'-processing). The 2,3,6-trichloro-substituted compound (**25**) was the most active, with  $\text{IC}_{50}$  of about  $\sim 4\text{ }\mu\text{M}$  against the ST reaction. This suggests a binding site pocket with high affinity for halogenated phenyl rings. This might be the site that binds the fluorophenyl group common in potent IN inhibitors like the clinically used drug raltegravir.


We next turned our attention to examining the following. (1) *The effect of a bulky aryl B groups:* benzyloxy and chloro substituted phenoxy in *ortho*, *meta* and *para*-positions of phenyl ring B enhanced activity for both 3'-P and ST steps; whereas an *ortho* furan substituent (compound **50**) increased the inhibitory potency against ST but decreased potency against 3'-P; the increase in activity was most pronounced for the *o*-benzyloxy substituent (compound **42**), with the 3'-P and ST  $\text{IC}_{50}$  values lowered to 21 and  $9\text{ }\mu\text{M}$ , respectively, compared to corresponding  $\text{IC}_{50}$  values of 75 and  $25\text{ }\mu\text{M}$  for the parent phenyl compound **10**. (2) *The effects of electron donating or withdrawing groups:* introduction of electron donating groups such as methoxy, methyl and cyclopentoxy (compounds **19**, **39** and **46**) decreased integrase inhibitory activity; and, increasing the number of electron donating groups resulted in further decrease in activity (compare compounds **19** and **20**). Introduction of strong electron withdrawing groups such as trifluoromethyl and nitro also diminished activity as exemplified by compounds **52** and **53**, respectively; (3) *Boisosteric replacement of phenyl ring B with polycyclic and heteroaromatic rings:* replacement of phenyl with thiophene or benzothiophene (compounds **45** and **49**, respectively) displayed improved activity. Naphthalene derivatives on the other hand decreased activity (compounds **47** and **48**). It thus appears that heteroaromatic B rings may improve integrase inhibitory activity.

### 2.2.3. Reduction of the $\alpha,\beta$ -unsaturated double bond of the chalcones or replacement with an amide moiety

To explore the importance of the  $\alpha,\beta$ -unsaturated bond of the chalcones on inhibitory activity, we reduced it to a single bond. Our first attempt at this was with compound **25**, but reduction also led to a loss of the Br substituent. However, reduction (Scheme 2) of analog **36**, having a fluorine substituent, which is not as good a




**Scheme 3.** Reagents and conditions: (a)  $\text{CHCl}_3$ ,  $\text{NaOH}$ ,  $\text{H}_2\text{O}$ ; (b)  $\text{NBS}$ ,  $\text{CH}_3\text{CN}$ , 2 h (c) 4-F benzylamine,  $\text{EDCI}$ ,  $\text{HOBT}$ ,  $\text{Et}_3\text{N}$ ; (d)  $\text{KMnO}_4$ ,  $\text{H}_2\text{O}$ /pyridine,  $90^\circ\text{C}$ .

**Table 1**  
Structure and HIV-1 integrase inhibitory activities of chalcones


Compd	Ar	X	R <sub>1</sub>	Activity IC <sub>50</sub> (μM)	
				3'-P <sup>a</sup>	ST <sup>b</sup>
10	Ph	COOH	Br	75 ± 14	25 ± 5
11	2-Cl-Ph	COOH	Br	52 ± 13	14 ± 4
12	2-F-Ph	COOH	Br	74 ± 25	32 ± 3
13	3-Cl-Ph	COOH	Br	45 ± 10	9 ± 3
14	4-Cl-Ph	COOH	Br	81 ± 28	12 ± 1
15	4-Br-Ph	COOH	Br	11 ± 4	5 ± 2
16	4-I-Ph	COOH	Br	69	35
17	2,4-di-Cl-Ph	COOH	Br	28 ± 12	7 ± 4
18	2,3-di-Cl-Ph	COOH	Br	58 ± 13	19 ± 11
19	2,3,di-MeO-Ph	COOH	Br	62 ± 14	34 ± 6
20	2,3,4-tri-MeO-Ph	COOH	Br	85 ± 20	51 ± 16
21	2,5-di-Cl-Ph	COOH	Br	47 ± 14	18 ± 4
22	2,6-di-Cl-Ph	COOH	Br	52 ± 12	16 ± 8
23	3,4-di-Cl-Ph	COOH	Br	27 ± 11	14 ± 4
24	2,3,5-tri-Cl-Ph	COOH	Br	20 ± 12	13 ± 5
25	2,3,6-tri-Cl-Ph	COOH	Br	23 ± 12	<3.7
26	4-Cl-Ph	COOH	Cl	85 ± 13	53 ± 21
27	2,4-di-Cl-Ph	COOH	Cl	>100	>100
28	2,3-di-Cl-Ph	COOH	Cl	100	92 ± 14
29	2,5-di-Cl-Ph	COOH	Cl	75	25
30	2,3,5-tri-Cl-Ph	COOH	Cl	55 ± 40	17 ± 3
31	2,3,6-tri-Cl-Ph	COOH	Cl	77 ± 20	30 ± 5
32	3,4-di-Cl-Ph	COOH	Cl	87 ± 13	41 ± 9
33	4-Cl-Ph	COOH	F	36 ± 13	17 ± 3
34	2,4-di-Cl-Ph	COOH	F	52 ± 13	24 ± 9
35	2,3-di-Cl-Ph	COOH	F	55 ± 18	30 ± 9
36	2,5-di-Cl-Ph	COOH	F	50	29 ± 6
37	2,3,5-tri-Cl-Ph	COOH	F	33 ± 14	15 ± 4
38	2,3,6-tri-Cl-Ph	COOH	F	82 ± 16	39 ± 6
39	3,5-di-Me-Ph	COOH	Br	>100	58
40	2-Cl-3,4-di-MeO-Ph	COOH	Br	>100	76
41	6-F-3,4-di-MeO-Ph	COOH	Br	>100	76
42	2-BzO-Ph	COOH	Br	21	9
43	3-(4-Cl-PhO)-Ph	COOH	Br	25	18
44	4-(4-Cl-PhO)-Ph	COOH	Br	22	16
45	3-Thienyl	COOH	Br	30 ± 12	18 ± 1
46	3-(Cyclopentyloxy)-Ph	COOH	Br	100	55
47	3-(2-OEt-naphthyl)	COOH	Br	92	46
48	5-MeO-naphthyl	COOH	Br	54	54 ± 25
49	5-Benzothiophene	COOH	Br	23 ± 3	11 ± 1
50	2-Furan-2-yl-Ph	COOH	Br	>100	16 ± 2
51	3-Cl-4,5-Methylenedioxy-Ph	COOH	Br	79	63
52	3-OCF <sub>3</sub> -Ph	COOH	Br	>100	52
53	3-NO <sub>2</sub> -Ph	COOH	Br	49 ± 8	32 ± 6
54	3-Br-Ph	COOH	Br	19 ± 4	18 ± 1
55	3-F-Ph	COOH	Br	71 ± 1	59 ± 2
56	2,3,6-tri-Cl-Ph	CN	Br	>100	44 ± 2
57	2,3,6-tri-Cl-Ph	COOH	CH <sub>3</sub>	>100	50 ± 9
58	2,3,5-tri-Cl-Ph	NO <sub>2</sub>	Br	>333	68 ± 5
59	2,3,6-tri-Cl-Ph	NO <sub>2</sub>	Br	36 ± 4	21 ± 3
S1360 <sup>c</sup>	N/A	N/A	N/A	11 ± 2	0.6 ± 0.1

<sup>a</sup> 3-P = 3'-processing.<sup>b</sup> ST = strand transfer.<sup>c</sup> S1360 = standard.

leaving group as Br, was successful in achieving our objective. The reduction product, compound **60**, was less active than unsaturated analog, showing that the reduction of the double bond in the chalcone template is detrimental to activity. Bioisosteric replacement of the α,β-unsaturated carbonyl system in the chalcone with an

**Table 2**  
Structure and HIV-1 integrase inhibitory activities of synthesized compounds



Compd	Ar	R <sub>2</sub>	R <sub>1</sub>	X	Activity (IC <sub>50</sub> , μM)	
					3'-P <sup>a</sup>	ST <sup>b</sup>
8	4-F-Ph	NH	Br	CHO	>100	8 ± 1
60	2,5-di-Cl-Ph	CH <sub>2</sub>	F	COOH	>100	57 ± 5
61	4-F-Ph	NH	Br	COOH	>100	15 ± 3
62	4-F-Ph	NH	H	CH <sub>3</sub>	>100	>100
63	4-F-Ph	NH	Br	H	>100	>100
S1360 <sup>c</sup>	N/A	N/A	N/A	N/A	11 ± 2	0.6 ± 0.1

<sup>a</sup> 3-P = 3'-processing.<sup>b</sup> ST = strand transfer.<sup>c</sup> S1360 = standard.


amide moiety (compound **61**, see Table 2) was well tolerated with retention of comparative inhibitory activity but an increase in ST selectivity. Interestingly, the aldehyde intermediate compound **8**, which was obtained during the synthesis of the amide analog compound **61**, was found to be more active than the carboxylic acid product **61**, indicating that an aldehyde group can substitute for the carboxylic acid group in these compounds, opening an opportunity for exploring other bioisosteric substitutions.

### 2.3. Inhibition of HIV replication in cell culture


To evaluate the potential of these new IN inhibitors as antiviral agents, five potent analogs were tested for the ability to inhibit HIV-1 replication in primary human peripheral blood mononuclear cells (PBMCs) infected with the virus. The toxicity of the compounds was also tested against PBMC and human CEM lymphoblastic leukemia cells. All five compounds tested inhibited HIV replication (Table 3). The most potent compounds, **25** and **15** showed comparable EC<sub>50</sub> values of 7.3 and 8.3 μM, respectively. Compound **49** (integrase ST IC<sub>50</sub> = 11 μM) exhibited an EC<sub>50</sub> of 10 μM, while analog **17** (integrase ST IC<sub>50</sub> = 7.0 μM) displayed an EC<sub>50</sub> of 13.9 μM. Thus, in general the biological activity correlated well with the IN inhibitory potency. Moreover, the two most potent IN inhibitory compounds of the set tested, **15** and **25**, were also found to have the highest activity against HIV replication, suggesting that IN might indeed be their antiviral target. These results demonstrate moderate antiviral activity, which is not very different from early members of the diketo acid IN inhibitors, which served as important lead compounds such as L-706,908 which had an EC<sub>50</sub> of 5.7 ± 4.7 μM for inhibition of HIV replication.<sup>31</sup> The reason that compound **8** (integrase ST IC<sub>50</sub> = 8.0 μM), in which the carboxylic acid is replaced by an aldehyde group showed a lower antiviral activity (EC<sub>50</sub> = 30.8 μM) than expected is not apparent, but may have to do with solubility, absorption, binding to serum proteins in cell culture media and/or cellular distribution factors. The compounds showed mild cytotoxicity, with selectivity index up to fourfold for compound **49**. It may be noteworthy that compound **49** is the only compound among the antiviral test set that had a 5-benzothiophene for ring B; this may offer clues for improving selectivity index. These results demonstrate reasonable antiviral activity when compared to the antiviral activity of reported diketo acid inhibitors like L-706,908 which has an EC<sub>50</sub> of 5.7 ± 4.7 μM<sup>31</sup> for inhibition of HIV replication. To put data in perspective with regard to AZT's activity, in the same assay, AZT had an EC<sub>50</sub> of 0.001 μM.<sup>31</sup>



L-706,908



L-870,810



S-1360

### 3. Molecular modeling

#### 3.1. Generation of a pharmacophore model using the PHASE program

Knowledge of the structure of target protein inhibitor interactions is limited as no crystal structure of human integrase–ligand–DNA complex is available. Therefore, to explore the bioactive conformation of these novel IN inhibitors, the PHASE<sup>32</sup> (v3.0) pharmacophore mapping program implemented in Maestro modeling program package (v8.5; Shrodinger Inc.) was used. The 3D structures of molecules were constructed in and imported from the SYBYL modeling program (v8.3, Tripos).

Conformers of each molecule were generated using the OPLS\_2005 force field in the macromodel conformational analysis program of PHASE. A set of pharmacophoric sites based on features defined in PHASE were assigned to the molecules. These included H-bond acceptor (A), H-bond donor (D), hydrophobic group (H), negatively charged group (N) and aromatic ring (R). The three most potent compounds of the series, compound **15**, **17** and **25** were selected as actives for use in identifying a common pharmacophore hypothesis using a tree-based partitioning technique of PHASE. The resulting pharmacophores were then scored and ranked.

The top twenty pharmacophore hypotheses were analyzed and validated by partial least square (PLS) regression-based PHASE 3D-

QSAR. Random external test sets were generated automatically to determine the predictive ability of the models. The hypothesis that afforded the best PLS statistics (Table 4) and the highest predicted ability consisted of six features viz., two H-bond acceptors, two hydrophobic groups, a negative group and one aromatic ring (Fig. 2a). Two H-bond acceptor features mapped onto the oxygen of the hydroxyl and the 3-keto moiety and negative feature mapped onto the carboxylate group of the salicylic acid pharmacophore. The two hydrophobic features mapped to the halogen substituent of ring A and the *para*-substituted halogen in aryl ring B. The aromatic pharmacophoric feature mapped onto aromatic ring A. Residuals of the training and the test set are listed in Tables 5 and 6, respectively. The prediction curve for the test set is shown in Figure 2b. As can be seen, the hypothesis gave a good predictive model with a predictive  $r^2$  of 0.57 for an external test set of ten

**Table 4**  
PLS Statistics of PHASE 3D-QSAR model

$q^2$	0.57
$r^2$	0.74
SD	0.17
F	28.3
PLS Factors	4
Pearson R	0.75

**Table 3**  
Anti-HIV activity of selected novel potent derivatives


Compd	Anti-HIV-1 activity in PBMCs <sup>a</sup>		Slope	R	Cytotoxicity (IC <sub>50</sub> , $\mu$ M)	
	EC <sub>50</sub> <sup>b</sup> ( $\mu$ M)	EC <sub>90</sub> <sup>b</sup> ( $\mu$ M)			PBMCs	CEM
<b>15</b>	8.7	21.0	2.5 $\pm$ 0.61	0.97	24.4	27.5
<b>17</b>	13.9	26.0	3.5	1.0	22.1	23.8
<b>25</b>	7.3	20.3	2.2 $\pm$ 0.96	0.91	22.7	19.0
<b>49</b>	10.1	24.3	2.5 $\pm$ 0.89	0.94	39.1	20.7
<b>8</b>	30.8	52.7	4.1	1.0	44.0	40.2
AZT	0.0025	0.01	1.6 $\pm$ 0.30	0.96	>100	56.1
L-706,908	5.7 $\pm$ 4.7 <sup>c</sup>	N/A	N/A	N/A	N/A	N/A

<sup>a</sup> Human peripheral blood mononuclear cells (PBMCs).

<sup>b</sup> EC<sub>50</sub> and EC<sub>90</sub> are the effective concentrations inhibiting 50% and 90% HIV replication, respectively.

<sup>c</sup> In the human leukemic T-cell line MOLT-4; in that assay the EC<sub>50</sub> of AZT was 0.001  $\mu$ M.





**Figure 2.** The best pharmacophore hypothesis (AAHHNR) mapped onto (a) compound **17** (b) alignment of actives and (c) inactives from PHASE pharmacophore mapping. Pharmacophore features are: red spheres and vectors for H-bond acceptor (A), brown rings for aromatic groups (R), dark red spheres refer to a negative feature (N) and green spheres indicate hydrophobic groups (H). Atoms in the ligands are represented as O, red; Br, wine red; Cl, dark green; H, white; C, gray. (d) PHASE 3D-QSAR predictions of the test set.

compounds. It was also able to distinguish active and inactive compounds, with the latter group not aligning well to the features (Fig. 2c and d).

### 3.2. 3D-QSAR analysis

#### 3.2.1. CoMFA and CoMSIA 3D-QSAR models

Statistical analysis by PLS<sup>33,34</sup> was done using CoMFA and CoMSIA descriptors as independent variables and biological activity in the form of  $pIC_{50}$  values as dependent variable. The molecular alignment used for CoMFA and CoMSIA 3D-QSAR modeling was derived from PHASE pharmacophore mapping (Shrodinger, Inc.). PHASE derived pharmacophore hypotheses have been successfully used to derive 'bioactive conformations' for 3D-QSAR modeling in previous work from our laboratory<sup>35</sup> and elsewhere (Telvekar et al.).<sup>36</sup> Thus, 'bioactive conformations' of all the molecules from PHASE conformational analysis were superimposed using the MATCH alignment tool in SYBYL (Fig. 3). Fifty three (53) synthesized target compounds were used for the 3D-QSAR analyses. The Leave-One-Out (LOO) method of cross-validation was used for initial assessment of the predictive abilities of the models with the training sets. The optimal number of components used in the final QSAR models was that which gave the smallest standard error of prediction. Reliability of the models was tested by prediction of 10 compounds selected as an external test set using factor analysis. The PLS analysis results from both the CoMFA and CoMSIA models are summarised in Table 7. Cross-validated  $q^2$  values of 0.43 and 0.54 were obtained for CoMFA and CoMSIA, respectively. Results of rigorous statistical testing using group cross-validation (20 groups), bootstrapping and randomization of activity data demonstrated the robustness of the QSAR models. Both CoMSIA and CoMFA identified six outliers, which were excluded on the basis of their extreme residual values (Table 8). Out of these, five outliers were common to both CoMFA and CoMSIA. These were compounds **16**,

**27**, **28**, **55** and **57**. The reasons for compound **55** being an outlier are not apparent. With compound **16** (iodo substituted) and compound **57** ( $R_1 = CH_3$ ), their outlier status may stem from structural uniqueness, whereas the outlier status of compounds **27** and **28** (least active analogs,  $R_1 = Cl$ ) may drive from different binding modes. The CoMFA and CoMSIA predictions for the training and the test sets are shown in Figure 4a and b, respectively. The CoMFA and CoMSIA residual values for the training set are given in Table 8 and the corresponding values for the test set in Table 9. In concordance with the higher  $q^2$  value, the CoMSIA model performed better than CoMFA model, and predicted all test set compounds within 0.56 log unit of the actual  $pIC_{50}$  values.

#### 3.2.2. CoMFA and CoMSIA contour maps

The CoMFA and CoMSIA contour maps derived from the PLS coefficients are displayed in Figure 5. Steric CoMFA maps (Fig. 5a) show green contours around the *ortho* and *para*-positions on aryl ring B indicating that bulky groups are favored at these positions. This is validated by the fact that a benzyl group substitution at the *ortho*-position resulted in higher activity (compound **42**). Compounds **44** and **49**, which have a bulky chloro phenoxy and thiophene groups around the green contour at the *para*-position also displayed better inhibitory activities. A large yellow contour surrounds a small green contour at the *meta*-position on aryl ring B suggesting that only a limited bulk is favorable for activity. This is apparent by the loss of inhibitory activity observed with compound **46**, the bulky cyclopentyl ring of which occupies the sterically disfavored yellow contour region. There is also a green contour evident at the  $R_1$ -position on the aromatic ring A. This may be the reason why among the three halogens at that position, bromine, which is the bulkiest, afforded the highest activity. At the aryl ring B, CoMFA electrostatic map reveals a red contour at the 3'-position suggesting that electronegative groups could increase activity at this position. This is most exemplified by the 3'-Cl


**Table 5**  
Residuals of training set by PHASE 3D-QSAR

Compd	Actual ST pIC <sub>50</sub>	Predicted ST pIC <sub>50</sub>	Residuals
10	4.60	4.85	−0.25
11	4.85	4.59	0.26
12	4.49	4.53	−0.04
14	4.92	4.84	0.08
15	5.30	4.94	0.36
16	4.45	4.85	−0.4
17	5.15	5.02	0.13
18	4.72	4.66	0.06
19	4.46	4.59	−0.13
20	4.29	4.29	0.00
21	4.74	4.87	−0.13
22	4.79	4.83	−0.04
24	4.88	4.82	0.06
25	5.40	5.40	0.00
26	4.27	4.49	−0.22
27	4.00	4.21	−0.21
28	4.03	4.48	−0.45
32	4.39	4.44	−0.05
33	4.76	4.63	0.13
34	4.62	4.33	0.29
35	4.52	4.44	0.08
36	4.54	4.63	−0.09
37	4.82	4.59	0.23
39	4.23	4.42	−0.19
40	4.12	3.94	0.18
41	4.12	4.15	−0.03
42	5.04	4.99	0.05
43	4.74	4.79	−0.05
44	4.79	4.79	0.00
45	4.74	4.85	−0.11
46	4.26	4.32	−0.06
47	4.33	4.16	0.17
48	4.26	4.28	−0.02
49	4.96	4.86	0.10
53	4.49	4.46	0.03
54	4.74	4.60	0.14
55	4.23	4.32	−0.09
56	4.40	4.24	0.16
57	4.30	4.37	−0.07
58	4.16	4.10	0.06
59	4.68	4.80	−0.12
61	4.82	4.83	−0.01
63	4.00	4.26	−0.26

**Table 6**  
Residuals of test set by PHASE 3D-QSAR

Compd	Actual ST pIC <sub>50</sub>	Predicted ST pIC <sub>50</sub>	Residual
13	5.04	4.86	0.18
23	4.85	4.78	0.07
29	4.60	4.71	−0.11
30	4.77	4.63	0.14
31	4.52	4.25	0.27
38	4.41	4.28	0.13
50	4.80	4.91	−0.11
51	4.20	4.51	−0.31
52	4.28	4.46	−0.18
62	4.00	4.35	−0.35

substitution in compound **13** which had an IC<sub>50</sub> value of 9 μM. However, for compound **55**, a decrease in activity was observed. Red contours are also present at the 5'- and the 6'-positions. The dichloro-substituted compounds **21** and **22** having Cl at the 5'- and 6'-positions showed improved activities. There are also distant blue contours around aryl ring B, which predicts that electropositive groups should increase activity at these positions. Electronegative oxygen atom of the methoxy substituents in compounds **20**, **40** and **41** points towards the blue contours and thus may explain the loss of activity observed by the presence of methoxy groups in



**Figure 3.** Superimposition of the conformation of molecules obtained from PHASE pharmacophore analysis. Atoms are represented as O, red; Br, dark green; Cl, green; F, bright green; H, cyan; C, white; S, yellow; N, blue.

**Table 7**  
PLS Statistics of CoMFA and CoMSIA 3D-QSAR

PLS statistics	CoMFA	CoMSIA
$q^2$	0.43	0.54
$r^2$	0.94	0.94
s	0.09	0.09
F	78.32	78.69
PLS components	6	6
Field contribution		
Steric	0.474	0.100
Electrostatic	0.526	0.428
Hydrophobic		0.283
Donor		0.146
Acceptor		0.044
Bootstrapping $r^2$ (20 runs)	0.96 ± 0.01	0.96 ± 0.01
Randomization $q^2$ (20 runs)	0.33	0.39
Cross-validation $q^2$ (20 runs)	0.44	0.50

these compounds. The red contours around the salicylic acid moiety in aryl ring A suggest that a negative charge is important for activity in this region, possibly for chelation with the active site Mg<sup>2+</sup> ions. This is in agreement with the proposed mechanism of action of these compounds as HIV-1 integrase inhibitors. There is a red contour around the 4'-position and both blue and red contours at the 5'-position (R<sub>1</sub>). This suggests that polarizability effect might be important at R<sub>1</sub> and could explain the increased activity conferred by bromine substitution over other halogens and the similar-sized hydrophobic methyl group at this position.

The CoMSIA steric contour map (Fig. 5b) also shows a large green contour around aryl ring B. Like in the CoMFA, the electrostatic contours of CoMSIA also show red contours at the 3' and the 4'-positions on ring B, and at the 5'-position (R<sub>1</sub>) on ring A. There is a blue contour located at the position of the hydrogen atom of the hydroxyl group of the salicylic acid moiety. The hydrophobic, H-bond donor and acceptor contours are shown in Figure 5c. There is a white contour located near the R<sub>1</sub> substituent position which suggests hydrophobic groups increase activity at this region. This might be the reason why compound **57** with a hydrophobic methyl group at this position was less active than the corresponding halogen compounds. The improved activity observed with bromine at R<sub>1</sub>-position may be the result because of a combination of polarizability effects. White contours are also evident at the 2' and the 5'-positions on the B ring, indicating that hydrophilic groups are favored for activity. This might be the explanation, or at least in part, for the decrease in potency of compounds **19** and **39**, which have hydrophobic methoxy and methyl groups at these re-

**Table 8**  
Residuals of training set of CoMFA and CoMSIA models

Compd	Actual ST pIC <sub>50</sub>	Residuals	
		CoMFA	CoMSIA
10	4.60	−0.13	−0.12
11	4.85	0.12	0.12
12	4.49	−0.18	−0.19
13	5.04	0.05	0.03
14	4.92	−0.15	−0.08
15	5.30	0.16	0.11
16 <sup>a</sup>	4.45	−0.62	−0.96
17	5.15	0.21	0.19
18	4.72	0.10	−0.07
19	4.46	0.06	−0.04
20	4.29	−0.05	−0.003
21	4.74	0.01	0.04
22	4.79	−0.03	−0.03
23	4.85	−0.13	−0.03
24	4.88	0.07	0.11
25	5.40	−0.05	−0.05
26	4.27	−0.02	0.09
27 <sup>a</sup>	4.00	−0.62	−0.89
28 <sup>a</sup>	4.03	−0.79	−0.74
29	4.60	0.02	0.005
30	4.77	0.01	0.11
31	4.52	0.03	0.06
33	4.76	−0.009	0.04
34	4.62	−0.02	−0.13
35	4.52	−0.03	−0.12
36	4.54	0.06	0.04
38	4.41	−0.001	−0.04
39	4.23	−0.05	−0.05
40	4.12	−0.02	−0.02
43	4.74	−0.07	−0.01
44 <sup>c</sup>	4.79	−0.07	0.56
46	4.26	−0.03	0.00
47	4.33	0.07	0.09
48	4.26	0.002	−0.03
49	4.96	0.10	0.04
51	4.20	0.01	−0.07
52 <sup>b</sup>	4.28	−0.84	−0.09
55 <sup>a</sup>	4.23	−0.41	−0.31
56	4.35	−0.05	−0.01
57 <sup>a</sup>	4.30	−1.03	−1.03
58	4.16	−0.01	0.11
62	4.00	0.07	0.04
63	4.00	−0.047	−0.05

<sup>a</sup> Represents common outliers.<sup>b</sup> Outlier only for CoMFA.<sup>c</sup> Outlier only for CoMSIA.

gions. Another white contour appears at the 3-keto oxygen again signifying the need for hydrophilic groups at this position. There is a yellow contour close to the 4'-position on ring B, suggesting that hydrophobic character is favored for activity. The 3'-keto func-


**Table 9**  
Residuals of the test set predictions given by CoMFA and CoMSIA models

Compd	Actual ST pIC <sub>50</sub>	Residuals	
		CoMFA	CoMSIA
32	4.39	−0.41	−0.28
37	4.82	0.35	0.30
41	4.12	−1.00	−0.32
42	5.04	0.25	0.35
45	4.74	0.14	0.37
50	4.79	0.09	0.09
53	4.49	−0.10	0.22
54	4.74	0.19	0.05
59	4.67	0.11	0.56
61	4.82	0.76	0.40


tionality perfectly fits the magenta contour within its vicinity indicating that they may act as H-bond acceptors. A cyan contour appears close to, and merges into the magenta contour, suggesting that the OH group is a H-bond donor at this position. It may be involved in H-bonding interactions with the acceptor carboxylate groups of the acidic catalytic triad residues of integrase active site. The general agreement between the maps and the observed biological activity trends suggests that these models could aid in the design of more active compounds.

#### 4. Conclusion

In summary, we have successfully synthesized and evaluated the HIV-1 integrase inhibitory activities of a series of chalcone derivatives and related amides harboring a 3-keto salicylic acid moiety as a diketo acid isostere. Consistent with the DKA class of IN inhibitors, these series of compounds were also generally selective against the IN strand transfer catalytic step. Introduction of a Br substituent at the R<sub>1</sub>-position and an  $\alpha,\beta$ -unsaturated carbonyl linker connecting aryl rings A and B resulted in significant enhancement of inhibitory activity. The most active compound has 2,3,6-trichloro substitution on aryl ring B. In the absence of a crystal structure of the complete human integrase, inhibitor and DNA complex, bioactive conformational hypothesis was obtained and validated by PHASE pharmacophore mapping and 3D-QSAR analysis. Statistically significant CoMFA and CoMSIA 3D-QSAR models were obtained using an alignment derived from the pharmacophore analysis, and may be useful for the design and prediction of the activities of novel related compounds as HIV-1 integrase inhibitors. Last but not the least, most of the potent compounds also inhibited HIV replication in cell culture with moderate antiviral activity, which supports further development of the series.

**Figure 4.** Predictions of ST activities of the training (blue diamonds) and test set (red triangles) compounds by (a) CoMFA and (b) CoMSIA





**Figure 5.** (a) CoMFA and (b) CoMSIA steric and electrostatic contour plots. The green contours indicate regions where bulky groups increase activity, whereas yellow contours indicate regions where bulky groups decrease activity. Blue contours indicate regions where electropositive groups increase activity, whereas red contours depict regions where electronegative groups increase activity. (c) CoMSIA hydrophobic, H-bond donor and H-bond acceptor contour plots; white contours indicate regions where hydrophilic groups increase activity, whereas yellow contours indicate regions where hydrophobic groups increase activity, cyan indicates region where H-donors increase activity and magenta contour indicate region where H-bond acceptor groups increase activity.

## 5. Experimental section

All reagents and solvents were purchased from the Aldrich Chemical Company and used without further purification. Progress of reactions was monitored by TLC on silica gel GHLF-250 micron plates (Analtech, Inc.). Fisher scientific Da visil grade 1740 (170–400 mesh) silica gel was used for flash column chromatography.  $^1\text{H}$  NMR spectra were recorded on Bruker AR, 300 or 500 MHz spectrometer; chemical shifts are expressed in  $\delta$  values (ppm) and coupling constants ( $J$ ) in Hertz. Mass spectral data were determined on a Bruker-HP Esquire-LC spectrometer (ESI-MS). Melting points of final products were determined using a Fisher-Johns melting point apparatus and are reported uncorrected. HPLC analysis of final compounds was carried out using a reverse phase SUPELCOSIL 5  $\mu\text{m}$  C-18 column of dimensions  $25 \times 4.6$  cm. Area% purity was detected at 254 nm. An initial isocratic method comprising 40% water (solvent A) and 60% methanol (solvent B) for 20 min at a flow-rate of 1.2 mL/min was optimized to a linear gradient elution with water (A) and methanol (B) at a flow-rate of 1.7 mL/min. Gradient elution of the mobile phase was either (a) 80% A to 35% A from 0 to 5 min, 35% A, 65% B for 5–12 min, and 35% A to 80% A from 12 to 15 min or (b) 60% A to 20% A from 0 to 4 min, 20% A and 80% B for 4–12 min and 20% A to 60% A from 12 to 15 min. The purity of all, but two, compounds was found to be  $\geq 95\%$ .

### 5.1. General procedure for the synthesis of acetylated intermediates (5a–5e)

5-Halogenated salicylic acids (46 mmol) were stirred with acetic anhydride (20 mL) and 80  $\mu\text{L}$  of concd  $\text{H}_2\text{SO}_4$  was then added. After a few (20) min the reaction mixture solidified and was poured into cold water. The solid so obtained was filtered and extracted with ethyl acetate, washed with brine and dried over

$\text{Na}_2\text{SO}_4$ . The solvent was evaporated under reduced pressure to obtain a crude solid which was recrystallized from ethyl acetate/hexane to afford white crystals (5a–5c).

#### 5.1.1. 2-Acetoxy-5-bromobenzoic acid (5a)

Yield 88%; mp 155–157  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  13.42 (br s, 1H, COOH), 8.0 (d, 1H,  $J = 2.4$  Hz, ArH), 7.85 (dd, 1H,  $J_1 = 2.7$  Hz,  $J_2 = 8.7$  Hz, ArH), 7.20 (d, 1H,  $J = 8.7$  Hz, ArH), 2.2 (s, 3H,  $\text{OCOCH}_3$ ); MS (ESI):  $m/z$  216.6  $[\text{M}-\text{COCH}_3]^-$ .

#### 5.1.2. 2-Acetoxy-5-fluorobenzoic acid (5b)

Yield 85%; mp 133–134  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  13.42 (br s, 1H, COOH), 7.67 (dd, 1H,  $J_1 = 3.3$  Hz,  $J_2 = 9.0$  Hz, ArH), 7.51 (ddd, 1H,  $J_1 = 8.7$  Hz,  $J_2 = 8.1$  Hz,  $J_3 = 3.3$  Hz, ArH), 7.26 (dd, 1H,  $J_1 = 4.8$  Hz,  $J_2 = 8.7$  Hz, ArH), 2.24 (s, 3H,  $\text{OCOCH}_3$ ); MS (ESI):  $m/z$  154.7  $[\text{M}-\text{COCH}_3]^-$ .

#### 5.1.3. Acetoxy-5-chlorobenzoic acid (5c)

Yield 85%; mp 150–152  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  7.88 (d, 1H,  $J = 2.7$  Hz, ArH), 7.72 (dd, 1H,  $J_1 = 2.7$  Hz,  $J_2 = 9.0$  Hz, ArH), 7.27 (d, 1H,  $J = 8.4$  Hz, ArH), 2.24 (s, 3H,  $\text{OCOCH}_3$ ); MS (ESI):  $m/z$  170.7  $[\text{M}-\text{COCH}_3]^-$ .

#### 5.1.4. Acetic acid-4-bromo-2-cyanophenyl ester (5d)

Yield 76%; mp 60–61  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.26 (d, 1H,  $J = 2.7$  Hz, ArH), 8.0 (dd, 1H,  $J_1 = 2.4$  Hz,  $J_2 = 8.7$  Hz, ArH), 7.44 (d, 1H,  $J = 8.7$  Hz, ArH), 2.29 (s, 3H,  $\text{OCOCH}_3$ ); MS (ESI):  $m/z$  197.5  $[\text{M}-\text{COCH}_3]^-$ .

#### 5.1.5. 2-Acetoxy-5-methylbenzoic acid (5e)

Yield 75%; mp 158–160  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.01 (d, 1H,  $J = 1.5$  Hz, ArH), 7.50 (dd, 1H,  $J_1 = 1.8$  Hz,  $J_2 = 8.4$  Hz, ArH), 7.12 (d, 1H,  $J = 8.1$  Hz, ArH), 2.43 (s, 3H,  $\text{OCOCH}_3$ ), 2.27 (s, 3H,  $\text{CH}_3$ ); MS (ESI):  $m/z$  150.7  $[\text{M}-\text{COCH}_3]^-$ .

## 5.2. General procedure for the synthesis of 3-acetyl-5-halo salicylic acids (6a–6e)

The acetylated intermediates (38.61 mmol) and  $\text{AlCl}_3$  (120 mmol) were mixed in a three-necked flask and heated to  $160^\circ\text{C}$  under mechanical stirring. After 3 h, the reaction mixture was cooled to room temperature and poured into ice containing 20 mL concentrated HCl. The slurry was extracted with ethyl acetate, acidified with 1 M HCl, washed with brine and dried over  $\text{Na}_2\text{SO}_4$ . The solvent was evaporated to give the crude product, which was washed with dichloromethane for removal of impurities, filtered and dried to give 3-acetyl salicylic acids as pale brown powders.

### 5.2.1. 2-Acetoxy-3-acetyl-5-bromobenzoic acid (6a)

Yield 34%; mp  $182\text{--}185^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.06 (d, 1H,  $J = 2.7$  Hz, ArH), 7.97 (d, 1H,  $J = 2.7$  Hz, ArH), 2.60 (s, 3H,  $\text{COCH}_3$ ); MS (ESI):  $m/z$  258.7  $[\text{M}-\text{H}]^-$ .

### 5.2.2. 2-Acetoxy-3-acetyl-5-fluorobenzoic acid (6b)

Yield 31%; mp  $135\text{--}136^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  7.78 (dd, 1H,  $J_1 = 3.3$  Hz,  $J_2 = 8.1$  Hz, ArH), 7.72 (dd, 1H,  $J_1 = 3.3$  Hz,  $J_2 = 8.7$  Hz, ArH), 2.62 (s, 3H,  $\text{COCH}_3$ ); MS (ESI):  $m/z$  196.7  $[\text{M}-\text{H}]^-$ .

### 5.2.3. 2-Acetoxy-3-acetyl-5-chlorobenzoic acid (6c)

Yield 32%; mp  $160\text{--}163^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  7.95 (d, 1H,  $J = 3.0$  Hz, ArH), 7.85 (d, 1H,  $J = 2.7$  Hz, ArH), 2.61 (s, 3H,  $\text{COCH}_3$ ); MS (ESI):  $m/z$  212.7  $[\text{M}-\text{H}]^-$ .

### 5.2.4. 3-Acetyl-5-bromo-2-hydroxybenzonitrile (6d)

Yield 26%; mp  $75\text{--}77^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  13.17 (s, 1H, OH), 8.41 (d, 1H,  $J = 2.4$  Hz, ArH), 8.32 (d, 1H,  $J = 2.4$  Hz, ArH), 2.68 (s, 3H,  $\text{COCH}_3$ ); MS (ESI):  $m/z$  239.6  $[\text{M}-\text{H}]^-$ .

### 5.2.5. 3-Acetyl-2-hydroxy-5-methylbenzoic acid (6e)

Yield 28%; mp  $>180^\circ\text{C}$  (dec);  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  7.80 (s, 1H, ArH), 7.64 (s, 1H, ArH), 2.59 (s, 3H,  $\text{COCH}_3$ ), 2.24 (s, 3H,  $\text{CH}_3$ ); MS (ESI):  $m/z$  192.7  $[\text{M}-\text{H}]^-$ .

## 5.3. General procedure for the synthesis of chalcones (10–57)

Equimolar amounts of aromatic aldehydes and intermediates **6a–6e** were dissolved in 10 mL of ethanol, and 5 mL of 25% aq NaOH was added. More solvent was added to help stirring in cases of excessive precipitation. The reaction mixture was stirred at rt for 1–5 days or until completion of reaction (6 h for compound **56**). The mixture was then poured into ice and acidified with 3 N HCl to pH 4. The yellow precipitate formed was filtered and washed with water. The crude solid was purified by column chromatography (10–25% ethyl acetate/hexane) and/or recrystallized with ethanol or THF/water to yield pure chalcones.

### 5.3.1. 5-Bromo-2-hydroxy-3-(3-phenyl-acryloyl)benzoic acid (10)

Yellow solid, yield 58%; mp  $210\text{--}212^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{MeOH}-d_4$ ):  $\delta$  8.08 (d, 1H,  $J = 2.4$  Hz, ArH), 8.04 (d, 1H,  $J = 2.7$  Hz, ArH), 7.81 (dd, 2H,  $J_1 = 2.1$  Hz,  $J_2 = 7.5$  Hz, ArH), 7.68 (s, 2H, H- $\alpha$  and H- $\beta$ ), 7.42–7.44 (m, 3H, ArH); MS (ESI):  $m/z$  346.6  $[\text{M}-\text{H}]^-$ ; HPLC:  $t_R$  7.71 min, purity 99.10%.

### 5.3.2. 5-Bromo-3-[3-(2-chlorophenyl)-acryloyl]-2-hydroxybenzoic acid (11)

Yellow solid, yield 60%; mp  $211\text{--}214^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.08 (d, 1H,  $J = 2.7$  Hz, ArH), 8.03 (d, 1H,  $J = 2.7$  Hz, ArH), 8.01 (dd, 1H,  $J_1 = 1.5$  Hz,  $J_2 = 7.5$  Hz, ArH), 7.94 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.78 (d, 1H,  $J = 15.9$  Hz, H- $\alpha$ ), 7.59 (dd, 1H,

$J_1 = 1.8$  Hz,  $J_2 = 8.4$  Hz, ArH), 7.46 (ddd, 2H,  $J_1 = 1.8$  Hz,  $J_2 = 7.2$  Hz,  $J_3 = 7.5$  Hz, ArH); MS (ESI):  $m/z$  378.6, 380.6  $[\text{M}-\text{H}]^-$  HPLC:  $t_R$  8.05 min, purity 98.82%.

### 5.3.3. 5-Bromo-3-[3-(2-fluorophenyl)-acryloyl]-2-hydroxybenzoic acid (12)

Yellow solid, yield 47%; mp  $185\text{--}187^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.05 (d, 1H,  $J = 2.1$  Hz, ArH), 8.0 (s, 1H, H- $\beta$ ), 7.93 (t, 1H,  $J = 7.2$  Hz, ArH), 7.74 (s, 1H, H- $\alpha$ ), 7.73 (d, 1H,  $J = 2.7$  Hz, ArH), 7.51 (d, 1H,  $J = 6.9$  Hz, ArH), 7.31 (dd, 1H,  $J_1 = 7.2$  Hz,  $J_2 = 8.7$  Hz, ArH), 7.29 (t, 1H,  $J = 7.2$  Hz, ArH); MS (ESI):  $m/z$  362.7, 364.7  $[\text{M}-\text{H}]^-$  HPLC:  $t_R$  7.24 min, purity 99.39%.

### 5.3.4. 5-Bromo-3-[3-(3-chlorophenyl)-acryloyl]-2-hydroxybenzoic acid (13)

Yellow solid, yield 56%; mp  $234\text{--}236^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.06 (d, 1H,  $J = 2.7$  Hz, ArH), 8.01 (d, 1H,  $J = 2.7$  Hz, ArH), 7.92 (s, 1H, ArH), 7.76 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.75 (t, 1H,  $J = 7.8$  Hz, ArH), 7.63 (d, 1H,  $J = 15.9$  Hz, H- $\alpha$ ), 7.50 (dd, 2H,  $J_1 = 1.5$  Hz,  $J_2 = 8.7$  Hz, ArH); MS (ESI):  $m/z$  378.6, 380.6  $[\text{M}-\text{H}]^-$ . HPLC:  $t_R$  7.67 min, purity 99.30%.

### 5.3.5. 5-Bromo-3-[3-(4-chlorophenyl)-acryloyl]-2-hydroxybenzoic acid (14)

Yellow solid, yield 62%; mp  $238\text{--}241^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.04 (d, 1H,  $J = 15.3$  Hz, H- $\beta$ ), 7.91 (d, 1H,  $J = 2.4$  Hz, ArH), 7.74 (d, 2H,  $J = 8.4$  Hz, ArH), 7.71 (d, 1H,  $J = 3.0$  Hz, ArH), 7.58 (d, 1H,  $J = 15.3$  Hz, H- $\alpha$ ), 7.50 (d, 2H,  $J = 8.4$  Hz, ArH); MS (ESI):  $m/z$  380.6  $[\text{M}-\text{H}]^-$ . HPLC:  $t_R$  8.1 min, purity 99.10%.

### 5.3.6. 5-Bromo-3-[3-(4-bromophenyl)-acryloyl]-2-hydroxybenzoic acid (15)

Yellow solid, yield 34%; mp  $236\text{--}238^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.06 (d, 1H,  $J = 2.7$  Hz, ArH), 8.02 (d, 1H,  $J = 2.7$  Hz, ArH), 7.74 (d, 2H,  $J = 8.7$  Hz, ArH), 7.69 (s, 1H, H- $\beta$ ), 7.67 (d, 2H,  $J = 8.7$  Hz, ArH), 7.65 (s, 1H, H- $\alpha$ ); MS (ESI):  $m/z$  424.6  $[\text{M}-\text{H}]^-$ . HPLC:  $t_R$  6.05 min, purity: 99.02%.

### 5.3.7. 5-Bromo-2-hydroxy-3-[3-(4-iodophenyl)-acryloyl]benzoic acid (16)

Yellow solid, yield 48%; mp  $237\text{--}239^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.06 (d, 1H,  $J = 2.7$  Hz, ArH), 8.01 (d, 1H,  $J = 2.7$  Hz, ArH), 7.83 (d, 1H,  $J = 8.4$  Hz, ArH), 7.73 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.69 (d, 1H,  $J = 8.4$  Hz, ArH), 7.60 (d, 1H,  $J = 15.3$  Hz, H- $\alpha$ ), 7.59 (d, 2H,  $J = 8.4$  Hz, ArH); MS (ESI):  $m/z$  472.6  $[\text{M}-\text{H}]^-$ . HPLC:  $t_R$  6.46 min, purity 94.70%.

### 5.3.8. 5-Bromo-3-[3-(2,4-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (17)

Yellow solid, yield 54%; mp  $234\text{--}235^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.08 (d, 1H,  $J = 2.7$  Hz, ArH), 8.05 (d, 1H,  $J = 8.4$  Hz, ArH), 8.05 (d, 1H,  $J = 2.7$  Hz, ArH), 7.87 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.76 (d, 1H,  $J = 15.9$  Hz, H- $\alpha$ ), 7.76 (d, 1H,  $J = 2.1$  Hz, ArH), 7.52 (dd, 1H,  $J_1 = 1.5$  Hz,  $J_2 = 8.4$  Hz, ArH); MS (ESI):  $m/z$  414.8  $[\text{M}-\text{H}]^-$ . HPLC:  $t_R$  10.91 min, purity 98.23%.

### 5.3.9. 5-Bromo-3-[3-(2,3-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (18)

Yellow solid, yield 50%; mp  $226\text{--}228^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.08 (d, 1H,  $J = 2.7$  Hz, ArH), 8.03 (d, 1H,  $J = 2.4$  Hz, ArH), 7.98 (d, 1H,  $J = 8.1$  Hz, ArH), 7.91 (s, 1H, H- $\beta$ ), 7.77 (d, 1H,  $J = 15.6$  Hz, H- $\alpha$ ), 7.74 (d, 1H,  $J = 7.8$  Hz, ArH), 7.46 (t, 1H,  $J = 7.8$  Hz, ArH); MS (ESI):  $m/z$  414.8  $[\text{M}-\text{H}]^-$ . HPLC:  $t_R$  6.46 min, purity: 99.08%.

**5.3.10. 5-Bromo-3-[3-(2,3-dimethoxyphenyl)-acryloyl]-2-hydroxybenzoic acid (19)**

Yellow solid, yield 51%; mp 195–197 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.02 (s, 1H, ArH), 7.92 (s, 1H, ArH), 7.86 (d, 1H, *J* = 16.2 Hz, H-β), 7.75 (d, 1H, *J* = 15.9 Hz, H-α), 7.40 (dd, 1H, *J*<sub>1</sub> = 3.0 Hz, *J*<sub>2</sub> = 5.7 Hz, ArH), 7.13–7.15 (m, 2H, ArH), 3.83 (s, 3H, OCH<sub>3</sub>), 3.77 (s, 3H, OCH<sub>3</sub>); MS (ESI): *m/z* 406.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.03 min, purity: 99.7%.

**5.3.11. 5-Bromo-2-hydroxy-3-[3-(2,3,4-trimethoxyphenyl)-acryloyl]-benzoic acid (20)**

Yellow solid, yield 49%; mp 194–195 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.05 (d, 1H, *J* = 1.8 Hz, ArH), 8.02 (d, 1H, *J* = 1.8 Hz, ArH), 7.80 (d, 1H, *J* = 15.9 Hz, H-β), 7.63 (d, 1H, *J* = 8.7 Hz, ArH), 7.10 (d, 1H, *J* = 15.9 Hz, H-α), 6.92 (d, 1H, *J* = 8.7 Hz, ArH), 3.87 (s, 1H, OCH<sub>3</sub>), 3.84 (s, 1H, OCH<sub>3</sub>), 3.77 (s, 1H, OCH<sub>3</sub>); MS (ESI): *m/z* 436.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.88 min, purity: 99.36%.

**5.3.12. 5-Bromo-3-[3-(2,5-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (21)**

Yellow solid, yield 51%; mp 232–234 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.16 (d, 1H, *J* = 2.4 Hz, ArH), 8.11 (d, 1H, *J* = 2.7 Hz, ArH), 8.0 (d, 1H, *J* = 2.7 Hz, ArH), 7.87 (s, 2H, H-α and H-β), 7.65 (d, 1H, *J* = 8.4 Hz, ArH), 7.57 (dd, 1H, *J*<sub>1</sub> = 2.4 Hz, *J*<sub>2</sub> = 8.4 Hz, ArH); MS (ESI): *m/z* 414.7 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 8.03 min, purity 97.30%.

**5.3.13. 5-Bromo-3-[3-(2,6-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (22)**

Yellow solid, yield 54%; mp 235–237 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.08 (d, 1H, *J* = 2.7 Hz, ArH), 7.96 (d, 1H, *J* = 2.7 Hz, ArH), 7.79 (d, 1H, *J* = 16.5 Hz, H-β), 7.67 (d, 1H, *J* = 16.5 Hz, H-α), 7.60 (d, 2H, *J* = 8.1 Hz, ArH), 7.41–7.46 (t, 1H, *J* = 8.7 Hz, ArH); MS (ESI): *m/z* 414.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.96 min, purity 96.60%.

**5.3.14. 5-Bromo-3-[3-(3,4-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (23)**

Yellow solid, yield 48%; mp 218–221 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.07 (d, 1H, *J* = 15.9 Hz, H-β), 8.0 (s, 1H, ArH), 7.93 (d, 1H, *J* = 2.7 Hz, ArH), 7.72 (d, 1H, *J* = 2.7 Hz, ArH), 7.70 (d, 2H, *J* = 7.5 Hz), 7.56 (d, 1H, *J* = 15.9 Hz, H-α); MS (ESI): *m/z* 414.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.25 min, purity 97.0%.

**5.3.15. 5-Bromo-2-hydroxy-3-[3-(2,3,5-trichlorophenyl)-acryloyl]benzoic acid (24)**

Yellow solid, yield 54%; mp 212–214 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.10 (d, 1H, *J* = 2.4 Hz, ArH), 8.08 (d, 1H, *J* = 2.7 Hz, ArH), 8.05 (d, 1H, *J* = 2.7 Hz, ArH), 7.96 (d, 1H, *J* = 2.4 Hz, ArH), 7.84 (br s, 2H, H-α and H-β); MS (ESI): *m/z* 448.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.41 min, purity 98.21%.

**5.3.16. 5-Bromo-2-hydroxy-3-[3-(2,3,6-trichlorophenyl)-acryloyl]benzoic acid (25)**

Yellow solid, yield 57%; mp 220–223 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.07 (d, 1H, *J* = 2.7 Hz, ArH), 7.97 (d, 1H, *J* = 2.4 Hz, ArH), 7.73 (d, 1H, *J* = 8.7 Hz, ArH), 7.71 (d, 1H, *J* = 16.2 Hz, H-β), 7.61–7.64 (m, 2H, H-α and ArH); MS (ESI): *m/z* 448.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.13 min, purity 99.20%.

**5.3.17. 5-Chloro-3-[3-(4-chlorophenyl)-acryloyl]-2-hydroxybenzoic acid (26)**

Yellow solid, yield 43%; mp 213–215 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 7.95 (d, 1H, *J* = 2.1 Hz, ArH), 7.93 (d, 1H, *J* = 2.7 Hz, ArH), 7.84 (dd, 2H, *J*<sub>1</sub> = 1.5 Hz, *J*<sub>2</sub> = 6.9 Hz, ArH), 7.66 (s, 2H, H-β and H-α), 7.51 (dd, 2H, *J*<sub>1</sub> = 1.5 Hz, *J*<sub>2</sub> = 6.9 Hz, ArH); MS (ESI): *m/z* 336.9 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.13 min, purity 99.55%.

**5.3.18. 5-Chloro-3-[3-(2,4-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (27)**

Yellow solid, yield 48%; mp 248–250 °C; <sup>1</sup>H NMR (300 MHz, MeOH-*d*<sub>4</sub>): δ 8.05 (d, 1H, *J* = 15.9 Hz, H-β), 7.93 (d, 1H, *J* = 8.7 Hz, ArH), 7.84 (d, 1H, *J* = 2.4 Hz, ArH), 7.80 (d, 1H, *J* = 15.3 Hz, H-α), 7.73 (d, 1H, *J* = 2.1 Hz, ArH), 7.65 (d, 1H, *J* = 2.7 Hz, ArH), 7.52 (dd, 1H, *J*<sub>1</sub> = 2.1 Hz, *J*<sub>2</sub> = 8.4 Hz, ArH); MS (ESI): *m/z* 370.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.76 min, purity 99.14%.

**5.3.19. 5-Chloro-3-[3-(2,3-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (28)**

Yellow solid, yield 41%; mp 250–252 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.10 (d, 1H, *J* = 15.3 Hz, H-β), 7.85–7.89 (m, 3H, H-α and ArH), 7.70 (d, 1H, *J* = 7.2 Hz, ArH), 7.62 (s, 1H, ArH), 7.45 (t, 1H, *J* = 7.5 Hz, ArH); MS (ESI): *m/z* 370.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.06 min, purity 96.61%.

**5.3.20. 5-Chloro-3-[3-(2,5-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (29)**

Yellow solid, yield 42%; mp 247–249 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.20 (d, 1H, *J* = 1.2 Hz, ArH), 8.04 (s, 2H, H-β and ArH), 7.90 (s, 2H, H-α and ArH), 7.67 (d, 1H, *J* = 8.4 Hz, ArH), 7.61 (dd, 1H, *J*<sub>1</sub> = 1.5 Hz, *J*<sub>2</sub> = 8.4 Hz, ArH); MS (ESI): *m/z* 370.7 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 8.39 min, purity 99.0%.

**5.3.21. 5-Chloro-2-hydroxy-3-[3-(2,3,5-trichlorophenyl)-acryloyl]benzoic acid (30)**

Yellow solid, yield 37%; mp 235–237 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.05 (d, 1H, *J* = 15.9 Hz, H-β), 7.91 (d, 2H, *J* = 1.2 Hz, ArH), 7.82 (d, 1H, *J* = 3.0 Hz, ArH), 7.75 (d, 1H, *J* = 15.6 Hz, H-α), 7.60 (d, 1H, *J* = 2.7 Hz, ArH); MS (ESI): *m/z* 404.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.18 min, purity 98.81%.

**5.3.22. 5-Chloro-2-hydroxy-3-[3-(2,3,6-trichlorophenyl)-acryloyl]benzoic acid (31)**

Yellow solid, yield 49%; mp 229–231 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 7.99 (d, 1H, *J* = 16.2 Hz, H-β), 7.86 (d, 1H, *J* = 2.1 Hz, ArH), 7.71 (d, 1H, *J* = 2.7 Hz, ArH), 7.69 (d, 1H, *J* = 7.8 Hz, ArH), 7.65 (d, 1H, *J* = 8.4 Hz, ArH), 7.60 (d, 1H, *J* = 16.2 Hz, H-α); MS (ESI): *m/z* 404.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.58 min, purity 95.66%.

**5.3.23. 5-Chloro-3-[3-(3,4-dichlorophenyl)-acryloyl]-2-hydroxybenzoic acid (32)**

Yellow solid, yield 54%; mp >240 °C (dec); <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.05 (d, 1H, *J* = 15.9 Hz, H-β), 7.99 (s, 1H, ArH), 7.80 (d, 1H, *J* = 3.0 Hz, ArH), 7.71 (d, 2H, *J* = 6.0 Hz, ArH), 7.58 (d, 1H, *J* = 3.0 Hz, ArH), 7.56 (d, 1H, *J* = 15.9 Hz, H-α); MS (ESI): *m/z* 370.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.78 min, purity 99.60%.

**5.3.24. 3-[3-(4-Chlorophenyl)-acryloyl]-5-fluoro-2-hydroxybenzoic acid (33)**

Yellow solid, yield 56%; mp 198–200 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 7.91 (d, 1H, *J* = 15.9 Hz, H-β), 7.60 (d, 1H, *J* = 15.9 Hz, H-α), 7.68 (dd, 1H, *J*<sub>1</sub> = 3.3 Hz, *J*<sub>2</sub> = 8.4 Hz, ArH), 7.76 (d, 2H, *J* = 8.4 Hz, ArH), 7.49 (d, 2H, *J* = 8.7 Hz, ArH), 7.54 (dd, 1H, *J*<sub>1</sub> = 3.3 Hz, *J*<sub>2</sub> = 9.3 Hz, ArH); MS (ESI): *m/z* 318.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.11 min, purity 99.11%.

**5.3.25. 3-[3-(2,4-Dichlorophenyl)-acryloyl]-5-fluoro-2-hydroxybenzoic acid (34)**

Yellow solid, yield 41%; mp 260–262 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.12 (d, 1H, *J* = 15.6 Hz, H-β), 7.93 (d, 1H, *J* = 8.4 Hz, ArH), 7.81 (d, 1H, *J* = 15.9 Hz, H-α), 7.74 (d, 1H, *J* = 2.1 Hz, ArH), 7.65 (dd, 1H, *J*<sub>1</sub> = 3.3 Hz, *J*<sub>2</sub> = 8.7 Hz, ArH), 7.52 (dd, 1H,

$J_1 = 2.1$  Hz,  $J_2 = 8.4$  Hz, ArH), 7.43 (dd, 1H,  $J_1 = 3.3$  Hz,  $J_2 = 9.3$  Hz, ArH); MS (ESI):  $m/z$  352.8 [M–H]<sup>–</sup>. HPLC:  $t_R$  6.51 min, purity 99.5%.

### 5.3.26. 3-[3-(2,3-Dichlorophenyl)-acryloyl]-5-fluoro-2-hydroxybenzoic acid (35)

Yellow solid, yield 47%; mp 222–224 °C; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  7.94 (d, 1H,  $J = 7.8$  Hz, ArH), 7.90 (s, 2H, H- $\alpha$  and H- $\beta$ ), 7.77 (dd, 1H,  $J_1 = 3.0$  Hz,  $J_2 = 8.5$  Hz, ArH), 7.73 (d, 1H,  $J = 8.0$  Hz, ArH), 7.46 (t, 1H,  $J = 8.0$  Hz, ArH), 7.66 (dd, 1H,  $J_1 = 2.5$  Hz,  $J_2 = 9.0$  Hz, ArH); MS (ESI):  $m/z$  352.8 [M–H]<sup>–</sup>. HPLC:  $t_R$  5.68 min, purity 97.20%.

### 5.3.27. 3-[3-(2,5-Dichlorophenyl)-acryloyl]-5-fluoro-2-hydroxybenzoic acid (36)

Yellow solid, yield 59%; mp 238–240 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  8.07 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.93 (d, 1H,  $J = 2.4$  Hz, ArH), 7.76 (d, 1H,  $J = 15.9$  Hz, H- $\alpha$ ), 7.66 (dd, 1H,  $J_1 = 3.6$  Hz,  $J_2 = 8.4$  Hz, ArH), 7.60 (d, 1H,  $J = 8.7$  Hz, ArH), 7.52 (dd, 1H,  $J_1 = 2.4$  Hz,  $J_2 = 8.7$  Hz, ArH), 7.42 (dd, 1H,  $J_1 = 3.6$  Hz,  $J_2 = 9.3$  Hz, ArH); MS (ESI):  $m/z$  352.8 [M–H]<sup>–</sup>. HPLC:  $t_R$  6.02 min, purity 98.43%.

### 5.3.28. 5-Fluoro-2-hydroxy-3-[3-(2,3,5-trichlorophenyl)-acryloyl]-benzoic acid (37)

Yellow solid, yield 53%; mp 231–233 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  8.02 (d, 1H,  $J = 2.4$  Hz, ArH), 7.95 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.92 (d, 1H,  $J = 2.4$  Hz, ArH), 7.79 (d, 1H,  $J = 15.6$  Hz, H- $\alpha$ ), 7.74 (dd, 1H,  $J_1 = 3.3$  Hz,  $J_2 = 8.4$  Hz, ArH), 7.64 (dd, 1H,  $J_1 = 3.3$  Hz,  $J_2 = 9$  Hz, ArH); MS (ESI):  $m/z$  388.6 [M–H]<sup>–</sup>. HPLC:  $t_R$  8.11 min, purity 97.86%.

### 5.3.29. 5-Fluoro-2-hydroxy-3-[3-(2,3,6-trichlorophenyl)acryloyl]benzoic acid (38)

Yellow solid, yield 46%; mp 235–237 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  8.11 (d, 1H,  $J = 16.2$  Hz, H- $\beta$ ), 7.70 (d, 1H,  $J = 8.7$  Hz, ArH), 7.62 (d, 1H,  $J = 8.7$  Hz, ArH), 7.58 (d, 1H,  $J = 16.2$  Hz, H- $\alpha$ ), 7.66 (dd, 1H,  $J_1 = 3.6$  Hz,  $J_2 = 8.4$  Hz, ArH), 7.44 (dd, 1H,  $J_1 = 3.6$  Hz,  $J_2 = 9.3$  Hz, ArH); MS (ESI):  $m/z$  388.6 [M–H]<sup>–</sup>. HPLC:  $t_R$  7.57 min, purity 97.57%.

### 5.3.30. 5-Bromo-3-[3-(3,5-dimethylphenyl)-acryloyl]-2-hydroxybenzoic acid (39)

Yellow solid, yield 51%; mp 223–225 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  8.06 (d, 1H,  $J = 2.7$  Hz, ArH), 7.99 (d, 1H,  $J = 1.8$  Hz, ArH), 7.61 (d, 1H,  $J = 16.5$  Hz, H- $\beta$ ), 7.58 (d, 1H,  $J = 16.5$  Hz, H- $\alpha$ ), 7.40 (s, 2H, H<sub>z</sub>, ArH), 7.10 (s, 1H, ArH), 2.32 (s, 6H, CH<sub>3</sub>); MS (ESI):  $m/z$  374.9 [M–H]<sup>–</sup>. HPLC:  $t_R$  6.59 min, purity 99.0%.

### 5.3.31. 5-Bromo-3-[3-(2-chloro-3,4-dimethoxyphenyl)-acryloyl]-2-hydroxybenzoic acid (40)

Yellow solid, yield 21%; mp 217–220 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  8.05 (d, 1H,  $J = 2.7$  Hz, ArH), 8.00 (d, 1H,  $J = 2.7$  Hz, ArH), 7.90 (d, 1H,  $J = 15.6$  Hz, H- $\beta$ ), 7.81 (d, 1H,  $J = 8.7$  Hz, ArH), 7.68 (d, 1H,  $J = 15.6$  Hz, H- $\alpha$ ), 7.16 (d, 1H,  $J = 8.7$  Hz, ArH), 3.90 (s, 3H, OCH<sub>3</sub>), 3.77 (s, 3H, OCH<sub>3</sub>); MS (ESI):  $m/z$  440.8 [M–H]<sup>–</sup>. HPLC:  $t_R$  6.46 min, purity 99.0%.

### 5.3.32. 5-Bromo-3-[3-(2-fluoro-4,5-dimethoxyphenyl)-acryloyl]-2-hydroxybenzoic acid (41)

Yellow solid, yield 32%; mp 193–195 °C; <sup>1</sup>H NMR (300 Hz, DMSO-*d*<sub>6</sub>):  $\delta$  8.03 (d, 1H,  $J = 2.7$  Hz, ArH), 7.93 (d, 1H,  $J = 2.7$  Hz, ArH), 7.66 (s, 2H, H- $\beta$  and H- $\alpha$ ), 7.38 (d, 1H,  $J = 7.2$  Hz, ArH), 7.00 (d, 1H,  $J = 12.3$  Hz, ArH), 3.84 (s, 3H, OCH<sub>3</sub>), 3.82 (s, 3H, OCH<sub>3</sub>); MS (ESI):  $m/z$  424.8 [M–H]<sup>–</sup>. HPLC:  $t_R$  7.19 min, purity 97.51%.

### 5.3.33. 3-[3-(2-Benzyloxyphenyl)-acryloyl]-5-bromo-2-hydroxybenzoic acid (42)

Yellow solid, yield 26%; mp >165 °C (dec); <sup>1</sup>H NMR (300 MHz, MeOH-*d*<sub>4</sub>):  $\delta$  7.92–7.95 (m, 2H, H- $\beta$  and ArH), 7.72 (d, 1H,  $J = 7.5$  Hz, ArH), 7.65 (d, 1H,  $J = 2.7$  Hz, ArH), 7.47 (d, 2H,  $J = 7.2$  Hz, ArH), 7.30–7.42 (m, 5H, H- $\alpha$ , and ArH), 7.20 (d, 1H,  $J = 8.1$  Hz, ArH), 7.0 (t, 1H,  $J = 7.5$  Hz, ArH), 5.22 (s, 2H, CH<sub>2</sub>); MS (ESI):  $m/z$  452.8 [M–H]<sup>–</sup>. HPLC:  $t_R$  7.83 min, purity 95.65%.

### 5.3.34. 5-Bromo-3-[3-(4-chlorophenoxy)-phenyl]acryloyl]-2-hydroxybenzoic acid (43)

Yellow solid, yield 42%; mp 162–165 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  7.94 (s, 2H, ArH), 7.74 (d, 1H,  $J = 6.9$  Hz, ArH), 7.70 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.40–7.46 (m, 6H, H- $\alpha$  and ArH), 7.20 (d, 1H,  $J = 7.8$  Hz, ArH), 7.0 (t, 1H,  $J = 6.9$  Hz, ArH); (ESI):  $m/z$  472.6 [M–H]<sup>–</sup>. HPLC:  $t_R$  5.86 min, purity 96.8%.

### 5.3.35. 5-Bromo-3-[3-(4-(4-chlorophenoxy)phenyl)acryloyl]-2-hydroxybenzoic acid (44)

Yellow solid, yield 35%; mp 219–220 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  8.06 (d, 1H,  $J = 2.1$  Hz, ArH), 8.02 (d, 1H,  $J = 2.1$  Hz, ArH), 7.84 (d, 2H,  $J = 8.4$  Hz, ArH), 7.67 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.59 (d, 1H,  $J = 15.9$  Hz, H- $\alpha$ ), 7.48 (d, 2H,  $J = 8.7$  Hz, ArH), 7.13 (d, 2H,  $J = 8.7$  Hz, ArH), 7.07 (d, 2H,  $J = 8.7$  Hz, ArH); MS (ESI):  $m/z$  472.6 [M–H]<sup>–</sup>. HPLC:  $t_R$  7.23 min, purity 99.44%.

### 5.3.36. 5-Bromo-2-hydroxy-3-(3-thiophen-3-yl-acryloyl)-benzoic acid (45)

Yellow solid, yield 54%; mp 207–210 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  7.97 (s, 2H, ArH), 7.75 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 7.65 (s, 1H, ArH), 7.61 (s, 1H, ArH), 7.58 (d, 1H,  $J = 15.6$  Hz, H- $\alpha$ ), 7.48 (d, 1H,  $J = 4.2$  Hz, ArH); MS (ESI):  $m/z$  352.6 [M–H]<sup>–</sup>. HPLC:  $t_R$  6.91 min, purity 99.71%.

### 5.3.37. 5-Bromo-3-[3-(3-cyclopentyloxy-4-methoxyphenyl)-acryloyl]-2-hydroxybenzoic acid (46)

Yellow solid, yield 31%; mp 233–235 °C; <sup>1</sup>H NMR (300 MHz, MeOH-*d*<sub>4</sub>):  $\delta$  7.91 (d, 1H,  $J = 2.7$  Hz, ArH), 7.82 (d, 1H,  $J = 15.6$  Hz, H- $\beta$ ), 7.67 (d, 1H,  $J = 3.0$  Hz, ArH), 7.59 (d, 1H,  $J = 15.9$  Hz, H- $\alpha$ ), 7.26 (d, 1H,  $J = 8.7$  Hz, ArH), 7.25 (s, 1H, ArH), 6.99 (d, 1H,  $J = 8.1$  Hz, ArH), 4.86 (p, 1H,  $J = 5.7$  Hz, cyclopentyl), 3.79 (s, 3H, CH<sub>3</sub>) 1.87–1.92 (m, 2H, cyclopentyl), 1.66–1.72 (m, 4H, cyclopentyl), 1.57–1.60 (m, 2H, cyclopentyl); MS (ESI):  $m/z$  460.8 [M–H]<sup>–</sup>. HPLC:  $t_R$  7.19 min, purity 97.51%.

### 5.3.38. 5-Bromo-3-[3-(2-ethoxynaphthalen-1-yl)acryloyl]-2-hydroxybenzoic acid (47)

Yellow solid, yield 23%; mp 216–218 °C; <sup>1</sup>H NMR (300 MHz, MeOH-*d*<sub>4</sub>):  $\delta$  8.48 (d, 1H,  $J = 15.9$  Hz, H- $\beta$ ), 8.30 (d, 1H,  $J = 8.4$  Hz, ArH), 8.19 (d, 1H,  $J = 2.7$  Hz, ArH), 8.17 (d, 1H,  $J = 15.9$  Hz, H- $\alpha$ ), 7.92 (d, 1H,  $J = 9.0$  Hz, ArH), 7.88 (d, 1H,  $J = 2.7$  Hz, ArH), 7.84 (d, 1H,  $J = 8.4$  Hz, ArH), 7.55 (t, 1H,  $J = 7.2$  Hz, ArH), 7.37–7.44 (m, 2H, ArH), 4.33 (q, 2H,  $J = 6.9$  Hz, CH<sub>2</sub>), 1.55 (t, 3H,  $J = 6.9$  Hz, CH<sub>3</sub>); MS (ESI):  $m/z$  440.9 [M–H]<sup>–</sup>. HPLC:  $t_R$  8.53 min, purity 95.0%.

### 5.3.39. 5-Bromo-2-hydroxy-3-[3-(6-methoxynaphthalen-2-yl)-acryloyl]benzoic acid (48)

Yellow solid, yield 15%; mp 239–241 °C; <sup>1</sup>H NMR (300 Hz, MeOH-*d*<sub>4</sub>):  $\delta$  8.17 (d, 1H,  $J = 2.4$  Hz, ArH), 8.01 (s, 1H, H- $\beta$ ), 7.86 (d, 1H,  $J = 7.9$  Hz, ArH), 7.84 (d, 2H,  $J = 8.1$  Hz, ArH), 7.83 (d, 1H,  $J = 2.4$  Hz, ArH), 7.80 (s, 2H, H- $\alpha$ , and ArH), 7.26 (d, 1H,  $J = 2.4$  Hz, ArH), 7.16 (dd, 1H,  $J_1 = 8.7$  Hz,  $J_2 = 2.4$  Hz, ArH), 3.93 (s, 3H, OCH<sub>3</sub>); MS (ESI):  $m/z$  426.8 [M–H]<sup>–</sup>. HPLC:  $t_R$  6.07 min, purity 98.93%.

**5.3.40. 3-(3-Benzo[b]thiophen-5-yl-acryloyl)-5-bromo-2-hydroxybenzoic acid (49)**

Yellow solid, yield 21%; mp 235–237 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.21 (s, 1H, ArH), 8.1 (d, 1H, *J* = 15.9 Hz, H-β), 8.0 (s, 1H, ArH), 7.92 (d, 1H, *J* = 2.7 Hz, ArH), 7.82 (d, 1H, *J* = 5.4 Hz, ArH), 7.73 (d, 1H, *J* = 15.9 Hz, H-α), 7.72 (d, 1H, *J* = 3 Hz, ArH), 7.72 (s, 1H, ArH), 7.53 (d, 1H, *J* = 5.1 Hz, ArH); MS (ESI): *m/z* 402.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 5.53 min, purity 98.31%.

**5.3.41. 5-Bromo-3-[3-(2-furan-2-yl-phenyl)-acryloyl]-2-hydroxybenzoic acid (50)**

Yellow solid, yield 36%; mp 208–209 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 7.92 (m, 3H, H-β and ArH), 7.87 (s, 1H, H-α), 7.81 (d, 1H, *J* = 2.7 Hz, ArH), 7.70 (s, 1H, ArH), 7.68 (d, 1H, 9.6 Hz, ArH), 7.48 (t, 2H, *J* = 7.5 Hz, ArH), 6.66 (s, 1H, ArH), 6.63 (s, 1H, ArH); MS (ESI): *m/z* 412.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 5.77 min, purity 95.02%.

**5.3.42. 5-Bromo-3-[3-(6-chlorobenzo[1,3]dioxol-5-yl)-acryloyl]-2-hydroxybenzoic acid (51)**

Yellow solid, yield 18%; mp 228–231 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.10 (s, 1H, ArH), 8.07 (s, 1H, ArH), 7.90 (d, 1H, *J* = 15 Hz, H-β), 7.66 (s, 1H, ArH), 7.63 (d, 1H, *J* = 14.7 Hz, H-α), 7.21 (s, 1H, ArH), 6.16 (s, 2H, CH<sub>2</sub>); MS (ESI): *m/z* 424.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.82 min, purity 99.0%.

**5.3.43. 5-Bromo-2-hydroxy-3-[3-(3-trifluoromethoxyphenyl)-acryloyl]benzoic acid (52)**

Yellow solid, yield 44%; mp 217–219 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.07 (d, 1H, *J* = 2.7 Hz, ArH), 8.03 (d, 1H, *J* = 2.7 Hz, ArH), 7.80–7.84 (br s, 2H, H-α and H-β), 7.72 (s, 1H, ArH), 7.70 (s, 1H, *J* = 4.0 Hz, ArH), 7.58 (t, 1H, *J* = 7.8 Hz, ArH), 7.45 (d, 1H, *J* = 7.5 Hz, ArH); MS (ESI): *m/z* 430.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.06 min, purity 99.30%.

**5.3.44. 5-Bromo-2-hydroxy-3-[3-(3-nitrophenyl)-acryloyl]benzoic acid (53)**

Yellow solid, yield 62%; mp 200–202 °C; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 8.62 (s, 1H, ArH), 8.26 (s, 2H, ArH), 8.07 (s, 1H, H-β), 8.02 (s, 1H, H-α), 7.81 (d, 2H, *J* = 7.0 Hz, ArH), 7.75 (t, 1H, *J* = 7.0 Hz, ArH); MS (ESI): *m/z* 391.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 11.47 min, purity 95.59%.

**5.3.45. 5-Bromo-3-[3-(3-bromophenyl)-acryloyl]-2-hydroxybenzoic acid (54)**

Yellow solid, yield 31%; mp 235–237 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.0 (d, 1H, *J* = 15.9 Hz, H-β), 7.92 (d, 1H, *J* = 2.7 Hz, ArH), 7.90 (s, 1H, ArH), 7.72 (d, 1H, *J* = 6.6 Hz, ArH), 7.70 (d, 1H, *J* = 2.4 Hz, ArH), 7.59 (d, 1H, *J* = 8.4 Hz, ArH), 7.55 (d, 1H, *J* = 15.9 Hz, H-α), 7.39 (t, 1H, *J* = 7.8 Hz, ArH); MS (ESI): *m/z* 424.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.14 min, purity 98.81%.

**5.3.46. 5-Bromo-3-[3-(3-fluorophenyl)-acryloyl]-2-hydroxybenzoic acid (55)**

Yellow solid, yield 59%; mp 190–193 °C; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 8.0 (d, 1H, *J* = 2.0 Hz, ArH), 7.93 (s, 1H, ArH), 7.80 (d, 1H, *J* = 16.0 Hz, H-β), 7.60–7.77 (m, 3H, H-α and ArH), 7.48 (d, 1H, *J* = 6.5 Hz, ArH), 7.28 (d, 1H, *J* = 8.0 Hz, ArH); MS (ESI): *m/z* 364.7 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 5.39 min, purity 92.5%.

**5.3.47. 5-Bromo-2-hydroxy-3-[3-(2,3,6-trichlorophenyl)-acryloyl]benzonitrile (56)**

Orange solid, yield 74%; mp 139–141 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.34 (d, 1H, *J* = 16.2 Hz, H-β), 7.69 (d, 1H, *J* = 3.0 Hz, ArH), 7.66 (d, 1H, *J* = 9.0 Hz, ArH), 7.58 (d, 1H, *J* = 8.7 Hz, ArH),

7.46 (d, 1H, *J* = 2.4 Hz, ArH), 7.43 (d, 1H, *J* = 16.2 Hz, H-α); MS (ESI): *m/z* 431.6 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.37 min, purity: 95.69%.

**5.3.48. 2-Hydroxy-5-methyl-3-[3-(2,3,6-trichlorophenyl)-acryloyl]benzoic acid (57)**

Yellow solid, yield 58.5%; mp 205–208 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 7.87 (s, 1H, ArH), 7.74 (s, 1H, ArH), 7.70 (d, 1H, *J* = 8.7 Hz, ArH), 7.59–7.64 (m, 3H, H-α, H-β and ArH), 2.29 (s, 3H, CH<sub>3</sub>); MS (ESI): *m/z* 384.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.01 min, purity 97.84%.

**5.4. General procedure for the synthesis of chalcones (58–59)**

A similar procedure to the one mentioned above was followed with appropriate aromatic aldehydes and commercially available 1-(5-bromo-2-hydroxy-3-nitro-phenyl)-ethanone starting material.

**5.4.1. 1-(5-Bromo-2-hydroxy-3-nitrophenyl)-3-(2,3,5-trichlorophenyl)propenone (58)**

Yellow solid, yield 65%; mp 175–178 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 8.13 (d, 1H, *J* = 15.6 Hz, H-β), 7.76 (d, 1H, *J* = 15.9 Hz, H-α), 8.05 (d, 1H, *J* = 2.7 Hz, ArH), 8.0 (d, 2H, *J* = 2.4 Hz, ArH), 7.92 (d, 1H, *J* = 2.4 Hz, ArH); MS (ESI): *m/z* 451.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 5.46 min, purity 97.30%.

**5.4.2. 1-(5-Bromo-2-hydroxy-3-nitrophenyl)-3-(2,3,6-trichlorophenyl)propenone (59)**

Yellow solid, yield 58%; mp 153–154 °C; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 8.32 (s, 1H, ArH), 8.24 (s, 1H, ArH), 7.77 (d, 1H, *J* = 16.5 Hz, H-β), 7.75 (d, 1H, *J* = 9.0 Hz, ArH), 7.65 (d, 1H, *J* = 15.0 Hz, H-α), 7.63 (d, 1H, *J* = 8.0 Hz, ArH); MS (ESI): *m/z* 451.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 6.53 min, purity 98.0%.

**5.5. General procedure for the synthesis of 3-[3-(2,5-dichlorophenyl)propionyl]-5-fluoro-2-hydroxybenzoic acid (60)**

Compound **36** (0.2 g, 0.563 mmol) was dissolved in methanol (10 mL) and Pd/C (10 mol %, 20 mg) was added. The resulting solution was stirred under hydrogen at atmospheric pressure for 2 h at room temperature. After the reaction was complete, the reaction mixture was filtered through celite, and evaporated to dryness. The resulting crude solid was recrystallized from ethanol to give a white solid, yield 43%; mp 163–164 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 7.79 (d, 1H, *J* = 8.4 Hz, ArH), 7.74 (d, 1H, *J* = 9.0 Hz, ArH), 7.49 (s, 1H, ArH), 7.45 (d, 1H, *J* = 8.7 Hz, ArH), 7.31 (d, 1H, *J* = 8.1 Hz, ArH), 3.39 (t, 2H, *J* = 6.9 Hz, CH<sub>2</sub>), 3.02 (t, 2H, *J* = 6.9 Hz, CH<sub>2</sub>); MS (ESI): *m/z* 356.9 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 8.34 min, purity 96.36%.

**5.6. General procedure for the synthesis of 5-bromo-N-(4-fluorobenzyl)-2-hydroxyisophthalamic acid (61)****5.6.1. 5-Bromo-3-formyl-2-hydroxybenzoic acid (7)**

*Method a:* NaOH (5.53 g, 0.138 mol) was added to a solution of 5-bromosalicylic acid (5 g, 23.0 mmol) in CHCl<sub>3</sub> (45 mL) and water (1 mL, 55.55 mmol) at room temperature. The reaction mixture was refluxed for 48 h, diluted with water and acidified with 6 N HCl to pH 1.0. The resulting solution was extracted with ethyl acetate washed with brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The organic layer was evaporated in vacuo to yield a crude product which was purified by column chromatography (25–35% methanol/dichloromethane) to give a white solid, yield 13%.

*Method b:* 3-Formyl salicylic acid hydrate (1 g, 6.0 mmol) was dissolved in dry acetonitrile (20 mL). *N*-Bromo succinimide



(1.02 g, 5.73 mmol) was added and the reaction mixture stirred at rt for 3 h. The white crystals formed were filtered and washed with water. The filtrate was extracted with ethyl acetate, and washed with brine. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness. The two fractions were combined to give intermediate **7** as white crystals, yield 90%; mp 172–174 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 10.27 (s, 1H, CHO), 8.12 (d, 1H, *J* = 2.7 Hz, ArH), 7.93 (d, 1H, *J* = 2.7 ArH); MS (ESI): *m/z* 244.7 [M–H]<sup>–</sup>.

#### 5.6.2. 5-Bromo-*N*-(4-fluorobenzyl)-3-formyl-2-hydroxybenzamide (**8**)

To a solution of intermediate **7** (0.4 g, 1.63 mmol), HOBT (0.441 g, 3.26 mmol) and Et<sub>3</sub>N (0.45 mL, 3.26 mmol) in DCM (20 mL) was added EDCI (0.344 g, 1.8 mmol) at 0 °C. 4-Fluorobenzylamine (0.20 mL, 1.8 mmol) was then added and the reaction mixture stirred overnight at room temperature. The reaction mixture was extracted with ethyl acetate, washed with dilute hydrochloric acid and sodium bicarbonate solution. The organic layer was further washed with brine and dried over sodium sulfate. The resulting crude was purified by recrystallization with ethyl acetate/hexane to give a yellow solid. Yield 41%; mp 135–137 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 10.27 (s, 1H, CHO), 9.90 (s, 1H, NH), 8.41 (s, 1H, ArH), 7.87 (s, 1H, ArH), 7.40 (d, 2H, *J* = 8.1 Hz, ArH), 7.18 (t, 2H, *J* = 8.1 Hz, ArH), 4.52 (s, 2H, CH<sub>2</sub>); MS (ESI): *m/z* 351.9 [M–H]<sup>–</sup>.

#### 5.6.3. 5-Bromo-*N*-(4-fluorobenzyl)-2-hydroxyisophthalamic acid (**61**)

Intermediate **8** (0.2 g, 0.568 mmol) was added to a mixture of water (2.5 mL) and pyridine (2.5 mL). The reaction mixture was heated to 90 °C, KMnO<sub>4</sub> (0.18 g, 1.13 mmol) was added to the hot solution and the heating continued overnight at 90 °C. After completion, the mixture was filtered through celite and evaporated to dryness. The resulting residue was purified by recrystallization from ethyl acetate/hexane. Yield 80%; mp 208–210 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 10.20 (s, 1H, NH), 7.98 (s, 1H, ArH), 7.85 (s, 1H, ArH), 7.13–7.34 (m, 4H, ArH), 4.50 (s, 2H, CH<sub>2</sub>); MS (ESI): *m/z* 367.9 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 4.21 min, purity 97.7%.

#### 5.7. General procedure for the synthesis of *N*-(4-fluoro-benzyl)-2-hydroxy-3-methylbenzamide (**62**) and 5-bromo-*N*-(4-fluoro-benzyl)-2-hydroxybenzamide (**63**)

A similar procedure to the one described for the synthesis of intermediate **8** was followed by condensing 4-fluorobenzyl amine and commercially available 2-hydroxy-3-methylbenzoic acid (**3a**) and 5-bromosalicylic acid (**3b**) to give compounds **62** and **63**, respectively.

##### 5.7.1. *N*-(4-Fluoro-benzyl)-2-hydroxy-3-methylbenzamide (**62**)

Yield 76%; mp 165–166 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 13.10 (s, 1H, OH), 9.43 (t, 1H, *J* = 6.0 Hz, NH), 7.72 (d, 1H, *J* = 7.2 Hz, ArH), 7.37 (d, 1H, *J* = 8.7 Hz, ArH), 7.35 (d, 1H, *J* = 8.4 Hz, ArH), 7.31 (d, 1H, *J* = 7.2 Hz, ArH), 7.15 (t, 2H, *J* = 9.0 Hz, ArH), 6.78 (t, 1H, *J* = 7.5 Hz, ArH), 4.48 (d, 1H, *J* = 6.0 Hz, CH<sub>2</sub>), 2.15 (s, 3H, CH<sub>3</sub>); MS (ESI): *m/z* 257.9 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 8.63 min, purity 99.11%.

##### 5.7.2. 5-Bromo-*N*-(4-fluoro-benzyl)-2-hydroxybenzamide (**63**)

Yield 62%; mp 83–84 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 12.47 (s, 1H, OH), 9.37 (t, 1H, *J* = 5.7 Hz, NH), 8.07 (d, 1H, *J* = 2.4 Hz, ArH), 7.55 (dd, 1H, *J*<sub>1</sub> = 2.4 Hz, *J*<sub>2</sub> = 8.7 Hz, ArH), 7.36 (dd, 2H, *J*<sub>1</sub> = 5.7 Hz, *J*<sub>2</sub> = 8.1 Hz, ArH), 7.15 (t, 2H, *J* = 9.0 Hz, ArH), 6.89 (d, 1H, *J* = 8.7 Hz, ArH), 4.47 (d, 2H, *J* = 5.7 Hz, CH<sub>2</sub>); MS (ESI): *m/z* 323.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 9.7 min, purity 98.66%.

#### 5.8. General procedure for the synthesis of 2-hydroxy-3-(3-phenylpropyl)benzoic acid (**9**)

Compound **10** (0.05 g, 0.195 mmol) was dissolved in ethanol (10 mL) and Pd/C (10 mol %, 5 mg) was added. The resulting solution was stirred under hydrogen at atmospheric pressure for 10 h at room temperature. The reaction mixture was then filtered through celite, and evaporated to dryness. The resulting crude solid was purified using preparative chromatography to give a white solid, yield 72%; mp 125–127 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>): δ 11.66 (s, 1H, OH), 7.67 (dd, 1H, *J*<sub>1</sub> = 1.5 Hz, *J*<sub>2</sub> = 7.8 Hz, ArH), 7.39 (dd, 1H, *J*<sub>1</sub> = 1.5 Hz, *J*<sub>2</sub> = 7.2 Hz, ArH), 7.16–7.30 (m, 5H, ArH), 6.85 (t, 1H, *J* = 7.8 Hz, ArH), 2.62 (t, 4H, *J* = 7.2 Hz, CH<sub>2</sub>), 1.81–1.91 (m, 2H, CH<sub>2</sub>); MS (ESI): *m/z* 254.8 [M–H]<sup>–</sup>. HPLC: *t*<sub>R</sub> 7.61 min, purity 98.05%.

#### 5.9. CoMFA and CoMSIA studies

Three-dimensional structure building and CoMFA and CoMSIA studies were performed on a Dell computer workstation with the Red Hat Linux operating system using the SYBYL program, version 8.0 (Tripos Associates Inc.). Partial atomic charges were assigned with MOPAC. The CoMFA descriptors, steric (Lennard-Jones 6–12 potential) and electrostatic (Coulombic potential) field energies, were calculated using Tripos force field with a distance-dependent dielectric and employing the SYBYL default parameters: 2 Å grid points spacing, an sp<sup>3</sup> carbon probe atom with +1 charge and vdW radius of 1.52 Å, and energy cut-off of 30 kcal/mol. The five similarity indices in CoMSIA, that is, steric, electrostatic, hydrophobic, H-bond donor and H-bond acceptor descriptors, were calculated<sup>37</sup> using a C<sup>1+</sup> probe atom with a radius of 1.0 Å placed at regular grid spacing of 2 Å. A Gaussian type distance dependence was used between the grid point *q* and each atom *i* of the molecule. The default value of 0.3 was used as the attenuation factor (*R*). Column filtering was set to 2.0 kcal/mol. CoMSIA steric indices are related to the third power of the atomic radii, electrostatic descriptors were derived from atomic partial charges, hydrophobic fields were derived from atom-based parameters,<sup>38</sup> and H-bond donor and acceptor indices were obtained by a rule-based method based on experimental results.<sup>39</sup> The CoMFA and CoMSIA descriptors were used as independent variables, and pIC<sub>50</sub> (–log IC<sub>50</sub>) values were used as the dependent variable in PLS regression analyses to derive 3D-QSAR models. The predictive value of the models was evaluated by leave-one-out (LOO) cross-validation. The cross validated coefficient, *q*<sup>2</sup>, was calculated using Eq 1:

$$q^2 = 1 - \frac{\sum (Y_{\text{predicted}} - Y_{\text{actual}})^2}{\sum (Y_{\text{observed}} - Y_{\text{mean}})^2} \quad (1)$$

where *Y*<sub>predicted</sub>, *Y*<sub>actual</sub>, and *Y*<sub>mean</sub> are predicted, actual and mean values of the target property (pIC<sub>50</sub>), respectively.  $\sum (Y_{\text{predicted}} - Y_{\text{actual}})^2$  is the predictive sum of squares (PRESS). Conventional correlation coefficient *r*<sup>2</sup> and its standard error, *s*, were also computed for the final PLS models. CoMFA and CoMSIA coefficient maps were generated by interpolation of the pair wise products between the PLS coefficients and the standard deviations of the corresponding CoMFA or CoMSIA descriptor values.

#### 5.10. Biological assays

##### 5.10.1. Materials, chemicals, and enzymes

All compounds were dissolved in DMSO and the stock solutions were stored at –20 °C. γ[<sup>32</sup>P]-ATP was purchased from either Amersham Biosciences or ICN. The expression systems for wild-type IN were a generous gift from Dr. Robert Craigie, Laboratory

of Molecular Biology, NIDDK, NIH, Bethesda, MD. Cell lines were obtained from American Type Culture Collection (ATCC, Rockville, MD).

### 5.10.2. Preparation of oligonucleotide substrates

The oligonucleotides 21top, 5'-GTGTGGAAATCTCTAGCAGT-3', and 21-bot, 5'-ACTGCTAGAGATTTCCACAC-3', were purchased from the Norris Cancer Center Microsequencing Core Facility (University of Southern California) and purified by UV shadowing on polyacrylamide gel. To analyze the extent of 3'-end processing and strand transfer using 5'-end labeled substrates, 21top was 5'-end labeled using T<sub>4</sub> polynucleotide kinase (Epicentre, Madison, WI) and  $\gamma$ [<sup>32</sup>P]-ATP (Amersham Biosciences or ICN). The kinase was heat-inactivated and 21-bot was added in 1.5 M excess. The mixture was heated at 95 °C, allowed to cool slowly to room temperature, and run through a G25 mini spin column (USA Scientific) to separate double-stranded oligonucleotide from unincorporated material.

### 5.10.3. Integrase assays

To determine the extent of 3'-processing (3'-P) and strand transfer (ST), wild-type IN was preincubated at a final concentration of 200 nM with the inhibitor in reaction buffer (50 mM NaCl, 1 mM HEPES, pH 7.5, 50  $\mu$ M EDTA, 50  $\mu$ M dithiothreitol, 10% glycerol (w/v), 7.5 mM MnCl<sub>2</sub>, 0.1 mg/mL bovine serum albumin, 10 mM 2-mercaptoethanol, 10% dimethylsulfoxide, and 25 mM MOPS, pH 7.2) at 30 °C for 30 min. An aliquot (5  $\mu$ L) was electrophoresed on a denaturing 20% polyacrylamide gel (0.09 M Tris-borate, pH 8.3, 2 mM EDTA, 20% acrylamide, and 8 M urea). Gels were dried, exposed in a PhosphorImager cassette, and analyzed using a Typhoon 8610 Variable Mode Imager (Amersham Biosciences) and quantitated using ImageQuant 5.2. Percent inhibition (%I) was calculated using the following equation:

$$\%I = 100 * [1 - (D - C)/(N - C)]$$

where C, N, and D are the fractions of 21-mer substrate converted to 19-mer (3'-P product) or strand transfer products for DNA alone, DNA plus IN, and IN plus compound, respectively. The IC<sub>50</sub> values were determined by plotting the logarithm of compound concentration versus percent inhibition to obtain the concentration that produced 50% inhibition.

### 5.11. Anti-HIV-1 activity assay

The testing of the ability of potent compounds to inhibit HIV replication in cell culture was done according to a previously reported procedure.<sup>40</sup> PBMC (107 cells/T25flask) were stimulated with phytohemagglutinin for 3 days and infected with a wild-type HIV-1 strain (strain LAI) at 100 50% tissue culture infective doses, as described previously.<sup>40</sup> The cultures were kept for 5 days in the presence of test compounds at serial 1-log dilutions. Subsequently, human PBMC were removed from the culture supernatant by centrifugation (400 g, 10 min, 4 °C). This clarified supernatant was tested by a reverse transcriptase assay.

### 5.12. Cytotoxicity assays

The cytotoxicity of compounds was evaluated using uninfected PBMC and CEM leukemia cells according to a previous method.<sup>41</sup> PBMC were obtained from whole blood of healthy individuals, while CEM were obtained from the ATCC (Rockville, MD). The PBMC and CEM cells were cultured in the presence or absence of compound for 6 days. After this time period, cells were stained with Trypan blue dye, and counted for cell proliferation and viability according to the previously reported procedure.<sup>42</sup>

### Acknowledgments

Financial support from the National Institute of Allergy and Infectious Diseases (NIAID), NIH Grant No. AI084710 and the Department of Pharmaceutical Sciences is gratefully acknowledged. RFS is also supported by NIH CFAR grant 2P30-AI-050409 and the Department of Veterans Affairs.

### References and notes

1. *RetroViruses*; Coffin, J. C., Hughes, S. H., Vermus, H. E., Eds.; Cold Spring Harbor Laboratory Press: Plainview, NY, 1999.
2. Sweeney, Z. K.; Klumpp, K. *Curr. Opin. Drug Discov. Dev.* **2008**, *11*, 458.
3. Von Hentig, N. *Drugs Today* **2008**, *44*, 103.
4. MacArthur, R. D.; Novak, R. M. *Clin. Infect. Dis.* **2008**, *47*, 236.
5. Manfredi, R.; Sabbatani, S. *Curr. Med. Chem.* **2006**, *13*, 2369.
6. De Clercq, E. *J. Med. Chem.* **1995**, *38*, 2491.
7. Zhang, L.; Ramratnam, B.; Tenner-Racz, K.; He, Y.; Vesanen, M.; Lewin, S.; Talal, A.; Racz, P.; Perelson, A. S.; Korber, B. T.; Markowitz, M.; Ho, D. D. *N. Eng. J. Med.* **1999**, *340*, 1605.
8. Richman, D. D. *Nature* **2001**, *410*, 995.
9. Esposito, D.; Craigie, R. *Adv. Virus. Res.* **1999**, *52*, 319.
10. Katz, R. A.; Skalka, A. M. *Annu. Rev. Biochem.* **1994**, *63*, 133.
11. La Femina, R. L.; Schnider, C. L.; Robbins, H. L.; Callahan, P. L.; Legro, K.; Roth, E.; Schleif, W. A.; Emini, E. A. *J. Virol.* **1992**, *66*, 7414.
12. Sakai, H.; Kawamura, M.; Sakuragi, S.; Shibata, R.; Ishimoto, A.; Ono, N.; Ueda, S.; Adachi, A. *J. Virol.* **1993**, *67*, 1169.
13. Farnet, C. M.; Wang, B.; Lipford, J. R.; Bushman, F. D. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, *93*, 9742.
14. Asante-Appiah, E.; Skalka, A. M. *Adv. Virus Res.* **1999**, *52*, 351.
15. Brown, P. O. *Integration*; Cold Spring Harbor Press: Cold Spring Harbor, NY, 1999.
16. Billich, A. *Curr. Opin. Invest. Drugs* **2003**, *4*, 206.
17. Hazuda, D. J.; Anthony, N. J.; Gomez, R. P.; Jolly, S. M.; Wai, J. S.; Zhuang, L.; Fisher, T. E.; Embrey, M.; Guare, J. P., Jr.; Egbertson, M. S.; Vacca, J. P.; Huff, J. R.; Felock, P. J.; Witmer, M. V.; Stillmock, K. A.; Danovich, R.; Grobler, J.; Miller, M. D.; Espeseth, A. S.; Jin, L.; Chen, I. W.; Lin, J. H.; Kassahun, K.; Ellis, J. D.; Wong, B. K.; Xu, W.; Pearson, P. G.; Schleif, W. A.; Cortese, R.; Emini, E.; Summa, V.; Holloway, M. K.; Young, S. D. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 11233.
18. Sato, M.; Kawakami, H.; Motomura, T.; Aramaki, H.; Matsuda, T.; Yamashita, M.; Ito, Y.; Matsuzaki, Y.; Yamataka, K.; Ikeda, S.; Shinkai, H. *J. Med. Chem.* **2009**, *52*, 4869.
19. Sorbera, L. A.; Serradell, N. *Drugs Future* **2006**, *31*, 310.
20. Boros, E. E.; Edwards, C. E.; Foster, S. A.; Fuji, M.; Fujiwara, T.; Garvey, E. P.; Golden, P. L.; Hazen, R. J.; Jeffrey, J. L.; Johns, B. A.; Kawasuji, T.; Kiyama, R.; Koble, C. S.; Kurose, N.; Miller, W. H.; Mote, A. L.; Murai, H.; Sato, A.; Thompson, J. B.; Woodward, M. C.; Yoshinaga, T. *J. Med. Chem.* **2009**, *52*, 2754.
21. Dubois, M.; Bailly, F.; Mbemba, G.; Mouscadet, J. F.; Debyser, Z.; Witvrouw, M.; Cotellet, P. *J. Med. Chem.* **2008**, *51*, 2575.
22. Pace, P.; Francesco, M. E. D.; Gardelli, C.; Harper, S.; Muraglia, E.; Nizi, E.; Orvieto, F.; Petrocchi, A.; Poma, M.; Rowley, M.; Scarpelli, R.; Laufer, R.; Gonzalez, P.; Monteagudo, E.; Bonelli, F.; Hazuda, D. J.; Stillmock, K. A.; Summa, V. *J. Med. Chem.* **2007**, *50*, 2225.
23. Pasquini, S.; Mugnaini, C.; Tintori, C.; Botta, M.; Trejos, A.; Arvela, R. K.; Larhed, M.; Witvrouw, M.; Michiels, M.; Christ, F.; Debyser, Z.; Corelli, F. *J. Med. Chem.* **2008**, *51*, 5125.
24. Plewe, M. B.; Butler, S. L.; Dress, K. R.; Hu, Q.; Johnson, T. W.; Kuehler, J. E.; Kuki, A.; Lam, H.; Liu, W.; Nowlin, D.; Peng, Q.; Rahavendran, S. V.; Tanis, S. P.; Tran, K. T.; Wang, H.; Yang, A.; Zhang, J. *J. Med. Chem.* **2009**, *52*, 7211.
25. Ramkumar, K.; Serrao, E.; Odde, S.; Neamati, N. *Med. Res. Rev.* **2010**, *30*, 890.
26. Summa, V.; Petrocchi, A.; Bonelli, F.; Crescenzi, B.; Donghi, M.; Ferrara, M.; Fiore, F.; Gardelli, C.; Pazz, G. O.; Hazuda, D. J.; Jones, P.; Kinzel, O.; Laufer, R.; Monteagudo, E.; Muraglia, E.; Nizi, E.; Orvieto, F.; Pace, P.; Pescatore, G.; Scarpelli, R.; Stillmock, K. A.; Witmer, M. V.; Rowley, M. *J. Med. Chem.* **2008**, *51*, 5843.
27. Al-Mawasawi, L. Q.; Al-Safi, R. I.; Neamati, N. *Expert Opin. Emerg. Drugs* **2008**, *13*, 213.
28. Patil, S.; Kamath, S.; Sanchez, T.; Neamati, N.; Schinazi, R. F.; Buolamwini, J. K. *Bioorg. Med. Chem.* **2007**, *15*, 1212.
29. Zwaagstra, M. E.; Timmerman, H.; Abdoelgafoer, R. S.; Zhang, M. Q. *Eur. J. Med. Chem.* **1996**, *31*, 861.
30. Zwaagstra, M. E.; Timmerman, H.; Tamura, M.; Tohma, T.; Wada, Y.; Onogi, K.; Zhang, M. Q. *J. Med. Chem.* **1997**, *40*, 1075.
31. Pannecouque, C.; Pluymsers, W.; Van Maele, B.; Tetz, V.; Cherepanov, P.; De Clercq, E.; Witvrouw, M.; Debyser, Z. *Curr. Biol.* **2002**, *12*, 1169.
32. Dixon, S.; Smondyrev, A.; Knoll, E.; Rao, S.; Shaw, D.; Friesner, R. *J. Comput. Aided Mol. Des.* **2006**, *20*, 647.
33. Kuo, C. L.; Assefa, H.; Kamath, S.; Brzozowski, Z.; Slawinski, J.; Saczewski, F.; Buolamwini, J. K.; Neamati, N. *J. Med. Chem.* **2004**, *47*, 385.
34. Clark, M.; Cramer, I. R. D. *Quant. Struct. Act. Relat.* **1993**, *12*, 137.

35. Zhu, Z.; Buolamwini, J. K. *Bioorg. Med. Chem.* **2008**, *16*, 3848.
36. Telvekar, V.; Kundaikar, H.; Patel, K.; Chaudhari, H. *QSAR Comb. Sci.* **2008**, *27*, 1193.
37. Klebe, G.; Abraham, U.; Mietzner, T. *J. Med. Chem.* **1994**, *37*, 4130.
38. Viswanadhan, V. N.; Ghose, A. K.; Revenkar, G. R.; Robins, R. J. *Chem. Inf. Comput. Sci.* **1989**, *29*, 163.
39. Klebe, G. *J. Mol. Biol.* **1994**, *237*, 212.
40. Stuyver, L. J.; Lostia, S.; Adams, M.; Mathew, J. S.; Pai, B. S.; Grier, J.; Tharnish, P. M.; Choi, Y.; Chong, Y.; Choo, H.; Chu, C. K.; Otto, M. J.; Schinazi, R. F. *Antimicrob. Agents Chemother.* **2002**, *46*, 3854.
41. Schinazi, R. F.; Sommadossi, J. P.; Saalman, V.; Cannon, D. L.; Xie, M. Y.; Hart, G. C.; Smith, G. A.; Hahn, E. F. *Antimicrob. Agents Chemother.* **1990**, *34*, 1061.
42. Sommadossi, J. P.; Carlisle, R.; Schinazi, R. F.; Zhou, Z. *Antimicrob. Agents Chemother.* **1988**, *32*, 997.