

Improved synthesis of trifluoromethyl sulfones used as intermediates for the preparation of di- or tri-substituted olefins

Fabrice Eugene, Bernard Langlois* and Eliane Laurent

Université Claude Bernard-Lyon I, Lab. de Chimie Organique 3, associé au CNRS, 43 Bd du 11 Novembre 1918, F-69622 Villeurbanne (France)

(Received January 18, 1993; accepted March 25, 1993)

Abstract

Primary and secondary trifluoromethyl sulfones (triflones) are efficiently obtained from easily available sodium trifluoromethanesulfinate (triflinate) and alkyl bromides in *N,N*-dimethylacetamide. This technique is more powerful than the potassium triflinate/acetonitrile system. Ethyl aconitate can be also produced in one step from ethyl bromoacetate and diisopropylethylamine, sodium triflinate being a catalyst.

Introduction

Sulfones are very popular tools in organic synthesis [1] because they increase to a large extent the acidity of hydrogens in α -positions since their conjugated bases are stabilised by a strong inductive effect [1c, 2]. These anions can be further halogenated, nitrated, alkylated or enter Michael, Claisen or Knoevenagel reactions. The sulfonyl moiety is then removed by sodium or aluminium amalgam [1c, 3], in most cases, or, in a few instances, by β -eliminations under strong basic conditions [4].

Replacement of alkyl or aryl sulfonyl moieties by the trifluoromethanesulfonyl ('triflyl') moiety constituted a major progress in this type of chemistry and attractive syntheses from trifluoromethyl sulfones ('triflones') [1c, 1f, 5, 6] and perfluoroalkyl sulfones [7] have been already been reported.

The advantage of the triflyl group lies in its strong electron-withdrawing effect, higher than those of mesyl, benzenesulfonyl and even nitro substituents [8]. Thus, hydrogens in positions α to CF_3SO_2 are very acidic, all the more so since the conjugated anions are also stabilised by conjugative effects [9]. Furthermore, the triflyl moiety can be removed by milder reducers than other sulfonyl groups (for instance with zinc/ethanol or Raney nickel [6]) or, more often, through β -elimination or $\text{S}_{\text{N}}2$ processes [6], since this substituent, although a poorer leaving group than halogens, is a far better leaving group than aryl- or alkyl-sulfonyl [10]. A thermal 1,2-elimination of triflinic acid ($\text{CF}_3\text{SO}_2\text{H}$)

can also be achieved [6], whereas this process is not possible from other sulfones [11].

Triflones can be produced by oxidation of trifluoromethyl thioethers [12], Friedel–Crafts condensations with trifluoromethanesulfonyl chloride [13] and reaction of organometallics upon trifluoromethanesulfonic anhydride [13, 14]. Substitution of alkyl bromides by potassium trifluoromethanesulfinate (potassium 'triflinate') in acetonitrile seems to be the most general method [5, 6] but, because of the low nucleophilicity of the triflinate anion, only primary bromides and not secondary ones are converted into triflones through clean but very slow reactions.


Results and discussion

Recently, a cheap and efficient manufacture of sodium triflinate has been reported from bromotrifluoromethane [15]. Hence, it seemed interesting to examine the reactivity of this easily available reagent in media other than acetonitrile. Of the usual dipolar aprotic solvents known to enhance the reactivity of anions, some suffer from severe drawbacks: hexamethylphosphoric triamide (HMPT) is carcinogenic; the high-boiling sulfolane is very tedious to recover; *N,N*-dimethylformamide (DMF) can formylate anionised triflones [5]; and dimethylsulfoxide (DMSO) is not quite stable upon prolonged heating. *N,N*-Dimethylacetamide (DMAc) is a good candidate, better than acetonitrile since it solvates cations better ($DN_{\text{DMAc}} > DN_{\text{CH}_3\text{CN}}$) and anions lesser ($AN_{\text{DMAc}} < AN_{\text{CH}_3\text{CN}}$) [16] (DN = donor number, AN = acceptor number).

*Author to whom correspondence should be addressed.

Formation of triflones in DMAc

Sodium triflinate has been reacted with several primary and secondary carbonylated alkyl bromides in DMAc. A comparison with acetonitrile as solvent has been undertaken for one of the substrates:



[R = CH₂CO₂Et (**1a**), CH(CH₃)CO₂Et (**1b**), (CH₂)₂-CO₂Et (**1c**), CH₂COPh (**1d**), CH₂-CO-Bu^t (**1e**), CH(CH₃)COPh (**1f**)]


The results obtained are summarised in Table 1.

From these results, it can be seen that replacement of acetonitrile by DMAc (entries 1–3) dramatically enhanced the reactivity of the triflinate anion since comparable results were obtained from **1a** in DMAc under milder conditions and in a shorter time than in acetonitrile.

The results from Table 1 also show that, under these conditions, activated substrates such as α-bromoketones, even secondary ones (**1f**, entry 8) which are not reactive in acetonitrile [6], formed triflones within a few hours in excellent yield [it should be noted that **1d** needed 48 h at 80 °C to produce **2d** in acetonitrile [6] in contrast to 7 h at 50 °C in DMAc (entry 6)]. Secondary α-bromoesters (e.g. **1b**), which are less reactive in nucleophilic substitution, also provided triflones in medium yield within 0.5 d only at temperatures as low as 65 °C. The advantage was also noticeable with simple primary bromides with no carbonyl group in the α-position (**1c**): they were completely converted in c. 2

d at 65 °C in DMAc instead of 7 d at 80 °C in acetonitrile [6].


In contrast, the formation of benzyl triflone from benzyl bromide and alkaline triflinates did not appear to be sensitive to solvent and cation effects: the same yield (91–95%) was obtained, at 80–90 °C for 6 h with 0.1 equiv. potassium iodide as catalyst, from potassium triflinate in acetonitrile [6] or sodium triflinate in DMAc. Such an observation indicates that benzyl triflone probably does not result from an S_N2 process but from an S_N1 substitution which is known to be less affected by the nature of the solvent and cation in aprotic media [17]. This presumption is reinforced by the fact that, contrary to other bromides, benzyl bromide reacted easily with sodium triflinate in a solid–liquid system using toluene as an apolar aprotic solvent and tris(3,6-dioxahexyl)amine N[(CH₂CH₂O)₂CH₃]₃ (TDA-1) as the phase-transfer catalyst:



Formation of olefins substituted by carboxy groups from ethyl bromoacetate in the presence of sodium triflinate

During the synthesis of ethyl (trifluoromethanesulfonyl)acetate (**2a**) in DMAc, methyl triflone (**3**) and ethyl fumarate (**4**) were formed as by-products. Their amounts increased with temperature and reaction time (Table 2). This phenomenon did not occur in acetonitrile.

TABLE 1. Synthesis of triflones from bromides in DMAc



Entry No.	R	x	Solvent	θ (°C)	Time (h)	Conv. 1 ^a (%)	Yield of 2 ^b (%)	
							NMR analysis	Isolated
1	EtO ₂ C-CH ₂	(1a) 1	CH ₃ CN	80	40	74	76	68
2	EtO ₂ C-CH ₂	(1a) 1	DMAc	60	2.25	66	61	
3	EtO ₂ C-CH ₂	(1a) 3	DMAc	60	3.25	76	62	
4	CH ₃ O ₂ C-CH(CH ₃)	(1b) 1.5	DMAc	65	13.5	62	58	52
5	EtO ₂ C-(CH ₂) ₂	(1c) 1.2 + 0.4 ^c	DMAc	65	52	92 ^d	65	57
6	Ph-CO-CH ₂	(1d) 1	DMAc	50	7	89	87	82
7	Bu ^t -CO-CH ₂	(1e) 1	DMAc	60	1.5	54	100	78
8	Ph-CO-CH(CH ₃)	(1f) 2	DMAc	70	9	76 ^d	96	91

^aFrom the NMR spectra of the crude product.

^bYields versus % converted **1**.

^c1.2 equiv. at t = 0 and 0.4 equiv. at t = 30 h.


^dFrom isolated products.

TABLE 2. Influence of the parameters in the reaction of ethyl bromoacetate and sodium triflate


$$\text{BrCH}_2\text{CO}_2\text{Et} \xrightarrow[\text{DMAc, } \theta \text{ } ^\circ\text{C}]{\text{CF}_3\text{SO}_2\text{Na (x equiv.)}} \text{CF}_3\text{SO}_2\text{CH}_2\text{CO}_2\text{Et} + \text{CF}_3\text{SO}_2\text{CH}_3 + \text{EtO}_2\text{C}-\text{CH}=\text{CH}-\text{CO}_2\text{Et}$$

(1a) (2a) (3) (4) (E-isomer)









Entry No.	x (equiv.)	θ ($^\circ\text{C}$)	Time (h)	Conv. 1a (%) ^a	Product yield (%)		
					2a ^{a, b}	3 ^{a-c, e}	4 ^{a, b, d}
1	1	60	2.25	66	61	14	0
2	3	60	1.25	60	50	6	0
3	3	60	3.25	76	62	18	0
4	1 ^f	100	9	94	7	31	39

^aFrom NMR analysis of the crude product.^bYields versus % converted 1a.^cAccording to the stoichiometry: 1 mol 1a → 1 mol 3.^dAccording to the stoichiometry: 2 mol 1a → 1 mol 4.^eA part of 3 probably lost during work-up.^fConv. CF₃SO₂Na = 100%.

As DMAc dissolves and activates bromide anions well, it could be suspected that 3 and 4 resulted from 2a and Br⁻. Thus, pure 2a was treated with tetrabutylammonium bromide in DMAc: 2a was almost completely consumed, 3 and 4 appeared as major products and ethyl β-(trifluoromethanesulfonyl)propionate (5) as the minor one.



This reaction has been rationalised as indicated in Scheme 1.




Scheme 1.

As bromide is known to remove alkoxycarbonyl groups from esters [18], the first step is assumed to be a decarboxylation which delivers the stabilised anion 3⁻, the conjugated base of 3. The pK_a value of 3 has been reported as c. 18 [9]. As Cameo simulation led us to estimate the pK_a value of 2a to be c. 6–8, we propose that an acid/base reaction occurs between 2a and 3⁻ and delivers 2a⁻, the conjugated base of 2a, which is sufficiently nucleophilic to substitute the triflyl group of 2a and leads to the secondary triflone 6. Triflone 6 can eliminate trifluoromethanesulfinic acid ('triflinic acid') to deliver ethyl fumarate (4). The triflinic acid then decomposes thermally into fluorocarbon and sulfur dioxide. This assumption is in accordance with the observation that sodium triflate was completely converted when ethyl fumarate became an important product (Table 2, entry 4). Ethyl β-(trifluoromethanesulfonyl)propionate (5) can also be formed, in a side process, through decarboxylation of 6.

As the above-described formation of ethyl fumarate could be a good model for the preparation of olefins

bearing electron-withdrawing substituents from the corresponding alkyl bromides and triflate, this reaction has been investigated more thoroughly.


In the proposed Scheme 1, 1 mol alkyl bromide and 1 mol triflate are consumed to produce 1 mol of the non-valuable methyl triflate (3). In addition, another mol of triflate is destroyed through reaction (11). In order to circumvent these drawbacks, ethyl bromoacetate was reacted with sodium triflate in DMAc in the presence of a strong non-nucleophilic base. Diisopropylethylamine (DIEA, $pK_a \approx 11$ [19]) was chosen for this purpose:



The results obtained are reported in Table 3.

In fact, when ethyl bromoacetate (1a) and sodium triflate were reacted in DMAc at 100 °C in the presence of diisopropylethylamine, no methyl triflate

TABLE 3. Synthesis of ethyl aconitate from ethyl bromoacetate diisopropylethylamine and sodium triflate








Entry No.	θ (°C)	Time (h)	x (equiv.)	Conv. Tf ⁻ (%) ^a	Product yield (%)			
					4 ^b	7 ^b	8 ^b	9 ^b
1	100	9	2	39	19	77	<1	<1
2	100	14.5	3	48	0	47	6	24
3	80	14.5	3	28	9	67	<1	<1

^aTf⁻ = CF₃SO₂⁻.

^bFrom the ¹H NMR spectra.

(3) was obtained. Ethyl fumarate (4) was formed in only small amounts and ethyl aconitate (7) was the major product. This latter compound 7 became the only significant one at a lower temperature (80 °C) (Table 3, entry 3). Such a quasi-selective reaction, in which the triflate anion acts as a catalyst (conv. CF₃SO₂Na = 28%) could be a good model for the preparation of trisubstituted olefins from α -bromocarbonylated substrates.

In the presence of DIEA, the mechanism proposed in Scheme 1 must be modified as indicated in Scheme 2.




(Tf = CF₃SO₂; E = CO₂Et)

Scheme 2.


In the first step, an acid-base reaction between 2a and DIEA matches the decarboxylation of 2a so that 2a⁻, the conjugated base of 2a, is obtained without the occurrence of 3. As 2a⁻ is formed as soon as 2a is produced, 2a⁻ can co-exist with ethyl bromoacetate (1a) which is more sensitive to nucleophilic displacement than 2a [10]. Thus, 6 is readily formed and deprotonated by DIEA to 6⁻, the conjugated base of 6. Then, 6⁻ can substitute ethyl bromoacetate to deliver the tertiary triflate 10 which, in the presence of DIEA, leads to ethyl aconitate and the triflate anion (which is thus recovered).

In order to reinforce the proposed hypotheses and to detect intermediates, the reaction has been performed under milder conditions:



The occurrence of **6** and **7** has thus been confirmed.

The two first entries of Table 3 show that, under high temperature and long reaction time, ethyl succinate (**8**) and ethyl tricarallylate (**9**) arose as by-products alongside **4** and **7**. Their amounts also increased when potassium iodide was omitted as the catalyst:



Thus, the system $\text{CF}_3\text{SO}_2\text{Na}/\text{DIEA}/\text{KI}$ in DMAc at 80 °C seems to provide the best conditions for obtaining ethyl aconitate directly from ethyl bromoacetate.

Experimental

All NMR analyses were undertaken with deuteriochloroform as solvent. ^1H NMR spectra were recorded at 60 MHz on a Varian EM 360 spectrometer; chemical shifts (δ) are given in ppm with TMS as internal reference. ^{19}F NMR spectra were recorded either at 56.4 MHz on a Varian EM 360 spectrometer or at 75.2 MHz on a Bruker WP 80 one or at 188.2 MHz on a Bruker AC 200 apparatus; chemical shifts are given in ppm with CFCl_3 as reference (δ positive upfield). ^{13}C NMR spectra were recorded at 15.1 MHz with TMS as internal reference. The following abbreviations are used: s (singlet), d (doublet), t (triplet), q (quadruplet), m (multiplet). Coupling constants (J) are given in Hz. Quantitative NMR analyses of the organic phases were obtained with benzotrifluoride as internal standard. Unreacted sodium triflate was estimated, after work-up, by ^{19}F NMR ($\delta = -87.8$ ppm) spectroscopic analysis of the aqueous phase with sodium trifluoroacetate as internal standard.

Reactions were monitored by gas-phase chromatography on a Varian 3300 apparatus fitted with a thermal conductivity detector and a 15-m length semicapillary column (internal diameter, 0.25 mm); the stationary phase was either DB1 or DBwax. Helium was the carrier gas.

In some cases, IR spectra, coupled with GPC, were recorded on a Bruker IFS 85 apparatus. IR frequencies are given in cm^{-1} . Mass spectrometry has been performed on a VG 305 spectrometer.

Flash chromatographic separations were performed on Merck 60H silica. The products were eluted with petroleum ether (abbr.: PE), pure or mixed with diethyl ether (abbr.: E) or methylene chloride.

Pure commercially available acetonitrile (SDS-Chromasol) was stored over 3 Å molecular sieves. *N,N*-Dimethylacetamide (DMAc) (Aldrich-GC) was distilled at atmospheric pressure prior to use and dried over 4 Å molecular sieves for 24 h. Sodium trifluoromethanesulfinate was generously provided by Rhône-Poulenc Co. and other commercially available substrates (Aldrich or Janssen) were used as received.

Reaction of sodium trifluoromethanesulfinate with ethyl bromoacetate

In a 50 ml flask, fitted with a reflux condenser, a thermometer and a magnetic stirrer, were placed consecutively, under nitrogen, the required quantities of sodium triflate, solvent and ethyl bromoacetate. The stirred reaction mixture was then heated under nitrogen at the desired temperature and maintained at this temperature for the requisite time. After reaction, 30 ml of water were added. The resulting mixture was extracted with 3×20 ml of diethyl ether. The ethereal phase was then washed twice with 15 ml of water, dried over magnesium sulfate, filtered and concentrated at room temperature under reduced pressure. The crude organic residue was either analysed by ^1H and ^{19}F NMR spectroscopy with benzotrifluoride as the internal standard or purified by flash chromatography. The gathered aqueous phases were analysed by ^{19}F NMR spectroscopy using sodium trifluoroacetate as the internal standard. Details for each experiment are given in Table 4.

From 5.93 g of the crude product resulting for the experiment reported in Table 4 (entry 1) were obtained, by flash chromatography, 0.68 g $\text{BrCH}_2\text{CO}_2\text{Et}$ (**1a**) (PE/E=94:6) and 3.00 g $\text{CF}_3\text{SO}_2\text{CH}_2\text{CO}_2\text{Et}$ (**2a**) (PE/E=90:10). Spectroscopic data for **2a**: ^1H NMR δ : 1.30 (t, 3H, CH_3 , $^3J=8$ Hz); 4.30 (q, 2H, OCH_2 , $^3J=8$ Hz); 4.35 (s, 2H, CH_2CO) (lit. value [6]: 4.35) ppm. ^{19}F NMR (56.4 MHz) δ : -77.7 (s) ppm. IR (cm^{-1}): 2991; 1769; 1400; 1277; 1223; 1130; 1030; 717; 621.

From 1.04 g of the crude product resulting from the experiment reported in Table 4 (entry 5) were obtained, by flash chromatography, 0.050 g $\text{BrCH}_2\text{CO}_2\text{Et}$ (**1a**) (PE/E=93:7) and 0.210 g ethyl fumarate (**4**) (PE/

TABLE 4. Experimental data for the reaction of $\text{CF}_3\text{SO}_2\text{Na}$ and ethyl bromoacetate (**1a**)

Entry No.	$\text{CF}_3\text{SO}_2\text{Na}$ [g (mmol)]	1a [g (mmol)]	Solvent (ml)	θ ($^\circ\text{C}$)	Time (h)	Crude product (g)	NMR analysis [g (%)] ^a		
1	4.22 (27.1)	4.52 (27.1)	MeCN (30)	80	40	5.93	1a	1.20	(26)
							2a	3.33	(76)
2	5.54 (35.5)	1.94 (11.6)	DMAc (15)	60	1.25	2.00	1a	0.77	(40)
							2a	0.77	(50)
							3	0.06	(6)
3	1.84 (11.8)	1.99 (11.9)	DMAc (15)	60	2.25	2.07	1a	0.68	(34)
							2a	1.03	(61)
							3	0.16	(14)
4	5.50 (35.2)	1.94 (11.6)	DMAc (15)	60	3.25	2.10	1a	0.46	(24)
							2a	1.20	(62)
							3	0.24	(18)
5	2.12 (13.6)	2.26 (13.5)	DMAc (20)	100	9	2.01	1a	0.14	(6)
							2a	0.19	(7)
							3	0.58	(31)
							4	0.43	(39)

^aVersus introduced **1a** for remaining **1a**; versus converted **1a** for other compounds.

E=93:7). Spectroscopic data for **4**: ^1H NMR δ : in accordance with the literature [20]. IR (cm^{-1}): 2988; 2947; 1744; 1647; 1472; 1394; 1321; 1298; 1259; 1155; 1098; 1043; 902; 862; 662.

Because of its rather high volatility, methyl trifluoromethyl sulfone (**3**) ($E_{b760}=128\text{ }^\circ\text{C}$ [21]) could not be isolated in this way. However, its occurrence has been demonstrated in the above crude product via the following spectroscopic characteristics. ^1H NMR δ : 3.11 (s) (lit. value [5m, 21]: 3.11) ppm. ^{19}F NMR (56.4 MHz) δ : -81 (s) (lit. value [21]: 3.6 (versus $\text{CF}_3\text{CO}_2\text{H}$)) ppm. IR (cm^{-1}): 1389; 1329; 1232; 1202; 1136; 957; 771; 735.

Synthesis of triflones **2b–f**

The procedure was the same as above, except that the bromo compounds **1b–f** were used as substrates instead of ethyl bromoacetate. *N,N*-Dimethylacetamide only was used as solvent. The experimental data are given in Table 5.

Methyl 2-(trifluoromethanesulfonyl)propionate (**2b**)

From the total crude product resulting from **1b** (Table 5, entry 1), flash chromatography delivered 1.76 g $\text{BrCH}(\text{CH}_3)\text{CO}_2\text{CH}_3$ (**1b**) (PE/E=93:7) and 2.00 g $\text{CF}_3\text{SO}_2\text{CH}(\text{CH}_3)\text{CO}_2\text{CH}_3$ (**2b**) (PE/E=80:20). NMR characteristics for **2b**: ^1H NMR δ : 1.77 (d, 3H, $\text{CH}-\text{CH}_3$, $^3J=7\text{ Hz}$); 3.87 (s, 3H, CO_2CH_3); 4.30 (q, 1H, $\text{CH}-\text{CH}_3$, $^3J=7\text{ Hz}$) ppm. ^{19}F NMR (188.2 MHz) δ : -75.12 (s) ppm.

Ethyl 3-(trifluoromethanesulfonyl)propionate (**2c**)

From 4.00 g of the crude product resulting from **1c** (Table 5, entry 2), flash chromatography delivered 0.33 g $\text{BrCH}_2\text{CH}_2\text{CO}_2\text{Et}$ (**1c**) (PE/E=94:6) and 2.80 g $\text{CF}_3\text{SO}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$ (**2c**) (PE/E=90:10).

As the conversion rate of **1c** could not be determined from the ^1H NMR spectrum of the crude product, it has been estimated from the amount of **1c** recovered by chromatography: conv. **1c**=92%; yield **2c**=65% (crude), 57% (isolated) both versus converted **1c**. NMR characteristics for **2c**: ^1H NMR δ : 1.27 (t, 3H, $\text{CO}_2\text{CH}_2\text{CH}_3$, $^3J=7\text{ Hz}$); 2.87 (t, 2H, $\text{CH}_2\text{CO}_2\text{Et}$, $^2J=8\text{ Hz}$); 3.57 (t, 2H, $\text{CF}_3\text{SO}_2\text{CH}_2$, $^3J=8\text{ Hz}$); 4.17 (q, 2H, $\text{CO}_2\text{CH}_2\text{CH}_3$, $^3J=7\text{ Hz}$) ppm. ^{19}F NMR (56.4 MHz) δ : -79.0 (s) ppm.

α -(Trifluoromethanesulfonyl)acetophenone (**2d**)

From the total crude product resulting from **1d** (Table 5, entry 3), flash chromatography offered 0.30 g $\text{C}_6\text{H}_5-\text{CO}-\text{CH}_2\text{Br}$ (**1d**) (PE/ CH_2Cl_2 =80:20) and 3.72 g $\text{C}_6\text{H}_5-\text{CO}-\text{CH}_2\text{SO}_2\text{CF}_3$ (**2d**) (PE/ CH_2Cl_2 =65:35). NMR characteristics for **2d**: ^1H NMR δ : 4.9 (s, 2H, CH_2) (lit. value [6]: 5.0); 7.3–7.9 (m, 5H, C_6H_5) ppm.

1-Trifluoromethanesulfonyl-3,3-dimethyl-2-butanone (**2e**)

From the total crude product resulting from **1e** (Table 5, entry 4), flash chromatography delivered 0.35 g $\text{Bu}^t-\text{CO}-\text{CH}_2-\text{Br}$ (**1e**) (PE/ CH_2Cl_2 =80:20) and 0.62 g $\text{Bu}^t-\text{CO}-\text{CH}_2\text{SO}_2\text{CF}_3$ (**2e**) (PE/ CH_2Cl_2 =70:30). NMR characteristics for **2e**: ^1H NMR δ : 1.23 (s, 9H,

TABLE 5. Experimental data for the synthesis of triflones **2b–f**

Entry No.	CF ₃ SO ₂ Na [g (mmol)]		RBr (1) [g (mmol)]		DMAC (ml)	θ (°C)	Time (h)	Crude product (g)	NMR analysis [g (%)] ^a		
1	6.72	(43.1)	1b	4.69 (28.1)	70	65	13.5	5.41	CF ₃ SO ₂ Na	2.91 (43)	
									1b	1.76 (38)	
									2b	2.26 (58)	
2	4.23 + 1.50	(27.2) (9.6) ^b	1c	4.11 (22.7)	38	65	52	4.45	CF ₃ SO ₂ Na	0	
									1c ^c		
									2c	3.19 (60) ^d	
3	3.80	(24.3)	1d	4.02 (20.2)	35	50	7	5.36	CF ₃ SO ₂ Na	0	
									1d	0.44 (11)	
									2d	3.94 (87)	
4	1.09	(7.0)	1e	1.13 (6.3)	10	60	1.5	1.58	CF ₃ SO ₂ Na	0	
									1e	0.53 (47)	
									2e	0.79 (100)	
5	6.47	(41.5)	1f	4.39 (20.6)	60	70	9	4.85	CF ₃ SO ₂ Na	0	
									1f ^c		
									2f	4.0 (73) ^d	

^aVersus introduced reagents for CF₃SO₂Na and **1b–f**; versus converted **1b–f** for other compounds.^b1.50 g CF₃SO₂Na added after 30 h.^c¹H NMR spectra of **1** and **2** too close to estimate **1** in the crude product.^dVersus introduced **1**.

C(CH₃)₃; 4.5 (s, 2H, COCH₂) ppm. ¹⁹F NMR (75.2 MHz) δ : -78.2 (s) ppm.

α -(Trifluoromethanesulfonyl)propiophenone (**2f**)

From the total crude product resulting from **1f** (Table 5, entry 5), flash chromatography delivered 1.05 g BrCH(CH₃)COC₆H₅ (**1f**) (PE/CH₂Cl₂=80:20) and 3.8 g CF₃SO₂CH(CH₃)COC₆H₅ (**2f**) (PE/CH₂Cl₂=60:40).

As the conversion of **1f** could not be estimated from the ¹H NMR spectrum of the crude product, it has been calculated from the quantity of **1f** recovered by chromatography: conv. **1f** = 76%; yield **2f** = 96% (crude), 91% (isolated) both versus converted **1f**. NMR characteristics for **2f**: ¹H NMR δ : 1.73 (d, 3H, CH₃, ³J = 7 Hz); 5.19 (q, 1H, CH–CH₃, ³J = 7 Hz); 7.40–8.15 (m, 5H, C₆H₅) ppm. ¹⁹F NMR (56.4 MHz) δ : -75.7 (s) ppm.

Synthesis of benzyltriflone (**2g**)

In a homogeneous phase (DMAC)

In a 100 ml flask, fitted with a reflux condenser, a thermometer and a mechanical stirrer, were placed, consecutively, under nitrogen, 7.49 g (48 mmol) of sodium triflinate, 0.63 g (3.8 mmol) of potassium iodide, 40 ml of DMAC and, finally, 6.84 g (40 mmol) of benzyl bromide (**1g**). The stirred reaction mixture was heated, under nitrogen, at 90 °C in an oil bath and maintained at this temperature for 6 h. After cooling, GPC analysis indicated a 91% yield of **2g**. The reaction mixture was then filtered and concentrated under reduced pressure.

The solid obtained was recrystallized in a mixture of water and methanol (1:3 v/v), washed with an aqueous solution of sodium thiosulfate and dried under vacuum. Pure **2g** (6.43 g) was thus obtained (72% yield).

Under solid–liquid phase-transfer conditions

In the same apparatus as above were introduced consecutively 4.68 g (30 mmol) of sodium triflinate, 0.46 g (2.8 mmol) of potassium iodide, 25 ml of toluene, 1.58 g (4.9 mmol) of tris-(3,6-dioxahexyl)amine and 4.28 g (25 mmol) of **1g**. The mixture was vigorously stirred and, under nitrogen, maintained at 80 °C for 10 h. After reaction and filtration, GPC and ¹⁹F NMR analyses indicated a 72% yield of **2g**. The same work-up as above afforded 3.00 g of pure **2g** (54% yield).

Benzyl triflone (**2g**): M.p. 104 °C (H₂O–MeOH). ¹H NMR δ : 4.41 (s, 2H, CH₂) (lit. value [6]: 4.49); 7.4 (m, 5H, C₆H₅) ppm. ¹³C NMR δ : 56.2 (s, CCH₂); 119.7 (q, CCF₃, ²J(C–F) = 328 Hz); 123.2 (s, C–CH₂); 129.2 (s, C meta); 130.0 (s, C para); 131.2 (s, C ortho) ppm. IR (cm⁻¹): 3000; 2960; 1500; 1460; 1415; 1400; 1360; 1350; 1325; 1290; 1225; 1200; 1190; 1120; 1075; 1030; 780; 720; 700; 640; 560; 525; 505. MS *m/z*: 224 (M⁺).

Reaction of **2a** with tetrabutylammonium bromide

In the same apparatus as described above for its preparation, 0.51 g (2.3 mmol) of **2a** were mixed with a solution of 0.78 g (2.4 mmol) of tetrabutylammonium bromide in 5 ml of DMAC. This medium was kept at 100 °C, under nitrogen, for 1.5 h. After the usual work-

up, 0.35 g of a crude mixture was obtained and analysed by ^1H and ^{19}F NMR spectroscopy, with benzo-trifluoride as internal standard. By comparison with the spectra of isolated compounds, **2a** (45 mg), **3** (125 mg), **5** (19 mg) and ethyl fumarate (**4**) (42 mg) were found.

Reaction of sodium triflinate with ethyl bromoacetate in the presence of diisopropylethylamine (DIEA)

The procedure was the same as that already described for reactions between **1a** and $\text{CF}_3\text{SO}_2\text{Na}$, except that DIEA was added just after the other components, prior to heating, and the reaction medium was hydrolysed with 30 ml of 10% aqueous hydrochloric acid instead of pure water. The experimental data are given in Table 6.

Entry 1: From the whole crude product, compounds **6**, **7** and **10** were separated by flash chromatography, in the following order of elution: 0 g $\text{BrCH}_2\text{CO}_2\text{Et}$ (**1a**); 0.30 g (14%) $\text{CF}_3\text{SO}_2\text{CH}(\text{CO}_2\text{Et})\text{CH}_2\text{CO}_2\text{Et}$ (**6**) (PE/E = 96:4); 0.13 g (7%) $\text{CF}_3\text{SO}_2\text{CH}(\text{CO}_2\text{Et})(\text{CH}_2\text{CO}_2\text{Et})_2$ (**10**) (PE/E = 96:4); and 0.61 g (48%) $(E)\text{-EtO}_2\text{CCH}_2\text{C}(\text{CO}_2\text{Et})=\text{CHCO}_2\text{Et}$ (**7**) (PE/E = 96:4).

Diethyl 2-(trifluoromethanesulfonyl)succinate (**6**): ^1H NMR δ : 1.35 (t, 6H, OCH_2CH_3 , $^3J=7$ Hz); 3.25 [dd, 2H, $\text{CH}_a\text{--H}_b\text{--CH}_c$, $^3J(\text{H}_a\text{--H}_c)=6$ Hz, $^3J(\text{H}_b\text{--H}_c)=9$ Hz]; 4.20 (q, 4H, OCH_2CH_3 , $^3J=7$ Hz); 4.57 [dd, 1H, $\text{CH}_c\text{--CH}_a\text{H}_b$, $^3J(\text{H}_c\text{--H}_a)=6$ Hz, $^3J(\text{H}_c\text{--H}_b)=9$ Hz] ppm. ^{19}F NMR (56.4 MHz) δ : -76.0 (s) ppm.

Ethyl aconitate (**7**): ^1H NMR δ : in accordance with the literature [20]. IR (cm^{-1}): 2988; 2949; 1755; 1738; 1653; 1472; 1418; 1371; 1321; 1273; 1171; 1097; 1040; 968.

Diethyl 3-(ethoxycarbonyl)-3-(trifluoromethanesulfonyl)pentane-1,5-dioate (**10**): ^1H NMR δ : 1.35 (t, 9H, OCH_2CH_3 , $^3J=7$ Hz); 3.70 (s, 4H, CH_2CO); 4.00 (q, 6H, OCH_2CH_3 , $^2J=7$ Hz) ppm. ^{19}F NMR (56.4 MHz) δ : -71.3 (s) ppm.

Entry 2: The crude product was analysed by ^1H and ^{19}F NMR spectroscopy as well as by IR spectroscopy coupled with GPC. The results were as follows: 0 g ethyl bromoacetate (**1a**); 0.09 g (8%) ethyl fumarate (**4**); 0.48 g (40%) ethyl aconitate (**7**); 0.36 g (30%)

ethyl succinate (**8**); and 0.23 g (19%) ethyl tricarballylate (**9**).

Entry 3: NMR analysis and flash chromatography of the whole crude product indicated complete conversion of ethyl bromoacetate and afforded **4** and **7** in the following yields: ethyl fumarate (**4**): crude 0.21 g (19%), isolated 0.11 g (10%) (PE/E = 90:10); and ethyl aconitate (**7**): crude 0.84 g (77%), isolated 0.65 g (60%) (PE/E = 80:20).

Entry 4: The same procedure indicated complete conversion of **1a** and provided the following results: ethyl aconitate (**7**): crude 0.73 g (47%), isolated 0.73 g (47%) (PE/E = 80:20); ethyl succinate (**8**): crude 0.15 g (10%), isolated 0.10 g (6%) (PE/E = 80:20); and ethyl tricarballylate (**9**): crude 0.37 g (24%), isolated 0.37 g (24%) (PE/E = 80:20). The ^1H NMR spectra of ethyl succinate (**8**) and tricarballylate **9** were in accordance with the literature [20].

Entry 5: NMR analysis of the crude product indicated again complete conversion of **1a** and delivered the following figures: 0.15 g (9%) ethyl fumarate (**4**) and 1.11 g (67%) ethyl aconitate (**7**).

Synthesis of triflone 6 and triflone 10 from ethyl (trifluoromethanesulfonyl)acetate (2a)

In order to confirm the structures of triflones **6** and **10**, isolated from **1a**, sodium triflinate and DIEA (Table 6, entry 1), these compounds have been prepared via another route from **2a** and ethyl bromoacetate (**1a**). Thus, 0.161 g (6.7 mmol) of sodium hydride, washed with 3×5 ml of petroleum ether, were added to 6 ml of *N,N*-dimethylformamide (DMF). Compound **2a** (1.06 g, 4.8 mmol) was dropped, under stirring, onto this suspension at -10°C within 10 min. The reaction mixture was stirred at room temperature for 14 h then 0.60 g (3.6 mmol) of ethyl bromoacetate, dissolved in 5 ml of DMF, was added. Stirring was continued at room temperatures for 24 h. After addition of 30 ml of water, extraction with 3×30 ml of ether and decantation, the organic phase was washed twice with 20 ml of water, dried over MgSO_4 , filtered and concentrated under reduced pressure. The resulting material (1.04 g) was separated by flash chromatography to yield: 0.62

TABLE 6. Experimental data for the reaction of $\text{CF}_3\text{SO}_2\text{Na}$ with ethyl bromoacetate (**1a**) in the presence of DIEA

Entry No.	$\text{CF}_3\text{SO}_2\text{Na}$ [g (mmol)]	1a [g (mmol)]	KI [g (mmol)]	DIEA [g (mmol)]	DMAC (ml)	θ ($^\circ\text{C}$)	Time (h)	Crude product (g)	Unconverted $\text{CF}_3\text{SO}_2\text{Na}$ [g (%)] (^{19}F NMR)
1	2.38 (15.3)	2.60 (15.6)		2.20 (17.1)	25	60	4	1.07	0.71 (30)
2	1.08 (6.9)	2.33 (14.0)		2.00 (15.5)	10	100	9	1.30	0.40 (35)
3	0.98 (6.3)	2.11 (12.6)	0.20 (1.2)	1.93 (15.0)	15	100	9	1.41	0.60 (61)
4	0.92 (5.9)	3.00 (18.0)	0.20 (1.2)	2.44 (18.9)	15	100	14.5	1.77	0.48 (52)
5	1.00 (6.4)	3.24 (19.4)	0.25 (1.5)	2.74 (21.2)	15	80	14.5	1.66	0.72 (72)

g triflone **6** (PE/E=95:5); 0.13 g triflone **10** (PE/E=95:5); 0.13 g ethyl bromoacetate (**1a**) (PE/E=94:6); and 0.07 g triflone **2a** (PE/E=94:6).

The ^{19}F and ^1H NMR spectra of triflones **6** and **10** prepared in this way were identical with those of the triflones obtained from ethyl bromoacetate, sodium triflinate and DIEA (Table 6, entry 1).

The new compounds described in this paper are as follows: methyl 2-(trifluoromethanesulfonyl)propionate (**2b**), ethyl 3-(trifluoromethanesulfonyl)propionate (**2c**), 1-(trifluoromethanesulfonyl)-3,3-dimethyl-2-butanone (**2e**), α -(trifluoromethanesulfonyl)propiophenone (**2f**), diethyl 2-(trifluoromethanesulfonyl)succinate (**6**) and diethyl 3-(ethoxycarbonyl)-3(trifluoromethanesulfonyl)-pentane-1,5-dioate (**10**).

Conclusions

Though primary triflones can be obtained cleanly in acetonitrile from potassium triflinate and alkyl bromides, this reaction is very slow and not suitable for the preparation of secondary triflones. The use of the readily available sodium triflinate in a more basic aprotic solvent such as *N,N*-dimethylacetamide allows the preparation of primary as well as secondary triflones under mild conditions. Such an improvement widens the synthetic usefulness of triflones.

The unique properties of the trifluoromethanesulfonyl group also allows the one-pot synthesis of ethyl aconitate from ethyl bromoacetate with sodium triflinate as a catalyst, provided that a stoichiometric amount of a non-nucleophilic tertiary amine like diisopropylethylamine is used.

Acknowledgements

The authors (especially F.E.) thank Rhône-Poulenc Chimie Co. for its financial support and generous grant of sodium triflinate, and Dr Laurent Gilbert for providing them with the results of the Cameo simulations.

References

- 1 For a review, see (a) J. March, *Advanced Organic Chemistry*, 3rd edn., John Wiley, New York, 1985; (b) T. Durst, in D. Barton and W.D. Ollis (eds.), *Comprehensive Organic Chemistry*, Pergamon, Oxford, 1979, Vol. 3, p. 171; (c) R.P. Magnus, *Tetrahedron*, **33** (1977) 2019; (d) L. Field, *Synthesis*, (1978) 713; (e) P. Caubère, *Utilisation des Dérivés du Soufre en Chimie Organique*, Masson, Paris, 1984, p. 50; (f) S. Patai, Z. Rappoport

- and C.M.J. Sterling, *The Chemistry of Sulphones and Sulphoxides*, John Wiley, Chichester, 1988; (g) K. Schank, in *Houben-Weyl's Methoden der Organischen Chemie*, Thieme, Stuttgart, 1985, Vol. E/11, 1129 ff; (h) N.S. Simpkins, *Sulphones in Organic Synthesis*, Pergamon Press, Oxford, 1992.
- 2 Ref. 1a, p. 154, and references therein; S. Wolfe, A. Rank and I.G. Csizmadia, *J. Am. Chem. Soc.*, **91** (1969) 1567.
- 3 Ref. 1a, pp. 413, 440 and 818, and references therein; M. Julia, M. Launey, J.P. Stacino and J.N. Verpeaux, *Tetrahedron Lett.*, **23** (1982) 2465.
- 4 J.L. Fabre, M. Julia and J.N. Verpeaux, *Tetrahedron Lett.*, **23** (1982) 2469; J. Bremner, M. Julia, M. Launay and J.P. Stacino, *Tetrahedron Lett.*, **23** (1982) 3265; M. Sellen, J.E. Backvall and P. Helquist, *J. Org. Chem.*, **56** (1991) 835.
- 5 (a) J.B. Hendrickson, K.W. Bair, R. Bergeron, A. Giga, P.L. Skipper, D.D. Sternbach and J.A. Wareing, *Org. Prep. Proced. Int.*, **9** (1977) 173; (b) J.B. Hendrickson, D.D. Sternbach and K.W. Bair, *Acc. Chem. Res.*, **10** (1977) 306; (c) H.R. Snyder, H.V. Anderson and D.P. Hallada, *J. Am. Chem. Soc.*, **73** (1951) 3258; (d) C.S. Rondestvedt Jr. and J.C. Wygant, *ibid.*, **73** (1951) 5785; (e) R.S. Glass and D.L. Smith, *J. Org. Chem.*, **39** (1974) 3712; (f) M. Hanack and F.M. Massa, *Tetrahedron Lett.*, (1977) 661; (g) M. Hanack and K. Laping, *Tetrahedron Lett.*, (1977) 4493; (h) *ibid.*, (1979) 1309; (i) Z. Wrogl and M. Makosza, *Org. Prep. Proced. Int.*, **22** (1990) 575; (j) J.B. Hendrickson, G.J. Boudreaux and P.S. Palumbo, *Tetrahedron Lett.*, **25** (1984) 4617; (k) J.B. Hendrickson and P.S. Palumbo, *ibid.*, **26** (1985) 2849; (l) J.B. Hendrickson and P.S. Palumbo, *J. Org. Chem.*, **50** (1985) 2110; (m) J.B. Hendrickson and P.L. Skipper, *Tetrahedron*, **32** (1976) 1627.
- 6 J.B. Hendrickson, A. Giga and J. Wareing, *J. Am. Chem. Soc.*, **96** (1974) 2275.
- 7 R. Sodoyer, E. Abad, E. Rouvier and A. Cambon, *J. Fluorine Chem.*, **22** (1983) 401.
- 8 W.A. Sheppard, *J. Am. Chem. Soc.*, **85** (1963) 1314.
- 9 F.G. Bordwell, N.R. Vanier, W.S. Matthews, J.B. Hendrickson and P.L. Skipper, *J. Am. Chem. Soc.*, **97** (1975) 7160.
- 10 X. Creary, *J. Org. Chem.*, **50** (1985) 5080.
- 11 Ref. 1a, p. 913.
- 12 T. Nguyen, M. Rubinstein and C. Wakselman, *J. Fluorine Chem.*, **21** (1982) 437; J.F. Harris Jr., *J. Org. Chem.*, **32** (1967) 2063.
- 13 J.B. Baudin, M. Julia, C. Rolando and J.N. Verpeaux, *Tetrahedron Lett.*, **25** (1984) 3203.
- 14 M. Hanack and F.W. Massa, *Tetrahedron Lett.*, **22** (1981) 557; R.J. Koshar and R.A. Mitsch, *J. Org. Chem.*, **38** (1973) 3358; L.M. Yagupolskii and L.M. Yagupolskaya, *Proc. Acad. Sci. USSR (Eng. Trans.)*, **134** (1960) 1207.
- 15 M. Tordeux, B. Langlois and C. Wakselman, *J. Org. Chem.*, **54** (1989) 2452.
- 16 C. Reichardt, *Solvents and Solvent Effects in Organic Chemistry*, 2nd revised edn., VCH, Weinheim, 1990, pp. 17–25.
- 17 Ref. 16, p. 209.
- 18 S. Takei and Y. Kawando, *Tetrahedron Lett.*, (1975) 4389; R.V. Stevens and A.W.M. Lee, *J. Am. Chem. Soc.*, **101** (1979) 7032.
- 19 Estimated value from: J.W. Smith, in S. Patai (ed.), *The Chemistry of Amino Groups*, John Wiley, New York, 1968, Chap. IV, p. 170.
- 20 C.J. Pouchert, *The Aldrich Library of NMR Spectra*, 2nd edn., Aldrich Chemical Co., Milwaukee, WI, 1983, p. 544A.
- 21 Q.Y. Chen, G.Y. Yang and S.W. Wu, *J. Fluorine Chem.*, **55** (1991) 291.