ChemComm

COMMUNICATION

Check for updates

Cite this: Chem. Commun., 2021, 57, 8268

Received 1st June 2021, Accepted 20th July 2021

DOI: 10.1039/d1cc02888j

rsc.li/chemcomm

Rhodium(III)-catalyzed asymmetric [4+1] spiroannulations of O-pivaloyl oximes with α -diazo compounds[†]

Lincong Sun,^a Bingxian Liu, ^b Yanlian Zhao,^b Junbiao Chang,^a Lingheng Kong, ^c Fen Wang,^c Wei-Qiao Deng ^b and Xingwei Li ^{*}

Chiral Rh^{III} catalysts can catalyze the asymmetric [4+1] spiroannulation of O-pivaloyl oximes with α -diazo homophthalimides under redox-neutral and acid/base-neutral conditions, leading to formation of chiral spirocyclic imines as a result of C-H activation and N-O cleavage. The reaction proceeded with high efficiency and features broad substrate scope, mild reaction conditions, and high to excellent enantioselectivities.

Spirocycles are commonly encountered structural motifs that have found wild applications in synthetic chemistry. Their unique 3D structures may impart diversified biological, chemical, and photochemical properties in various natural and synthetic compounds. Thus, they have also been used as key intermediates during synthesis and development of drug molecules.¹ Chiral spirocycles have also attracted much attention as prevalent scaffolds of chiral ligands.² A number of synthetic strategies have been documented to access spirocycles.³ They can be prepared through alkylation, metal-catalyzed reactions, and cycloaddition reactions. Despite these synthetic routes, many of them suffer from employment of highly functionalized reagents and lengthy procedures with generation of waste. Therefore, atom- and step-economic catalytic synthesis of spirocycles with a quaternary carbon center in a regioand stereocontrolled fashion remains an ongoing theme.

In the past decade, metal-catalyzed and chelation-assisted C-H bond functionalization has been developed as an increasingly important strategy for the construction of complex

(hetero)cyclic molecules, including spirocyclic scaffolds.⁴ Of note, rhodium(III) cyclopentadienyl complexes have been established as one of most efficient catalysts for synthesis of spirocyclic compounds via a C-H activation pathway.⁵ By following this strategy, arenes bearing a heteroatom directing group have been successfully employed for the synthesis of various spirocyclic skeletons via coupling with unsaturated reagents such as alkynes, alkenes, and α -diazocarbonyl compounds.⁶ Meanwhile, the discovery of chiral rhodium catalysts has opened a new avenue to access chiral polycyclic scaffolds.7-11 In 2015, You and coworkers developed the first enantioselective dearomatization reaction of 2-naphthols for the synthesis of spirocyclic enones in excellent enantioselectivities (Scheme 1a).6b By using enol as a coordinating group, the Lam group realized Rh(m)catalyzed enantioselective synthesis of spirocyclic indenes (Scheme 1a).¹² In 2019, the Waldmann group disclosed a related enantioselective [3+2] spiroannulation via formal C(sp³)-H bond activation (Scheme 1a).^{6c} Very recently, we realized oxidative [4+1] spiroannulation of nitrone and guinone diazides, which occurred via initial C-H arylation followed by axial-to-central chirality transfer (Scheme 1b).¹³ Thus, these spirocyclic products were mostly constructed under oxidative conditions. On the other hand, in 2014 Cramer reported the [4+1] annulation of N-OPiv benzamides and diazo compounds for enantioselective synthesis of lactams.¹⁴ Later, the arene substrate was extended to indoles by Song and coworkers.^{11a} Our objective was to construct aza-spirocycles on different platforms via C-H activation assisted by other readily available oxidizing directing groups. As integration of eco-friendly synthesis of spirocycles and asymmetric C-H bond activation, we now report Rh(m)-catalyzed enantioselective [4+1] spirocyclization between O-pivaloyl oximes and α -diazo homophthalimides (Scheme 1c).

We initiated our studies using *O*-pivaloyl oxime **1a** as a model substrate, and its [4+1] spiroannulation with α -diazo homophthalimides was explored using the Cramer-type Cp^{*X*}Rh(m) catalysts (Table 1).^{7c,d} The desired product (*R*)-**3** was obtained with generally good enantioselectivity and efficiency when (*R*)-**Rh2** was used in various solvents (entries 1–5), and



View Article Online

^a Henan Key Laboratory of Organic Functional Molecule and Drug Innovation, Key Laboratory of Green Chemical Media and Reactions, Ministry of Education, School of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang 453007, China. E-mail: lixw@snnu.edu.cn

^b Institute of Molecular Science and Engineering, Institute of Frontier and Interdisciplinary Sciences, Shandong University, Qingdao, China

^c School of Chemistry and Chemical Engineering, Shaanxi Normal University, Xi'an 710062, China

[†] Electronic supplementary information (ESI) available: Experimental procedures and characterization data of all compounds. CCDC 2069779 (44), 2073889 (52) and 2073891 (53). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1cc02888j

(a) Oxidative [3+2] Spiroannulation



DCE seemed to outperform other solvents. The reaction temperature and additives were also screened (entries 5–8 and 9–14), and the enantioselectivity only slightly varied. To our delight, both high enantioselectivity and yield were maintained when the catalyst loading was lowered to 4 mol% even in the absence of any acid/base additive (entry 15). In contrast, replacing the catalyst with (*R*)-**Rh1** and (*R*)-**Rh3** gave lower enantioselectivities (see the ESI[†]).

Table 1	Optimization	studies ^a
Table 1	Optimization	studies

$ \begin{array}{c} \begin{array}{c} \begin{array}{c} & & & \\ & & $							
Entry	Additive	Solvent	T (°C)	$\operatorname{Yield}^{b}(\%)$	ee ^c (%)		
1	Li ₂ CO ₃	PhCl	40	72	89		
2	Li ₂ CO ₃	TFE	40	65	75		
3	Li_2CO_3	Et_2O	40	81	87		
4	Li_2CO_3	EtOAc	40	80	80		
5	Li_2CO_3	DCE	40	83	90		
6 ^c	Li_2CO_3	DCE	25	83	90		
7 ^c	Li_2CO_3	DCE	10	84	91		
8 ^c	Li_2CO_3	DCE	0	75	92		
9 ^c	LiOAc	DCE	10	72	88		
10^{c}	K_2CO_3	DCE	10	50	90		
11 ^c	Cs_2CO_3	DCE	10	NR	—		
12^c	HOAc	DCE	10	80	84		
13 ^c	PivOH	DCE	10	81	89		
14^{c}	—	DCE	10	83	91		
15^{cd}	_	DCE	10	85	91		

^{*a*} Reaction Conditions: **1a** (0.06 mmol), **2a** (0.05 mmol), (*R*)-**Rh2** (5 mol%), AgSbF₆ (20 mol%), additive (1.0 equiv.), solvent (1 mL), under Ar for 24 h. ^{*b*} Isolated yield. ^{*c*} Under Ar for 48 h. ^{*d*} (*R*)-**Rh2** (4 mol%) and AgSbF₆ (16 mol%) were used.

With the establishment of the optimal reaction conditions, we next investigated the scope of the O-pivaloyl oximes (Scheme 2). It was observed that introduction of various electron-donating, -withdrawing, and halogen groups at the para position of the phenyl ring gave consistently good yields and excellent enantioselectivities (3-13, 87-91% ee). Of note, a 1.0 mmol scale synthesis of product 3 also proved to be successful, with no deterioration of the vield or enantioselectivity. We also found that oxime esters bearing a *meta* methyl, methoxy, and halogen group also reacted efficiently (14-17) in comparably excellent enantioselectivity (70-90% yields, 90-94% ee). In contrast, ortho-fluorine substituted O-pivaloyl oxime (1s) underwent the C-H annulation with low regioselectivity and slightly lower enantioselectivity (18 and 18'). The presence of methyl and halogen group at the *ortho* position was also tolerated (19-22). No deterioration of the enantioselectivity was observed with different alkyl substitutions (23-26). Our studies further revealed that symmetrical benzophenonederived O-pivaloyl oximes were tolerated giving moderate to good yields and excellent enantioselectivities (27-31, 94-97%) ee). In addition, cyclic O-pivaloyl oximes also reacted smoothly with 2a, affording products 32 and 33 with 88% and 92% ee, respectively. When 2-naphthyl-substituted O-pivaloyl oxime was employed, the corresponding spirocyclic product 34 was also obtained (97% yield, 89% ee). Furthermore, heteroatomic



Scheme 2 Scope of O-pivaloyl oximes in spiro compounds synthesis. Reaction conditions: 1 (0.12 mmol), 2 (0.10 mmol), (R)-Rh2 (4 mol%), and AgSbF₆ (16 mol%) in DCE (2 mL) under Ar for 48 h, isolated yield.

substrate such as thienvl ring, afforded the corresponding products with lower enantioselectivity (35 and 36).

Next, we investigated the scope of the diazo compounds. As shown in Scheme 3, introduction of various electron-donating, -withdrawing, and halogen groups to the N-phenyl ring allowed the coupling to gave the desired products in good to excellent yields and high enantioselectivities (37-46, 86-96% ee), which revealed that the reaction efficiency was slightly affected by the electronic effect. Meanwhile, the absolute configuration of the product 44 was assigned as (R) by X-ray crystallography (CCDC⁺ 2069779). Moreover, variation of the N-substituent to -Bn (47) and 2-Naph (49) group all led to high yields (82% and 85%, respectively) and enantioselectivities (90% ee and 91% ee, respectively). However, lower enantioselectivity (79% ee) was obtained for the N^{-n} Bu (48) diazo compound, indicative of the electronic effect. Examination of the 6-substituent in the diazo compound revealed compatibility of bromo group (50, 91% ee).

To demonstrate the synthetic utility of the reaction system, the synthetic transformations of 3 was conducted (Scheme 4). Treatment of 3 with *m*-CPBA afforded the oxidation product N-oxide 51 in 64% yield without loss of the enantiopurity. Pd/ C-catalyzed reduction of 3, surprisingly, afforded a ringopening and ring-expansion product 52 (CCDC[†] 2073889) in 88% yield (4:1 dr), which may occur via C=N reduction and C-N reductive cleavage. Treatment of 3 with NaBH₄ in EtOH led to ring-opening of the imide group with incorporation of the solvent, affording hemiaminal 53 in 55% yield, and the structure of 53 (CCDC⁺ 2073891) was determined by X-ray crystallography.

To shed light on the mechanism of this spirocyclization reaction, we carried out several experimental studies (Scheme 5). First, H/D exchange of 1a was carried out in the presence of D₂O, the reactant was recovered in 92% yield and the ¹H NMR analysis revealed slight deuteration at the *ortho* positions of the phenyl ring, suggesting reversibility of the C-H cleavage. Second, another H/D exchange in the presence of 2a was conducted under the standard conditions, affording compound 3- D_{y} (the D_{y} is a statistical average value of compound $3-D_{\nu}$), and the deuterium incorporation was observed at



Scheme 3 Scope of diazo compounds in spiro compounds synthesis (see Scheme 2 for reaction conditions).



Scheme 4 Synthetic transformations of compounds 3



the methyl group probably due to acidity of the α-methyl group.¹⁵ Two parallel reactions of **1a** and **1a**-D₅ with **2a** were performed for kinetic isotope effect, and the $k_{\rm H}/k_{\rm D}$ = 2.2 was obtained at a low conversion under the standard conditions, suggesting that the ortho C-H activation event might be involved in the turnover-limiting step.

On the basis of the previous reports,^{11a,14} the enantioselectivity for the (R)-3 formation may be explained by the following stereochemical model (Scheme 6). Following the C-H bond activation and formation of the carbene species, both intermediates A and B can be possible. Regarding the subsequent migratory insertion, the intermediate B is preferred likely because of the reduced steric interaction between the OPiv group and the carbene ligand as we have previously proposed in the [4+1] annulation of quinone diazides,¹³ eventually leading



Scheme 6 Plausible stereochemical model of the chiral induction.

to the major product (*R*)-3. Indeed, our DFT studies revealed that the intermediate **B** is 5.0 kcal mol⁻¹ more stable, and this may also suggest a lower kinetic barrier during the migratory insertion of the intermediate **B**.

In summary, we have developed an efficient and mild protocol for the enantioselective synthesis of spirocyclic imines through a procedure of rhodium-catalyzed enantioselective C-H functionalization and N-O cleavage. This asymmetric [4+1] spiroannulation occurred under redox-neutral and acid/ base additive-free conditions *via* C-H bond activation of the oxime ester with cyclic α -diazo compounds as a coupling reagent. The reaction proceeded in high efficiency and features broad substrate scope, mild and redox-neutral reaction conditions, and high to excellent enantioselectivities. Further development of new reaction modes of rhodium-catalyzed enantioselective C-H activation reactions is underway in our laboratory.

We acknowledge the financial support for this work from NSFC (21525208 and 21801066) and the Shaanxi Normal University.

Conflicts of interest

The authors declare no competing financial interest.

Notes and references

- For reviews on the synthesis of selected spirocyclic natural products and drug discovery, see: (a) S. Kotha, A. Deb, K. Lahiri and E. Manivannan, Synthesis, 2009, 165; (b) V. A. D'yakonov, O. A. Trapeznikova, A. de Meijere and U. M. Dzhemilev, Chem. Rev., 2014, 114, 5775; (c) Y. Zheng, C. M. Tice and S. B. Singh, Bioorg. Med. Chem. Lett., 2014, 24, 3673. For individual reports, see; (d) T. Doi, Y. Iijima, M. Takasaki and T. Takahashi, J. Org. Chem., 2007, 72, 3667; (e) A. Bolognese, G. Correale, M. Manfra, A. Esposito, E. Novellino and A. Lavecchia, J. Med. Chem., 2008, 51, 8148; (f) S. A. Snyder, T. C. Sherwood and A. G. Ross, Angew. Chem., Int. Ed., 2010, 49, 5146; (g) T. Trieselmann, H. Wagner, K. Fuchs, D. Hamprecht, D. Berta, P. Cremonesi, R. Streicher, G. Luippold, A. Volz, M. Markert and H. Nar, J. Med. Chem., 2014, 57, 8766.
- 2 For selected reviews, see: (a) J.-H. Xie and Q.-L. Zhou, Acc. Chem. Res., 2008, 41, 581; (b) S.-F. Zhu and Q.-L. Zhou, Acc. Chem. Res., 2012, 45, 1365; (c) K. Ding, Z. Han and Z. Wang, Chem. - Asian J., 2009, 4, 32; (d) Q.-L. Zhou and J.-H. Xie, Top. Organomet. Chem., 2011, 36, 1. For individual reports, see: (e) V. B. Birman, A. L. Rheingold and K.-C. Lam, Tetrahedron: Asymmetry, 1999, 10, 125; (f) M. A. Arai, T. Arai and H. Sasai, Org. Lett., 1999, 1, 1795; (g) Z. Freixa, M. S. Beentjes, G. D. Batema, C. B. Dieleman, G. P. F. van Strijdonck, J. N. H. Reek, P. C. J. Kamer, J. Fraanje, K. Goubitz and P. W. N. M. van Leeuwen, Angew. Chem., Int. Ed., 2003, 42, 1284; (h) Z. Han, Z. Wang, X. Zhang and K. Ding, Angew. Chem., Int. Ed., 2009, 48, 5345; (i) J. Li, G. Chen, Z. Wang, R. Zhang, X. Zhang and K. Ding, Chem. Sci., 2011, 2, 1141; (j) X. Wang, Z. Han, Z. Wang and K. Ding, Angew. Chem., Int. Ed., 2012, 51, 936; (k) Z. Zheng, Y. Cao, Q. Chong, Z. Han, J. Ding, C. Luo, Z. Wang, D. Zhu, Q.-L. Zhou and K. Ding, J. Am. Chem. Soc., 2018, 140, 10374.

- 3 A. Ding, M. Meazza, H. Guo, J. W. Yang and R. Rios, *Chem. Soc. Rev.*, 2018, 47, 5946.
- 4 For selected reviews of transition-metal-catalyzed C-H functionalization, see: (a) L. McMurray, F. O'Hara and M. J. Gaunt, Chem. Soc. Rev., 2011, 40, 1885; (b) S. H. Cho, J. Y. Kim, J. Kwak and S. Chang, Chem. Soc. Rev., 2011, 40, 5068; (c) L. Ackermann, Chem. Rev., 2011, 111, 1315; (d) B.-J. Li and Z.-J. Shi, Chem. Soc. Rev., 2012, 41, 5588; (e) P. B. Arockiam, C. Bruneau and P. H. Dixneuf, Chem. Rev., 2012, 112, 5879; (f) Z. Shi, D. C. Koester, M. Boultadakis-Arapinis and F. Glorius, J. Am. Chem. Soc., 2013, 135, 12204; (g) G. Song and X. Li, Acc. Chem. Res., 2015, 48, 1007; (h) T. Gensch, M. N. Hopkinson, F. Glorius and J. Wencel-Delord, Chem. Rev., 2016, 45, 2900; (i) Y. Yang, J. Lan and J. You, Chem. Rev., 2017, 117, 8787; (j) J. R. Hummel, J. A. Boerth and J. A. Ellman, Chem. Rev., 2017, 117, 9163; (k) G. Cera and L. Ackermann, Top. Curr. Chem., 2016, 374, 191; (l) A. S. K. Hashmi, Acc. Chem. Res., 2014, 47, 864.
- ⁵ For individual reports, see: (a) A. Seoane, N. Casanova, N. Quiñones, J. L. Mascareñas and M. Gulías, J. Am. Chem. Soc., 2014, 136, 7607; (b) M.-B. Zhou, R. Pi, M. Hu, Y. Yang, R.-J. Song, Y. Xia and J.-H. Li, Angew. Chem., Int. Ed., 2014, 53, 11338; (c) D. J. Burns, D. Best, M. D. Wieczysty and H. W. Lam, Angew. Chem., Int. Ed., 2015, 54, 9958; (d) B. Liu, P. Hu, Y. Zhang, Y. Li, D. Bai and X. Li, Org. Lett., 2017, 19, 5402.
- 6 For selected examples of synthesis of spirocyclic compounds, see: (a) T. Nishimura, Y. Ebe and T. Hayashi, J. Am. Chem. Soc., 2013, 135, 2092; (b) J. Zheng, S.-B. Wang, C. Zheng and S.-L. You, J. Am. Chem. Soc., 2015, 137, 4880; (c) H. Li, R. Gontla, J. Flegel, C. Merten, S. Ziegier, A. P. Antonchick and H. Waldmann, Angew. Chem., Int. Ed., 2019, 58, 307; (d) S. Guo, L. Sun, Y. Liu, N. Ma, X. Zhang and X. Fan, Org. Lett., 2019, 21, 4082; (e) Z. Song, Z. Yang, P. Wang, Z. Shi, T. Li and X. Cui, Org. Lett., 2020, 22, 6272.
- 7 For selected reviews, see: (a) C. G. Newton, D. Kossler and N. Cramer, J. Am. Chem. Soc., 2016, 138, 3935. Selected examples:
 (b) B. Ye and N. Cramer, Science, 2012, 338, 504; (c) B. Ye and N. Cramer, J. Am. Chem. Soc., 2013, 135, 636; (d) B. Ye, P. A. Donets and N. Cramer, Angew. Chem., Int. Ed., 2014, 53, 507; (e) Y. Sun and N. Cramer, Angew. Chem., Int. Ed., 2017, 56, 364; (f) Y. Sun and N. Cramer, Angew. Chem., Int. Ed., 2018, 57, 15539; (g) M. Brauns and N. Cramer, Angew. Chem., Int. Ed., 2019, 58, 8902; (h) S.-G. Wang, Y. Liu and N. Cramer, Angew. Chem., Int. Ed., 2019, 58, 18136.
- 8 T. K. Hyster, L. Knörr, T. R. Ward and T. Rovis, *Science*, 2012, 338, 500.
- 9 (a) J. Zheng and S.-L. You, Angew. Chem., Int. Ed., 2014, 53, 13244;
 (b) J. Zheng, W.-J. Cui, C. Zheng and S.-L. You, J. Am. Chem. Soc., 2016, 138, 5242.
- 10 (a) F. Wang, Z. Qi, Y. Zhao, S. Zhai, G. Zheng, R. Mi, Z. Huang, X. Zhu, X. He and X. Li, *Angew. Chem., Int. Ed.*, 2020, **59**, 13288;
 (b) J. Sun, W. Yuan, R. Tian, P. Wang, X. Zhang and X. Li, *Angew. Chem., Int. Ed.*, 2020, **59**, 22706.
- (a) X. Chen, S. Yang, H. Li, B. Wang and G. Song, ACS Catal., 2017, 7, 2392; (b) T. Li, C. Zhou, X. Yan and J. Wang, Angew. Chem., Int. Ed., 2018, 57, 4048; (c) E. A. Trifonova, N. M. Ankudinov, A. A. Mikhaylov, D. A. Chusov, Y. V. Nelyubina and D. S. Perekalin, Angew. Chem., Int. Ed., 2018, 57, 7714; (d) S. Satake, T. Kurihara, K. Nishikawa, T. Mochizuki, M. Hatano, K. Ishihara, T. Yoshino and S. Matsunaga, Nat. Catal., 2018, 1, 585; (e) H. Li, X. Yan, J. Zhang, W. Guo, J. Jiang and J. Wang, Angew. Chem., Int. Ed., 2019, 58, 6732.
- 12 S. R. Chidipudi, D. J. Burns, I. Khan and H. W. Lam, *Angew. Chem., Int. Ed.*, 2015, **54**, 13975.
- 13 L. Kong, X. Han, S. Liu, Y. Zou, Y. Lan and X. Li, *Angew. Chem., Int. Ed.*, 2020, **59**, 7188.
- 14 B. Ye and N. Cramer, Angew. Chem., Int. Ed., 2014, 53, 7896.
- 15 D. Zhao, F. Lied and F. Glorius, Chem. Sci., 2014, 5, 2869.