

## Selective Piperidine Synthesis Exploiting Iodine-Catalyzed Csp<sup>3</sup>-H Amination under Visible Light

Hongwei Zhang, and Kilian Muniz

ACS Catal., Just Accepted Manuscript • Publication Date (Web): 10 May 2017

Downloaded from <http://pubs.acs.org> on May 10, 2017

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



# Selective Piperidine Synthesis Exploiting Iodine-Catalyzed C<sub>sp</sub><sup>3</sup>-H Amination under Visible Light

Hongwei Zhang<sup>1</sup> and Kilian Muñoz\*<sup>1,2</sup>

<sup>1</sup> Institute of Chemical Research of Catalonia (ICIQ), The Barcelona Institute of Science and Technology, 16 Avgda. Països Catalans, 43007 Tarragona, Spain.

<sup>2</sup> ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain


**ABSTRACT:** A route to selective piperidine formation through intramolecular catalytic C<sub>sp</sub><sup>3</sup>-H amination is described. This hydrocarbon amination reaction employs a homogeneous iodine catalyst derived from halogen coordination between molecular iodine and a terminal oxidant. It relies on visible light initiation and proceeds within two catalytic cycles that comprise a radical C-H functionalization and an iodine-catalyzed C-N bond formation. Under these conditions, the commonly observed preference for pyrrolidine synthesis based on halogenated nitrogen intermediates within the Hofmann-Löffler domain is effectively altered in favor of a free radical promoted piperidine formation. The protocol is demonstrated for a total of 30 applications.

**KEYWORDS:** amination, C-H functionalization, halogen bonding, iodine, light initiation.

Intramolecular amination of remote aliphatic C-H bonds is of particular conceptual interest as it streamlines existing protocols for the preparation of saturated N-heterocycles.<sup>1</sup> These compounds are usually accessible by classic radical chemistry,<sup>2</sup> in which modified Hofmann-Löffler reactions have demonstrated a unique potential.<sup>3</sup> We recently initiated exploration into iodine-catalyzed Hofmann-Löffler reactions<sup>4</sup> that provide the expected access to pyrrolidines from position-selective C-H functionalization based on intramolecular 1,5-H abstraction<sup>3,5</sup> through a nitrogen centered radical pathway (Figure 1, top).


In contrast, the related C-H amination strategy toward the piperidine core is significantly more challenging as the required 1,6-H abstraction from nitrogen-centered radicals is kinetically disfavored.<sup>6</sup> Consequently, a C-H amination strategy towards piperidines has remained elusive.<sup>7</sup> Piperidines represent important structural subunits in molecules of pharmaceutical, biological and medicinal interest, and exercise important pharmacophoric properties.<sup>8</sup> In fact, a recent analysis on the occurrence of nitrogen heterocycles in FDA approved pharmaceuticals identified the piperidine core as the most frequent member.<sup>9</sup> As a result, piperidine synthesis within intramolecular amination of remote C-H bonds would constitute an important synthetic advance. We here report on conditions for such a selective synthesis for the first time (Figure 2, bottom).

To override the given “innate” preference for pyrrolidine formation, we decided to pursue conditions that would preferentially generate free radicals outside the amidyl radical manifold involved in the Hofmann-Löffler pathway.<sup>3</sup> Within such a scenario, free radical hydrogen atom abstraction should address the weakest C-H bond and could be predicted by the introduction of carefully pre-organized substitution.<sup>10,11</sup>




**Figure 1.** Position-selective intramolecular C-H amination for pyrrolidine and piperidine synthesis.

We previously reported that iodinated reagents such as NIS **1b** effectively provide intermediates for exclusive Hofmann-Löffler reactions.<sup>12</sup> As a consequence of this observation and in order to prevent potential background reactions, we turned to less reactive bromine-based reagents. Catalytic amounts of molecular iodine<sup>13</sup> were pursued to generate low amounts of free radicals as the reaction carriers and thus to minimize potential side reactions. These halide reagent combinations provide the desired gateway to the elusive piperidine formation. Table 1 provides insight in the optimization of catalytic reaction conditions that allow for selective piperidine synthesis within C-H amination.

**Table 1.** Iodine-catalyzed piperidine formation: optimization.



entry	catalyst	oxidant (equiv)	time [h]	yield [%] <sup>a</sup>
1	I <sub>2</sub> (5 mol%)	<b>1a</b> (1.2)	4	47
2	I <sub>2</sub> (5 mol%)	<b>1a</b> (1.6)	4	62
3	I <sub>2</sub> (5 mol%)	<b>1a</b> (2.0)	2	80
4	I <sub>2</sub> (5 mol%)	<b>1b</b> (2.0)	12	34
5	I <sub>2</sub> (5 mol%)	<b>1c</b> (2.0)	12	n.d. <sup>b</sup>
6	I <sub>2</sub> (5 mol%)	<b>1d</b> (2.0)	12	72
7	-	<b>1a</b> (2.0)	12	0
8 <sup>c</sup>	I <sub>2</sub> (5 mol%)	<b>1a</b> (2.0)	12	0
9	KI (10 mol%)	<b>1a</b> (2.0)	12	50
10	TBAI (10 mol%)	<b>1a</b> (2.0)	12	30
11 <sup>d</sup>	I <sub>2</sub> (5 mol%)	<b>1a</b> (2.0)	12	73
12 <sup>e</sup>	I <sub>2</sub> (5 mol%)	<b>1a</b> (2.0)	12	67
13 <sup>f</sup>	I <sub>2</sub> (5 mol%)	<b>1a</b> (2.0)	12	0



<sup>a</sup>Isolated yield after purification. <sup>b</sup>n.d. = not determined (observation of trace amounts of **3a** in the crude NMR). <sup>c</sup>Reaction in the dark lab. <sup>d</sup>Reaction in CH<sub>2</sub>Cl<sub>2</sub>. <sup>e</sup>Reaction in MeCN. <sup>f</sup>Reaction in THF.

The reaction was developed with **2a** as substrate and departed from the observation that a combination of visible light exposure, 5 mol% molecular iodine and *N*-bromo succinimide (NBS) **1a** provided a selective transformation to the desired piperidine **3a** (entry 1). Subsequent rise of the amount of oxidant to 2 equivalents provided **3a** in 80% yield (entries 2,3) without detection of the corresponding pyrrolidine. This observation proves that the current conditions are capable of overriding conventional Hofmann-Löffler chemistry. Related iodo and chloro derivatives NIS **1b** and NCS **1c** provided significantly decreased reactivity (entries 4,5) accompanied by formation of the undesired pyrrolidine, while *N*-bromo phthalimide **1d** gave a comparable yield of 72% (entry 6). Control experiments verify that no formation of **3a** is obtained without the iodine catalyst or in the absence of light (entries 7,8). Alternative iodine catalyst sources and reactions in alternative solvents gave lower yields (entries 9-13).

Under the optimized conditions, the scope of the reaction was explored (Scheme 1). For tosylamide **2a**, the reaction was extended to a 1g-scale. Several additional sulfonimides **2b-e** including mesyl, nosyl, trimethylsilylethylsulfonylethyl (SES) and 2-bromophenyl sulfonyl also provide the corresponding piperidination products **3b-e**. Use of different substituents in the chain is demonstrated for **3f,g** and the reaction could be extended to the synthesis of the unsubstituted 2-phenyl piperidine **3h**.




**Scheme 1.** Piperidine Formation from C-H Amination: Scope (0.2 mmol scale). Yields refer to isolated material after purification. All reactions proceed with >90% selectivity in favor of piperidine formation (>95% yield based on recovered starting material). <sup>a</sup>1.6 equiv. of NBS. <sup>b</sup>20 mol% I<sub>2</sub>, white LED.

Common organic substituents are well tolerated on the arene group as demonstrated for derivatives **3i-3v**. These examples include 2-, 3- and 4-disubstitution patterns as well as higher substitution and as for **3q,r** also include carbonyl derivatives, which are non-compatible with the corresponding light induced iodine-catalyzed Hofmann-Löffler reaction.<sup>4</sup> The reaction also proceeds for heteroaromatic (**3w,x**) and dibenzylic derivatives (**3y**), and yields diastereomerically pure piperidine **3z** from cyclic stereocontrol, while acyclic stereocontrol is not possible under the reaction conditions (**3aa** and **3ab**). The C-H amination scope also includes related heterocycles such as the pharmaceutically relevant piperazine core **3ac**. For compounds **3c**, **3m** and **3z** their constitution was unambiguously established by single crystal X-ray analyses.<sup>14</sup>

It is noteworthy that potentially competing pyrrolidine formation was not observed in any of these cases. While the reactions were usually conducted with tosylamides as the representative sulfonamide groups, the use of SES and Ns enables a convenient approach to the corresponding free piperidines.<sup>14</sup> The reaction conditions could also be extended to the selective formation of a seven-membered derivative **5**, using an increased catalyst loading. It demonstrates the inherent potential of the current methodology for the synthesis of more advanced nitrogen heterocycles such as azepanes as well.


This novel C-H amination reaction is rationalized by the following merger of two catalytic cycles (Figure 2).



**Figure 2. Position-selective intramolecular C-H amination.**

It initiates from visible light-assisted homolytic cleavage of the N-Br bond in NBS.<sup>15</sup> The N-centered succinimidoyl radical then abstracts a hydrogen atom at the benzylic position of the substrate **2**. The respective benzylic C-H bonds are of low-

er BDE than the competing ones, which provides the required selectivity within this free radical reaction step.<sup>16</sup> Control experiments with deuterated starting materials **2h-d<sub>1</sub>** and **2h-d<sub>2</sub>** provide intra- and intermolecular kinetic isotope effects of 2.6 and 2.7, respectively (Scheme 2, eq. 1,2).<sup>14</sup>



**Scheme 2. Control Experiments.**

These results suggest the intramolecular radical C-H abstraction to be the slow step of the reaction, which is further corroborated for a Hammett correlation with a  $\rho$ -value of -2.9 (eq. 3). The intermediary benzylic radical **A** abstracts an iodine atom from a halogen-bonded<sup>17</sup>  $I_2$ -NBS adduct **B** to generate IBr **6** and the intermediary benzyl iodide **D** and regenerates the succinimidoyl radical.<sup>18</sup> The formation of the latter closes the catalytic cycle of radical C-H functionalization. The benzylic iodide undergoes nucleophilic substitution to the pyrrolidine product **3**.<sup>19</sup> The liberated HI regenerates the molecular iodine catalyst upon reaction with IBr **6**. This compound could potentially engage in radical halide formation itself, but is unproductive under the current conditions as demonstrated by a control experiment (Scheme 2, eq. 4). Molecular iodine recoordinates NBS **1a** within a halogen bonding mode,<sup>17</sup> which closes the second catalytic cycle of iodine catalysis. Closely related species such as  $I_2$ -N-chlorophthalimide had been invoked previously by Ishihara in iodolactonization reactions.<sup>20,21</sup> The postulation of a free radical mechanism outside the classical N-centered radical from N-halogenation as in the Hofmann-Löffler scenario is in agreement with the observation that molecular iodine does not convert tosylamides into their N-iodinated derivatives.<sup>4</sup> However, upon polarization by halogen bonding with the N-bromo succinimide **1a** the radical pathway to C-H functionalization is switched on. The postulated pathways do not involve direct benzylic bromination with NBS **1a**. This is in agreement with the observation that benzyl bromide derivative **7** does not cyclize to piperidine **3e** under the reaction conditions (Scheme 2, eq. 5) and thus con-

stitutes a dead-end. Obviously, successful nucleophilic piperidine formation requires the more reactive benzyl iodide intermediate **D**.<sup>14</sup>

In summary, we have identified mild and uniform conditions for a selective iodine-catalyzed C-H amination of 2-aryl substituted piperidines. This reaction overrides the common preference for pyrrolidine formation within the Hofmann-Löffler manifold and significantly enlarges both the scope of light-induced iodine-catalysis and position-selective C-H amination reactions. In addition, it diversifies the chemical space of piperidines.

## AUTHOR INFORMATION

### Corresponding Author

\* kmuniz@iciq.es

### Notes

The authors declare no competing financial interest.

## ASSOCIATED CONTENT

**Supporting Information.** Experimental details, control experiments and compound characterization (PDF), and details on the X-ray analyses (CIF). The Supporting Information is available free of charge on the ACS Publications website.

## ACKNOWLEDGMENT

Financial support for this project was provided from the Spanish Ministry for Economy and Competitiveness and FEDER (CTQ2014-56474R grant to K. M., and Severo Ochoa Excellence Accreditation 2014-2018 to ICIQ, SEV-2013-0319), and the ICIQ-COFUND Program (fellowship to H. Z.). The authors are grateful to the CERCA Programme of the Government of Catalonia and to COST Action CA15106 “C-H Activation in Organic Synthesis” (CHAOS).


## ABBREVIATIONS

NBS, N-bromo succinimide; DCE, dichloroethane.

## REFERENCES

- (1) (a) Cabrele, C.; Reiser, O. *J. Org. Chem.* **2016**, *81*, 10109-10125. (b) Eicher, T.; Hauptmann, S.; Speicher, A. *Chemistry of Heterocycles: Structure, Reactions, Synthesis and Applications*, Wiley-VCH, New York, 2013; (c) Ricci, A. ed., *Amino Group Chemistry: From Synthesis to the Life Sciences*, Wiley-VCH, Weinheim, 2007; (d) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K. *Angew. Chem. Int. Ed.* **2012**, *51*, 8960-9009.
- (2) Yan, M.; Lo, J. C.; Edwards, J. T.; Baran, P. S. *J. Am. Chem. Soc.* **2016**, *138*, 12692-12741.
- (3) (a) Wolff, M. E. *Chem. Rev.* **1963**, *63*, 55-64. (b) Stella, L. *Angew. Chem. Int. Ed.* **1983**, *22*, 337-350. (c) Jeffrey, J. L.; Sarpong, R. *Chem. Sci.* **2013**, *4*, 4092-4106.
- (4) Martínez, C.; Muñoz, K. *Angew. Chem. Int. Ed.* **2015**, *54*, 8287-8291.
- (5) ((a) Hu, X.-Q.; Chen, J.-R.; Xiao, W.-J. *Angew. Chem. Int. Ed.* **2017**, *56*, 1960-1962. (b) Xiong, T.; Zhang, Q. *Chem. Soc. Rev.* **2016**, *45*, 3069-3087. (c) Chen, J.-R.; Hu, X.-Q.; Lu, L.-Q.; Xiao, W.-J. *Chem. Soc. Rev.* **2016**, *45*, 2044-2056.
- (6) Nechab, M.; Mondal, S.; Bertrand, M. P. *Chem. Eur. J.* **2014**, *20*, 16034-16059.
- (7) For exceptions under stoichiometric reaction conditions: (a) Wawzonek, S.; Nelson Jr., M. F.; Thelen, P. J. *J. Am. Chem. Soc.* **1951**, *73*, 2806-2808. (b) Ban, Y.; Kimura, M.; Oishi, T. *Chem. Pharm. Bull.* **1976**, *24*, 1490-1496. (c) Freire, R.; Martín, A.; Pérez-Martín, I.; Suárez, E. *Tetrahedron Lett.* **2002**, *43*, 5113-5116. (d) Cherney, E. C.; Lopchuk, J. M.; Green, J. C.; Baran, P. S. *J. Am. Chem. Soc.* **2014**, *136*, 12592-12595. (e) Chen, K.; Richter, J. P.; Baran, P. S. *J. Am. Chem. Soc.* **2008**, *130*, 7247-7249.
- (8) (a) Horton, D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, *103*, 893-930. (b) Felpin, F. X.; Lebreton, J. *Eur. J. Org. Chem.* **2003**, 3693-3712. (c) O'Hagan, D. *Nat. Prod. Rev.* **2000**, *17*, 435-446. (d) Ritchie, T. J.; Macdonald, S. J. F.; Young, R. J.; Pickett, S. D. *Drug Discovery Today* **2011**, *16*, 164-171.
- (9) Vitaku, E.; Smith, D. T.; Njardarson, J. T. *J. Med. Chem.* **2014**, *57*, 10257-10274.
- (10) General review on catalysis of radical reactions: Studer, A.; Curran, D. P. *Angew. Chem. Int. Ed.* **2016**, *55*, 58-102.
- (11) General review on hydrogen atom transfer: Mayer, J. M. *Acc. Chem. Res.* **2011**, *44*, 36-46.
- (12) O'Broin, C. Q.; Fernández, P.; Martínez, C.; Muñoz, K. *Org. Lett.* **2016**, *18*, 436-439.
- (13) For general concepts of iodine-catalysed C-H/N-H coupling: (a) Finkbeiner, P.; Nachtsheim, B. *Synthesis* **2013**, *45*, 979-999. (b) Uyanik, M.; Ishihara, K. *ChemCatChem* **2012**, *4*, 177-185. (c) Li, J.; Lear, M. J.; Kawamoto, Y.; Umehiya, S.; Wong, A. R.; Kwon, E.; Sato, I.; Hayashi, Y. *Angew. Chem. Int. Ed.* **2015**, *54*, 12986-12990.
- (14) See Supporting Information for details.
- (15) (a) Saikia, I.; Borah, A. J.; Phukan, P. *Chem. Rev.* **2016**, *11*, 6837-7042. (b) Djerassi, C. *Chem. Rev.* **1948**, *48*, 271-317. (c) Skell, P. S.; Day, J. C. *Acc. Chem. Res.* **1978**, *11*, 381-387.
- (16) For related iodine catalyzed intermolecular benzylic amination: (a) Xue, Q.; Xie, J.; Li, H.; Cheng, X.; Zhu, C. *Chem. Commun.* **2013**, *49*, 3700-3702. (b) Zhang, X.; Wang, M.; Li, P.; Wang, L. *Chem. Commun.* **2014**, *50*, 8006-8009. However, these conditions are not compatible with the requirements for the present intramolecular reaction.<sup>14</sup>
- (17) a) Cavallo, G.; Metrangolo, P.; Milani, R.; Pilati, T.; Priimagi, A.; Resnati, G.; Terraneo, G. *Chem. Rev.* **2016**, *116*, 2478-2601. (b) Metrangolo, P.; Resnati, G. *Halogen Bonding Vol I: Impact on materials chemistry and life sciences*, Springer, Berlin, 2015. (c) Metrangolo, P.; Resnati, G. *Halogen Bonding Vol II: Impact on materials chemistry and life sciences*, Springer, Berlin, 2015. (d) Breugst, M.; Detmar, E.; von der Heiden, D. *ACS Catalysis* **2016**, *6*, 3203-3212.
- (18) Photochemical initiation of the reaction may also occur via homolysis at the I<sub>2</sub>-NBS stage **B**.
- (19) Hydrogen bonding between the benzyliodide **D** and IBr may also be involved as an accelerator in the C-N bond formation to **3**: Tsuji, N.; Kobayashi, Y.; Takemoto, Y. *Chem. Commun.* **2014**, *50*, 13691-13694.
- (20) (a) Nakatsuji, H.; Sawamura, Y.; Sakakura, A.; Ishihara, K. *Angew. Chem. Int. Ed.* **2014**, *53*, 6974-6977. (b) Sakakura, A.; Ishihara, K. *Chem. Rec.* **2015**, *15*, 728-742.
- (21) For additional support on interaction between NBS and molecular iodine: Chaikovskii, V. K.; Skorokhodov, V. I.; Filimonov, V. D. *Russ. J. Org. Chem.* **2001**, *37*, 1503-1504.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- selective piperidine formation
- benign light induced reaction
- radical C-H functionalization
- iodine-catalyzed C-N bond formation