

Preparation and Properties of Tetrathio-derivatives of
Quinodimethanes Fused to a 1,2,5-Thiadiazole Unit and Related
Heterocycles

Yoshiro YAMASHITA* and Tsutomu MIYASHI

Department of Chemistry, Faculty of Science, Tohoku University,
Aramaki, Sendai 980

Wittig-Horner reaction of 2-dimethoxyphosphinyl-1,3-benzodithiolo with 1,4-benzoquinones fused to a 1,2,5-thiadiazole unit and related heterocycles gave a new type of donors which show reversible two-electron oxidation waves and formed conductive complexes with iodine and DDQ.

Recently bis(1,3-dithiolo) donors containing quinodimethane structures have received considerable attention since the cation radical states are stabilized by formation of a new aromatic sextet as well as extended conjugation.¹⁾ In this connection, anthraquinodimethane **1** is an interesting donor. Although the preparation is reported,²⁾ the properties have not been described. **1** seems to be a nonplanar molecule due to the steric interaction caused by the peri-hydrogens as found in tetracyanoanthraquinodimethane (TCNAQ).³⁾ The steric interaction in TCNAQ has been removed by the replacement of the benzene rings with 1,2,5-thiadiazole rings.⁴⁾ It seems also possible to remove the steric interaction in donor **1** by the replacement to give mono-substituted derivative **2a** and bis-substituted derivative **3a**. Another advantage of the replacement is that S---N interactions can be expected in **2a** and **3a**, which lead to high-dimensional structures in their complexes and ion radical salts.⁵⁾ From these viewpoints, we have prepared donors **2a**, **3a**, and related heterocycles **2b**, **3b**, and investigated their properties.

Bis(1,3-dithiolo) donors seem to be prepared by the Wittig reaction of the corresponding diones, which is the most straight-forward and simple method, however such methods have not been used except for the preparation of **1**. We have succeeded in preparing the donors **2** and **3** by using the Wittig-Horner reaction. Thus, the reaction of 2-dimethoxyphosphinyl-1,3-benzodithiolo^{2,6)} (**4**) with naphtho[2,3-c][1,2,5]thiadiazole-4,9-dione⁷⁾ in the presence of n-BuLi gave **2a** (mp 367-370 °C) in 90% yield with a small amount of ketone **5a** (mp 289-291 °C). The similar reaction of naphtho[2,3-c][1,2,5]selenadiazole-4,9-dione⁸⁾ gave **2b** (mp 345-347 °C) in 69% yield with a small amount of ketone **5b** (mp 281-283 °C). The ketones **5a,b** were mainly obtained when one equivalent of the Wittig reagent was used. Ketone **6** (mp 186-189 °C) was also obtained along with **1** in the reaction of **4** with anthraquinone. Bis-substituted derivative **3a** (mp >400 °C) and **3b** (mp 376-




Table 1. Absorption maxima in the electronic spectra

Compound	λ_{\max}/nm ($\log \epsilon$) in CH_2Cl_2
1	237(4.85), 348(4.29), 398sh(4.42), 415(4.19)
2a	235(4.58), 360(4.21), 422(4.64)
2b	233(4.57), 305(4.19), 414(4.60)
3a	240sh, 257sh, 302, 380sh, 400, 420
3b	248(4.60), 335sh(4.36), 374(4.59)
5a	254(4.42), 280sh(4.24), 306sh(3.97), 404(4.39), 480(4.01)
5b	252(4.29), 266sh(4.26), 322(4.01), 334sh(3.99), 407(4.42), 492(3.59)
6	248(4.67), 272sh(4.18), 347(3.95), 452(4.19)
8a	251(4.49), 320(4.28), 520(3.91)




Table 2. Decomposition points, molar ratios,^{a)} and electrical resistivities, ρ ,^{b)} of complexes of donors **1**, **2**, and **3** with iodine and DDQ

Donor	I complex			DDQ complex		
	Decomp/°C	Ratio	ρ/Ω cm	Decomp/°C	Ratio	ρ/Ω cm
1	302-305	1:5.9	$>10^8$	360-363	0.5:1:H ₂ O	8.8×10^7
2a	300-310	1:2.6	8.1×10	>300	0.7:1	1.4×10^2
2b	343-346	1:2.6	1.8×10^2	>230	0.7:1	4.8×10^2
3b	>400	1:2.6	3.5×10	--- ^{c)}		

a) Determined on the basis of elemental analyses. b) Measured by a two-probe technique on compaction samples. c) No attempt.

378 °C) were also prepared by the similar reaction of the corresponding quinones^{8,9)} in 50% and 38% yields, respectively. These results show that the Wittig-Horner reaction of 2-dimethoxyphosphinyl-1,3-benzodithiole with 1,4-benzoquinones fused to aromatic rings provides a useful method to prepare highly-conjugated bis(1,3-dithiole) donors.

The absorption maxima in the electronic spectra of the bis(1,3-dithiole) donors **1**, **2a,b**, **3a,b**, and ketones **5a,b**, **6** are shown in Table 1. The absorptions of the ketones are red-shifted as compared with those of the corresponding bis(1,3-dithiole) donors. This can be attributed to the contribution of the dipolar structure **7**. For a comparison, dicyanomethylene compound **8a** (mp > 250 °C), which is considered to be more polarized, was synthesized by the reaction of **4** with **8b**, and was found to have an absorption maximum at 520 nm.

The cyclic voltammetry of **1**, **2a**, and **2b** in acetonitrile¹⁰⁾ showed reversible two-electron oxidation waves at 0.39, 0.65, 0.70 V vs. SCE, respectively, indicating that Coulombic repulsion is decreased in the dications of these conjugated molecules. Two-electron oxidation waves are also observed in analogous bis(1,3-dithiole) donors with a naphthalene^{1f)} or 1,6-methano[10]annulene moiety.^{1d)} The higher values in **2a** and **2b** reflect the electron withdrawing property of a 1,2,5-thiadiazole ring and a 1,2,5-selenadiazole ring.¹¹⁾ On the other hand, the oxidation potential of **3b** in dichloromethane was 0.78 V vs. SCE (irreversible).¹²⁾

Anthraquinodimethane (**1**) formed complexes with iodine and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) whose electrical resistivities are high as shown in Table 2. On the other hand, heterocycles **2a** and **2b** gave better conductive complexes with iodine and DDQ than **1**, in which the molar ratios are also different from those of the complexes of **1**. In addition, electrochemical oxidation of a solution containing **2a** or **2b** (10^{-3} mol dm⁻³) and tetraethylammonium perchlorate (0.06 mol dm⁻³) in dichloromethane using platinum electrodes at 10 μ A/cm² gave perchlorate salts of the cation radicals. The molar ratios were 1:1.2 (2:ClO₄⁻) in both salts based on elemental analyses, indicating that these are novel salts

in which the dication states are involved.¹³⁾ The electrical resistivities of these salts of **2a** and **2b** were 7.1×10^2 and $7.4 \times 10^2 \Omega \text{ cm}$, respectively.

In the case of bis-substituted derivative **3a**, the low solubility in solvents prevented the study on complexes with acceptors. However, the iodine complex of bithieno derivative **3b** exhibited higher conductivity as shown in Table 2, indicating that the bis-substituted derivatives of **1** are promising electron donors to provide highly conductive compounds.

References

- 1) a) Y. Ueno, A. Nakayama, and M. Okawara, *J. Chem. Soc., Chem. Commun.*, 1978, 74; b) M. Sato, M. V. Lakshmikantham, M. P. Cava, and A. F. Garito, *J. Org. Chem.*, 43, 2084 (1978); c) J.-M. Fabre, E. Torreilles, and L. Giral, *Tetrahedron Lett.*, 1978, 3703; d) R. Neidlein, D. Droste-Tran-Viet, A. Gieren, M. Kokkinidis, R. Wilckens, H.-P. Geserich, and W. Ruppel, *Helv. Chim. Acta.*, 67, 574 (1984); e) M. R. Bryce, *J. Chem. Soc., Perkin Trans. 1*, 1985, 1675; f) M. R. Bryce, *Mol. Cryst. Liq. Cryst.*, 120, 305 (1985).
- 2) K. Akiba, K. Ishikawa, and N. Inamoto, *Bull. Chem. Soc. Jpn.*, 51, 2674 (1978).
- 3) U. Schubert, S. Hünig, and A. Aumüller, *Liebigs, Ann. Chem.*, 1985, 1216; C. Kabuto, Y. Fukazawa, T. Suzuki, Y. Yamashita, T. Miyashi, and T. Mukai, *Tetrahedron Lett.*, 27, 925 (1986).
- 4) Y. Yamashita, T. Suzuki, T. Mukai, and G. Saito, *J. Chem. Soc., Chem. Commun.*, 1985, 1044.
- 5) G. C. Papavassilliou, S. Y. Yiannopoulos, and J. S. Zambounis, *J. Chem. Soc., Chem. Commun.*, 1986, 821; T. Suzuki, C. Kabuto, Y. Yamashita, and T. Mukai, *Bull. Chem. Soc. Jpn.*, 60, 2109 (1987); T. Suzuki, C. Kabuto, Y. Yamashita, T. Mukai, T. Miyashi, and G. Saito, *ibid.*, in press.
- 6) K. Ishikawa, K. Akiba, and N. Inamoto, *Tetrahedron Lett.*, 1976, 3695.
- 7) R. Neeff and O. Bayer, *Chem. Ber.*, 90, 1137 (1957).
- 8) R. Neidlein, D. Tran-Viet, A. Gieren, M. Kokkinidis, R. Wilckens, H.-P. Geserich, and W. Ruppel, *Chem. Ber.*, 115, 2898 (1982).
- 9) D. W. H. MacDowell and J. C. Wiscowaty, *J. Org. Chem.*, 37, 1712 (1972).
- 10) Measured at a platinum electrode with 0.1 mol dm^{-3} of tetraethylammonium perchlorate as a supporting electrolyte; scan rate 100 mV s^{-1} .
- 11) T. Suzuki, C. Kabuto, Y. Yamashita, and T. Mukai, *Chem. Lett.*, 1987, 1129.
- 12) The oxidation potential of **3b** could not be measured in acetonitrile due to the low solubility. The irreversible potential was calculated as E_{pa} (anodic peak potential) - 0.03. That of **1** in dichloromethane was 0.68 V vs. SCE (irreversible). That of **3a** could not be measured even in dichloromethane due to the low solubility.
- 13) The salts were obtained as fine black needles and the analytical data were reproducible.

(Received January 22, 1988)