

Direct Oxidative Conversion of
Methylarenes into Aromatic Nitriles


Daisuke Tsuchiya, Yuhsuke Kawagoe, Katsuhiko Moriyama, and Hideo Togo*

Graduate School of Science, Chiba University, Yayoi-cho 1-33, Inage-ku,
Chiba 263-8522, Japan

togo@faculty.chiba-u.jp

Received July 8, 2013

ABSTRACT



A variety of methylarenes were successfully converted into the corresponding aromatic nitriles in good to moderate yields by the treatment with NBS or DBDMH in the presence of a catalytic amount of AIBN or BPO, followed by the reaction with molecular iodine in aq NH₃ in a one-pot procedure. The present reaction is a useful and practical transition-metal-free method for the preparation of aromatic nitriles from methylarenes.

Study of the practical transition-metal free methods for the preparation of aromatic nitriles is very important since they are precursors for the preparation of amides, esters, primary amines, carboxylic acids, aldehydes, ketones, and nitrogen-containing heterocycles, such as tetrazoles and oxazoles.¹ They have also great importance in the synthesis of agrochemicals, therapeutic drugs, functional materials,

natural products, dyes, and pigments.^{1,2} Today, the most conventional methods for the preparation of aromatic nitriles are the dehydration of primary aromatic amides^{3,4} with SOCl₂, TsCl/Py, P₂O₅, POCl₃, COCl₂, (EtO)₃P/I₂, Ph₃P/CCl₄, or (COCl)₂/DMSO and the Sandmeyer reaction of aromatic diazonium ion with toxic CuCN.^{4,5} Recently, the direct transformation of aromatic bromides into the corresponding aromatic nitriles was actively studied with CuCN in *N,N*-dimethylformamide (DMF) at 153 °C (the Rosenmund–von Braun reaction),^{6a} and related reactions with palladium and metal cyanide were

(1) (a) Sandler, S. R.; Karo, W. Nitriles (Cyanides). In *Organic Functional Group Preparations*; Wasserman, H. H., Ed.; Academic Press: San Diego, 1983; Vol. 12-I of Organic Chemistry, Chapter 17. (b) Fabiani, M. E. *Drug News Perspect.* **1999**, 12, 207.

(2) (a) Friedrich, K.; Wallensfels, K. *The Chemistry of the Cyano Group*; Rappoport, Z., Ed.; Wiley-Interscience: New York, 1970. (b) North, M. *Comprehensive Organic Functional Group Transformation*; Katritzky, A. R., Meth-Cohn, O., Rees, C. W., Eds.; Pergamon: Oxford, 1995. (c) Murahashi, S.-I. Synthesis from Nitriles with Retention of the Cyano Group. *Science of Synthesis*; Georg Thieme Verlag: Stuttgart, 2004; Vol. 19, pp 345–402. (d) Collier, S. J.; Langer, P. Application of Nitriles as Reagents for Organic Synthesis with Loss of the Nitrile Functionality. *Science of Synthesis*; Georg Thieme Verlag: Stuttgart, 2004; Vol. 19, pp 403–425.

(3) (a) Reisner, D. B.; Coring, E. C. *Organic Synthesis*; Wiley: New York, 1963; Collect. Vol. IV, p 144. (b) Krynitsy, J. A.; Carhart, H. W. *Organic Synthesis*; Wiley: New York, 1963; Collect. Vol. IV, p 436. (c) Humber, J. G.; Davis, M. A. *Can. J. Chem.* **1966**, 44, 2113. (d) Yamato, E.; Sugawara, S. *Tetrahedron Lett.* **1970**, 4348. (e) Lehnert, W. *Tetrahedron Lett.* **1971**, 1501. (f) Campagna, F.; Caroti, A.; Casini, G. *Tetrahedron Lett.* **1977**, 1813. (g) Mai, K.; Patal, G. *Tetrahedron Lett.* **1986**, 27, 2203. (h) Claremon, D. A.; Phillips, B. *Tetrahedron Lett.* **1988**, 29, 2155. (i) Nemoto, H.; Kubota, Y.; Yamamoto, Y. *J. Org. Chem.* **1990**, 55, 4515. (j) Sznajdman, M. L.; Crasto, C.; Hecht, S. M. *Tetrahedron Lett.* **1993**, 34, 1581. (k) Xue, C. B.; Degrado, W. F. *Tetrahedron Lett.* **1995**, 61, 6486 and references cited therein. (l) Heck, M.; Wagner, A.; Mioskowski, C. *J. Org. Chem.* **1996**, 61, 6486. (m) Nakajima, N.; Ubukata, M. *Tetrahedron Lett.* **1997**, 38, 2099. (n) Bose, D. S.; Jayalakshmi, B.; Goud, P. R. *Synthesis* **1999**, 1724. (o) Bose, D. S.; Jayalakshmi, B. *Synthesis* **1999**, 64. (p) Kuo, C.; Zhu, J.; Wu, J.; Chu, C.; Yao, C.; Shia, K. *Chem. Commun.* **2007**, 301.


(4) Larock, R. C. *Comprehensive Organic Transformation*; VCH Publishers: Weinheim, 1989; pp 976–993.

(5) Sandmeyer, T. *Ber.* **1884**, 17 (1633), 2650.

(6) (a) Sharman, W. M.; Van Lier, J. E. In *Porphyrin Handbook*; Kadish, E., Smith, K. M., Guillard, R., Eds.; Academic Press: New York, 2003; Vol. 15, p 1. (b) Weissman, S. A.; Zewge, D.; Chen, C. *J. Org. Chem.* **2005**, 70, 1508. (c) Littke, A.; Soumeillant, M.; Kaltenbach, R. F., III; Cherney, R. J.; Tarby, C. M.; Kiau, S. *Org. Lett.* **2007**, 9, 1711. (d) Martin, M. T.; Liu, B.; Cooley, B. E., Jr.; Eaddy, J. F. *Tetrahedron Lett.* **2007**, 48, 2555. (e) Nandurkar, N. S.; Bhanage, B. M. *Tetrahedron* **2008**, 64, 3655. (f) Iqbal, Z.; Lyubimtsev, A.; Hanack, M. *Synlett* **2008**, 2287. (g) Chen, G.; Weng, J.; Zheng, Z.; Zhu, X.; Cai, Y.; Cai, J.; Wan, Y. *Eur. J. Org. Chem.* **2008**, 3524. (h) Schareina, T.; Zapf, A.; Cotte, A.; Müller, N.; Beller, M. *Synthesis* **2008**, 3351. (i) Buono, F. G.; Chidambaram, R.; Mueller, R. H.; Waltermire, R. E. *Org. Lett.* **2008**, 10, 5325. (j) Chattopadhyay, K.; Dey, R.; Ranu, B. C. *Tetrahedron Lett.* **2009**, 50, 3164. (k) DeBlase, C.; Leadbeater, N. E. *Tetrahedron* **2010**, 66, 1098. (l) Yan, G.; Kuang, C.; Zhang, Y.; Wang, J. *Org. Lett.* **2010**, 12, 1052. (m) Yu, H.; Richey, R. N.; Miller, W. D.; Xu, J.; May, S. A. *J. Org. Chem.* **2011**, 76, 665. (n) Ushkov, A. V.; Grushin, V. V. *J. Am. Chem. Soc.* **2011**, 133, 10999. (o) Yeung, P. Y.; Tsang, C. P.; Kwong, F. Y. *Tetrahedron Lett.* **2011**, 52, 7038. (p) Hajipour, A. R.; Rafiee, F.; Ruoho, A. E. *Tetrahedron* **2012**, 53, 526. (q) Zhang, D.; Sun, H.; Zhang, L.; Zhou, Y.; Li, C.; Jiang, H. *Chem. Commun.* **2012**, 48, 2909.

reported.^{6b–q} However, all of these reactions require toxic metal cyanides. Typical examples of cyanide-free and transition-metal-free methods for the preparation of aromatic nitriles are the reaction of electron-rich aromatics with chlorosulfonylisocyanate to form *N*-chlorosulfonyl amides and the subsequent treatment with DMF to give aromatic nitriles, together with generation of SO₃ and HCl,^{7a,b} and the reaction of indoles or pyrroles with triphenylphosphine–thiocyanogen (TPPT).^{7c} Three methods reported by us recently are the reaction of bromoarenes, methoxybenzene, dimethoxybenzenes, and 1,3-difluorobenzene with *n*-BuLi and subsequently DMF, followed by the reaction with molecular iodine and aq NH₃,^{8a,b} the reaction of electron-rich aromatics, such as dimethoxybenzenes, with POCl₃–DMF, followed by the reaction with molecular iodine and aq NH₃,^{8c,d} and the reaction of bromoarenes with Mg and then DMF, followed by the reaction with molecular iodine and aq NH₃.^{8e,f} On the other hand, studies of the direct oxidative conversion of methylarenes into aromatic nitriles are extremely limited, as shown in Scheme 1. One conversion method requires high temperatures (360–460 °C) using ammonia and oxygen in the presence of (VO)₂P₂O₇ catalyst^{9a} (Scheme 1, eq 1).

Scheme 1. Established Methods for Transformation of Methylarene to Aromatic Nitrile



Another conversion method involves the reaction of methylarenes with excess amounts of NaN₃ and (diacetoxyiodo)benzene (DIB) in the presence of CuSO₄ catalyst at room temperature,^{9b} which requires toxic NaN₃ and is effective for only electron-rich methylarenes (Scheme 1, eq 2). Here, we would like to report a cyanide-free and transition-metal-free one-pot conversion of methylarenes into aromatic nitriles by treatment with *N*-bromosuccinimide (NBS) or 1,3-dibromo-5,5-dimethylhydantoin (DBDMH) in the presence of a

catalytic amount of benzoyl peroxide (BPO) or 2,2'-azobis(isobutyronitrile) (AIBN), followed by the reaction with molecular iodine in aq NH₃ (Scheme 1, eq 3) because we know for a fact that benzylic halides can be effectively converted into the corresponding aromatic nitriles with molecular iodine and aq NH₃ (60 °C for a few hours).¹⁰ Thus, 4-bromotoluene **1a** was treated with NBS or DBDMH in the presence of a catalytic amount of BPO or AIBN in carbon tetrachloride at 80 °C, followed by the treatment with molecular iodine and aq NH₃ for 4 h at 60 °C, as the Wohl-Ziegler reaction is generally carried out in carbon tetrachloride.¹¹ After optimization studies, we found that BPO and AIBN showed the same reactivity as a radical initiator (entries 1 vs 2), DBDMH had higher reactivity than NBS (entries 1 vs 3), and acetonitrile exhibited higher reactivity than carbon tetrachloride in the Wohl-Ziegler reaction (entries 1 vs 4), to give 4-bromobenzonitrile **2a** in good yields, as shown in Table 1. Then, using the optimum conditions (Table 1, entry 7),

Table 1. One-Pot Transformation of 4-Bromotoluene to 4-Bromobenzonitrile

entry	first step			second step		
	radical initiator	"Br"	time (h)	I ₂ (equiv)	time (h)	yield (%)
1	BPO	DBDMH	2	2.5	4	86
2	AIBN	DBDMH	2	2.5	4	86
3 ^a	BPO	NBS	2	2.5	4	52
4 ^b	BPO	DBDMH	2	2.5	4	55
5 ^c	BPO	DBDMH	2	2.5	4	76
6	BPO	DBDMH	2	4.0	4	83
7	BPO	DBDMH	3	2.5	12	87
8		DBDMH	2	2.5	4	11

^a NBS (1.1 equiv) was used. ^b Reaction was carried out in CCl₄. ^c Reaction was carried out at 90 °C.

the conversion of 1-methylnaphthalene **11** into 1-cyanonaphthalene **21** with DBDMH in the presence of BPO, followed by treatment with molecular iodine and aq NH₃, was performed. However, 4-bromo-1-methylnaphthalene was obtained in 84% yield, together with 1-cyanonaphthalene in 9% yield, as shown in entry 1 of Table 2. Thus, under warming conditions, the polar bromination of 1-methylnaphthalene occurred to give mainly 4-bromo-1-methylnaphthalene. After optimization studies to suppress the polar bromination of aromatics, it was found that

(7) (a) Gerhard, L. *Ber.* **1967**, 100, 2719. (b) Gerhard, L. *Org. Synth.* **1970**, 50, 52. (c) Tamura, Y.; Kita, Y.; Kawasaki, T. *J. Synth. Org. Chem.* **1980**, 38, 891.

(8) (a) Ushijima, S.; Togo, H. *Synlett* **2010**, 1562. (b) Ushijima, S.; Moriyama, K.; Togo, H. *Tetrahedron* **2011**, 67, 958. (c) Ushijima, S.; Togo, H. *Synlett* **2010**, 1067. (d) Ushijima, S.; Moriyama, K.; Togo, H. *Tetrahedron* **2012**, 68, 4588. (e) Ishii, G.; Moriyama, K.; Togo, H. *Tetrahedron Lett.* **2011**, 52, 2402. (f) Ishii, G.; Harigae, R.; Moriyama, K.; Togo, H. *Tetrahedron* **2013**, 69, 1462.


(9) (a) Review: Lücke, B.; Narayana, K. V.; Martin, A.; Jähnisch, K. *Adv. Synth. Catal.* **2004**, 346, 1407 and references cited therein. (b) Zhou, W.; Zhang, L.; Jiao, N. *Angw. Chem. Int. Ed.* **2009**, 48, 7094.

(10) (a) Iida, S.; Togo, H. *Synlett* **2008**, 1639. (b) Iida, S.; Ohmura, R.; Togo, H. *Tetrahedron* **2009**, 65, 6257.

(11) (a) Wohl, A. *Ber.* **1919**, 52, 51. (b) Ziegler, K. *Ann.* **1942**, 551, 30. *Organic Synthesis*; Wiley: New York, 1973; Collect. Vol. V, p 145.

irradiation of the mixture of 1-methylnaphthalene **1l** and NBS in carbon tetrachloride with a tungsten lamp (200 W, *W-hv*) gave 1-cyanonaphthalene **2l** in 88% yield (Table 2, entry 6). Thus, those two reaction systems, namely, DBDMS with BPO in refluxing acetonitrile (method A) for electron-deficient methylarenes and NBS under irradiation with a tungsten lamp in carbon tetrachloride at room temperature (method B) for electron-rich methylarenes, are recommended for the conversion of methylarenes **1** into aromatic nitriles **2**.
























Table 2. One-Pot Transformation of 1-Methylnaphthalene to 1-Cyanonaphthalene

<div>  </div>						
first step						
entry	"Br" (mol %)	solvent	temp (°C)	time (h)	yield (%)	
1	BPO	DBDMH (55)	MeCN	80	2	9 (84) ^b
2	AIBN	DBDMH (55)	CCl ₄	60	2	54
3	<i>hν</i> ^a	NBS (110)	CCl ₄	rt	1	80
4	<i>hν</i> ^a	NBS (110)	CCl ₄	rt	1	52
5	<i>hν</i> ^a	DBDMH (55)	CCl ₄	rt	1	4
6	<i>hν</i> ^a	NBS (110)	CCl ₄	rt	2	88

^a The first step reaction was carried out under tungsten lamp irradiation. ^b The number in parenthesis indicates the yield of 4-bromo-1-methylnaphthalene.

Then, these two methods (methods A and B) were used for the conversion of various methylarenes **1** into aromatic nitriles **2**, as shown in Table 3. For electron-deficient methylarenes **1**, such as 2-bromotoluene **1b**, 3-bromotoluene **1c**, 4-chlorotoluene **1d**, 4-iodotoluene **1e**, 4-benzoyltoluene **1f**, 4-methanesulfonyltoluene **1g**, 4-cyanotoluene **1h**, and 4-nitrotoluene **1i**, both methods A and B furnished the corresponding aromatic nitriles **2** in good to moderate yields (entries 1–9), while the same treatment of methyl 4-methylbenzoate **1j** using methods A and B gave 4-cyanobenzamide **2j** in good yields instead of methyl 4-cyanobenzoate (entry 10). This is due to the occurrence of amidation of formed methyl 4-cyanobenzoate by aq NH₃ at 60 °C. For toluene **1k**, 2-methylnaphthalene **1m**, 4-methylbiphenyl **1n**, and 4-methyl-*tert*-butylbenzene **1s**, both methods A and B gave the corresponding aromatic nitriles **2k**, **2m**, **2n**, and **2s** in good to moderate yields (entries 11–14 and 19). It should be noted that method A could not be used for 4-methoxytoluene **1o** and 2-methoxytoluene **1p** because of polar bromination of the aromatic rings. However, using method B, 4-methoxytoluene **1o** and 2-methoxytoluene **1p** were smoothly transformed into 4-methoxybenzonitrile **2o** and 2-methoxybenzonitrile **2p** in good yields, respectively (entries 15 and 16). Moreover, when method B was used for 2-methoxytoluene **1p** and


Table 3. One-Pot Transformation of Methylarenes to Aromatic Nitriles

<div> <div>Method A: BPO (10 mol %) DBDMH (55 mol %) CH₃CN, 80 °C</div> <div>Method B: NBS (110 mol %) CCl₄, rt, W-hv</div> </div>		<div> <div>I₂ (2.5 equiv) aq NH₃</div> <div>60 °C</div> </div>		Ar-CN 2	
entry	product	yield (%) (time ^{1st} (h), time ^{2nd} (h))	Method A	Method B	
1		2a	87 (3, 12)	88 ^{a,b} (2, 4)	
2		2b	84 ^c (4, 48)	56 (2, 4)	
3		2c	71 ^c (4, 48)	69 ^b (2, 4)	
4		2d	80 ^d (21, 12)	63 (2, 4)	
5		2e	82 ^c (2, 18)	68 ^b (2, 4)	
6		2f	74 ^d (12, 8)	60 (2, 5)	
7		2g	60 (12, 4)	75 ^b (2, 18)	
8		2h	87 (2, 15)	85 (2, 15)	
9		2i	71 ^d (13, 12)	61 (4, 12)	
10		2j	75 ^e (2, 4)	78 ^e (4, 18)	
11		2k	80 ^{c,f} (4, 4)	61 (2, 4)	
12		2l	71 ^{c,g,h,i} (2, 4)	88 (2, 4)	
13		2m	60 ^{c,g} (2, 4)	77 (2, 4)	
14		2n	82 ^{c,j} (2, 16)	53 (2, 4)	
15		2o	0 (2, 4)	90 (2, 4)	
16		2p	-	92 (2, 4)	
17		2q	-	71 ^{a,k} (5, 4)	
18		2r	-	72 ^k (6, 4)	
19		2s	76 ^c (2, 4)	82 (2, 4)	
20		2t	0 (2, 4)	40 ^l (2, 4)	
21		2u	0 (2, 4)	48 ^{b,m} (4, 18)	
22		2v	64 ^{c,n} (2, 4)	74 ^{b,n} (2, 4)	
23		2w	63 ^{c,n} (2, 12)	59 ⁿ (2, 12)	

^a CH₃CN was used instead of CCl₄. ^b DBDMH (1.0 equiv) was added instead of NBS. ^c CCl₄ was used instead of CH₃CN. ^d DBDMH (0.75 equiv) was added. ^e Methyl 4-methylbenzoate was used. ^f The reaction was carried out on a 3 mmol scale. ^g The reaction was carried out at 60 °C. ^h NBS (1.1 equiv) was added instead of DBDMH. ⁱ AIBN (0.1 equiv) was added instead of BPO. ^j DBDMH (1.0 equiv) was added. ^k NBS (2.2 equiv) was added. ^l The reaction was carried out in diluted conditions (0.1 M). ^m I₂ (3.5 equiv) was added at the second step. ⁿ NBS (2.2 equiv) was added at the first step, and I₂ (5.0 equiv) and aq NH₃ (6 mL) were added at the second step.


4-methyl-1,2-dimethoxybenzene **1r** with 2.2 equiv of NBS, brominated aromatic nitriles, such as 5-bromo-2-methoxybenzonitrile **2q** and 2-bromo-4,5-dimethoxybenzonitrile

Scheme 2. Plausible Reaction Mechanism for Transformation of Methylarenes into Nitriles



2r were selectively obtained (entries 17 and 18). *N*-Benzenesulfonyl-3-methylindole **1t** and 3-methylbenzothiophene **1u** were also converted into *N*-benzenesulfonyl-3-cyanoindole **2t** and 3-cyanobenzothiophene **2u** in moderate yields, respectively, by method **B** (entries 20 and 21). Furthermore, *p*-xylene **1v** and *m*-xylene **1w** were also converted into terephthalonitrile **2h** and isophthalonitrile **2w** in moderate yields, respectively, by methods **A** and **B** using twice the amounts of DBDMS or NBS reagent (entries 22 and 23). The present reaction pathway is shown in Scheme 2. At the initial step, the Wohl–Ziegler reaction of methylarenes with DBDMH or NBS in the presence of BPO or AIBN under warming conditions or irradiation conditions with a tungsten lamp occurs to form the corresponding arylmethyl bromides. By adding molecular iodine and aq NH₃ to the reaction mixture, the S_N2 reaction of arylmethyl bromides by NH₃ occurs to form arylmethylamines. Once arylmethylamines are formed, they are smoothly converted into the corresponding aromatic nitriles via the formation of imines and *N*-iodoimines.¹² Finally, the present method was applied to the preparation of Febuxostat precursor **5**, a nonpurine selective inhibitor of xanthine oxidase.¹³ 2-Methoxytoluene was treated with NBS (method **B**) under irradiation conditions to form 5-bromo-2-methoxybenzonitrile **2q**, as shown in Scheme 3. Then, compound **2q** was demethylated with BBr₃ and treated with 1-bromo-2-methylpropane to form

Scheme 3. Synthesis of Febuxostat



5-bromo-2-isobutoxybenzonitrile **4**. Compound **4** was coupled with 4-methyl-5-*t*-butoxycarbonylthiazole in the presence of Ni(OAc)₂ and *t*-BuOLi to generate Febuxostat precursor **5**.¹⁴ Once precursor **5** is formed, Febuxostat was easily obtained in 93% yield by treatment with CF₃CO₂H.

In conclusion, a variety of methylarenes were smoothly converted into the corresponding aromatic nitriles in good to moderate yields by the treatment with NBS or DBDMH in the presence of a catalytic amount of AIBN or BPO, followed by the reaction with molecular iodine in aq NH₃ in a one-pot procedure. The present reaction is a useful, practical, and transition-metal-free method for the preparation of aromatic nitriles from methylarenes.

Acknowledgment. Financial support in the form of a Grant-in-Aid for Scientific Research (No. 25105710) from the Ministry of Education, Culture, Sports, Science, and Technology in Japan, Iodine Research Project in Chiba University, is gratefully acknowledged.

Supporting Information Available. Experimental procedures, spectral data, and copies of NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(12) (a) Iida, S.; Togo, H. *Synlett* **2006**, 2633. (b) Iida, S.; Togo, H. *Synlett* **2007**, 407. (c) Iida, S.; Togo, H. *Tetrahedron* **2007**, 63, 8274.

(13) Okamoto, K.; Eger, B. T.; Nishuno, T.; Kondo, S.; Pai, E. F.; Nishino, T. *J. Biol. Chem.* **2003**, 278, 1848.

(14) Canivel, J.; Yamaguchi, J.; Ban, I.; Itami, K. *Org. Lett.* **2009**, 11, 1733.

The authors declare no competing financial interest.