

Variation of the C₁ Component in the Silicon-Induced Domino Synthesis of Cyclopentanes

Nico Bräuer, Tycho Michel and Ernst Schaumann*

*Institut für Organische Chemie, Technische Universität Clausthal
Leibnizstraße 6, D-38678 Clausthal-Zellerfeld, Germany*


Received 20 April 1998; revised 3 July 1998; accepted 19 July 1998

Abstract

The reaction of mono- or bisilylated thioanisole derivatives **2b–e** with 3,4-epoxybutyl tosylate (**3**) affords the cyclopentanol **6a–c**. Migratory aptitudes of two different silyl groups in the Brook 1,4-rearrangement is examined starting from **2d, e** giving the order $\text{SiMe}_2\text{Ph} > \text{SiMe}_3 > \text{SiMePh}_2$. The relative configuration of the resulting 1,3-functionalized cyclopentanes is established by NOE experiments after derivatization of the alcohol to the corresponding pivalates **7a–c**. © 1998 Elsevier Science Ltd. All rights reserved.



Keywords: Carbanion; Oxiranes; Rearrangements; Silicon and compounds

Recently, we have reported the one-pot synthesis of 1,3-difunctionalized cyclopentanes by a silicon-induced domino reaction [1,2]. This reaction proceeds via 1,4-shift of the silyl group from carbon to oxygen [3] after nucleophilic attack of a silyl-substituted thioacetal anion on an epoxyhomoallyl tosylate, followed by intramolecular displacement of the tosylate group (Scheme 1, Route B) [4]. Remarkably, the approach is not limited to a migrating SiMe_3 group, but the bulky SiMe_2^tBu group migrates too [5,6,7,8], but then oxetane formation becomes a competing process (Scheme 1, Route A).




Scheme 1

Here we wish to report on the extension of this reaction by using the C_1 bis-anion equivalents **1**, **2a-f** which should allow higher flexibility in subsequent transformations [9]. Furthermore, the modifications should provide information on the relative migratory aptitudes of different silyl groups in the cyclization reaction.




- 2a:** $R^1 = \text{H}$, $n = 0$
b: $R^1 = \text{SiMe}_3$, $n = 0$
c: $R^1 = \text{SiMe}_2\text{Ph}$, $n = 1$
d: $R^1 = \text{SiMe}_3$, $n = 1$
e: $R^1 = \text{SiMe}_3$, $n = 2$
f: $R^1 = \text{SiMe}_3$, $n = 3$

The C_1 building blocks **1** and **2a,b** have been reported before [10,11,12]. The C_1 components **2c-f** were synthesized by deprotonation and subsequent silylation of thioanisole using the procedure of Seebach (Scheme 2) [12].




Scheme 2

Reaction of the lithiated O,S-acetal **1** with 3,4-epoxybutyl tosylate (**3**) gave vinyl sulfide **4** in only 14% yield. This indicates that the desired domino reaction does occur, but only in low yields and with elimination of methoxide after the cyclization step (Scheme 3).



Scheme 3


Use of **2a** as C₁ component yielded the oxetane **5** as a mixture of diastereomers (Scheme 4).



Scheme 4

No cyclopentane could be isolated [13]. Insufficient stabilization of the carbanion after 1,4-carbon-oxygen shift seems to be the reason for this failure leading to the alternative oxetane formation (cf. Scheme 1, Route A).

The introduction of a second silyl group into **2a** should favour cyclopentane formation because of its additional stabilizing effect on the carbanion after 1,4-migration. Indeed, reaction of lithiated **2b** with **3** led to the 1,3-difunctionalized cyclopentane, which was isolated as cyclopentanol **6a** after purification by column chromatography (SiO₂) in good yield (Table 1, entry 1).



Scheme 5

Table 1.

Cyclopentane Formation by Silicon-Induced Domino Reaction and Diastereomer Ratios (d.r.)^a


Entry	C ₁ Component	~R ¹ (d.r.)	~SiMe _{3-n} Ph _n (d.r.)
1	2b	6a 76% (>20:1) ^b	---
2	2c	6b 62% (8.1:1)	---
3	2d	6b 35% (6.1:1)	6a 40% (>20:1) ^b
4	2e	6c 28% (10.3:1)	6a 16% (>20:1) ^b
5	2f	---	---

a) Diastereomer ratios were determined by gas chromatography after derivatization to the corresponding pivalate **7a-c**.

b) No other diastereomer could be detected in the 200 MHz ¹H NMR spectrum.

As judged by the yields of products **6a** vs. **6b** from **2d**, the SiMe₂Ph group migrates a little better than the SiMe₃ group (Table 1, entry 3). Obviously, the phenyl ring has a stabilizing effect on the migrating group and this electronic effect compensates steric hindrance. If a second methyl group on silicon is replaced by a phenyl ring as in **2e** the overall yield of **6** decreases, indicating that approach of **2e** on **3** is sterically hindered. Interestingly, **6c** resulting from migration of the SiMe₃ group is the main product, and not **6a**. This implies that the bulkiness of SiMePh₂ overrules the favourable electronic effect (Table 1, entry 4). Starting from **2f**, no cyclization product could be observed (Table 1, entry 5). Steric hindrance of the initial epoxide ring opening seems to be the reason for this failure.

The relative configuration of the major diastereomer was determined by NOE experiments after derivatization of the alcohol to the corresponding pivalate **7a-c** (62–92%). A 1,3-*trans*-arrangement of pivalate and silyl group was confirmed in each case.



Experimental Section

The ¹H and ¹³C NMR spectra were recorded using a Bruker DPX-200 (200 MHz) as dilute solutions in CDCl₃ with TMS as internal standard. Chemical shift values are given in parts per million (ppm). NOE measurements were carried out on a Bruker ARX-400 (400 MHz). The infrared spectra were recorded using a Bruker Vektor 22 spectrometer as neat films or as KBr-pellet. Elemental analyses were determined by the Institut für Pharmazeutische Chemie, Technische Universität Braunschweig. High resolution mass spectrum was measured by the Hans-Knöll-Institut für Naturstoff-Forschung, Jena. Column chromatography was carried out using Merck silica gel 6 (70–230 mesh). Silylated thioanisoles **1**, **2a,b** [6] and the epoxide **3** [1] are described in the literature.

General procedure for the preparation of silylated thioanisols 2d-f: A solution of **2a** in dry THF (1 ml/mmol) was cooled to -78°C , one equiv. of *n*-BuLi (1.6 M in hexane) was added dropwise and the mixture was stirred at 0°C for 2 h. At -78°C the appropriate silane (1.1 equiv.) was added to this solution. The solution was allowed to warm to room temperature and stirred for 12 h. The reaction mixture was hydrolyzed with diethyl ether/saturated NH_4Cl solution (1:1). The aqueous phase was separated and extracted twice with diethyl ether. The combined organic extracts were dried and the solvent evaporated. The crude product was purified by column chromatography (PE).

Phenyldimethylsilyl(phenylthio)(trimethylsilyl)methane (2d): Preparation from 10.86 g (55.3 mmol) of **2a** gave 12.93 g (71%) as a colourless oil. ^1H NMR: δ , -0.02 (s, 9H, SiMe_3), 0.38 and 0.45 (each s, 3H, SiMe_2Ph), 1.73 (s, 1H, CH), 7.05–7.68 (m, 10H, SiMe_2Ph , SPh); ^{13}C NMR: δ , -2.2, -1.8 (SiMe_2Ph), -0.4 (SiMe_3), 18.7 (CH), 125.0, 127.6, 127.9, 128.5, 129.1, 133.9, 138.7, 140.0 (SPh, SiMe_2Ph); IR: 3070, 3019, 2955, 2897, 1582, 1478, 1438, 1427, 1250, 1114, 1086, 1024, 995, 839, 782, 734, 669 cm^{-1} ; $\text{C}_{18}\text{H}_{26}\text{SSi}_2$ (330.62): calcd. C 65.39, H 7.93, S 9.70; found C 64.99, H 8.06, S, 9.71.

Diphenylmethylsilyl(phenylthio)(trimethylsilyl)methane (2e): Preparation from 12.62 g (64.2 mmol) of **2a** gave 14.92 g (59%) as a colourless oil. ^1H NMR: δ , 0.01 (s, 9H, SiMe_3), 0.79 (s, 3H, SiMePh_2), 2.21 (s, 1H, CH), 7.05–7.72 (m, 15H, SiMePh_2 , SPh); ^{13}C NMR: δ , -3.4 (SiMe_3), -0.3 (SiMePh_2), 17.2 (CH), 125.6, 127.5, 127.7, 128.1, 128.4, 129.2, 129.3, 134.7, 134.8, 136.4, 136.8, 139.6 (SPh, SiMePh_2); IR: 3069, 3050, 3018, 2955, 2896, 1582, 1478, 1427, 1250, 989, 838, 790, 735, 698 cm^{-1} ; $\text{C}_{23}\text{H}_{28}\text{SSi}_2$ (392.69): calcd. C 70.34, H 7.19, S 8.16; found C 70.33, H 7.12, S 7.95.

Phenylthio(trimethylsilyl)(triphenylsilyl)methane (2f): Preparation from 12.4 g (63.2 mmol) of **2a** gave 16.06 g (56%) as colourless solid (Fp: 64°C). ^1H NMR: δ , 0.00 (s, 9H, SiMe_3), 2.61 (s, 1H, CH), 7.12–7.89 (m, 20H, SiPh_3 , SPh); ^{13}C NMR: δ , 0.3 (SiMe_3), 16.0 (CH), 125.1, 127.6, 127.9, 128.5, 129.5, 130.1, 134.7, 134.9, 135.1, 135.4, 136.1, 139.7 (SiPh_3 , SPh); IR: 3070, 3050, 3014, 2954, 2926, 2854, 1726, 1632, 1582, 1479, 1428, 1249, 1188, 1167, 1109, 1024, 998, 854, 795, 738, 699 cm^{-1} ; $\text{C}_{28}\text{H}_{30}\text{SSi}_2$ (454.75): calcd. C 73.95, H 6.65, S 7.05; found C 73.96, H 6.83, S 7.03.

Bis(phenyldimethylsilyl)(phenylthio)methane (2c): A solution of 8.82 g (71 mmol) of thioanisol in dry THF (1 ml/mmol) was cooled to -78°C and one equiv. of *n*BuLi (1.6M in hexane) and one equiv. of TMEDA were added. The solution was stirred at 0°C for 2 h. At -78°C 1.1 equiv. of chlorophenyldimethylsilane was added. The reaction mixture was allowed to warm to room temperature and stirred for 12 h. After aqueous workup the resulting phenyldimethylsilyl phenylthiomethane was deprotonated and silylated again, repeating the general procedure described above. Column chromatography gave 15.11 g (54%) of **2c** as a colourless solid (Fp: 34°C). ^1H NMR: δ , -0.01 (s, 12H, SiMe_2Ph), 1.66 (s, 1H, CH), 6.81–7.32 (m, 15H, SiMe_2Ph , SPh); ^{13}C NMR: δ , -2.2 (SiMe_2Ph), 19.1 (CH), 125.2, 127.6, 128.4, 128.5, 129.1, 133.9, 138.5, 139.7 (SPh, SiMe_2Ph); IR: 3069, 3050, 2956, 1582, 1478, 1427, 1250, 1113, 991, 836, 733, 699 cm^{-1} ; $\text{C}_{23}\text{H}_{28}\text{SSi}_2$ (392.69): calcd. C 70.34, H 7.19, S 8.16; found C 70.15, H 7.22, S 8.12.

General procedure for the preparation of compounds 4,5 and 6a-c by the silicon-induced domino reaction: To a solution of 1.2 equiv. of silylated thioanisole **1**, **2a-f** in dry THF (4 ml/mmol) 1.1 equiv. of *n*-BuLi (1.6 M in hexane) was added at -78 °C. The solution was stirred at 0 °C for 2 h and cooled again to -78 °C. One equiv. of **3** in dry THF (4 ml/mmol) was added dropwise and the reaction mixture was allowed to warm to 0 °C and stirred for 12 h. The mixture was hydrolyzed with diethyl ether/ saturated NH₄Cl solution (1:1), the aqueous phase was extracted twice with diethyl ether, the combined organic layers were dried and the solvent evaporated. The crude products were purified by column chromatography (PE/EE 20:1→10:1).

1-Phenylthio-3-(trimethylsilyloxy)cyclopent-1-ene (4): 1.21 g (5 mmol) of **3** gave 187 mg (14%) as yellow oil. ¹H NMR: δ, 0.03 (s, 9H, SiMe₃), 1.84-2.00 and 2.84-3.13 (each *m*, 2H, CH₂), 5.08 (*dt*, *J*= 4.6, 7.4 Hz, 1H, 3-*H*), 6.56 (*d*, *J*= 7.4Hz, 1H, 2-*H*), 6.90-7.35 (*m*, 5H, *SPh*); ¹³C NMR: δ, -0.8 (SiMe₃), 29.6, 36.5 (C-4, C-5), 69.2 (C-3), 129.0, 129.1, 129.5 (*SPh*), 135.6, 137.2 (C-1, *SPh*), 153.6 (C-2); IR: 3420, 3060, 2940, 1710, 1580, 1480, 1440, 1250, 1020, 840, 740, 690 cm⁻¹; no elemental analysis could be obtained because of its decomposition character.

2-(2-Phenylthioethyl-2-trimethylsilyl)oxetane (5): 1.21 g (5 mmol) of **3** gave 538 mg (40%) of two diastereomers (1.5:1*) as colourless oil. ¹H NMR: δ, -0.05 (s, 9H*, SiMe₃), -0.02 (s, 9H, SiMe₃), 1.64 [*ddd*, *J*= 4.4, 11.4, 14.4 Hz, 1H, PhS(SiMe₃)CHCH₂], 1.84-2.20 and 2.21-2.50 (each *m*, 9H), 4.20-4.48 (*m*, 4H, 4-*H**, 4-*H*), 4.74-4.90 (*m*, 2H, 2-*H**, 2-*H*), 6.92-7.25 (*m*, 10H, *SPh**, *SPh*); ¹³C NMR: δ, -2.8*, -2.4 (SiMe₃), 27.6, 27.7* (C-3), 29.1, 29.5* (PhS(SiMe₃)CH), 39.9, 40.7* (PhS(SiMe₃)CHCH₂), 67.7*, 67.8 (C-4), 80.5*, 80.9 (C-2), 125.5, 125.7, 127.3, 128.6, 128.7, 129.0, 138.0, 138.6 (*SPh*, *SPh**); IR: 2950, 2880, 1580, 1480, 1250, 980, 840, 740, 690 cm⁻¹; C₁₄H₂₂OSSi (266.45): calcd. C 63.10, H 8.32, S 12.03; found C 62.69, H 8.08, S 11.89.

3-Phenylthio-3-(trimethylsilyl)cyclopentanol (6a): 242 mg (1 mmol) of **3** and **2b** as C₁ component gave 201 mg (76%) as a colourless oil. ¹H NMR: δ, 0.00 (s, 9H, SiMe₃), 1.63-2.11 (*m*, 6H, CH₂), 2.31 (*br s*, 1H, OH), 4.21 (*qui*, *J*= 4.4 Hz, 1H, 1-*H*), 7.20-7.65 (*m*, 5H, *SPh*); ¹³C NMR: δ, -3.0 (SiMe₃), 33.3, 35.9 (C-4, C-5), 43.7 (C-2), 44.3 (C-3), 74.9 (C-1), 128.7, 128.9, 133.1, 137.1 (*SPh*); IR: 3380, 2960, 1440, 1250, 1070, 840, 750, 690 cm⁻¹; C₁₄H₂₂OSSi (266.45): calcd. C 63.10 H 8.32 S 12.03; found C 62.67, H 8.34, S 11.92.

3-Phenyldimethylsilyl-3-(phenylthio)cyclopentanol (6b): 1.21 g (5 mmol) of **3** and **2c** as C₁ component gave 1.02 g (62%) as a colourless oil. ¹H NMR: δ, 0.12, 0.18 (each *s*, 3H, SiMe₂Ph); 1.33-1.48 (*m*, 2H), 1.62-2.06 (*m*, 5H), 3.76-3.89 (*m*, 1H, 1-*H*), 6.99-7.47 (*m*, 10H, *SPh*, SiMe₂Ph); ¹³C NMR: δ, -4.6, -4.5 (SiMe₂Ph), 33.6, 35.7 (C-4, C-5), 44.2 (C-2), 44.5 (C-3), 74.7 (C-1), 127.7, 128.7, 128.9, 129.3, 132.8, 134.5, 136.6, 137.0 (*SPh*, SiMe₂Ph); IR: 3396, 3069, 2956, 2861, 1473, 1427, 1409, 1250, 1069, 1111, 1026, 831, 818, 774, 749, 737, 702 cm⁻¹; C₁₉H₂₄OSSi (328.52): EI-HRMS: *m/z* = 328.1326 (Δ = -0.9 mmu).

3-Diphenylmethylsilyl-3-(phenylthio)cyclopentanol (6c): 1.21 g (5 mmol) of **3** and **2e** as C₁ component gave as second major fraction 537 mg (28%) as a colourless oil. ¹H NMR: δ, 0.53 (*s*, 3H, SiMePh₂), 1.57–1.71 (*m*, 1H, CH₂), 2.01 (*br s*, 1H, OH), 2.14–2.52 (*m*, 5H, CH₂), 3.98–4.12 (*m*, 1H, 1-*H*), 7.13–7.76 (*m*, 15H, SPh, SiMePh₂); ¹³C NMR: δ, -5.2 (SiMePh₂), 34.4, 35.5 (C-4, C-5), 44.0 (C-3), 44.8 (C-2), 74.6 (C-1), 127.7, 127.8, 128.5, 129.0, 129.3, 129.4, 132.3, 134.9, 135.1, 135.4, 135.5, 137.4 (SPh, SiMePh₂); IR: 3386, 3070, 3049, 2956, 2860, 1473, 1427, 1254, 1109, 1069, 1025, 789, 738, 726, 699 cm⁻¹; C₂₄H₂₆OSSi (390.58): calcd. C 73.80 H 6.71 S 8.21; found C 73.82, H 6.91, S 8.14.

General procedure for the pivaloylation of the alcohols 6a-c: One equiv. of the appropriate alcohol was dissolved in pyridine (2 ml/mmol) and the solution was cooled to -10 °C. 1.2 equiv. of pivaloyl chloride were added dropwise, the reaction mixture was allowed to warm to room temperature and stirred overnight. Pyridine was evaporated under reduced pressure using toluene as co-solvent. The resulting solid was dissolved in dichloromethane and was washed with saturated NaHCO₃ solution, water and brine. After drying, the solvent was evaporated and the obtained crude product was purified by column chromatography (PE/EE 20:1).

3-Pivaloyloxy-1-(phenylthio)-1-(trimethylsilyl)cyclopentane (7a): 828 mg (3.1 mmol) of **6a** gave 924 mg (85%) as a colourless oil. ¹H NMR: δ, -0.03 (*s*, 9H, SiMe₃), 1.05 (*s*, 9H, CMe₃), 1.43–1.64 (*m*, 1H, CH₂), 1.75–2.02 (*m*, 4H, CH₂), 2.36 (*dd*, *J* = 7.6, 15.0 Hz, 1H, 2-*H*), 4.90 (*m*, 1H, 3-*H*); ¹³C NMR: δ, -3.0 (SiMe₃), 27.0 (CMe₃), 31.8, 33.5 (C-4, C-5), 38.5 (CMe₃), 40.9 (C-2), 43.0 (C-1), 76.0 (C-3), 128.5, 128.6, 133.3, 137.2 (SPh), 178.4 (C=O); IR: 2960, 2905, 2871, 1725, 1479, 1439, 1284, 1250, 1160, 1035, 841, 750, 694 cm⁻¹; C₁₉H₃₀O₂SSi (350.56): calcd. C 65.09, H 8.63 S 9.15; found C 65.04, H 8.68, S 9.25.

3-Pivaloyloxy-1-(phenyldimethylsilyl)-1-(phenylthio)cyclopentane (7b): 562 mg (1.7 mmol) of **6b** gave 614 mg (87%) of a colourless oil as two diastereomers (8.1:1*). ¹H NMR: δ, 0.28, 0.29, 0.31*, 0.32* (each *s*, 3H, SiMe₂Ph), 0.99*, 1.02 (each *s*, 9H, CMe₃), 1.33–1.65 (*m*, 2H, 2H*, CH₂), 1.77–2.01 (*m*, 3H, 3H*, CH₂), 2.33 (*dd*, *J* = 6.8, 15.2 Hz, 1H, 2-*H**), 2.42 (*dd*, *J* = 7.6, 15.2 Hz, 1H, 2-*H*), 4.58–4.73 (*m*, 1H, 3-*H*), 4.86–4.99 (*m*, 1H, 3-*H**); ¹³C NMR (only one diastereomer could be detected): δ, -4.6 (SiMe₂Ph), 27.0 (CMe₃), 31.6, 33.9 (C-4, C-5), 38.0 (CMe₃), 41.2 (C-2), 43.4 (C-1), 75.9 (C-3), 127.8, 128.5, 128.6, 129.4, 133.1, 134.6, 136.6, 137.1 (SPh, SiMe₂Ph), 178.4 (C=O); IR: 3070, 2968, 2870, 1723, 1479, 1438, 1428, 1284, 1250, 1160, 1111, 832, 816, 775, 750, 737, 702 cm⁻¹; C₂₄H₃₂O₂SSi (412.62): calcd. C 69.86, H 7.82, S 7.77; found C 70.00, H 7.93, S 7.75.

3-Pivaloyloxy-1-(diphenylmethylsilyl)-1-(phenylthio)cyclopentane (7c): 80 mg (0.2 mmol) of **6c** gave 60 mg (62%) of a colourless oil as two diastereomers (10.3:1*). ¹H NMR: δ, 0.49, 0.56* (each *s*, 3H, SiMePh₂), 0.93*, 1.04 (each *s*, 9H, CMe₃), 1.35–1.66 (*m*, 2H, 2H*, CH₂), 1.93–2.10 (*m*, 3H, 3H*, CH₂), 2.49 (*dd*, *J* = 7.6, 15.6 Hz, 1H, 2-*H**), 2.62 (*dd*, *J* = 8.0, 15.6 Hz, 1H, 2-*H*), 4.55–4.70 (*m*, 1H, 3-*H*), 4.89–5.02 (*m*, 1H, 3-*H**); ¹³C NMR (only one diastereomer could be detected): δ, -5.2 (SiMePh₂), 26.9 (CMe₃), 31.4, 34.6 (C-4, C-5), 38.4 (CMe₃), 41.9 (C-2), 43.0 (C-1), 76.0 (C-3), 127.7, 128.4, 128.7, 129.4, 132.7, 134.9, 135.1, 135.4, 135.5, 137.5 (SPh, SiMePh₂), 178.3 (C=O); IR: 3070, 2968, 2870, 1723, 1585, 1477, 1429, 1396,

1367, 1285, 1256, 1159, 1108, 1033, 982, 789, 737, 699 cm^{-1} ; $\text{C}_{29}\text{H}_{34}\text{O}_2\text{SSi}$ (474.69): calcd. C 73.37, H 7.22, S 6.75; found C 73.66, H 7.51, S 6.74.

Acknowledgements

Support of this work by the Fonds der Chemischen Industrie, Frankfurt, and by the Deutsche Forschungsgemeinschaft (Scha 231/9-1) is gratefully acknowledged.

References

- [1] Fischer, M. R.; Kirschning, A.; Michel, T.; Schaumann, E.; *Angew. Chem.* **1994**, *106*, 220-221; *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 217-218.
- [2] Michel, T.; Kirschning, A.; Beier, C.; Bräuer, N.; Schaumann, E.; Adiwidjaja, A.; *Liebigs Ann.* **1996**, 1811-1821.
- [3] Jankowski, P.; Raubo, P.; Wicha, J.; *Synlett* **1994**, 985-992 and literature cited herein.
- [4] For use of halohydrines see Takeda, T.; Naito, S.; Ando, K.; Fujiwara, T.; *Bull. Chem. Soc. Jpn.* **1983**, *56*, 967-968.
- [5] Tietze, L. F.; Geissler, H.; Gewert, J. A.; Jakobi, U.; *Synlett* **1994**, 511-512.
- [6] Shinokubo, H.; Oshima, K.; Utimoto, K.; *Chem. Lett.* **1995**, 461-462.
- [7] Shinokubo, H.; Miura, K.; Oshima, K.; Utimoto, K.; *Tetrahedron* **1996**, *52*, 503-514.
- [8] Smith, A. B., III; Boldi, A. M.; *J. Am. Chem. Soc.* **1997**, *119*, 6925-6926.
- [9] For chemistry of mixed organosulfur-silicon compounds see Block, E.; Aslam, M.; *Tetrahedron* **1988**, *44*, 281-324.
- [10] Hackett, S.; Livinghouse, T.; *J. Chem. Soc., Chem. Comm.* **1986**, 75-76.
- [11] Hackett, S.; Livinghouse, T.; *J. Org. Chem.* **1986**, *51*, 879-885.
- [12] Gröbel, B.-T.; Seebach, D.; *Chem. Ber.* **1977**, *110*, 852-866.
- [13] For use of halohydrines see Coe, P. L.; Jones, A. S.; Kumar, A.; Walker, R. T.; *Tetrahedron Lett.* **1988**, *29*, 835-836.