ACS Medicinal Chemistry Letters

Letter

Subscriber access provided by DUESSELDORF LIBRARIES

Discovery of potent and selective inhibitors of Toxoplasma gondii thymidylate synthase for opportunistic infections

Nilesh Zaware, Hitesh Sharma, Jie Yang, Ravi K. V. Devambatla, Sherry F. Queener, Karen S. Anderson, and Aleem Gangjee

ACS Med. Chem. Lett., Just Accepted Manuscript • DOI: 10.1021/ml400208v • Publication Date (Web): 04 Oct 2013 Downloaded from http://pubs.acs.org on October 9, 2013

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Medicinal Chemistry Letters is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

7 8

9 10

11

12 13

14

15

16 17

18

19

20

21

26

27

28

29

30

31

32

33

34

35 36 37

38 39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59 60

Discovery of potent and selective inhibitors of *Toxoplasma* gondii thymidylate synthase for opportunistic infections.

Nilesh Zaware,^{a§}Hitesh Sharma,^{b§} Jie Yang,^a Ravi Kumar Vyas Devambatla,^a Sherry F. Queener,^c Karen S. Anderson^{b*} and Aleem Gangjee^{a*}

^aDivision of Medicinal Chemistry, Graduate School of Pharmaceutical Sciences, Duquesne University, 600 Forbes Avenue, Pittsburgh, PA 15282, United States

^bDepartment of Pharmacology, Yale University School of Medicine, New Haven, CT 06511, United States^cDepartment of Pharmacology and Toxicology, Indiana University School of Medicine, Indianapolis, IN 46202, United States.

§These authors contributed equally to this work.

KEYWORDS. Opportunistic infection; Toxoplasma gondii; Crystal structure; Thymidylate-synthase inhibitors; Activesite inhibitors.

ABSTRACT: Infection by the parasite *Toxoplasma gondii* (tg) can lead to toxoplasmosis in immunocompromised patients such as organ transplant, cancer and HIV/AIDS patients. The bifunctional thymidylate synthase-dihydrofolate reductase (TS-DHFR) enzyme is crucial for nucleotide synthesis in *T. gondii*, and represents a potential target to combat *T. gondii* infection. While species selectivity with drugs has been attained for DHFR, TS is much more conserved across species and specificity is significantly more challenging. We discovered novel substituted-*9H*-pyrimido[4,5-*b*]indoles **1-3** with single-digit nanomolar K_i for tgTS, two of which, **2** and **3**, are 28- and 122-fold selective over human TS (hTS). The synthesis of these compounds, and their structures in complex with tgTS-DHFR are presented along with binding measurements and cell culture data. These results show, for the very first time, that in spite of the high degree of conservation of active site residues between hTS and the parasite TS, specificity has been accomplished via novel structures and provides a new target (TS) for selective drug development against parasitic infections.

TEXT

According to the World Health Organization, the parasite Toxoplasma qondii (T. qondii, tg) has infected up to a third of the world's population.¹ Though the advent of antiretroviral therapy (ART) has decreased the incidence of T. gondii infection in AIDS patients, the infections still arise because of noncompliance with therapy, discontinuation of therapy because of toxicities with current treatments, and the emergence of drug-resistance strains. Late diagnosis of HIV and the rise of untreated HIV in developing countries also contributes to the persistence of T. gondii infections as a clinical issue.² T. gondii has been defined by the Centers for Disease Control and Prevention as an AIDS defining infection and a major cause of morbidity and indeed mortality in HIV patients as recently as 2009³ and 2013.⁴ Toxoplasma encephalitis (TE) is the principal opportunistic infection (OI) caused by T. gondii in AIDS patients. In addition to AIDS, other immune compromised patients such as those with organ transplants and those undergoing cancer chemotherapy are also highly susceptible to *T. gondii* infections.

Sulfadiazine (SDZ) with pyrimethamine (PM) (Fig. 1) is the principal drug combination for T. gondii infection along with folinic acid for rescue of host bone marrow toxicity from PM.² Alternate first line therapy with sulfamethoxazole (SMX) and trimethoprim (TMP) (Fig. 1) is also used.² Problems in chemoprophylaxis, treatment failures, high rates of adverse drug reactions and drug intolerance to these very similar first-line agents, particularly PM and the sulfonamide component,⁵ indicate an urgent need for new agents. The most clinically useful combinations, PM/SDZ (with folinic acid rescue) or TMP/SMX, act synergistically.⁶ The sulfonamide component inhibits dihydropteroate synthase (DHPS) which catalyzes an early step in the synthesis of folic acid. DHPS catalyzes the synthesis of *de novo* folate biosynthesis, a pathway absent in the human host. PM and TMP inhibit dihydrofolate reductase (DHFR). DHFR catalyzes the synthesis of tetra-

hydrofolate (THF) from folic acid as well as dihydrofolate (DHF). Thus the most successful drugs for T. qondii inhibit folate biosynthesis and metabolism at two separate steps to afford inhibition of purine and pyrimidine synthesis. Since Toxoplasma cannot salvage thymidine from the host,⁷ indirect inhibition of thymidylate synthase (TS) via DHFR inhibition is possible, but the direct inhibition of TS, the enzyme that catalyzes the conversion of dUMP to dTMP for DNA synthesis would be most logical. However, no selective inhibitors of T. gondii TS (tgTS) are known, up to now, that do not also potently inhibit human TS (hTS) as well.⁸ Unlike DHFR, where the human and parasite enzymes are somewhat divergent in amino acid sequence and structure allowing for selective inhibition of the parasite DHFR by 100-fold or greater, such divergence is much less in TS from parasites and humans.⁹ There is significant homology between the active site residues of tgTS and hTS. The significance of selective inhibition of tgTS is that this would inhibit the replication of T. gondii cells since T. gondii, unlike human, lacks salvage of thymidine thus TS function in T. gondii is obligatory for survival.¹⁰ Such selective inhibitors could be used alone or in combination with PM, TMP, SMX and other sulfonamides.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59 60 Based on the potent bicyclic hTS inhibitor nolatrexed reported by Webber et al.,¹¹ we designed tricyclic compounds 1-3 using as a novel scaffold for hTS inhibition and as potential anticancer agents. Compounds 1-3 (Figure 1) contain a tricyclic pyrimido[4,5-*b*]indole scaffold connected to a phenyl or a naphthalene moiety via a sulfur bridge. Compounds 1-3 were synthesized using Scheme 1. Compound 4 was prepared by a method developed by Gangjee et al.¹² Ullmann coupling¹³ of 4 with aryl thiols 5-7 in a microwave apparatus (Initiator[®] from Biotage) in the presence of potassium carbonate provided target compounds 1-3, respectively, in yields of 28-40%.

Inhibition assays were performed on isolated *T. gondii* TS-DHFR and hTS. Compounds **1-3** exhibited single-digit nanomolar K_i values (Table 1). Surprisingly compounds **2** and **3** showed unprecedented selectivity for tgTS over hTS of **28-** and **122-**fold respectively.

To determine the structural basis for the observed activity and selectivity, we determined the crystal structures of inhibitors 2 and 3 with T. gondii TS-DHFR (Figure 2). By superimposing the tgTS and hTS (Figure S₃), almost all of the residues in the active site were identical. It was surprising that a remarkable difference of 28- and 122-fold in the K_i values of 2 and 3 respectively were observed in spite of almost identical architectures of the active sites of TS of the two organisms. As shown in Fig. 2, the crystal structure of the tgTS-DHFR with 2 and 3 revealed extensive base stacking interactions between the inhibitor and the nucleotide dUMP as expected. Using co-crystal structures of 2 and 3 with tgTS-DHFR, we generated ligand interactions using MOE 2011.10.14 Compound 2 is bound to the active site by aromatic stacking of the pyrimido [4,5b]indole scaffold and the dUMP pyrimidine ring. Additionally, hydrogen bonds are formed between the N9 of **2** and carboxyl of Asn4o6, the Asp513 carboxylic oxygen with N3 and the 2-NH₂ group. The 2-NH₂ group forms additional hydrogen bond with the backbone oxygen of Ala6o9. The scaffold is also stabilized by hydrophobic interactions of the C-ring with Trp4o3. The phenyl ring is oriented almost at a right angle to the tricyclic scaffold and interacts with Ile4o2, Leu516, Phe52o and Met6o8. Compound **3** retains the interactions of the tricyclic scaffold and the hydrogen bonds with Asp513, Ala6o9 and Asn4o9. Similar to **2**, the 12-naphthyl ring of **3** is oriented almost at right angle to the tricyclic scaffold and interacts with Ile4o2, Leu516, Phe520 and Met6o8.

The structure of **3** in the TS active site suggested that the I^D-naphthalene moiety of **3** must be a key determinant of higher selectivity. The position of the compound in the active site orients the I^D-naphthalene moiety close to the C-terminal tail of the *T. gondii* TS domain. In fact, the residues of the C-terminal tail are well ordered only in *T. gondii* and not in the known X-ray crystal structures of human TS. The I^D-naphthalene substituent of **3** also makes hydrophobic interactions with M6o8 only in tgTS, since the C-terminal tail shields the active site from solvent, the tail could be a possible structural determinant that facilitates additional hydrophobic interactions with tgTS conferring selectivity for the parasite (Fig. 2).

Compounds 1-3 have also been evaluated against *T. gondii* cells in culture (Table 2) and 3, at 5 μ M, was found to be equivalent to PM at 3 μ M indicating a significant effect on *T. gondii* in culture comparable to a clinically used agent PM. The culture model requires that the experimental compounds rapidly kill exposed tachyzoites as they are released from cells and/or that the experimental compounds penetrate the cell and the vacuole containing tachyzoites where the compounds may kill the organism or prevent its replication.

We have discovered, for the first time, highly potent (single-digit nM) tgTS inhibitors 1-3 with up to a remarkable 122-fold selectivity for tgTS over hTS (K_i in Table 1). This level of selectivity is unprecedented for any parasite with regard to TS. For comparison, TMP selectively inhibits tgDHFR and is used extensively with SMX to treat Toxoplasma infection, but has an IC₅₀ = 2.7μ M and a selectivity of only 49-fold for tgDHFR over hDHFR.¹⁵ Further PM, a widely used gold standard tgDHFR inhibitor for TE and other *T. gondii* infections, only has a potency of IC₅₀ = 3.9 μ M and a selectivity of only 5.9 for tgDHFR over mammalian DHFR. In addition, we have also, for the very first time, solved the X-ray crystal structure of the bifunctional tgTS-DHFR enzyme with compounds 2 and 3.¹⁶ In cell cuture study, compound 3 showed similar potency to PM, a highly effective standard drug used as a control in these studies. We have thus demonstrated with 2 and 3 that structure variation affords increased selectivity while maintaining potency (compared to 1). Supplementary Figure 3 shows dUMP and compound 3 and the superposition of key active site residues (using tg numbering) in both human and tgTS. While all residues are identical between the two species, only M608 is present in the tgTS structure. Future studies will be directed toward solving the structure of compound 3 with human TS to more fully discern possible differences. TS is a completely new and novel target for *T. gondii*. There is no report, to our knowledge, of selective and potent inhibitors of tgTS as anti-OI agents. The ability to provide agents that are selective for tgTS on the basis of structure-based rational design is a major step towards providing novel clinically useful agents with tgTS inhibition.

FIGURES



Figure 1. DHFR inhibitors (PM and TMP) in the clinic for *T. gondii* infections and compounds **1-3**.



Figure 2. *T. gondii* TS active site with key residues highlighted in red. The inhibitors **2** and **3** and dUMP are colored yellow.

SCHEMES





TABLES

Table 1. Kinetic Evaluation o	of 1-3 Reveals Species Selectiv-
ity ^a	

Cmpd	IC ₅₀ (nM)		K _i (nM)		Selectivi- ty (hTS
	tgTS	hTS	tgTS	hTS	K _i / tgTS K _i)
1	21 ± 13	28.8 ± 21	1.8 ± 1.1	3.3 ± 2.4	1.83
2	23 ± 6.6	481 ± 162	2.0 ± 0.6	55.7 ± 18.8	27.9
3	35.8 ± 2.9	3250 ± 184	3.1 ± 0.25	378 ± 21	121.9

^{*a*}*T. gondii* TS-DHFR (25 nM) and human TS (50 nM) were preincubated with 100 μ M dUMP and inhibitor. The reaction was initiated with 100 μ M methyleneTHF. ^{*b*} Due to the concentrations of enzyme required for assay and the limits of detection, for those compounds having K_i values in the lower nanomolar range, the values reported represent an upper limit.

Table 2. Toxoplasma Cell Culture Study

Condition	<i>T. gondii</i> in media at Day 4	<i>T. gondii</i> in cells at Day 4	Estimated IC₅₀ value, µMª
No drug	3710000	1140000	
1 (5 μM)	2350000	340000	2.6
2 (5 μM)	100000	140000	3.7
3 (5 μM)	<25000	<2500	0.6
PM (3 μM)	<25000	<2500	0.57

^{*a*}Dose response curves using data from the three concentrations of test compounds gave estimates of IC_{50} values.

ACKNOWLEDGMENTS

This work was supported, in part, by NIH grants CA152316 (to A.G.), the Duquesne University Adrian Van Kaam Chair in Scholarly Excellence (A. G.) and AI083146 (to K.S.A.).

ASSOCIATED CONTENT

Supporting Information. Synthetic procedures, compound characterization, biological assays and X-ray structure determination. This material is available free of charge via the Internet at http://pubs.acs.org.

Accession Codes

The atomic coordinates of *T.gondii* TS-DHFR/Compound **2** and *T.gondii* TS-DHFR/Compound **3** described here have been deposited in the RCSB Protein Data Bank (www.pdb.org) with accession codes 4KY4 and 4KYA respectively.

AUTHOR INFORMATION

Corresponding Authors

*These authors contributed equally to this work: (A.G.) Tel.: +1-412-396-6070, fax: +1-412-396-5593, e-mail:

gangjee@duq.edu; (K.S.A.) Tel.: +1-203-785-4526, fax: +1-203-785-7670, e-mail: <u>karen.anderson@yale.edu</u>.

Notes

The authors declare no competing financial interest.

ABBREVIATIONS

tg, *T. gondii*; TS, thymidylate synthase; DHFR, dihydrofolate reductase; TE, toxoplasma encephalitis; OI, opportunistic infection; SDZ, sulfadiazine; PM, pyrimethamine; SMX, sulfamethoxazole; TMP, trimethoprim.

REFERENCES

- 1. Montoya, J. G.; Liesenfeld, O. Toxoplasmosis. *Lancet* 2004, 363, 1965–1976.
- Kaplan, J. E.; Benson, C.; Holmes, K. K.; Brooks, J. T.; Pau, A.; Masur, H. Guidelines for Prevention and Treatment of Opportunistic Infections in HIV-Infected Adults and Adolescents. Recommendations from CDC, The National Institutes of Health and the HIV Medicine Association of the Infectious Diseases Society of America. CDC, *MMWR*, April 10, 2009, *Vol* 58, NoRR-4.
- 3. CDC. <u>http://www.cdc.gov/parasites/toxoplasmosis/</u>. May 2013.
- 4. Dudrand, C. M.; Flexner, C. HIV Cure: Knocking on the Door. *Clin. Pharmacol. Ther.* **2013**, *93*, 382–384.
- Aspinall, T. V.; Joynson, D. H.; Guy, E.; Hyde, J. E.; Sims, P. F. The Molecular Basis of Sulfonamide Resistance in *Toxoplasma gondii* and Implications for the Clinical Management of Toxoplasmosis. *J. Infect. Dis.* 2002, *185*, 1637–1643.
- 6. Chulay, J. D.; Watkins, W. M.; Sixsmith, D. G. Synergistic Antimalarial Activity of Pyrimethamine and Sulfadoxine

against Plasmodium falciparum in vitro. Am. J. Trop. Med. Hyg. **1984**, 33, 325–330.

- 7. Hyde, J. E. Targeting Purine and Pyrimidine Metabolism in Human Apicomplexan Parasites. *Curr. Drug Targets* 2007, 8, 31-47.
- Pang, C. K.; De, S. K.; White, J.; Buckner, F. S.; Varani, G.; Rathod, P. K. Differential Drug Binding by the Highly Conserved *Plasmodium falciparum* Thymidylate Synthase. *Mol. Biochem. Parasitol.* 2005, *143*, 121–124.
- Ferrari, S.; Lasasso, V.; Costi, M. P. Sequence-Based Identification of Specific Drug Target Regions in the Thymidylate Synthase Enzyme Family. *Chem. Med. Chem.* 2008, 3, 392– 401.
- Fox, B. A.; Bzik, D. J. De novo Pyrimidine Biosynthesis is Required for Virulence of *Toxoplasma gondii*. *Nature* 2002, 415, 926–929.
- Webber, S. E.; Bleckman, T. M.; Attard, J.; Deal, J. G.; Kathardekar, V.; Welsh, K. M.; Webber, S.; Janson, C. A.; Matthews, D. A.; Smith, W. W.; Freer, S. T.; Jordan, S. R.; Bacquet, R. J.; Howland, E. F.; Booth, C. L. J.; Ward, R. W.; Hermann, S. M.; White, J.; Morse, C. A.; Hilliard, J. A.; Bartlett, C. A. Design of thymidylate synthase inhibitors using protein crystal structures: The synthesis and biological evaluation of a novel class of 5-substituted quinazolinones. *J. Med. Chem.* **1993**, *36*, 733–746.
- 12. Gangjee, A.; Zaware, N.; Raghavan, S.; Ihnat, M.; Shenoy, S.; Kisluik, R. L. Single Agents with Designed Combination Chemotherapy Potential: Synthesis and Evaluation of Substituted Pyrimido[4,5-b]indoles as Receptor Tyrosine Kinase and Thymidylate Synthase Inhibitors and as Antitumor Agents. J. Med. Chem. 2010, 53, 1563–1578.
- Sperotto, E.; van Klink, G. P. M.; de Vries, J. G.; van Koten, G. Ligand-Free Copper-Catalyzed C-S Coupling of Aryl Iodides and Thiols. J. Org. Chem. 2008, 73, 5625–5628.
- 14. Molecular Operating Environment (MOE 2011.10); Chemical Computing Group, Inc.: Montreal, Quebec, Canada, 2008; www.chemcomp.com.
- Piper, J. R.; Johnson, C. A.; Krauth, C. A.; Carter, R. L.; Hosmer, C. A.; Queener, S. F.; Borotz, S. E.; Pfefferkorn, E. R. Lipophilic Antifolates as Agents against Opportunistic Infections. 1. Agents Superior to Trimetrexate and Piritrexim against *Toxoplasma gondii* and *Pneumocystis carinii* in *in vitro* Evaluations. *J. Med. Chem.* 1996, 39, 1271–1280.
- 16. Sharma, H.; Landau, M. J.; Vargo, M. A.; Spasov, K. A.; Anderson, K. S. First Three-Dimensional Structure of *Toxoplasma gondii* Thymidylate Synthase-Dihydrofolate Reductase: Insights for Catalysis, Interdomain Interactions, and Substrate Channeling. *Biochemistry* 2013, *Submitted*.

4

60

HN H2N 3 H 1. gondii TS K _i = 3.1 nM human TS K _i = 378 nM	F520 L516 U512 V403 Y555 M608 N406 L486	Compound 3	