Benzoxazolamines and Benzothiazolamines: Potent, Enantioselective Inhibitors of Leukotriene Biosynthesis with a Novel Mechanism of Action

Edward S. Lazer,*,† Clara K. Miao,† Hin-Chor Wong,† Ronald Sorcek,† Denice M. Spero,† Alex Gilman,† Kollol Pal,† Mark Behnke,[†] Anne G. Graham,[‡] Jane M. Watrous,[‡] Carol A. Homon,[‡] Juergen Nagel,^{§,||} Arvind Shah,[§] Yvan Guindon,[⊥] Peter R. Farina,[‡] and Julian Adams[†]

Departments of Medicinal Chemistry, Biochemistry, and Analytical Chemistry, Boehringer Ingelheim Pharmaceuticals, Incorporated, 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877

Received December 10, 1993®

A series of benzoxazolamine and benzothiazolamine analogs that inhibit leukotriene (LT) biosynthesis are described. The initial lead, (S)-N-(benzothiazol-2-yl)phenylalanine ethyl ester (5a), was discovered in a screening program for inhibition of Ca-ionophore-A23187-induced LTB₄ release in human polymorphonuclear leukocytes (IC₅₀ $0.23 \,\mu$ M). Through structural modification, it was determined that hydrophobic substituents in the 5-position and replacement of the phenyl ring of phenylalanine with a cyclohexyl group greatly enhance potency. Several ester bioisosteres that retain potency and enantiomeric selectivity are described. Lead optimization culminated in (S)-N-[2-cyclohexyl-1-(2-pyridinyl)ethyl]-5-methyl-2-benzoxazolamine (43b), IC₅₀ 0.001 μ M. The compounds described are not inhibitors of 5-lipoxygenase but, rather, act at the level of arachidonic acid release.

Since the late 1970's, extensive research has been devoted to the discovery of compounds that block the biosynthesis or activity of leukotrienes (LTs). LTs are produced by the metabolism of arachidonic acid by 5-lipoxygenase (5-LO). Arachidonic acid is derived from membrane phospholipids upon hydrolysis by phospholipase A_2 (PLA₂). The now well-known biochemical pathway from arachidonic acid to LTs has been reviewed in the literature.^{1,2} Reviews describing the status of research on inhibitors of leukotriene biosynthesis^{3,4} and peptide leukotriene antagonists⁵ have also appeared recently.

Leukotriene B_4 (LTB₄) is a potent chemotactic agent for leukocytes and stimulates a number of proinflammatory responses.⁶ The peptide leukotrienes LTC_4 , LTD_4 , and LTE₄ are potent bronchoconstrictors of human lung in vitro⁷ and in vivo.⁸ The peptide LTs also promote mucus secretion⁹ and airway hyperresponsiveness in normal¹⁰ and asthmatic¹¹ subjects. These properties and the fact that LTs have been found in diseased tissues implicate LTs as major contributors in a number of diseases including asthma, rheumatoid arthritis, psoriasis, and inflammatory bowel disease.^{12,13} Several inhibitors of LT biosynthesis³ as well as LTD₄ receptor antagonists⁵ have been in clinical trials, and recent results are encouraging, especially in asthma.14-16

Inhibitors of LT biosynthesis can be generally divided into four classes. The majority are 5-LO inhibitors that inactivate 5-LO via their antioxidant properties, possibly by keeping the iron associated with the active site in the inactive reduced state. BI-L-239 (1)¹⁷ is a representative of this class. N-Hydroxyurea derivatives are exemplified by zileuton (A-64077) (2). The design of this class was originally based on the ability of the hydroxamate functionality to bind to the active-site ferric ion.⁴ Recent studies, however, indicate that representatives of this group, including 2, can reduce 5-LO as measured by the ability to enhance a pseudoperoxidase reaction,¹⁸ raising the possibility that antioxidant activity may also contribute to the mechanism of this class.

A series of enantioselective 5-LO inhibitors that possess neither redox nor iron-chelating properties have been described¹⁹ and have led to development candidate ICI D2138 (3). Finally, MK-886 (4)²⁰ was the first compound reported to block the activation of 5-LO by an 18-kDa membrane protein termed 5-lipoxygenase-activating protein (FLAP).



In this paper, we describe a series of benzoxazoles and benzothiazoles that do not act by any of the mechanisms described above but, rather, inhibit LT biosynthesis by affecting the availability of arachidonic acid.

In Vitro Optimization

This series evolved from a hit in a screening program to detect inhibitors of Ca-ionophore-induced LTB₄ biosynthesis in human neutrophils. Compound 5a was found to have an IC₅₀ of 0.23 μ M in this assay. For comparison, the antioxidant 5-LO inhibitor 1 has an IC₅₀ of 0.34 μ M in this assay. Compound 5a did not inhibit other calciummediated events (oxidative burst or enzyme release) and was not structurally related to known 5-LO inhibitors.

© 1994 American Chemical Society

[†] Department of Medicinal Chemistry.

[‡] Department of Biochemistry.

Department of Analytical Chemistry.

¹ Bio-Mega/Boehringer Ingelheim Research Inc., 2100 Rue Cunard, Laval, Quebec, Canada. | Present address: Boehringer Ingelheim KG, D6507 Ingelheim am

Rhein, Germany.
Abstract published in Advance ACS Abstracts, February 15, 1994.



compd no.	R/S	x	R	n	syn meth	mp (°C)	formulaª	IC ₅₀ ^b (µM) or % inh of LTB₄/conc
5a	S	NH	\mathbf{Et}	1	A	oil	$C_{18}H_{18}N_2O_2S$	0.23
5b	RS	NH	Et	1	В	137-139	$C_{18}H_{18}N_2O_2S$	0.36 (0.32–0.41)
6	RS	NH	Et	2	В	69.5-72	$C_{19}H_{20}N_2O_2S$	0.52 (0.38-0.67)
7	RS	NH	\mathbf{Et}	0	В	123 - 125	$C_{17}H_{16}N_2O_2S$	61%/3 µM
8	RS	NCH3	Et	1	В	oil	$C_{19}H_{20}N_2O_2S$	1.0 (0.62-1.4)
9	RS	CH_2NH	\mathbf{Et}	1	с	150-153	C ₁₉ H ₁₉ ClN ₂ O ₂ S·HCl	$55\%/3\mu M$
10	RS	CH_2	\mathbf{Et}	1	с	46-48	$C_{19}H_{19}NO_2S$	$56\%/1\mu M$
11	RS	NH	н	1	В	250-251	C ₁₆ H ₁₄ N ₂ O ₂ S·HCl ^d	$-19\%/3~\mu M$

^a All compounds gave satisfactory elemental analyses ($\pm 0.4\%$) for C, H, and N, unless otherwise noted. ^b IC₅₀s (the concentration required to inhibit the maximum response by 50%) were determined from $n \ge 3$ experiments by nonlinear regression analysis with 95% confidence bounds given in parentheses (SAS Software System, SAS Institute, Inc., Cary, NC). A Hill-type equation (the sigmoid E_{max} model) served as the model for analysis.^{27,28} Percent inhibition is the average of duplicates from n = 1 experiment. ^c See Experimental Section. ^d C: calcd, 57.39; found 57.97.

We therefore embarked on a program to optimize for LTB₄ biosynthesis inhibition by structure modification. The racemate **5b** was prepared and found to be less potent (IC₅₀ 0.36 μ M), giving us our first indication of enantiomeric selectivity (Table 1). The homophenylalanine analog **6** is slightly less active (IC₅₀ 0.52 μ M), while the phenylglycine derivative **7** is much less potent. Substitution on or replacement of the 2-amino group led to a reduction in activity (**8-10**). Compound **11** is representative of several carboxylic acid analogs of active esters that we prepared. All of the carboxylic acid derivatives were totally inactive, indicating that the ester is the active species and a metabolically stable bioisosteric replacement was needed.

Halogen substitution on the aromatic ring of the phenylalanine ester gave modest improvements in potency (12-14, Table 2). An alkyl or methoxy substituent in the 6-position of the benzothiazole had a minor effect on potency (15-17). We found that replacement of the benzothiazole with a benzoxazole generally improved activity (for example, compare 18a and 5a). We therefore switched to the benzoxazole nucleus for the majority of subsequent analogs.

The two most significant breakthroughs in in vitro potency are illustrated by compounds 19-23a. A comparison of 20 and 22 with 19 and 21 illustrates the striking enhancement in potency of an alkyl substituent in the 5-position, while in the 6-position there is relatively little effect in both a benzoxazole and a benzothiazole. Comparison of 23a with 18a shows that replacement of the R₂ phenyl with a cyclohexyl group also results in a dramatic increase in potency.

As mentioned above, we recognized the need to replace the ester with a metabolically stable group. Compounds 24-28 illustrate that the ester at R_3 could be replaced by a phenyl group and that potency can still be enhanced with a cyclohexyl group at R_2 and a substituent (in this case, Cl or OMe) in the 5-position at R_1 . Comparison of 26a-c shows that enantiomeric selectivity is retained. Compounds 29-33 show that various substituents in the 2-, 3-, or 4-position of the phenyl group at R_3 all reduce potency compared to the unsubstituted 26a, probably indicating steric limitations at that position. Several other functional groups or rings could be used to replace the ester and maintain activity, especially if in combination with a 5-substituent at R_1 or a cyclohexyl ring at R_2 . This is illustrated with amides (34 and 35), ethers (36-38), a thioether (39), and a ketone (40). Compounds 36-38 clearly show that increasing the size of the ether group reduces potency, again consistent with steric limitations. The 3-methyl-5-oxadiazole group, which has been reported as an ester bioisostere for antimuscarinic compounds,²¹ retained potent activity with a cyclohexyl at R_2 (41) and moderate activity with a 4-fluorophenyl at R_2 (42).

Compounds 43-45 illustrate the difference in potency of the pyridine isomers at R_3 . The racemic 2-pyridyl isomer 43a was 5 times more potent than the 3-isomer (44) which was more potent that the 4-isomer (45). We found similar potency ratios regardless of the substituents at R_1 and R_2 .

The effect on potency of modifications at R_1 is illustrated by compounds 46-63. Adding an isopropyl on the 5-position to compound 26a results in 46 and only a slight increase in potency. We believe this reflects a limit in hydrophobicity in the molecule beyond which no substantial increase in potency can be demonstrated. Among other racemic compounds with a cyclohexyl at R_2 and a phenyl at R_3 , 51 ($R_1 = 5$ -cyano) and 47 ($R_1 =$ ethoxycarbonyl) were the most potent followed by 53 ($R_1 = 5$ -acetamido). The 5-sulfonamide 54, 5-dimethylamide 49, and 5-carboxylic acid 48 had similar potencies with IC₅₀s in the 150-170 nM range. The 5-hydroxymethyl 52 and 5-carboxamide 50 were least potent.

The potencies of additional R_1 substituents are illustrated by a second set of compounds having a 4-fluorophenyl group at R_2 and a 2-pyridyl ring at R_3 (55–63). A number of substituents in the 5-position (CF₃, OMe, iPr, NO₂, and Cl) resulted in compounds with IC₅₀s of 20 nM or less.

Additional modifications at R_2 are illustrated by compounds 64-73. The 3-chlorophenyl substituent in 65 provides a more potent compound than the 4-chlorophenyl (64) or 4-fluorophenyl (57) groups. Compounds 66 and 67 show that a 4-methoxy substituent on the phenyl ring retains good activity but the 4-hydroxy substituent

Benzoxazolamines and Benzothiazolamines

drastically reduces potency. The effects of various heterocyclic substituents are illustrated with compounds 68-76. A piperidine (68a), morpholine (69), and thiomorpholine (70) each provide reasonable activity, whereas the thiomorpholine sulfone 71 and N-methylpiperazine 72 are inactive. A 2-methyl-4-thiazolyl group is active with either a 2-pyridiyl (73) or an ester (74) at R_3 . Histidine derivatives 75 and 76 are relatively inactive. Overall, these results suggest that an ionizable or hydrophilic group in the distal region of a group at R_2 has a deleterious effect on potency.

Stereoselectivity for LTB₄ biosynthesis inhibition is illustrated by several pairs of enantiomers in Table 2. Examples are given for compounds with a variety of substituents at R_2 and R_3 . Esters (18a,b and 23a,b) and phenyl (26b,c) and pyridyl bioisosteres at R_3 (43b,c and 59b,c) show selectivities of between 10- and 57-fold. When absolute configuration was determined or known, the more active enantiomer corresponded to the L-amino acid or the S-configuration in most cases. In compounds 68-72, this corresponds to the R-configuration. These results suggest that there is an enantioselective interaction of these compounds with a molecular target.

Compound 43b (BIRM 270) has been selected from this group for more extensive evaluation and is presently under development as a topically administered treatment for asthma. In a sheep model of allergen-induced bronchoconstriction and hyperresponsiveness,²² 43b blocked the late-phase response and the airway hyperresponsiveness when given by aerosol administration at doses of 1, 3, or 10 mg. These results will be published separately.

Chemistry

The preparation of benzothiazoles is illustrated in Scheme 1. Reaction of a phenyl isothiocyanate and the appropriately substituted amine or amino acid ester gave a thiourea, which was cyclized using SO_2Cl_2 (method A). Alternatively, amino acids could be alkylated with 2-chlorobenzothiazole using NaOH in DMSO and then esterified (method B). The 5-substituted compound 22 was prepared from 3-ethylphenyl isothiocyanate and required separation from the 7-isomer.

Benzoxazoles were prepared either by alkylation of the appropriately substituted 2-chlorobenzoxazole with an amine or an amino acid as described in method B above or by using diisopropylethylamine with an amine or an amino acid ester (method C, Scheme 2). The substituted 2-chlorobenzoxazoles were prepared by chlorination of the corresponding 2-mercaptobenzoxazole. These were prepared by reaction of the appropriate aminophenol with CS_2 .²³ If not available commercially, the aminophenols were prepared by nitration of the substituted phenol followed by reduction of the nitro group.²³

1,2-Diarylethylamines and 1-aryl-2-cyclohexylethylamines were prepared as described in Scheme 3. The Grignard reagent was prepared from the benzyl- or cyclohexylmethyl bromide and allowed to react with the appropriate arylnitrile. In situ reduction of the intermediate imine with NaBH₄ gave the amine (method D). Alternatively, reaction of the Grignard reagent with an aldehyde gave an alcohol, which was oxidized to a ketone followed by reductive amination (method E). The ketone could also be prepared by Friedel-Crafts acylation, where applicable.

Scheme 4 outlines the synthesis of a number of compounds containing R_3 modifications that were derived

from carboxylic acids or esters. Amides 34 and 35 were prepared by coupling of an amine with the carboxylic acid using CDI in DMF. Ketone 40 was prepared by reaction of the carboxylic acid with methyllithium. Methyl oxadiazole derivatives 41 and 42 were prepared from the corresponding esters with acetamide oxime and NaH as described in the literature.²¹ Ethers 36–38 were synthesized by alkylation of the corresponding alcohol. Thioether 39 was prepared by reaction of the corresponding mesylate with sodium thiomethoxide. In each case, the ether or thioether was introduced onto the protected amine prior to the coupling step. Reduction of the ester followed by reaction with MsCl gave the mesylate.

Several compounds were prepared by further transformation after the coupling step as illustrated in Scheme 5. Hydrolysis of ester 47 gave 48, which was converted to amides 49 and 50. Treatment of amide 50 with trifluoroacetic anhydride gave nitrile 51. Reduction of ester 47 gave alcohol 52. Acetamide 53 and sulfonamide 54 were prepared from the amine, which was obtained by reduction of the 5-nitro compound.

The synthesis of 68a-72 is illustrated in Scheme 6. Method F outlines the preparation of single enantiomers, starting with (R)-phenylglycinol. Protection of the amine as the tBOC derivative is followed by formation of the mesylate. Displacement with the appropriate heterocycle followed by deprotection gave the desired amine which upon reaction with 2-chloro-5-isopropylbenzoxazole gave 68-70 and 72. Racemic 71 was prepared (method G) by oxidation of morpholine followed by reaction with 2-chloroacetophenone. Reductive amination of the resulting ketone gave the desired amine, which was converted to 71.

Chiral compounds were prepared by either of three methods. Amino acid derivatives **5a**, and **18a,b** were prepared from the commercially available chiral phenylalanine ester. Catalytic reduction of the phenylalanine ester with Rh/Al_2O_3 gave the cyclohexylalanine esters used for **23a,b**. Enantiomers of **26a** and **59a** were prepared by separation of the racemates using chiral HPLC. Enantiomers of **43a** were prepared by chromatographic separation of diastereomeric amides obtained from the precursor racemic amine and L-valine as described in a separate publication.²⁴

Mechanism

Compound 43b and others from this series had little to no activity in a calcium-ionophore-A23187-stimulated assay for 5-LO inhibition in which exogenous [¹⁴C]arachidonic acid is provided to the neutrophils and radiolabeled 5-LO metabolites are measured. This procedure,²⁵ unlike the assay used to characterize the compounds presented here, bypasses PLA₂ and is therefore a measurement of 5-LO activity. The relative lack of activity in this system initially indicated that our compounds were not acting directly on 5-LO. This was later confirmed by a similar lack of activity against 5-LO from RBL-1 and neutrophil cytosolic preparations.

We found that the inhibition of LTB_4 biosynthesis by 43b could be overcome by addition of exogenous arachidonic acid. This suggested to us that the mechanism of action could involve limiting arachidonic acid availability. In fact, direct measurement of arachidonic acid and platelet-activating factor (PAF) released from calciumionophore-stimulated neutrophils showed that 43b in-

Table 2. Inhibition of Leukotriene Biosynthesis by Benzoxazole and Benzothiazole Derivatives



compd	חימ	37				syn			$IC_{50}^{b}(\mu M)$ or
<u> </u>	R/S RS	Y S	<u>к</u> 1 ч	4-ClCeH4	R ₃	B		CasHarClNaOaS	$\frac{1}{20}$ (0.14–0.27)
13	RS	š	н	$4-BrC_6H_4$	CO_2Et	B	104-106	$C_{18}H_{17}BrN_2O_2S$	0.15 (0.12-0.19)
14	RS	S	H	4-FC ₆ H ₄	CO ₂ Et	В	129-131	$C_{18}H_{17}FN_2O_2S$	0.33 (0.22-0.44)
15 16	RS RS	S	6-127 6-nBu	4-CIC ₆ H ₄ 4-CIC ₆ H ₄	CO ₂ Et	A	100-107	$C_{21}H_{23}CIN_2O_2S$ $C_{00}H_{01}CIN_2O_2S$	0.33(0.29-0.37) 0.18(0.14-0.21)
17	RS	š	6-OMe	4-ClC ₆ H ₄	CO_2Et	Ä	129-131	$C_{19}H_{19}ClN_2O_3S$	0.23 (0.16-0.30)
18a	S	0	н	C ₆ H₅	CO_2Et	В	oil	$C_{18}H_{18}N_2O_{3^{*1}/2}H_2O$	0.15 (0.13-0.17)
18D 19	R S	0	H 6-iPr	C ₆ H ₅ C ₆ H ₅	CO ₂ Et CO ₂ Et	B	oil	$C_{18}H_{18}N_2O_3$ $C_{02}H_{04}N_2O_3$	1.5 (0.63-2.3) 0.22 (0.15-0.28)
20	\tilde{s}	ŏ	5-iPr	C ₆ H ₅	CO_2Et	č	oil	$C_{21}H_{24}N_2O_3$	0.0005 (0.0004-0.0007)
21	S	S	6-Et	C ₆ H ₅	CO ₂ Et	A	102-104	$C_{20}H_{22}N_2O_2S$	0.24 (0.19-0.29)
22 23a	S	8	5-Et H	C ₆ H ₅ C ₆ H ₁₁	CO ₂ Et CO ₂ Me	A C	oil	$C_{20}H_{22}N_2O_2S^c$ $C_{17}H_{22}N_2O_2$	0.012 (0.0091-0.016)
23b	\tilde{R}	ŏ	н	C_6H_{11}	CO ₂ Me	č	oil	$C_{17}H_{22}N_2O_3^d$	0.10 (0.078-0.13)
24	RS	S	Н	C ₆ H ₅	C ₆ H ₅	B	54-56	$C_{21}H_{18}N_2S$	0.17 (0.15-0.19)
25 26a	RS	0	ਸ ਸ	C ₆ H ₅ C ₂ H ₁₁	C ₆ H ₅ C ₂ H ₂	В	154-156 131-139	$C_{21}H_{18}N_2O$ $C_{21}H_{21}N_2O$	0.051 (0.029-0.073)
26b	-	ŏ	Ĥ	C_6H_{11}	C_6H_5	e	oil	$C_{21}H_{24}N_2O$	0.006 (0.004-0.007)
26c	+	0	H	$C_{6}H_{11}$	C ₆ H ₅	e	oil	$C_{21}H_{24}N_2O$	0.34 (0.13-0.55)
27	RS	0	5-Cl 5-OMe	C ₆ H ₅	C_6H_5	C	149-151 148-150	$C_{21}H_{17}CIN_2O$	0.023 (0.012 - 0.034)
29	RS	ŏ	H	C_6H_5 C_6H_{11}	4-ClC ₆ H ₄	č	143 - 130 142 - 144	$C_{22}H_{20}C_{20}$ $C_{21}H_{23}ClN_2O$	0.009 (0.004-0.02)
30	RS	Õ	н	C6H11	$4-FC_6H_4$	Ċ	144-145	$C_{21}H_{23}FN_2O$	0.17 (0.12-0.23)
31	RS	0	H	C_6H_{11}	4-MeOC ₆ H ₄	C	63-67	$C_{22}H_{26}N_2O_2$	0.55 (0.38-0.71)
32 33	пS RS	ő	л Н	$C_{eH_{11}}$	2-ClCeH4	č	136-138 146-148	C ₂₂ H ₂₆ N ₂ O C ₂₁ H ₂₆ ClN ₂ O	0.30(0.28-0.33) 0.080(0.062-0.097)
34	S	ŏ	5-Me	C ₆ H ₁₁	C(O)NHMe	g	202-204	$C_{18}H_{25}N_3O_2$	0.072 (0.059-0.085)
35	RS	0	5-iPr	$4 - FC_6H_4$	$C(O)NC_5H_{10}$	g	109-110	$C_{24}H_{28}FN_3O_2$	0.3/
36 37	RS RS	0	5-1Pr 5-iPr	4-FC6H4 4-FC6H4	CH ₂ OMe CH ₂ OEt	g	oil	$C_{20}H_{23}FN_2O_2$ $C_{01}H_{02}FN_2O_2$	0.017 (0.010 - 0.023) = 0.041 (0.027 - 0.055)
38	RS	ŏ	5-iPr	4-FC ₆ H ₄	CH ₂ OnPr	g	oil	$C_{22}H_{27}FN_2O_2$	0.17 (0.12-0.23)
39	S	0	5-iPr	C ₆ H ₅	CH ₂ SMe	g	oil	$C_{20}H_{24}N_2OS \cdot 1/_4H_2O$	0.013 (0.0089-0.017)
40 41	S S	0	6-1PT 5-Me	C ₆ H ₅ C ₆ H ₁₁	C(O)Me 5-(3-MeCoNoO)	g	107-110	$C_{20}H_{22}N_2OS$ $C_{10}H_{24}N_4O_2$	0.35 (0.27-0.44)
42	$\tilde{R}S$	ŏ	5-iPr	4-FC ₆ H ₄	$5-(3-MeC_2N_2O)$	g	86-88	$C_{21}H_{21}FN_4O_2$	0.44 (0.17-0.71)
43a	RS	0	5-Me	C_6H_{11}	2-C₅H₄N	B	104-105	$C_{21}H_{25}N_3O$	0.004 (0.003-0.004)
43D 43c	S R	0	5-Me 5-Me	$C_{6}H_{11}$	2-C5H4N 2-C5H4N	C	80-80 75-78	$C_{21}H_{25}N_3U H_2U$ $C_{01}H_{02}N_9O$	0.001 (0.0007 - 0.001) 0.040 (0.033 - 0.047)
44	RS	ŏ	5-Me	C ₆ H ₁₁	3-C₅H₄N	B	150-151	$C_{21}H_{25}N_{3}O$	0.020 (0.017-0.022)
45	RS	0	5-Me	C_6H_{11}	4-C₅H₄N	B	188-189	$C_{21}H_{25}N_{3}O \cdot 1/_{2}H_{2}O$	$0.19 (*^{h}-0.42)$
40 47	пS RS	0	5-CO ₂ Et	C_6H_{11}	C ₆ H ₅	i	119-122 129-132	$C_{24}H_{30}N_{2}O^{4}/_{2}H_{2}O$	0.006 (0.005-0.008)
48	\widetilde{RS}	ŏ	5-CO ₂ H	C_6H_{11}	C ₆ H ₅	i	232-234	$C_{22}H_{24}N_2O_3$	0.15 (0.084-0.22)
49	RS	0	5-C(O)NMe ₂	C_6H_{11}	C ₆ H₅	i	194-196	$C_{24}H_{29}N_3O_2$	0.17 (0.13-0.20)
50 51	RS RS	0	$5-C(0)NH_2$ 5-CN	C_6H_{11} C_2H_{11}	C ₆ H ₅ C ₆ H ₅	ı i	181-183	$C_{22}H_{25}N_3O_2$ $C_{22}H_{25}N_3O_2$	35%/0.3 μM 0.015 (0.013–0.018)
52	RS	ŏ	5-CH₂OH	C_6H_{11}	C ₆ H ₅	i	187-189	$C_{22}H_{26}N_2O_2$	$41\%/0.3\mu M$
53	RS	0	5-NHC(O)Me	C_6H_{11}	C ₆ H ₅	i	145-147	$C_{23}H_{27}N_3O_2$	0.078 (0.055-0.10)
54 55	RS RS	0	5-NHSU2Me	C6H11 4-FCeH₄	C6H5 2-C₅H₄N	i B	188 - 190 105 - 107	$C_{22}H_{27}N_3O_3S$ $C_{21}H_{15}F_4N_3O$	0.16(0.12-0.20) 0.008(0.005-0.01)
56	\tilde{RS}	ŏ	5-OMe	4-FC ₆ H ₄	2-C ₅ H ₄ N	ē	111-113	$C_{21}H_{18}FN_3O_2$	0.011 (0.0056-0.016)
57	RS	0	5-iPr	4-FC ₆ H ₄	2-C₅H₄N	C	56-58	$C_{23}H_{22}FN_3O$	0.012 (0.0077-0.017)
58 598	RS RS	0	5-NU2 5-Cl	4-FC6H4 4-FC6H4	$2-C_5H_4N$ $2-C_5H_4N$	C C	174-175	$C_{20}H_{15}FIN_4O_3$ $C_{20}H_{15}CIFN_5O$	0.016 (0.010 - 0.021) 0.020 (0.015 - 0.025)
59b	-	ŏ	5-Cl	4-FC ₆ H ₄	2-C ₅ H ₄ N	e	oil	C ₂₀ H ₁₅ ClFN ₃ O	0.020 (0.014-0.027)
59c	+	0	5-Cl	4-FC ₆ H ₄	2-C₅H₄N	e	oil	C ₂₀ H ₁₅ ClFN ₃ O	0.32 (0.094-0.55)
60 61	RS RS	0	5-F 5.6-diF	4-FC6H4 4-FC6H4	2-C5H4N 2-C5H4N	B	117-119 143-144	$C_{20}H_{15}F_{2}N_{3}O$ $C_{20}H_{14}F_{3}N_{3}O$	$0.021 (0.015 - 0.027) \\ 0.030 (0.018 - 0.042)$
62	RS	Õ	5-CO ₂ Et	4-FC ₆ H ₄	$2-C_5H_4N$	Ĉ	151-152	$C_{23}H_{20}FN_3O_3$	0.027 (0.0080-0.047)
63 64	RS	0	5-SO ₂ NMe ₂	$4 - FC_6H_4$	2-C₅H₄N	C	175-175	$C_{22}H_{21}FN_4O_3S$	0.050 (0.024 - 0.075)
65	RS	ŏ	5-iPr	3-ClC6H4	2-C5H4I 2-C5H4N	č	51-52	$C_{23}H_{22}CIN_{3}O - 74H_{2}O$ $C_{23}H_{22}CIN_{3}O$	0.005 (0.002-0.007)
66	RS	Ó	5-NO ₂	4-OMeC ₆ H ₄	2-C₅H₄N	C	162-163	$C_{21}H_{18}N_4O_4$	0.014 (0.011-0.017)
67 689	RS R	0	o-NO2 5-iPr	4-0HC6H4 k	2-U5H4N CeHe	C F	153-154 180 dec	U20H16N4U4 CooHmNoO.2HCl.H-O	27%/0.3 μM 0.10 (0.084–0.12)
68b	\ddot{s}	ŏ	5-iPr	k	C ₆ H₅	F	185 dec	C ₂₃ H ₂₉ N ₃ O·2HCl·H ₂ O	1.2 (0.79–1.6)
69 70	R	0	5-iPr	l	C ₆ H ₅	F	135-137	$C_{22}H_{27}N_3O_2$	0.052 (0.044-0.061)
70	к RS	0	5-Cl	n n	$C_6 \Pi_5$ $C_6 H_5$	r G	189-191	$C_{19}H_{20}CIN_3O_3S$	$46\%/1 \mu M$
72	R	ŏ	5-C1	0	C ₆ H ₅	F	58-60	C ₂₀ H ₂₃ ClN ₄ O	$21\%/1\mu M$
73	RS	0	5-iPr	р	2-C₅H₄N	С	116-117	$C_{21}H_{22}N_4OS^q$	0.029 (0.029-0.030)

Table 2. (Continued)

compd no.	R/S	Y	R ₁	R_2	R3	syn meth	mp (°C)	formulaª	IC50 ^b (µM) or % inh of LTB4/conc
74	RS	0	5-iPr	р	CO ₂ Et	F	oil	C ₁₉ H ₂₃ N ₃ O ₃ S	0.016 (0.0057-0.026)
75	\boldsymbol{S}	0	5-iPr	r	$\rm CO_2Et$	C*	166-168	C17H20N4O3.2HCl	$26\%/1 \mu M$
76	\boldsymbol{s}	0	5-iPr	t	$\rm CO_2Et$	C*	143-146	C ₁₉ H ₂₄ N ₄ O ₃ ·2HCl ^u	59%/1 µM

^a See footnote a, Table 1. ^b See footnote b, Table 1. ^c FAB HRMS m/z 355.1474 (M + H)⁺; calcd 355.1492. ^d C: calcd, 67.53; found, 67.07. ^e Prepared by chiral HPLC of the racemate. See Experimental Section. ^f Estimate. No nonlinear regression analysis due to curve shape. ^g Scheme 4. ^h Lower bound was negative or equal to zero. ⁱ Scheme 5. ^j FAB HRMS m/z 376.1177 (M + H)⁺; calcd 376.1173. ^k N-Piperidinyl. ⁱ N-Morpholinyl. ^m N-Thiomorpholinyl. ⁿ N-S,S-Dioxothiomorpholinyl. ^o N-N'-Methylpiperizinyl. ^p 2-Methyl-4-thiazolyl. ^q FAB HRMS m/z 378.1493 (M + H)⁺; calcd 378.1517. ^r 4-Imidazolyl. ^s Due to insolubility in CH₂Cl₂, toluene with enough MeOH to dissolve was used as solvent. ^t 3-Ethyl-4-imidazolyl. ^a C: calcd, 53.15; found, 54.64.

Scheme 1^a



 a (a) Ether, 0 °C; (b) chlorobenzene, SO₂Cl₂, 0 °C (method A); (c) NaOH, DMSO (method B).

Scheme 2^a



^a (a) CS₂, KOH, MeOH, H₂O; (b) PCl₅, POCl₃; (c) NaOH, DMSO (method B); (d) iPr₂NEt, CH₂Cl₂ (method C).

Scheme 3^a



^a (a) Mg, ether; (b) R₃CN; (c) NaBH₄, MeOH (method D); (d) Mg, ether; (e) R₃CHO; (f) MnO₂; (g) (1) HC(O)NH₂, HC(O)OH, (2) 2 N HCl (method E); (h) R₃H, AlCl₃, CH₂Cl₂.

hibited both, and the dose–response curves were coincidental with LTB₄. This is consistent with a site of action at the level of arachidonic acid release from phospholipids. However, no direct effect on cytosolic PLA₂ was observed for this compound, suggesting an indirect effect on PLA₂ activity. These results are being published separately.²⁶

Experimental Section

Melting points were taken on a Buchi 510 melting point apparatus and are uncorrected. ¹H NMR were all consistent with molecular structures and were recorded on a Bruker 250

Scheme 4^a





^a (a) CDI, DMF, NHR₄R₅; (b) MeLi, THF; (c) NaH, CH₃C-(NOH)NH₂; (d) (Bu₄N)⁺HSO₄⁻, R₃I, NaOH, H₂O; (e) TFA, CH₂Cl₂; (f) method C, R₁ = iPr; (g) MsCl, Et₃N, CH₂Cl₂; (h) NaSCH₃.

Scheme 5^a



 a (a) 2 N NaOH, EtOH; (b) CDI, DMF, Me₂NH; (c) CDI, DMF, NH₃; (d) Et₃N, THF, [CF₃C(O)]₂O; (e) LAH, THF; (f) H₂, Pd/C, EtOH; (g) AcCl, Et₃N, DMF; (h) MsCl, Et₃N, DMF.

WM spectrometer. Elemental analyses were performed at Midwest Microlab, Indianapolis, IN, and were within 0.4% of the calculated values unless otherwise indicated.

Ethyl (R,S)-N-(Benzothiazol-2-ylmethyl)-4-chlorophenylalanine HCl (9). (R,S)-4-Chlorophenylalanine ethyl ester HCl (1.85 g, 7 mmol) and several 4A molecular sieves were added to a solution of 2-benzothiazolecarboxaldehyde²⁹ (1 g, 6.1 mmol) in 30 mL of EtOH. NaBH₃CN (0.8 g, 12.7 mmol) was added, and the reaction mixture was stirred at room temperature for 23 h. The reaction mixture was poured into 150 mL of H₂O, made basic with 2 N NaOH, and extracted with EtOAc. The combined EtOAc extracts were washed with brine, dried (Na₂SO₄), and

Scheme 6

Method F^a



Method G^b

^a (a) $O(CO_2tBu)_{2}$; (b) MsCl, Et₃N; (c) THF, reflux; (d) TFA, CH_2Cl_2 ; (e) method B or C. ^b (a) H_2O_2 , HOAc; (b) iPr_2NEt ; (c) NH_4OAc , $NaBH_3CN$, (d) iPr_2NEt , CH_2Cl_2 .

concentrated. The resulting oil was chromatographed on silica gel, eluting with 98 CH₂Cl₂:2MeOH, and the fractions containing the product were collected, concentrated, and treated with ethereal HCl. The HCl salt was recrystallized from EtOH giving 9 (0.4 g, 16%), mp 150–153 °C. ¹H NMR (CDCl₃): δ 8.0 (d, 1H, aromatic), 7.5 (d, 1H, aromatic), 7.3 (m, 7H, aromatic), 4.1 (q, 2H, CH₂), 3.3 (m, 5H, 2 CH₂S, 1 CH), 1.1 (t, 3H, CH₃). Anal. (C₁₉H₁₉ClN₂O₂S HCl) C, H, N.

Ethyl (R,S)-3-(Benzothiazol-2-yl)-2-benzylpropionate (10). A solution of ethyl hydrocinnamic acid (3.56 g, 20 mmol) in 20 mL of THF was added under N_2 to a solution of lithium diisopropylamide prepared by addition of 15 mL of 1.6 N n BuLi (24 mmol) to 2.42 g (24 mmol) of diisopropylamine in 100 mL of THF at -78 °C. The temperature was maintained between -40 and -15 °C for 30 min after addition. A solution of 3.26 g of 2-benzothiazolecarboxaldehyde²⁹ in 30 mL of THF was added over 15 min (-40 to -20 °C). The reaction mixture was allowed to warm slowly to room temperature. The reaction mixture was poured into a mixture of ice and water, made acidic with 2 N HCl, and extracted with EtOAc. The combined extracts were washed with brine, dried (Na₂SO₄), and concentrated giving an oil which was chromatographed on silica gel, eluting with 99 CH₂-Cl₂:1 MeOH. The fractions containing the product (a diastereometric pair) were combined and concentrated giving 4.2 g (61%)of ethyl 3-(benzothiazol-2-yl)-3-hydroxy-2-benzylpropionate as an oil. A sample of the first eluting diastereomer was recrystallized (ligroine) and characterized, mp 94-96 °C. ¹H NMR (CDCl₃): § 8.05 (d, 1H, aromatic), 7.95 (d, 1H, aromatic), 7.45 (m, 7H, aromatic), 5.1 (d, 1H, CH), 4.7 (d, 1H, OH), 4.1 (q, 2H, CH₂), 3.75 (m, 1H, CH), 3.3 (m, 2H, CH₂), 1.1 (t, 3H, CH₃).

The above hydroxy ester (4.2 g, 12.3 mmol) was dissolved in 20 mL of pyridine and cooled on an ice bath. POCl₃ (2.73 g, 17.8 mmol) was added, and the reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction mixture

was then heated to 90–100 °C for 2 h, cooled, and added to ice and water. The product was extracted with EtOAc, washed with 2 N HCl and brine, dried (Na₂SO₄), and concentrated. The residue was chromatographed on silica gel, eluting with CH₂Cl₂, giving 1.47 g (37%) of ethyl 3-(benzothiazol-2-yl)-2-benzylpropenoate as an oil. ¹H NMR (CDCl₃): δ 8.1 (d, 1H, aromatic), 7.95 (s, 1H, olefin), 7.95 (d, 1H, aromatic), 4.5 (s, 2H, CH₂), 4.25 (q, 2H, CH₂), 1.25 (t, 3H, CH₃).

71

A solution of 1.43 g (4.43 mmol) of the above olefin in 75 mL of EtOH was combined with 650 mg of 10% Pt/C and hydrogenated under 40 psi until TLC (CH₂Cl₂) showed no starting material remained (24 h). The reaction mixture was filtered through Celite and concentrated. The residue was recrystallized from ligroine giving 0.57 g (39.5%) of 10, mp 46–48 °C. ¹H NMR (CDCl₃): δ 8.0 (d, 1H, aromatic), 7.85 (d, 1H, aromatic), 7.3 (m, 7H, aromatic), 4.1 (q, 2H, CH₂), 3.45 (m, 1H, CH), 3.3 (m, 2H, CH₂), 3.05, (m, 2H, CH₂), 1.1, (t, 3H, CH₃). Anal. (C₁₉H₁₉NO₂S) C, H, N.

Method A. (R,S)-N-(6-Isopropylbenzothiazol-2-yl)-4chlorophenylalanine Ethyl Ester (15). (R,S)-4-Chlorophenylalanine ethyl ester HCl (5 g, 18.9 mmol) was converted to the free base with Et₃N and dissolved in 75 mL of ether. This solution was added to a solution of 4-isopropylphenylisothiocyanate (3.36 g, 19 mmol) in 150 mL of ether. The temperature was maintained at 0 °C or less during the addition by cooling in an ice-salt bath. The reaction mixture was stirred for 4.5 h on the ice bath and then concentrated. The foamy residue was triturated with petroleum ether resulting in 6.1 g (15.1 mmol, 80%) of the intermediate thiourea, mp 73-75 °C.

The thiourea (6.0 g, 14.8 mmol) was dissolved in 25 mL of chlorobenzene and cooled on an ice bath to 0 °C. Sulfuryl chloride (2.76 g, 20.4 mmol) in 5 mL of chlorobenzene was added dropwise. After 5.5 h, the reaction mixture was concentrated and the residue

General Procedure for Preparation of 2-Chlorobenzoxazoles. 2-Chloro-5-methylbenzoxazole. To a suspension of 1.0 g (6.05 mmol) of 5-methyl-2-mercaptobenzoxazole in 8.22 g (53.6 mmol) of POCl₃ at room temperature was added 1.26 g (7.2 mmol) of PCl_5 along with 5 mL of CH_2Cl_2 . The reaction mixture turned into a solution after the addition. After 1 h of stirring at room temperature, the reaction mixture was concentrated to remove excess POCl₃, and the residue was treated with Na₂CO₃ solution until pH8 was reached. The aqueous phase was extracted with CH₂Cl₂, and the combined CH₂Cl₂ extracts were washed with brine, dried $(MgSO_4)$, and concentrated to give 1.1 g of crude material after azeotroping with toluene. Short-column chromatography on silica gel starting with petroleum ether followed by 10% CH₂Cl₂/petroleum ether provided 820 mg (81% yield) of 2-chloro-5-methylbenzoxazole as an oil. ¹H NMR (CDCl₃): δ 2.45 (s, 3H, CH₃), 7.1–7.4 (m, 3H, Ar).

Method B. (R,S)-N-(Benzothiazol-2-yl)phenylalanine Hydrochloride (11). A 11.8-g (71.4 mmol) sample of (R,S)phenylalanine was added in portions to a suspension of 5.7 g of powdered NaOH (143 mmol) in 50 mL of DMSO and stirred under N₂ for 30 min. 2-Chlorobenzothiazole (11 g, 65 mmol) was added over 15 min at room temperature. The reaction mixture was heated on an oil bath set at 95 °C for 4 h. The cooled reaction mixture was poured into 200 mL of ice and water and the pH of the resulting solution adjusted to 1-2 by the addition of 10 N HCl. More ice was added, and the mixture was filtered. The white solid was dissolved in alkaline solution, stirred with Celite, filtered, and acidified with 2 N HCl, and the resulting white precipitate was filtered, rinsed with water and EtOH, and dried. This resulted in 6.47 g of 11 (30%), mp 250-251 °C. ¹H NMR $(DMSO-d_6): \delta 13.5 (br s, 2H, NH^{2+}), 10.1 (br s, 1H, CO_2H), 7.9-$ 7.1 (m, 9H, aromatic), 5.1 (m, 1H, CH), 3.25 (m, 2H, CH₂). Anal. (C₁₆H₁₄N₂O₂S·HCl) H, N; C: calcd, 57.39; found, 57.97.

(S)-N-(Benzoxazol-2-yl)cyclohexylalanine Methyl Ester (23a). To a solution of 10.0 g (46.36 mmol) of (S)-(-)-phenylalanine methyl ester hydrochloride in 75 mL of MeOH was added 1.4 g of Rh/Al₂O₃. The resulting mixture was hydrogenated on a Parr shaker overnight at room temperature under 50 psi of hydrogen. The catalyst was filtered off with the aid of Celite. The filtrate was concentrated to give a white solid which was recrystallized from CH₂Cl₂/ether to give 9.2 g (95% yield) of (S)-(-)-cyclohexylalanine methyl ester hydrochloride, mp 154-155 °C. ¹H NMR (CDCl₃): δ 0.8-1.7 (m, 13H, aliph), 3.74 (s, 3H, OCH₃), 4.0 (t, 1H, CH), 8.6 (br, 3H, NH₃⁺).

To a mixture of 2.21 g (10 mmol) of (S)-(-)-cyclohexylalanine methyl ester hydrochloride and 1.68 g (11 mmol) of 2-chlorobenzoxazole in 75 mL of CH₂Cl₂ was added 2.58 g (20 mmol) of diisopropylethylamine. The resulting mixture was refluxed overnight. The reaction mixture was diluted with CH₂Cl₂ and washed with H₂O. The CH₂Cl₂ phase was dried and concentrated to give an oil material weighing 3.05 g. Flash column chromatography on silica gel starting with 1:1 petroleum ether:CH₂Cl₂ followed by 100% CH₂Cl₂ and 1% MeOH in CH₂Cl₂ gat at at al of 1.78 g (59% yield) of **23a** as an oil. ¹H NMR (CDCl₃): δ 1.0–1.8 (m, 13H, aliph), 3.72 (s, 3H, CH₃), 4.65 (m, 1H, CH), 5.48 (br, 1H, NH), 7.02–7.38 (m, 4H, Ar). Anal. (C₁₇H₂₂N₂O₃) C, H, N.

Methods C and E. (R,S)-N-(2-Cyclohexyl-1-phenylethyl)-2-ben zoxazolamine (26a). Cyclohexylacetyl chloride³⁰ (10 g, 62.2 mmol) in 75 mL of CH₂Cl₂ was added over 30 min to 8.29 g (62.2 mmol) of AlCl₃ in 75 mL of CH₂Cl₂ and stirred under N₂ on an ice bath. After 40 min, 5.08 g (65 mmol) of benzene in 30 mL of CH₂Cl₂ was added over 25 min. The reaction mixture was stirred on ice for 30 min and then stirred for 2 h at room temperature, refluxed for 3 h, and stirred overnight at room temperature. The reaction mixture was poured into cold 2 N HCl, the organic phase was separated, and the aqueous phase was washed with more CH₂Cl₂. The combined organic phases were washed with 2 N HCl and brine, dried (Na₂SO₄), and concentrated. The residual oil (12 g) was chromatographed on silica gel, eluting with CH₂Cl₂, giving 1-phenyl-2-cyclohexylethanone (10.4 g, 51.4%). ¹H NMR (CDCl₃): δ 7.9 (m, 2H, phenyl), 7.5 (m, 3H, phenyl), 2.85 (d, 2H, CH₂), 2.0 (m, 1H, CH), 1.7 (m, 5H, cyclohexyl), 1.1 (m, 5H, cyclohexyl).

A mixture of the above ketone (10.37 g, 51.4 mmol) and formamide (11.57 g, 250 mmol) was stirred on an oil bath. The oil-bath temperature was raised to 170 °C as 5.9 g (128 mmol) of formic acid was added in portions. Additional formamide and formic acid were added at various intervals over the next 30 h, until TLC (98 CH₂Cl₂:2 MeOH) showed the reaction was over. The cooled reaction mixture was then dissolved in EtOAc, washed with water and brine, dried (Na₂SO₄), and concentrated. The resulting formamide (10.88 g, 91%) was suspended in 250 mL of 2 N HCl and heated to reflux. After 4 h, the reaction mixture was cooled and the precipitated amine hydrochloride filtered, rinsed with water, and dried giving 9.35 g (76% from ketone) of 1-phenyl-2-cyclohexylethylamine HCl, mp 288 °C. ¹H NMR (CDCl₃): δ 8.5 (br s, 3H, NH₃⁺), 7.4 (m, 5H, phenyl), 4.25 (br t, 1H, CH), 1.7 (m, 7H, CH, CH₂), 1.0 (m, 6H, CH₂).

A mixture of 1.48 g (7.3 mmol) of the free base of the above amine, 1.12 g of 2-chlorobenzoxazole (Aldrich), and 1.14 g (8.8 mmol) of diisopropylethylamine in 30 mL of CH₂Cl₂ was heated at reflux for 24 h. The reaction mixture was diluted with 50 mL of CH₂Cl₂ and washed with H₂O, 1 N HCl, saturated Na₂CO₃ solution, and brine, dried (Na₂SO₄), and concentrated. The resulting solid was recrystallized from EtOH giving 1.4 g (60%) of **26a**, mp 131–132 °C. ¹H NMR (CDCl₃): δ 7.3 (m,9H, aromatic), 6.3 (br s, 1H, NH), 5.0 (m, 1H, CH), 1.75 (m, 7H, CH, CH₂), 1.5–0.8 (m, 6H, aliphatic). Anal. (C₂₁H₂₄N₂O) C, H, N.

Method D. 1-(2-Pyridyl)-2-cyclohexylethylamine Dihydrochloride. Cyclohexylmethyl bromide (22.1 g, 0.125 mol) in 200 mL of anhydrous ether was added dropwise to 3.04 g of Mg turnings (0.125 mol) at a rate to maintain reflux. After the reaction subsided, the mixture was heated to reflux for 1 h and then cooled to 0 °C. 2-Cyanopyridine (10 g, 0.096 mol) in 100 mL of ether was then added, keeping the temperature under 10 °C. After addition, the reaction mixture was stirred for an additional 1 h on the ice bath and then 300 mL of MeOH was added slowly at first and then more quickly. A dark green gum settled which was stirred with a glass rod until it dissolved. After 30 min, 7.26 g of NaBH₄ was added in portions, keeping the temperature at about 10 °C. The reaction mixture was then allowed to warm to room temperature and stirred for 2 h. The solvent was then removed on a rotary evaporator, and the residue was treated with H₂O and then 2 N HCl. The mixture was extracted with CH₂Cl₂ and then made basic with 2 N NaOH and the product extracted into CH₂Cl₂, washed with brine, dried (Na₂-SO4), and concentrated. The resulting oil was dissolved in ether and filtered, and the dihydrochloride salt was precipitated with ethereal HCl giving 14.26 g (61%), mp 133-136 °C. NMR was identical to that of the sample prepared by method E.

(S)-3-Cyclohexyl-2-[(5-methylbenzoxazol-2-yl)amino]-Nmethylpropionamide (34). A solution of 1.08 g (3.57 mmol) of (S)-N-(5-methylbenzoxazolyl)cyclohexylalanine in 15 mL of CH₂-Cl₂ was cooled on an ice bath. Carbonyldiimidazole (0.88 g, 15.4 mmol) was added in portions. After the solution was stirred for 1 h, methylamine was bubbled in for 45 min. The reaction mixture was then diluted with CH₂Cl₂, washed with H₂O and brine, dried (Na₂SO₄), and concentrated. The residue was chromatographed on silica gel, eluting with 99 CH₂Cl₂:1 MeOH. After recrystallization from iPrOH, 0.2 g (18%) of 34 was obtained, mp 202-204 °C. ¹H NMR (CDCl₃): δ 7.15 (m, 2H, aromatic), 6.85 (d, 1H, aromatic), 6.55 (br d, 1H, NH), 5.9 (d, 1H, NH), 4.4 (m, 1H, CH), 2.8 (d, 3H, CH₃), 2.4 (s, 3H, CH₃), 1.9-0.85 (m, 13H, aliphatic). Anal. (C₁₈H₂₆N₃O₂) C, H, N.

(R,S)-5-Isopropyl-N-[3-(4-fluorophenyl)-1-(methoxymethyl)prop-2-yl]benzoxazolamine (36). To a solution of 6.5 g (33 mmol) of D,L-4-fluorophenylalanine methyl ester in 30 mL of dry THF was added 4.19 g (98.9 mmol) of LiCl followed by 3.74 g (98.9 mmol) of NaBH₄ and 60 mL of absolute EtOH under argon at room temperature. The reaction mixture was kept at 50-55 °C for 2 h and left overnight at room temperature. The reaction was quenched with saturated NH₄Cl and the mixture extracted with CH₂Cl₂. The combined CH₂Cl₂ extracts were dried (MgSO₄) and concentrated to give 5.1 g of oil. Trituration of the partially crystallized material with ether provided a first crop of D,L-4-fluorophenylalaninol (1.87 g) as a white crystalline solid, mp 160 °C. A second crop of 920 mg was obtained after flash column chromatography of the filtrate on silica gel with 2.5% MeOH in CH₂Cl₂. A total of 2.79 g (50% yield) was obtained. ¹H NMR (CDCl₃/DMSO- d_6): δ 2.83 (dd, 1H, CHPh), 3.0 (dd, 1H, CHPh), 3.3 (m, 1H, CH), 3.48 (dd, 1H, CHOH), 3.64 (dd, 1H, CHOH), 6.94 (t, 2H, Ar), 7.18 (dd, 2H, ArF).

D,L-N-(tert-Butoxycarbonyl)-p-fluorophenylalaninol was prepared by the same procedure described below in method F for (R)-(-)-N-(tert-butoxycarbonyl)-2-phenylglycinol. ¹H NMR (CDCl₃): δ 1.4 (s, 9H, tBu), 1.84 (br, 1H, OH), 3.5 (d, 2H, CH₂-Ph), 3.55 (dd, 1H, CHO), 3.67 (dd, 1H, CHO), 3.8 (m, 1H, CHN), 4.7 (br, 1H, NH), 7.0 (t, 2H, Ar), 7.15 (dd, 2H, ArF).

A solution of 350 mg (13 mmol) of the above alcohol, 440 mg (1.3 mmol) of tetrabutylammonium hydrogen sulfate, and 920 mg (6.5 mmol) of iodomethane in 4 mL of THF and 3 mL of 50% NaOH/H₂O was stirred vigorously overnight at room temperature. The reaction was quenched with H₂O. The aqueous phase was extracted with ether, dried (MgSO₄), and concentrated to give 380 mg of oil which crystallized on standing. Flash column chromatography starting with petroleum ether followed by 5% and then 20% EtOAc in petroleum ether afforded 360 mg (98% yield) of the methyl ether, mp 66–67.5 °C. ¹H NMR (CDCl₃): δ 1.4 (s, 9H, tBu), 2.8 (m, 2H, CH₂Ar), 3.3 (dd, 2H, CH₂O), 3.5 (s, 3H, CH₃O), 3.85 (m, 1H, CH), 4.84 (br, 1H, NH), 6.95 (t, 2H, Ar), 7.15 (dd, 2H, ArF).

A solution of 350 mg (1.24 mmol) of the above methyl ether in 1 mL of CF₃CO₂H and 1 mL of CH₂Cl₂ was stirred under anhydrous conditions. The reaction was quenched with 2 N NaOH to pH 9, and the aqueous phase was extracted with CH₂-Cl₂, dried (MgSO₄), and concentrated to give 200 mg (88% yield) of D,L-4-fluorophenylalaninyl methyl ether as an oil which was used directly in the next step. ¹H NMR (CDCl₃): δ 1.5 (s, 2H, NH₂), 2.5 (dd, 1H, CHAr), 2.7 (dd, 1H, CHAr), 3.2 (m, 2H, CH₂O), 3.35 (dm, 4H, CH, OCH₃), 6.97 (t, 2H, Ar), 7.15 (dd, 2H, ArF).

The coupling of the 2-chloro-5-isopropylbenzoxazole and the above amine was done by using Method C to afford an 86% yield of 36 as an oil. ¹H NMR (CDCl₃): δ 1.28 (d, 6H, 2 Me), 3.0 (m, 3H, isoprCH, CH₂Ar), 3.37 (s, 3H, CH₃O), 3.37–3.4 (m, 2H, CH₂O), 4.2 (m, 1H, CH), 5.2 (br, 1H, NH), 6.9–7.28 (m, 7H, Ar). Anal. (C₂₀H₂₃FN₂O₂) C, H, N.

(S)-5-Isopropyl-N-[1-(methylthio)-3-phenylprop-2-yl]-2benzoxazolamine (39). The mesylate derivative of (S)-N-tBocphenylalaninol was prepared by the same procedure as described in method F for the mesylate of (R)-N-tBoc-phenylglycinol. ¹H NMR (CDCl₃): δ 1.4 (s, 9H, tBu), 2.9 (dd, 2H, CH₂Ph), 3.0 (s, 3H, CH₃SO₃), 4.1 (dd, 2H, CH₂OSO₂), 4.25 (m, 1H, CH), 4.7 (br, 1H, NH), 7.15-7.3 (m, 5H, Ar).

Sodium thiomethoxide (210 mg, 3.06 mmol) was added to a solution of 1.0 g (2.78 mmol) of the above methane sulfonate in 5 mL of DMF at 0 °C under Argon. The reaction mixture was stirred at 0 °C for 0.5 h and then at room temperature overnight. After being diluted with ether, the organic phase was washed with saturated NH₄Cl and H₂O. The combined aqueous phase was back-extracted with ether. The ether extracts were dried (MgSO₄) and concentrated to give 820 mg of residue. Flash column chromatography with petroleum ether followed by 5% EtOAc to 15% EtOAc in petroleum ether provided 560 mg (72% yield) of (S)-(-)-N-(tert-butoxycarbonyl)-1-(methylthio)-3-phenyl-2-propylamine, mp 82–90 °C. ¹H NMR (CDCl₃): δ 1.4 (s, 9H, tBu), 2.14 (s, 3H, CH₃S), 2.58 (d, 2H, CH₂S), 2.9 (d, 2H, CH₂Ph), 4.0 (m, 1H, CH), 4.65 (br, 1H, NH), 7.15–7.3 (m, 5H, Ar).

A solution of 200 mg (0.71 mmol) of the above N-protected thioether in 0.5 mL of CF_3CO_2H and 0.5 mL of CH_2Cl_2 was stirred at room temperature for 5 h under anhydrous conditions. The reaction was quenched with saturated Na₂CO₃ and the mixture extracted with CH_2Cl_2 . The combined CH_2Cl_2 extracts were dried and concentrated to give (S)-(-)-2-amino-3-phenylpropyl methyl thioether (100 mg, 78%) as an oil. ¹H NMR ($CDCl_3$): δ 1.75 (br, 2H, NH₂), 2.1 (s, 3H, CH_3S), 2.48 (dd, 1H, CHS), 2.64 (dd, 1H, CHS), 2.7 (dd, 1H, CHPh), 2.85 (dd, 1H, CHPh), 3.2 (m, 1H, CH), 7.2–7.35 (m, 5H, Ar).

Via method C with 2-chloro-5-isopropylbenzoxazole and the above amine, **39** was obtained in 59% yield as an oil. ¹H NMR (CDCl₃): δ 1.28 (d, 6H, 2 Me), 2.14 (s, 3H, CH₃S), 2.76 (d, 2H, CH₂S), 2.97 (m, 1H, CHMe₂), 3.1 (dd, 2H, CH₂Ph), 4.3 (m, 1H,

CH), 5.0 (d, 1H, NH), 6.9 (d, 1H, Ar), 7.1 (d, 1H, Ar), 7.3 (m, 6H, Ar). Anal. $(C_{20}H_{24}N_2OS^{-1}/_4H_2O)$ C, H, N.

(S)-3-[(6-Isopropylbenzothiazol-2-yl)amino]-4-phenyl-2butanone (40). According to a general procedure for methyl ketones,³¹ a solution of (S)-N-(6-isopropylbenzothiazol-2-yl)phenylalanine (2 g, 5.9 mmol) in 60 mL of THF was stirred at -5 °C under N₂ on an ice-salt bath. A solution of 26 mL of 1.4 N MeLi in ether (36.4 mmol) was added via syringe, keeping the temperature below 10 °C. After the solution was stirred for 2 h at 0 °C, 8.56 g of chlorotrimethylsilane (78 mmol) was added, the reaction mixture was warmed to room temperature, and 45 mL of 1 N HCl was added. After stirring 30 min, the aqueous phase was extracted with ether, and the combined organic extracts were washed with water and brine, dried (Na₂SO₄), and concentrated. After chromatography on silica gel with CH₂Cl₂, the product was recrystallized from EtOH giving 0.71 g of 40 (36%), mp 107-110 °C. ¹H NMR (CDCl₃): δ 7.5 (m, 2H, aromatic), 7.2 (m, 6H, aromatic), 5.75 (d, 1H, NH), 5.0 (m, 1H, CH), 3.3 (m, 2H, CH₂), 2.95 (m, 1H, CH), 2.25 (s, 3H, CH₃), 1.25 (d, 6H, CH₃). Anal. (C20H22N2OS) C, H, N.

Methods B and E. (R.S)-N-[2-Cyclohexyl-1-(2-pyridinyl)ethyl]-5-methyl-2-benzoxazolamine (43a). A solution of 17.7 g (0.1 mol) of cyclohexylmethyl bromide in 50 mL of ether was added dropwise to a mixture of 2.43 g (0.1 mol) of Mg turnings in 25 mL of ether. The reaction mixture started to reflux immediately. At the end of the addition when the refluxing subsided, the reaction mixture was heated to reflux for an additional 0.5 h. After the mixture had cooled to room temperature, 5.36 g (0.05 mol) of 2-pyridinecarboxyaldehyde was added slowly to the freshly generated Grignard reagent. The reaction was exothermic during the addition. The resulting mixture was stirred at room temperature for 3 h before it was poured into ice- H_2O . The aqueous phase was treated with dilute HCl until pH 7 was reached. The aqueous phase was extracted with ether, and the combined ether extracts were washed with brine, dried (MgSO₄), filtered, and concentrated to give 9.1 g of an oily residue. Crystallization from CH₂Cl₂/petroleum ether provided a total of 6.05 g (59% yield) of 2-cyclohexyl-1-(2-pyridyl)ethanol, mp 72-74 °C. ¹H NMR (CDCl₃): δ 0.95-1.8 (m, 12H, 6 CH₂), 1.99 (m, 1H, CH), 4.05 (d, 1H, OH), 4.8 (m, 1H, CH), 7.2 (m, 2H, pyr), 7.7 (t, 1H, pyr), 8.5 (d, 1H, pyr).

To a solution of 4.63 g (22.5 mmol) of the above alcohol in 50 mL of CH_2Cl_2 was added a total of 8.88 g (0.1 mol) of MnO_2 in portions over a period of 5 h under anhydrous conditions. The resulting mixture was stirred at room temperature overnight. The catalyst was filtered off with the aid of Celite. The filtrate was concentrated to give an oil. Chromatography on a short column of silica gel, eluting with petroleum ether followed by $20\% CH_2Cl_2$ in petroleum ether, provided 4.1 g (89% yield) of (cyclohexylnethyl)-2-pyridyl ketone as an oil. ¹H NMR (CDCl₃): $\delta 1.07-1.3$ (m, 5H, cyclohexyl), 1.74 (m, 5H, cyclohexyl), 2.0 (m, 1H, CH of cyclohexyl), 3.1 (d, 2H, CH₂), 7.45 (m, 1H, pyr), 7.85 (t, 1H, pyr), 8.05 (d, 1H, pyr), 8.7 (d, 1H, pyr).

A mixture of 1.8 g (8.85 mmol) of the above ketone and 2.22 g (49.2 mmol) of formamide was heated on an oil bath at 160 °C. To this was added 1.2 g (26.5 mmol) of formic acid in portions while keeping the temperature between 160 and 165 °C. The reaction was completed in 3 h. After cooling, 17 mL of 2 N HCl solution was added, and the resulting mixture was refluxed for 5 h. The reaction mixture was diluted with ether and treated with 10 N NaOH followed by saturated Na₂CO₃ until pH 9 was reached. The aqueous phase was extracted with ether, and the combined ether extracts were washed with brine, dried (MgSO₄), and concentrated to give a dark oil. Upon treatment of the oil with ethereal HCl, a solid was obtained. The solid was recrystallized from MeOH/CH₂Cl₂/petroleum ether to give 1.3 g (54%)yield) of 1-(2-pyridyl)-2-cyclohexylethylamine dihydrochloride. ¹H NMR (DMSO-d₆): δ 0.8–1.1 (m, 6H, aliph), 1.55–1.8 (m, 7H, aliph), 4.5 (m, 1H, CH), 7.5 (m, 1H, pyr), 7.67 (d, 1H, pyr), 7.95 (m, 1H, pyr), 8.6 (br, 2H, NH₂), 8.68 (d, 1H, pyr), 9.05 (br, 2H, 2NH+).

A mixture of 531 mg (13 mmol) of powdered NaOH in DMSO was stirred at room temperature for 10 min before 1.2 g (4.34 mmol) of the above amine dihydrochloride was added. After the mixture was stirred for 0.5 h at room temperature, 661 mg (3.95 mmol) of 2-chloro-5-methylbenzoxazole was added. The resulting

Benzoxazolamines and Benzothiazolamines

mixture was kept at 60 °C with stirring for 2 h. The reaction was quenched with H₂O. The aqueous phase (pH 9) was extracted with ether, dried, and concentrated to give an 1.3-g oily product. This residue was treated with ethereal HCl to give a gummy solid which was recrystallized from CH₂Cl₂/petroleum ether giving 1.1 g of product as the dihydrochloride salt (69% yield), mp > 140 °C. A portion (680 mg) of the salt was converted to the free base by dissolving in H₂O and adjusting the pH to 9. The aqueous phase was extracted with CH₂Cl₂, dried (MgSO₄), and concentrated to give 570 mg of oil. Upon trituration with petroleum ether, 458 mg of 43a (82% recovery) was obtained, mp 104-105 °C. ¹H NMR (CDCl₉): δ 1.0-1.8 (m, 13H, aliph), 2.35 (s, 3H, CH₃), 5.1 (dd, 1H, CHN), 5.95 (d, 1H, NH), 6.8 (m, 1H, Ar), 7.1 (d, 1H, Ar), 7.18 (m, 2H, Ar, pyr), 7.3 (d, 1H, pyr), 7.64 (m, 1H, pyr), 8.55 (d, 1H, pyr). Anal. (C₂₁H₂₅N₃O) C, H, N.

2-[(2-Cyclohexyl-1-phenylethyl)amino]-5-benzoxazolecarboxylic Acid Ethyl Ester (47). This compound was prepared by method C in 66% yield, mp 129–132 °C. NMR (CDCl₃): δ 8.0 (s, 1H, Ar), 7.8 (d, 1H, Ar), 7.3 (m, 6H, Ar), 6.75 (d br, 1H, NH), 5.0 (m, 1H, CH) 2.0–0.85 (m, 13H, CH₂, cyclohexyl). Anal. (C₂₄H₂₈N₂O₃) C, H, N.

2-[(2-Cyclohexyl-1-phenylethyl)amino]-5-benzoxazolecarboxylic Acid (48). A mixture of 1.8 g of 47 (4.6 mmol), 20 mL of 2 N NaOH, and 30 mL of EtOH was refluxed for 2.5 h. The solvent was then removed in vacuo and the residue treated with water and 2 N HCl giving a white suspension. After stirring for 2h, the product was filtered, rinsed with water, and recrystallized from MeOH giving 1.15 g of 48 (69%), mp 232-234 °C. ¹H NMR (DMSO): δ 12.75 (s, 1H, CO₂H), 8.7 (d, 1H, NH), 7.7 (m, 2H, Ar), 7.4 (m, 6H, Ar), 4.9 (m, 1H, CH), 1.9–0.8 (m, 13H, CH₂, cyclohexyl). Anal. (C₂₂H₂₄N₂O₃) C, H, N.

2-[(2-Cyclohexyl-1-phenylethyl)amino]-N,N-dimethyl-5benzoxazolecarboxamide (48). A solution of 1.6 g (4.4 mmol) of 48 in 10 mL of DMF was cooled on an ice bath. CDI (1.22 g, 7.5 mmol) was added in portions, and the reaction mixture was stirred an additional 20 min. Dimethylamine was bubbled through the reaction for 15 min. After stirring an additional 30 min, the reaction mixture was poured into 50 mL of ice water. The precipitated product was filtered, washed with H₂O, dried, and boiled with EtOAc and a trace of EtOH. The product was filtered and dried giving 1.13 g of 49 (66%), mp 194-196 °C. ¹H NMR (CDCl₃): δ 7.3 (m, 7H, Ar), 7.05 (d, 1H, Ar), 6.65 (d br, 1H, NH), 4.95 (m, 1H, CH), 3.0 (d, 6H, N, CH₃), 1.9–0.8 (m, 13H, CH₂, cyclohexyl). Anal. (C₂₄H₂₉N₃O₂) C, H, N.

2-[(2-Cyclohexyl-1-phenylethyl)amino]-5-benzoxazolecarboxamide (50). This was prepared by the same method as 49, using ammonia instead of dimethylamine (49%), mp 181– 183 °C. ¹H NMR (CDCl₃): δ 8.6 (d, 1H, NH), 7.9–7.0 (m, 10H, Ar, NH₂), 4.85 (m, 1H, CH), 1.9–0.8 (m, 13H, CH₂, cyclohexyl). Anal. (C₂₂H₂₅N₃O₂) C, H, N.

2-[(2-Cyclohexyl-1-phenylethyl)amino]-5-benzoxazolenitrile (51). A solution of 1.27 g of 50 (3.5 mmol) and 1.1 mL of Et₃N (7.7 mmol) was cooled to 5 °C, and 0.54 mL (3.85 mmol) of trifluoroacetic anhydride was added over 5 min. The reaction mixture was stirred at room temperature for 2 h, the reaction quenched with H₂O, the mixture concentrated, and the product taken up in EtOAc. After being washed with 1 N HCl, saturated Na₂CO₃ solution, and brine, the solution was dried (Na₂SO₄) and concentrated. The product was purified by flash chromatography on silica gel, eluting with CH₂Cl₂. After titration with hexane, 820 mg (68%) of 51 was obtained as a white solid, mp 166-168 °C. 'H NMR (CDCl₃): δ 7.35 (m, 8H, Ar), 6.7 (d br, 1H, NH), 5.0 (m, 1H, CH), 2.0-0.8 (m, 13H, CH₂, cyclohexyl). Anal. (C₂₂H₂₃N₃O) C, H, N.

N-(2-Cyclohexyl-1-phenylethyl)-5-(hydroxymethyl)benzoxazolamine (52). A solution of 785 mg 47 (2 mmol) in 10 mL of THF was added dropwise to a suspension of 0.17 g (4.5 mmol) of LiAlH₄ in 10 mL of THF. The reaction mixture was stirred at room temperature for 30 min and then heated to reflux for 3 h. The reaction mixture was cooled, and excess LiAlH₄ and Li salts were decomposed with a few drops of EtOH followed by THF-H₂O. The reaction mixture was filtered and the filtrate diluted with EtOAc, washed with H₂O and brine, dried (Na₂-SO₄), and concentrated. The product was stirred with petroleum ether and then filtered and dried giving 250 mg of 52 (34%), mp 187-189 °C. ¹H NMR (DMSO): δ 8.45 (d, 1H, Ar), 7.3 (m, 6H, Ar), 6.9 (d, 1H, Ar), 5.1 (t, 1H, OH), 4.85 (m, 1H, CH), 4.45 (d, 2H, CH₂), 1.9–0.85 (m, 13H, CH₂ cyclohexyl). Anal. $(C_{22}H_{26}N_2O_2)$ C, H, N.

5-Acetamido-N-[2-cyclohexyl-1-(2-pyridinyl)ethyl]-2-benzoxazolamine (53). A solution of 1.75 g (4.79 mmol) of 5-nitro-[2-cyclohexyl-1-(2-pyridinyl)ethyl]-2-benzoxazolamine in 17.5 mL of absolute EtOH with 200 mg of 10% Pd/C was hydrogenated on a Parr shaker overnight. The catalyst was filtered off with the aid of Celite. The filtrate was concentrated, and the residue was dissolved in ether and extracted with 2 N HCl. The acidic aqueous phase was washed with ether after which the pH was adjusted to 8 by the addition of saturated NaHCO₃. This aqueous phase was extracted with CH_2Cl_2 . The CH_2Cl_2 extracts were dried and concentrated to give a residue which was crystallized from EtOH/petroleum ether to afford 928 mg (58%) of 5-amino-[2-cyclohexyl-1-(2-pyridinyl)ethyl]-2-benzoxazolamine, mp 181-184 °C dec. ¹H NMR (CDCl₃): δ 0.97-1.8 (m, 13H, aliph), 3.6 (br, 2H, NH₂), 4.95 (m, 1H, CH), 6.35 (dd, 1H, Ar), 6.65 (d, 1H, Ar), 6.97 (d, 1H, Ar), 7.25–7.32 (m, 5H, Ph).

To a solution of 200 mg (0.6 mmol) of 5-amino-N-[2-cyclohexyl-1-(2-pyridinyl)ethyl]-2-benzoxazolamine and 66 mg (0.66 mmol) of Et₃N in 2 mL of DMF at 0 °C was added 52 mg (0.66 mmol) of acetyl chloride. After stirring at 0 °C for 1 h followed by 2 h at room temperature, the reaction mixture was diluted with EtOAc, washed with 2 N HCl, saturated NaHCO₃, and brine. The EtOAc phase was dried (Na₂SO₄) and concentrated to give 240 mg of a brown oil. Preparative TLC on silica gel in 5% MeOH/CH₂Cl₂(2×) gave 190 mg of solid. Upon recrystallization from EtOH/petroleum, 135 mg (60%) of 53 was obtained, mp 145-147 °C. ¹H NMR (DMSO-d₆): δ 0.9-1.8 (m, 13H, aliph), 2.0 (s, 3H, CH₃), 4.82 (m, 1H, CH), 7.1-7.5 (m, 7H, Ar), 8.45 (d, 1H, NH), 9.85 (s, 1H, NHCO). Anal. (C₂₃H₂₇N₃O₂) C, H, N.

N-[2-Cyclohexyl-1-(2-pyridinyl)ethyl]-5-methanesulfonamidobenzoxazolamine (54). To a solution of 200 mg (0.6 mmol) of 5-amino-N-[2-cyclohexyl-1-(2-pyridinyl)ethyl]-2-benzoxazolamine in 2 mL of DMF and 66 mg (0.66 mmol) of Et₃N at 0 °C was added 75 mg (0.66 mmol) of methane sulfonyl chloride. After stirring at 0 °C for 1 h followed by 2 h at room temperature, the reaction mixture was diluted with EtOAc. The EtOAc solution was washed with 2 N HCl, saturated NaHCO₃, and brine. The EtOAc phase was dried and concentrated to give 270 mg of brown solid. Preparative TLC on silica gel with 5% MeOH/ CH₂Cl₂ gave 180 mg of solid. Recrystallization from EtOH/ petroleum ether provided 166 mg (68% yield) of white crystalline product, mp 188-190 °C. ¹H NMR (DMSO-d_θ): δ 0.9-1.8 (m, 13H, aliph), 2.87 (s, 3H, CH₃), 4.85 (m, 1H, CH), 6.8 (d, 1H, Ar), 7.06 (d, 1H, Ar), 7.2-7.38 (m, 6H, Ar), 8.57 (d, 1H, NH), 9.46 (s, 1H, NHSO₂). Anal. (C₂₂H₂₇N₃O₃S) C, H, N, S.

Method F. (R)-5-Isopropyl-N-[1-phenyl-2-(N-piperidinyl)ethyl]-2-benzoxazolamine Dihydrochloride (68a). To a suspension of 9.55 g (69.6 mmol) of (R)-(-)-2-phenylglycinol in 50 mL of CH₂Cl₂ at 0 °C was added 19.2 mL (18.23 g, 83.5 mmmol) of di-tert-butyl dicarbonate under anhydrous conditions. The reaction mixture was removed from the ice bath and stirred for 2 h at room temperature. The reaction mixture was then concentrated, and the residue was treated with ether giving a white crystalline product which was filtered to afford a 13-g first crop and a 1.9-g second crop. A total of 14.8 g (90% yield) of (R)-(-)-N-(tert-butoxycarbonyl)-2-phenylglycinol was collected as a white crystalline solid, mp 135-138 °C. ¹H NMR (CDCl₃): δ 1.4 (s, 9H, tBu), 2.4 (br, 1H, OH), 3.83 (t, 2H, CH₂), 4.78 (m, 1H, CH), 5.3 (br, 1H, NH), 7.27 (m, 5H, Ar).

Methanesulfonyl chloride (4.78 g, 41.7 mmol) was added dropwise to a solution of 9.0 g (37.9 mmol) of (R)-(-)-N-(tertbutoxycarbonyl)-2-phenylglycinol and 5.8 mL (4.22 g, 41.7 mmol) of Et₃N in 90 mL of CH₂Cl₂at -10 °C under anhydrous conditions. After 1.5 h at 0 °C, the reaction was quenched with ice-H₂O. The aqueous phase was extracted with CH₂Cl₂. The combined CH₂-Cl₂ extracts were washed with dilute HCl, dilute Na₂CO₃, and brine and dried over anhydrous MgSO₄. After filtration, the filtrate was concentrated to afford 11.8 g of a white solid. The solid was recrystallized from CH₂Cl₂/petroleum ether to give 11.47 g (96%) of the mesylate as a white crystalline solid (96% yield), mp 112-114 °C. ¹H NMR (CDCl₃): δ 1.4 (s, 9H, tBu), 2.89 (s, 3H, CH₃SO₃), 4.41 (m, 2H, CH₂OSO₂), 5.0 (m, 1H, CH), 5.2 (br, 2H, NH), 7.3 (m, 5H, Ar).

A solution of 2.0 g (6.34 mmol) of the above mesylate and 2.7 g (31.7 mmol) of piperidine in 20 mL of freshly distilled THF was refluxed under argon for 5 h. The piperidinium sulfonate salt was precipitated by the addition of ether and removed by filtration. The filtrate was concentrated to give 1.9 g of oil. Flash column chromatography of the oil on silica gel and elution with 5% acetone in petroleum ether afforded 960 mg (50%) of (R)-N-(tert-butoxycarbonyl)-1-phenyl-2-piperidinylethylamine as a white crystalline solid, mp 83–84 °C. ¹H NMR (CDCl₃): δ 1.4 (m, 9H, tBu), 1.6 (m, 6H, 3 CH₂), 2.28 (m, 2H, CH₂N), 2.5 (m, 4H, CH₂NCH₂), 4.6 (m, 1H, CH), 5.5 (br, 1H, NH), 7.3 (m, 5H, Ar).

A solution of 940 mg (3.09 mmol) of the protected amine obtained above was stirred in 2 mL of TFA and 2 mL of CH₂Cl₂ at room temperature for 1.5 h under anhydrous conditions. The reaction was quenched with H₂O, and the pH was adjusted to 8 by the addition of saturated Na₂CO₃. The aqueous phase was extracted with CH₂Cl₂. The combined CH₂Cl₂ extracts were washed with brine, dried over anhydrous MgSO₄, filtered, and concentrated to give 600 mg (95%) of (R)-(-)-1-phenyl-2piperidinylethylamine as a yellow oil. Without further purification, this oily product was used directly in the following reaction. ¹H NMR (CDCl₃): § 1.4-1.6 (m, 6H, CH₂), 1.9 (s, 2H, NH₂), 2.3 (m, 4H, 2 CH₂N), 2.55 (m, 2H, CH₂N), 4.1 (dd, 1H, CH), 7.3 (m, 5H, Ar).

To a solution of 600 mg (2.94 mmol) of (R)-(-)-1-phenyl-2piperidinylethylamine in 3 mL of DMSO was added 130 mg (3.23 mmol) of powdered NaOH. After 10 min of stirring, 690 mg (3.52 mmol) of 2-chloro-5-isopropylbenzoxazole was added. The resulting mixture was stirred at 60 °C for 3 h. The reaction was quenched with H₂O after cooling. The aqueous phase was extracted with EtOAc. The combined EtOAc extracts were extracted with 2 N HCl. The combined HCl extracts were washed with EtOAc and then treated with 2 N NaOH to adjust the pH to 8–9. The alkaline aqueous phase was extracted with CH_2Cl_2 . The combined CH₂Cl₂ extracts were dried and concentrated to give 540 mg of an oily residue. Flash column chromatography on silica gel, eluting with 1% MeOH in CH₂Cl₂, provided 460 mg (46%) of oil. The oil was treated with ethereal HCl to give a precipitate which was recrystallized from CH₂Cl₂/ether to afford 500 mg of 68a (37%) as a white crystalline solid after drying in a vacuum at 50 °C overnight, mp 180 °C dec. ¹H NMR for the free base: δ 1.2 (d, 6H, 2 CH₃), 1.5-1.7 (m, 6H, 3 CH₂), 2.5-3.0 (m, 7H, CH of iPr, 3 CH₂N), 5.0 (dd, 1H, CH), 6.8-7.4 (m, 9H, Ar, NH). Anal. (C23H29N3O·2HCl·H2O) C, H, N.

The preparation of 68b (mp 185 °C dec) was carried out by following the same scheme and procedures, starting with (S)-(+)-phenylglycinol. The NMR spectrum was identical to that of 68a.

Method G. 5-Chloro-N-[1-phenyl-2-(S,S-dioxo-N-thiomorpholino)ethyl]benzoxazolamine (71). To a 0 °C solution of the HCl salt of thiomorpholine (4.72 g, 34 mmol) in 40 mL of AcOH was added 30% H₂O₂ (14 mL). After the addition was complete, the reaction mixture was warmed to 100 °C and stirred overnight. The mixture was concentrated under reduced pressure and then recrystallized from EtOH to afford the thiomorpholinedioxide (3.4 g, 40%), mp 250 °C dec. ¹H NMR (DMSO- d_6): δ 3.32 (d, 4H), 3.25 (d, 4H).

To a solution of the above thiomorpholinedioxide (0.40 g, 3.0 mmol) in 15 mL of toluene were added iPr₂NEt (0.34 g, 3.0 mmol) and α -chloroacetophenone (0.46 g, 3.0 mmol). The mixture was refluxed for 6 h and cooled and then eluted through a silica gel column with EtOAc:hexane (2:1) to obtain pure α -(S,S-dioxothiomorpholino)acetophenone (326.0 mg, 43%) as a white solid, a portion of which was taken on to the next reaction. ¹H NMR (CDCl₃): δ7.95 (d, 2H, ArH), 7.58–7.65 (m, 1H, ArH), 7.50 (t, 2H, ArH), 4.05 (s, 2H, CH₂CO), 3.13 (br q, 8H, CH₂).

To a solution of α -(S,S-dioxothiomorpholino)acetophenone (248 mg, 0.98 mmol) in MeOH (10 mL) and a few drops of CHCl₃ was added NH₄OAc (85.0 mg, 11.0 mmol) followed by a solution of NaCNBH₃ (200 mg, 3.2 mmol) in MeOH (1 mL). After the mixture was stirred for 2 days at room temperature, the reaction was quenched with concentrated HCl and the mixture basified with KOH (s) and diluted with H_2O and then extracted with $CHCl_3$ (5 × 20 mL). The combined organic layers were dried over Na_2SO_4 and evaporated and then eluted through a silica column with 10% MeOH-CH2Cl2 to yield 1-phenyl-2-(S,Sdioxothiomorpholino)ethylamine, 115 mg (46%). ¹H NMR (CDCl₃): § 7.30-7.41 (m, 5H, ArH), 4.20 (dd, 1H, CH), 2.90-3.21 (m, 8H), 2.55-2.72 (m, 2H), 2.0 (br s, 2H, NH₂).

A mixture of 2.5-dichlorobenzoxazole (44 mg, 0.23 mmol), 1-phenyl-2-(S,S-dioxothiomorpholino)ethylamine (60 mg, 0.24 mmol), and iPr2NEt (37 mg, 0.29 mmol) in 3 mL of dichloroethane was heated to reflux and stirred overnight. After cooling to room temperature, the mixture was concentrated, taken up in a small amount of CH₂Cl₂, and eluted through a silica gel column with EtOAc:hexane (3:2) to obtain an oil, which was triturated with Et₂O and dissolved in EtOAc and then precipitated with hexanes. Filtration and recrystallization from EtOAc afforded 44 mg (47%) of 71, mp 189-191 °C. ¹H NMR (CDCl₃): δ 7.26-7.42 (m, 6H, ArH), 7.21 (d, 1H, ArH), 7.0 (dd, 1H, ArH), 5.78 (d, 1H, NH), 4.98 (q, 1H, CH), 2.83-3.23 (m, 10H, CH₂). Anal. (C₁₉H₂₀ClN₃O₃S) C, H, N.

Chiral HPLC Separation of 26a. (-)- and (+)-N-(2-Cyclohexyl-1-phenylethyl)-2-benzoxazolamine (26b,c). An 80-mg sample of 26a was dissolved in 20 mL of mobile phase (hexane:iPrOH:Et₂NH, 990:10:1). Multiple injections of up to 5 mL on a semipreparative Chiralcel OG (Daicel) column (25 cm \times 20 mm) were made at ambient temperature, with a flow rate of 9.0 mL/min. The detector was set at 280 nM. The first fraction, (+)-26c, had a retention time of 11.7 min and was isolated in >99% purity (37 mg). The second fraction, (-)-26b, had a retention time of 14.87 min and was isolated in 99% purity (37 mg)

Chiral HPLC Separation of 58a. (-)- and (+)-5-Chloro-N-[2-(4-fluorophenyl)-1-(2-pyridyl)ethyl]-2-benzoxazolamine (58b,c). A 10-mg sample of 58a was dissolved in 5 mL of mobile phase (hexane:iPrOH:Et₂NH, 990:10:1). Multiple injections of up to 100 μ L on an analytical Chiralcel OD (Daicel) column (25 cm \times 4.6 mm) were made at ambient temperature, with a flow rate of 1.0 mL/min. The detector was set at 254 nM. The first fraction, (-)-58b, eluted at 22.35 min, while (+)-58c eluted at 29.78 min. A total of 3.8 and 3.0 mg, respectively, was isolated, both >99% pure.

Biological Methods. Inhibition of LTB, Biosynthesis in Human PMNs. Human polymorphonuclear leukocytes were obtained from the heparinized blood of healthy, medication-free donors by dextran sedimentation and Ficoll-Paque separation.³² Compound solutions were prepared from DMSO stock solutions and diluted into Dulbecco's phosphate-buffered saline without Ca²⁺ and Mg²⁺, by a robotic sample processor (Tecan/RSP 505).³³ Buffer solutions of compound and cells $(2.5 \times 10^6 \text{ cells/mL in})$ Dulbecco's PBS with Ca²⁺ and Mg²⁺) were incubated at 28 °C for 15 min with shaking prior to the addition of calcium ionophore A23187 (2.5 μ M final concentration). After 10 min, the reaction was terminated by adding EGTA (10 mM final concentration), the solution centrifuged (300g, 7 min, 10 °C), and the supernatant analyzed for LTB₄ by commercially available RIA reagents (Advanced Magnetics, Cambridge, MA, and New England Nuclear, Boston, MA).

Acknowledgment. We acknowledge Tracy Saboe and Scot Campbell for NMR support in structure determination, Regina Mucci for technical assistance in in vitro screening, and Edward Graham and Tapon Roy for statistical analysis.

References

- (1) Borgeat, P. Biochemistry of the Lipoxygenase Pathways in Neutrophils. Can. J. Physiol. Pharmacol. 1989, 67, 936-942.
- (2) Samuelsson, B.; Funk, C. D. Enzymes Involved in the Biosynthesis of Leukotriene B₄. J. Biol. Chem. 1989, 264, 19469-19472.
- (3) Musser, J. H.; Kreft, A. F. 5-Lipoxygenase: Properties, Pharmacology, and the Quinolinyl (bridged) aryl Class of Inhibitors. J. Med. Chem. 1992, 35, 2501-2513.
- Summers, J. B. Inhibitors of Leukotriene Biosynthesis. Drug News (4) Perspect. 1990, 3, 3517-3526. (5) Shaw, A.; Krell, R. D. Peptide Leukotrienes: Current Status of
- Research. J. Med. Chem. 1991, 34, 1235-1242. Ford-Hutchinson, A. W. LTB₄ in Inflammation. Crit. Rev.
- (6) Immunol. **1990**, 10, 1–12.
- Dahlen, S.-E.; Hedquist, P.; Hammerstrom, B.; Samuelsson, B. Leukotrienes are Potent Constrictors of Human Bronchi. Nature (London) 1980, 288, 484-486.

Benzoxazolamines and Benzothiazolamines

- (9) Marom, Z.; Shelhamer, J. H.; Bach, M. K.; Morton, D. R.; Kaliner, M. Slow Reacting Substances LTC₄ and LTD₄ Increase the Release of Mucus from Human Airways In Vitro. Am. Rev. Respir. Dis. 1982, 126, 449-451.
- (10) Kaye, M. G.; Smith, L. J. Effect of Inhaled Leukotriene (LT) D₄ on Airway Hyperreactivity in Normal Subjects: Comparison with Histamine and Platelet Activating Factor (PAF). *Clin. Res.* 1989, 37, 478A.
- (11) Arm, J. P.; Spur, B. W.; Lee, T. H. Leukotrienes C₄, D₄, and E₄ Enhance Histamine Responsiveness in Asthmatic Airways. Am. Rev. Resp. Dis. 1991, 144, 1053-1057.
- (12) Lewis, R. A.; Austen, K. F.; Soberman, R. J. Leukotrienes and Other Products of the 5-Lipoxygenase Pathway. Biochemistry and Relation to Pathobiology in Human Diseases. N. Engl. J. Med. 1990, 323, 645–654.
- (13) Wallace, J. L. 5-Lipoxygenase: A Rational Target for Therapy of Inflammatory Bowel Diseases? Trends Pharmacol. Sci. 1990, 11, 51-53.
- (14) Israel, E.; Dermarkarian, R.; Rosenberg, M.; Sperling, R.; Taylor, G.; Rubin, P.; Drazen, J. M. The Effects of a 5-Lipoxygenase Inhibitor on Asthma Induced by Cold, Dry Air. N. Engl. J. Med. 1990, 323, 1740-1744.
- Israel, E.; Drazen, J. M.; Perlman, H.; Cohn, J.; Rubin, P. A. Double-Blind Multicenter Study of Zileuton, a Potent 5-Lipoxygenase (5-LO) Inhibitor Versus Placebo in the Treatment of Spontaneous Asthma in Adults. J. Allergy Clin. Immunol. 1992, 89, 236.
 Hui, K. P.; Barnes, N. C. Lung Function Improvement in Asthma
- (16) Hui, K. P.; Barnes, N. C. Lung Function Improvement in Asthma with a Cysteinyl-Leukotriene Reception Antagonist. *Lancet* 1991, 337, 1062–1063.
- (17) Lazer E. S.; Farina, P. R.; Gundel, R. H.; Wegner, C. D. 2,6-Disubstituted-4-(2-arylethenyl)phenol 5-Lipoxygenase Inhibitors: Development and Profile of BI-L-239. Drugs Future 1991, 16, 641-646.
- (18) Reindeau, D.; Falgueyret, J.-P.; Guay, J.; Veda, N.; Yamamoto, S. Pseudoperoxidase activity of 5-Lipoxygenase Stimulated by Potent Benzofuranol and N-Hydroxyurea Inhibitors of the Lipoxygenase Reaction. *Biochem J.* 1991, 274, 287–292.
- (19) Crawley, G. C.; Dowell, R. I.; Edwards, P. N.; Foster, S. J.; McMillan, R. M.; Walker, E. R. H.; Waterson, D.; Bird, T. C. G.; Bruneau, P.; Girodeau, J. M. Methoxytetrahydropyrans. A New Series of Selective and Orally Potent 5-Lipoxygenase Inhibitors. J. Med. Chem. 1992, 35, 2600-2609.
- (20) Rouzer, C. A.; Ford-Hutchinson, A. W.; Morton, H. E.; Gillard, J. W. MK886, a Potent and Specific Leukotriene Biosynthesis Inhibitor Blocks and Reverses the Membrane Association of 5-Lipoxygenase in Ionophore-challenged Leukocytes. J. Biol. Chem. 1990, 265, 1436-1442.

- (21) Saunders, J.; MacLeod, K. M.; Merchant, K.; Showell, G. A.; Snow, R. J.; Street, L. J.; Baker, R. Ester Bioisosteres: Synthesis of Oxadiazolyl-1-azabicyclo[2.2.1.]heptanes as Muscarinic Agonists. J. Chem. Soc., Chem. Commun. 1988, 118-119.
- (22) Abraham, W. M. The Role of Eicosanoids in Allergen-Induced Early and Late Bronchial Responses in Allergic Sheep. Adv. Prostaglandin, Thromboxane, Leukotriene Res. 1990, 201–208.
- (23) Katz, L.; Cohen, M. S. Benzoxazole Derivatives. I. 2-Mercaptobenzoxazoles. J. Org. Chem. 1954, 19, 758-766.
- (24) Miao, C. K.; Sorcek, R.; Nagel, J. H. The Resolution of Racemic Amines by the Formation of Diastereomeric Amides with Amino Acids. Org. Prep. Proced. Int. 1992, 24, 87-91.
- (25) Lazer, E. Š.; Wong, H.-C.; Wegner, C. D.; Graham, A. G.; Farina, P. R. Effect of Structure on Potency and Selectivity in 2,6-Disubstituted 4-(2-arylethenyl)phenol Lipoxygenase Inhibitors. J. Med. Chem. 1990, 33, 1892-1898.
- (26) Farina, P. R.; Graham, A. G.; Hoffman, A. F.; Watrous, J. M.; Borgeat, P.; Nadeau, M.; Hansen, G.; Dinallo, R. M.; Adams, J.; Miao, C. K.; Lazer, E. S.; Parks, T. P.; Homon, C. A. BIRM 270: A Novel inhibitor of Arachidonate Release which Blocks Leukotriene B₄ and Platelet Activating Factor Biosynthesis in Human Neutrophils. J. Pharmacol. Exp. Ther., submitted for publication.
- (27) Holford, N. H. G.; Sheiner, L. B. Pharmacokinetic and Pharmacodynamic Modeling In Vivo. Crit. Rev. Bioeng. 1981, 5, 273-322.
- (28) Holford, N. H. G.; Sheiner, L. B. Kinetics of Pharmacologic Response. Pharmacol. Ther. 1982, 16, 143-166.
- (29) Campaigne, E.; Thompson, R. L.; Van Werth, J. E. Some Heterocyclic Aldehyde Thiosemicarbazones Possessing Anti-Viral Activity. J. Med. Chem. 1959, 1, 577-599.
- (30) Mihina, J. S.; Herbst, R. M. The Reaction of Nitriles with Hydrazoic Acid: Synthesis of Monosubstituted Tetrazoles. J. Org. Chem. 1950, 15, 1082-1092.
- (31) Rubottom, G. M.; Kim, C. Preparation of Methyl Ketones by the Sequential Treatment of Carboxylic Acids with Methyllithium and Chlorotrimethylsilane. J. Org. Chem. 1983, 48, 1550–1552.
- (32) Henson, P. M.; Zanolari, B.; Schwartzman, N. A.; Hong, S. R. Intracellular Control of Human Neutrophil Secretion. I. C5a-Induced Stimulus-Specific Desensitization and the Effects of Cytochalasin B. J. Immunol. 1978, 121, 851-855.
- (33) An alternate method for compound dilution if a robotic sample processor is unavailable is described as follows. Test compounds were solubilized in DMSO and diluted in Dulbecco's PBS without Ca²⁺ and Mg²⁺. Dilutions were carried out in siliconized glass tubes (Surfasil, Pierce) with solution transferred using siliconized pipette tips (Costar). To reduce precipitation of compounds, the initial dilution from the DMSO stock was placed into prewarmed buffer, with mixing by gentle inversion. Subsequent dilutions were mixed by vortexing.