

Accepted Manuscript

Synthesis of new chiral ferrocenyl P,N-ligands with a benzoxazole ring and their application in Ag-catalyzed asymmetric [3+2] cycloaddition

Sheng Yan, Cheng Zhang, Ya-Hui Wang, Zhong Cao, Zhuo Zheng, Xiang-Ping Hu

PII: S0040-4039(13)00751-X

DOI: <http://dx.doi.org/10.1016/j.tetlet.2013.05.003>

Reference: TETL 42903

To appear in: *Tetrahedron Letters*



Please cite this article as: Yan, S., Zhang, C., Wang, Y-H., Cao, Z., Zheng, Z., Hu, X-P., Synthesis of new chiral ferrocenyl P,N-ligands with a benzoxazole ring and their application in Ag-catalyzed asymmetric [3+2] cycloaddition, *Tetrahedron Letters* (2013), doi: <http://dx.doi.org/10.1016/j.tetlet.2013.05.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.


Graphical Abstract

To create your abstract, type over the instructions in the template box below.
Fonts or abstract dimensions should not be changed or altered.

Synthesis of new chiral ferrocenyl *P,N*-ligands with a benzoxazole ring and their application in Ag-catalyzed asymmetric [3+2] cycloaddition

Leave this area blank for abstract info.

Sheng Yan, Cheng Zhang, Ya-Hui Wang, Zhong Cao,* Zhuo Zheng and Xiang-Ping Hu*





Tetrahedron Letters
journal homepage: www.elsevier.com

Synthesis of new chiral ferrocenyl *P,N*-ligands with a benzoxazole ring and their application in Ag-catalyzed asymmetric [3+2] cycloaddition

Sheng Yan^{a,b}, Cheng Zhang^b, Ya-Hui Wang^b, Zhong Cao^{a,*}, Zhuo Zheng^b and Xiang-Ping Hu^{b,*}

^a School of Chemistry and Biological Engineering, Changsha University of Science and Technology, Changsha 410004, China

^b Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

Keywords:

P,N-ligand

benzoxazole

asymmetric catalysis

silver

cycloaddition

ABSTRACT

New chiral ferrocenyl *P,N*-ligands with a benzoxazole ring as the *N*-donor group have been synthesized from 2-aminophenol through a three-step transformation and successfully employed in the Ag-catalyzed asymmetric [3+2] cycloaddition of azomethine ylides with electron-deficient alkenes. High diastereoselectivities, excellent enantioselectivities and good yields have been achieved for a variety of substrates, demonstrating the potential of these new *P,N*-ligands in asymmetric catalysis.

2009 Elsevier Ltd. All rights reserved.

Asymmetric catalysis is one of the most powerful methods for the preparation of enantiomerically pure compounds. In this context, a number of metal complexes with hybrid chiral *P,N*-ligands bearing an oxazoline as the *N*-donor group have been recognized as powerful catalysts in a variety of asymmetric catalytic reactions such as hydrogenation, allylic alkylation, and so on.¹ In sharp contrast, few chiral *P,N*-ligands with an oxazole/benzoxazole ring as the *N*-donor moiety have been found to be efficient in catalytic asymmetric reactions presumably due to the lack of α -central chirality close to the *N*-donor atom as that in oxazoline ligands (Figure 1).² However, the increased rigidity caused by the presence of a benzoxazole ring in the ligand backbone may be advantageous in some cases to create an efficient chiral environment around the central metal. The most important task for the development of an efficient chiral phosphine-benzoxazole ligand should be to search for an appropriate chiral skeleton.





Figure 1. Structure of *P*-oxazoline and *P*-benzoxazole ligands


In our recent work, we have developed a series of chiral *P,N*-ligands based on chiral 1-ferrocenylethylamine skeleton. These ligands displayed excellent enantio-induction in the Pd-catalyzed asymmetric allylic alkylation,³ Ru-catalyzed asymmetric cyclopropanation,⁴ Cu-catalyzed asymmetric propargylic amination,⁵ Cu-catalyzed asymmetric [3+3] cycloaddition,⁶ and so on.⁷ The success of these ligands in asymmetric catalysis demonstrated the potential of 1-ferrocenylethylamine as the chiral ligand backbone. As an effort toward further exploration of the application of this chiral skeleton in asymmetric catalysis,⁸ herein we report the synthesis of a new type of chiral *P,N*-ligands with a benzoxazole ring as the *N*-donor moiety from 1-ferrocenylethylamine derivatives, which were found to be highly effective in the Ag(I)-catalyzed asymmetric [3+2] cycloaddition of azomethine ylides with some electron-deficient alkenes.



* Corresponding author. Tel.: +86-411-84379276; fax: +86-411-84684746; e-mail: xiangping@dicp.ac.cn

Scheme 1. Synthesis of chiral ferrocenyl *P,N*-ligands (R_c,S_a)-**1a-b**. Reagents and conditions: (a) KOH, EtOH/H₂O (5/1), refluxing, 3 h; (b) POCl₃/PCl₅, CH₂Cl₂, rt, 12 h; (c) Ac₂O, 110 °C, NH₃ or MeNH₂, MeOH, 80 °C; (d) *i*-Pr₂NEt, DMF, rt-50 °C, 12 h.

Table 1. Catalytic asymmetric [3+2] cycloaddition reaction of *N*-(4-chlorobenzylidene)glycine ethyl ester **6a** with *N*-phenylmaleimide **7**^a



entry	ligand	Ag salt	base	solvent	temperature	Yield (%) ^b	Endo/exo	Ee (%) ^d
1	1a	AgOAc	Et ₃ N	toluene	rt	85	>95/5	50
2	1b	AgOAc	Et ₃ N	toluene	rt	93	95/5	95
3	1c	AgOAc	Et ₃ N	toluene	rt	90	>95/5	87
4	1b	AgOTf	Et ₃ N	toluene	rt	97	90/10	27
5	1b	AgBF ₄	Et ₃ N	toluene	rt	82	92/8	64
6	1b	AgOAc	Et ₃ N	CH ₂ Cl ₂	rt	93	90/10	97
7	1b	AgOAc	Et ₃ N	THF	rt	93	93/7	74
8	1b	AgOAc	Et ₃ N	CH ₃ CN	rt	78	95/5	97
9	1b	AgOAc	Et ₃ N	Et ₂ O	rt	92	93/7	95
10	1b	AgOAc	-	toluene	rt	94	95/5	93
11	1b	AgOAc	<i>i</i> -Pr ₂ NEt	toluene	rt	92	>95/5	95
12	1b	AgOAc	DBU	toluene	rt	87	>95/5	31
13	1b	AgOAc	DABCO	toluene	rt	91	95/5	94
14	1b	AgOAc	Et ₃ N	toluene	40	94	95/5	94
15	1b	AgOAc	Et ₃ N	toluene	0	93	95/5	96
16 ^e	1b	AgOAc	Et ₃ N	toluene	-10	89	>95/5	97
17 ^f	1b	AgOAc	Et ₃ N	toluene	-20	94	>95/5	98

^a Reactions were performed in 3 mL of solvent with 0.5 mmol of substrate **6a**, 10 mol% of base additives, and 3 mol% of catalyst prepared in situ from Ag salts and 1.1 equiv of ligand at indicated temperature for 2 h


^b Isolated yields.

^c *endo/exo*-Ratio was determined by ¹H NMR.

^d Enantiomeric excesses were determined by chiral HPLC.

^e Reaction time, 6 h.

^f Reaction time, 12 h.



Scheme 2. Synthesis of chiral ferrocenyl *P,N*-ligand (R_c,S_a)-**1c**. Reagents and conditions: (a) Ac₂O, 110 °C, 3 h, and then LiAlH₄, THF, reflux for 12 h; (b) *n*-BuLi, Et₂O, and then reflux for 12 h.

Chiral phosphine-benzoxazole ligands **1a-b** were synthesized from 2-aminophenol according to the procedure shown in Scheme 1. Initially, 2(3*H*)-benzoxazolethione (**3**) was prepared as a white needle in 82% yield by the reaction of 2-aminophenol with CS₂ in refluxing aqueous EtOH for 3 hours.⁹ The treatment of 2(3*H*)-benzoxazolethione with POCl₃ and PCl₅ in CH₂Cl₂ at room temperature generated 2-chlorobenzoxazole **4** as a colorless oil in 60% yield.¹⁰ Finally, 2-chlorobenzoxazole **4** was converted


into the desired phosphine-benzoxazole ligands **1a-b**¹¹ in high yields by the reaction with chiral phosphine-amine intermediates, (R_c,S_p)-PPFNHR,¹² under the basic condition.

For a comparison, (R_c,S_p)-**1c** with a oxygen linker was also prepared in a similar procedure as shown in Scheme 2.

With these newly developed ferrocenyl phosphine-benzoxazole ligands in hand, we then examined their efficiency in the Ag-catalyzed asymmetric [3+2] cycloaddition of azomethine ylides with *N*-phenylmaleimide.¹³ Initially, the cycloaddition of *N*-(4-chlorobenzylidene)glycine ethyl ester (**6a**) with *N*-phenylmaleimide **7** was selected as a model reaction. The reaction was performed in Et₂O at rt in the present of a catalytic amount of Et₃N (10 mol%) and the catalyst (3 mol%) *in situ* prepared from silver salts and 1.1 equiv of ferrocenyl phosphine-benzoxazole ligands **1a-c**,¹⁴ and the results are summarized in Table 1. In comparison with ligand **1a** bearing a *N*-H proton, ligand **1b** with a *N*-methyl substituent showed significantly improved enantioselectivity in this model reaction, giving the *endo*-cycloadduct **8a** in good diastereoselectivity (95/5) with

enantioselectivity of up to 95% ee (entry 1 *vers* entry 2). With (R_c, S_p)-**1c** as ligand, the reaction gave the cycloadducts in slightly decreased enantioselectivity (87% ee) but with the same configuration as that obtained with (R_c, S_p)-**1a** (entry 3). This result suggested that in this class of P, N -ligands, the N -atom in the benzoxazole ring rather than the one directly attached to the stereogenic carbon center is the actual N -donor atom. The reaction proved to be highly sensitive to the catalyst precursors. Thus, the use of AgOTf and AgBF₄ resulted in dramatically decreased enantioselectivities (entries 4 and 5). The nature of the solvent has some influence on the conversion, diastereoselectivity and enantioselectivity (entries 6-9). Further work on the screening of various bases indicated that Et₃N was the best choice (entries 10-13). The reaction temperature had some impact on the reactivity and enantioselectivity, and the reaction performed at lower temperature tended to give better enantioselectivity (entries 14-17). When the reaction temperature was carried out at -20 °C, up to 98% ee was achieved (entry 17).

Table 2. Catalytic asymmetric [3+2] cycloaddition of imino esters **6a-l** with N -phenylmaleimide **7**^a



Entry	Substrate (R)	Yield (%) ^b	<i>endo/exo</i> ^c	ee (%) ^d
1	6a : 4-ClC ₆ H ₄	94	95/5	98
2	6b : 3-ClC ₆ H ₄	94	>95/5	95
3	6c : 2-ClC ₆ H ₄	92	>95/5	98
4	6d : Ph	90	>95/5	97
5	6e : 4-FC ₆ H ₄	89	94/6	98
6	6f : 4-CF ₃ C ₆ H ₄	87	94/6	97
7	6g : 4-BrC ₆ H ₄	94	>95/5	95
8	6h : 4-MeC ₆ H ₄	85	92/8	>99
9	6i : 4-MeOC ₆ H ₄	80	90/10	98
10	6j : 1-naphthyl	85	95/5	98
11	6k : 2-thienyl	75	>95/5	88
12	6l : cyclohexyl	80	93/7	91

^a Reactions were performed in 3 mL of toluene with 0.5 mmol of substrate **6a-j**, 10 mol% of Et₃N, and 3 mol% of catalyst prepared *in situ* from AgOAc and 1.1 equiv of **1b** at -20 °C for 12 h

^b Isolated yields.

^c *endo/exo*-Ratio was determined by ¹H NMR.

^d Enantiomeric excesses were determined by chiral HPLC.

Under the optimized reaction conditions, the scope of the present catalytic system in the 1,3-dipolar cycloaddition was carried out. As shown in Table 2, various imino esters **6a-i** prepared from various substituted benzaldehydes reacted with N -phenylmaleimide **7** to give the corresponding *endo*-adducts **8a-i** in high yields and good to perfect enantioselectivities (95->99% ee) (entries 1-9). The results disclosed that the electronic property of the substituent on the phenyl ring had little effect in the enantioselectivities, but had some influence on the diastereoselectivities (entries 1, 4-9). The substrates bearing an electron-withdrawing substituent tended to give higher

diastereoselectivities than those with an electron-donating group (entries 8-9 *vers* entries 5-7). An azomethine ylide derived from 1-naphthylaldehyde also proved to be a suitable substrate in this transformation, producing *endo*-cycloadduct in 98% ee (entry 10). 2-Thienyl substrate **6k** was also efficient, giving *endo*-cycloadduct in 88% ee (entry 11). More importantly, the presence catalytic system showed good reactivity for alkyl substituted substrate **6l**, giving *endo*-cycloadduct in 91% ee (entry 12).

To further extend the scope of this catalytic system, the scope and generality of this catalytic system with regard to iminoesters and dipolarophiles were also investigated (Figure 1). The present catalytic system is remarkably tolerant and remains high in reactivity for azomethine ylides derived from amino esters other than glycinate. Thus, under the optimized reaction conditions, the azomethine ylide derived from phenylalanine successfully reacted with N -phenylmaleimide **7**, leading to perfect *endo*-selectivities (>95/5) and excellent enantioselectivities (96% ee). A variety of other dipolarophiles was also subjected to this reaction. With N -methylmaleimide, high *endo*-selectivity (*endo/exo* > 95/5) and good enantioselectivity (94% ee) was achieved. However, for dimethyl maleate and *tert*-butyl acrylate, the present catalytic system proved to be less efficient, displaying very low enantioselectivity.




Figure 1. Cycloadducts of the reaction between imino esters with other dipolarophiles.

In conclusion, we have developed a new family of chiral ferrocenyl P, N -ligands with a benzoxazole fragment through an efficient and simple synthesis route. These ligands were evaluated in the Ag(I)-catalyzed asymmetric [3+2] cycloaddition of azomethine ylides with various electronic-deficient alkenes, in which high *endo*-selectivities and good to excellent enantioselectivities (up to >99% ee) were achieved. Further applications of these new ligands in other type of asymmetric reactions are in progress.

Acknowledgments

Support for this research from the Dalian Institute of Chemical Physics (CAS). The authors also thank Prof. Hongchao Guo for providing several chiral materials (synthesized in the National Key Technologies R&D Program of China, 2012BAK25B03, CAU) for ligand synthesis.

References and notes

- (a) Bausch, C. C.; Pfaltz, A. In *Privileged Chiral Ligands and Catalysts*; Zhou, Q.-L., Ed.; Wiley-VCH: Weinheim, 2011; pp 221-256; (b) Tannert, R.; Pfaltz, A. In *Phosphorus (III) Ligands in Homogeneous Catalysis: Design and Synthesis*; Kamer, P. C. J.; van Leeuwen, W. N. M., Eds.; Wiley, 2012; pp 235-260.
- (a) Mazuela, J.; Paptchikhine, A.; Tolstoy, P.; Pamies, O.; Dieguez, M.; Andersson, P. G. *Chem. Eur. J.* **2010**, *16*, 620-638; (b) Mazuela, J.; Paptchikhine, A.; Pamies, O.; Andersson, P. G.; Dieguez, M. *Chem. Eur. J.* **2010**, *16*, 4567-4576; (c) Mazuela, J.; Tolstoy, P.; Pamies, O.; Andersson, P. G.; Dieguez, M. *Org. Biomol. Chem.* **2011**, *9*, 941-946.
- (a) Hu, X.; Dai, H.; Hu, X.; Chen, H.; Wang, J.; Bai, C.; Zheng, Z. *Tetrahedron: Asymmetry* **2002**, *13*, 1687-1693; (b) Hu, X.; Chen, H.; Hu, X.; Dai, H.; Bai, C.; Wang, J.; Zheng, Z. *Tetrahedron Lett.* **2002**, *43*, 9179-9182; (c) Hu, X.; Dai, H.; Bai, C.; Chen, H.; Zheng, Z. *Tetrahedron: Asymmetry* **2004**, *15*, 1065-1068; (d) Hu, X.-P.; Chen, H.-L.; Zheng, Z. *Adv. Synth. Catal.* **2005**, *347*, 541-548.
- Dai, H.; Hu, X.; Chen, H.; Bai, C.; Zheng, Z. *J. Mol. Catal., A: Chem.* **2004**, *211*, 17-21.
- Zhang, C.; Wang, Y.-H.; Hu, X.-H.; Zheng, Z.; Xu, J.; Hu, X.-P. *Adv. Synth. Catal.* **2012**, *354*, 2854-2858
- Zhang, C.; Hu, X.-H.; Wang, Y.-H.; Zheng, Z.; Xu, J.; Hu, X.-P. *J. Am. Chem. Soc.* **2012**, *134*, 9585-9588.
- (a) Dai, H.; Hu, X.; Chen, H.; Bai, C.; Zheng, Z. *Chin. J. Catal.* **2003**, *24*, 565-566; (b) Yu, S.-B.; Hu, X.-P.; Deng, J.; Huang, J.-D.; Wang, D.-Y.; Duan, Z.-C.; Zheng, Z. *Tetrahedron Lett.* **2008**, *49*, 1253-1256.
- (a) Hu, X.-P.; Zheng, Z. *Org. Lett.* **2004**, *6*, 3585-3588; (b) Hu, X.-P.; Zheng, Z. *Org. Lett.* **2005**, *7*, 419-422; (c) Deng, J.; Duan, Z.-C.; Huang, J.-D.; Hu, X.-P.; Wang, D.-Y.; Xu, X.-F.; Zheng, Z. *Org. Lett.* **2007**, *9*, 4825-4828; (d) Deng, J.; Hu, X.-P.; Huang, J.-D.; Yu, S.-B.; Wang, D.-Y.; Duan, Z.-C.; Zheng, Z. *J. Org. Chem.* **2008**, *73*, 2015-2017; (e) Wang, D.-Y.; Hu, X.-P.; Deng, J.; Yu, S.-B.; Duan, Z.-C.; Zheng, Z. *J. Org. Chem.* **2009**, *74*, 4408-4410; (f) Duan, Z.-C.; Hu, X.-P.; Zhang, C.; Zheng, Z. *J. Org. Chem.* **2010**, *75*, 8319-8321.
- Chen, W.-B.; Jin, G.-Y. *Heteroatom Chem.* **2001**, *12*, 151-155.
- Lazer, E. S.; Miao, C. K.; Wong, H.-C.; Sorcek, R.; Spero, D. M.; Gilman, A.; Pal, K.; Behnke, M.; Graham, A. G.; Watrous, J. M.; Homon, C. A.; Nage, J.; Shah, A.; Guindon, Y.; Farina, P. R.; Adams, J. J. *Med. Chem.* **1994**, *37*, 913-923.
- Selected physical and spectral data for (R_c, S_p)-**1a**: yellow solid; mp: 120-124 °C; $[\alpha]_D^{25}$: -304.6 (c 0.50, CHCl₃); ¹H NMR (400 MHz, CDCl₃): δ 1.53 (d, $J = 6.4$ Hz, 3H), 3.79 (s, 1H), 4.02 (s, 5H), 4.30 (s, 1H), 4.50 (s, 1H), 5.11-5.15 (m, 1H), 5.71 (br, 1H), 6.94-6.98 (m, 1H), 7.06-7.13 (m, 7H), 7.32-7.35 (m, 4H), 7.48-7.52 (m, 2H); ³¹P NMR (162 MHz, CDCl₃): δ -24.6; ¹³C NMR (100 MHz, CDCl₃): δ 22.4, 48.9 (d, $J = 4.1$ Hz), 69.2, 69.9, 70.2 (d, $J = 4.4$ Hz), 72.1 (d, $J = 4.2$ Hz), 95.1, 108.5, 116.2, 120.4, 123.5, 128.0, 128.1, 128.2, 129.1, 132.6 (d, $J = 18.8$ Hz), 134.9 (d, $J = 20.6$ Hz), 136.6 (d, $J = 7.6$ Hz), 138.7 (d, $J = 8.3$ Hz), 143.1, 148.3, 160.4; HRMS (m/z) calcd. for C₃₁H₂₇FeN₂OP: 530.1210, found: 530.1209.
Selected physical and spectral data for (R_c, S_p)-**1b**: yellow needle; mp: 154-156 °C; $[\alpha]_D^{25}$: -364.1 (c 0.50, CHCl₃); ¹H NMR (400 MHz, CDCl₃): δ 1.59 (d, $J = 7.2$ Hz, 3H), 2.33 (s, 3H), 3.76 (s, 1H), 4.13 (s, 5H), 4.32 (s, 1H), 4.50 (s, 1H), 5.81-5.83 (m, 1H), 6.81-6.86 (m, 4H), 6.89-6.93 (m, 1H), 7.00-7.09 (m, 3H), 7.28-7.30 (m, 4H), 7.44-7.48 (m, 2H); ³¹P NMR (162 MHz, CDCl₃): δ -25.7; ¹³C NMR (100 MHz, CDCl₃): δ 18.3, 28.6, 51.4 (d, $J = 8.9$ Hz), 68.7, 69.9 (d, $J = 3.5$ Hz), 70.0, 72.1 (d, $J = 4.5$ Hz), 77.8 (d, $J = 10.2$ Hz), 91.9, 92.1, 108.3, 115.7, 119.5, 123.3, 127.6 (d, $J = 7.2$ Hz), 128.0 (d, $J = 17.4$ Hz), 128.1, 128.7, 132.5, 132.7, 134.4, 134.6, 136.8 (d, $J = 9.9$ Hz), 137.8 (d, $J = 9.5$ Hz), 143.5, 148.6, 161.0. HRMS (m/z) calcd. for C₃₂H₂₉FeN₂OP: 544.1367, found: 544.1367.
Selected physical and spectral data for (R_c, S_p)-**1c**: yellow solid; mp: 196-198 °C; $[\alpha]_D^{25}$: -346.6 (c 0.53, CH₂Cl₂); ¹H NMR (400 MHz, CDCl₃): δ 1.85 (d, $J = 7.2$ Hz, 3H), 3.81 (s, 1H), 4.17 (s, 5H), 4.43 (s, 1H), 4.83 (s, 1H), 5.91-5.93 (m, 1H), 6.68-6.93 (m, 9H), 7.29-7.48 (m, 5H); ³¹P NMR (162 MHz, CDCl₃): δ -26.4; ¹³C NMR (100 MHz, CDCl₃): δ 16.9, 48.6 (d, $J = 8.0$ Hz), 69.0, 70.1 (d, $J = 3.5$ Hz), 70.7, 72.7 (d, $J = 4.5$ Hz), 77.8 (d, $J = 10.2$ Hz), 89.8, 90.1, 109.5, 109.7, 121.3, 122.7, 127.4 (d, $J = 6.9$ Hz), 127.8, 128.0 (d, $J = 7.6$ Hz), 129.1, 131.9 (d, $J = 20$ Hz), 134.7 (d, $J = 20$ Hz), 136.4 (d, $J = 9.1$ Hz), 136.8 (d, $J = 9.2$ Hz), 142.3, 152.7. HRMS (m/z) calcd. for C₃₁H₂₆FeNO₂P: 531.1051, found: 531.1047.
- (a) Hayashi, T.; Hayashi, C.; Uozumi, Y. *Tetrahedron: Asymmetry* **1995**, *6*, 2503-2506; (b) Boaz, N. W.; Debenham, S. D.; Mackenzie, E. B.; Large, S. E. *Org. Lett.* **2002**, *4*, 2421-2424.
- For recent reviews about 1,3-dipolar cycloaddition reactions of azomethine ylides, see: (a) Engels, B.; Christl, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 7968-7970; (b) Stanley, L. M.; Sibi, M. P. *Chem. Rev.* **2008**, *108*, 2887-2902; (c) Alvarez-Corral, M.; Munoz-Dorado, M.; Rodriguez-Garcia, I. *Chem. Rev.* **2008**, *108*, 3174-3198; (d) Adrio, J.; Carretero, J. C. *Chem. Commun.* **2011**, *47*, 6784-6794. For recent examples, see: (e) Kim, H. Y.; Li, J.-Y.; Kim, S.; Oh, K. *J. Am. Chem. Soc.* **2011**, *133*, 20750-20753; (f) He, L.; Chen, X.-H.; Wang, D.-N.; Luo, S.-W.; Zhang, W.-Q.; Yu, J.; Ren, L.; Gong, L.-Z. *J. Am. Chem. Soc.* **2011**, *133*, 13504-13518; (g) Reboredo, S.; Reyes, E.; Vicario, J. L.; Badia, D.; Carrillo, L.; C_ozar, A.; Cosst' o, F. P. *Chem.; Eur. J.* **2012**, *18*, 7179-7188; (h) Wang, M.; Wang, C.-J.; Lin, Z. *Organometallics* **2012**, *31*, 7870-7876; (i) Liu, T.-L.; He, Z.-L.; Tao, H.-Y.; Wang, C.-J. *Chem. Eur. J.* **2012**, *18*, 8042-8046; (j) Jing, X.; He, C.; Dong, D.; Yang, L.; Duan, C. *Angew. Chem. Int. Ed.* **2012**, *51*, 10127-10131; (k) He, Z.; Liu, T.; Tao, H.; Wang, C.-J. *Org. Lett.* **2012**, *14*, 6230-6233; (l) Han, M.-L.; Wang, D.-Y.; Zeng, P.-W.; Zheng, Z.; Hu, X.-P. *Tetrahedron: Asymmetry* **2012**, *23*, 306-312.
- General procedure for catalytic asymmetric 1,3-dipolar cycloaddition of azomethine ylides. Under nitrogen atmosphere, a solution of AgOAc (0.015 mmol) and ligand **1b** (0.0165 mmol) in 3 mL of toluene was stirred at room temperature for 1 h. The solution was cooled to -20 °C, and then added imino esters (0.5 mmol), Et₃N (7.5 μL, 0.05 mmol) and *N*-phenylmaleimide **7** (0.6 mmol) successively. After 12 h, the mixture was passed through a short column of silica gel and the diastereometric ratio (*endo/exo*) was determined by the NMR spectroscopic analysis of the crude product. The residue was then purified by column chromatography on silica gel, and was submitted to ee analysis by HPLC with a chiral column.