

Article

## Discovery of Novel Dual Inhibitors of Wild-type and the Most Prevalent Drug-Resistant Mutant, S31N, of M2 Proton Channel from Influenza A Virus

Jizhou Wang, Chunlong Ma, Jun Wang, Hyunil Jo, Belgin Canturk, Giacomo Fiorin, Lawrence H. Pinto, Robert A. Lamb, Michael L. Klein, and William F. DeGrado

*J. Med. Chem.*, **Just Accepted Manuscript** • DOI: 10.1021/jm301538e • Publication Date (Web): 25 Feb 2013

Downloaded from <http://pubs.acs.org> on March 25, 2013

### Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



**ACS Publications**  
High quality. High impact.

Journal of Medicinal Chemistry is published by the American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036  
Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

# Discovery of Novel Dual Inhibitors of Wild-type and the Most Prevalent Drug-Resistant Mutant, S31N, of M2 Proton Channel from Influenza A Virus

Jizhou Wang<sup>\*,a</sup>, Chunlong Ma<sup>b</sup>, Jun Wang<sup>c</sup>, Hyunil Jo<sup>c</sup>, Belgin Canturk<sup>d</sup>, Giacomo Fiorin<sup>e</sup>,  
Lawrence H. Pinto<sup>b</sup>, Robert A. Lamb<sup>f,g</sup>, Michael L. Klein<sup>e</sup>, and William F. DeGrado<sup>\*,c</sup>

<sup>a</sup> Influmedix Inc., 170 North Radnor-Chester Road, Suite 300, Radnor, PA 19087


<sup>b</sup>Department of Molecular Biosciences, Northwestern University, Evanston, Illinois 60208

<sup>c</sup>Department of Pharmaceutical Chemistry, University of California, San Francisco, CA 94158

<sup>d</sup>Department of Chemistry, University of Pennsylvania, 231 South 34th St., Philadelphia, PA 19104


<sup>e</sup>Institute for Computational Molecular Science and Department of Chemistry, Temple University, Philadelphia, Pennsylvania 19122-6078

<sup>f</sup>Department of Molecular Biosciences, and <sup>g</sup>Howard Hughes Medical Institute, Northwestern University, Evanston, Illinois 60208-3500



**Abstract:** Anti-influenza drugs, amantadine and rimantadine, targeting the M2 channel from influenza A virus are no longer effective because of widespread drug resistance. S31N is the predominant and amantadine-resistant M2 mutant, present in almost all of the circulating influenza A strains as well as in the pandemic 2009 H1N1 and the highly pathogenic H5N1 flu strains. Thus, there is an urgent need to develop second-generation

M2 inhibitors targeting S31N mutant. However, S31N mutant presents a huge challenge to drug discovery, and it has been considered undruggable for several decades. Using structural information, classical medicinal chemistry approaches and M2-specific biological testing, we discovered benzyl-substituted amantadine derivatives with activity against both S31N and WT, among which 4-(adamantan-1-ylaminomethyl)-benzene-1,3-diol is the most potent dual inhibitor. These inhibitors demonstrate that S31N is a druggable target and provide a new starting point to design novel M2 inhibitors that address the problem of drug-resistant influenza A infections.



**Figure 1.** Structures of four approved antiviral drugs for influenza A

## INTRODUCTION

Seasonal influenza infections as well as the emergence of life-threatening strains of influenza are a major worldwide health concern. Every year, influenza epidemics cause numerous deaths and millions of hospitalizations.<sup>1,2</sup> Besides the yearly epidemic outbreaks, influenza viruses are even more threatening pathogens due to their potency to cause pandemics, as occurred in 2009 with the emergence and worldwide spread of pandemic H1N1. Available prophylactic vaccines are not completely effective against emerging flu strains and thus effective anti-viral therapy is an essential component in the fight against influenza A infections. Currently, there are only four antivirals approved for use against influenza virus infections in humans, two targeting the M2 proton channel, amantadine (oral) and rimantadine (oral) and two targeting the neuraminidase, oseltamivir (oral), and zanamivir (inhaled) (Fig.1). Resistance to both classes is a problem: resistance to the


adamantane class of drugs is now so pervasive that the Centers for Disease Control and Prevention (CDC) has advised against its continued use<sup>2,3</sup> and widespread resistance to oseltamivir has been increasingly reported since 2007-2008.<sup>4-8</sup> Even more concerning are the reports of the emergence of flu strains with dual resistance to both adamantane and neuraminidase classes of antivirals.<sup>9</sup>

Resistance to M2 proton channel drugs is associated with mutations in the transmembrane domain of the M2 protein. The homotetrameric structure of the M2 channel places constraints on the types of drug-resistant mutations that can be accommodated. A single mutation to the pore-lining residues of the protein results in four changes to the residues lining the pore, where the drug binds. While many mutations can be tolerated with retention of function *in vitro*,<sup>10-12</sup> only a few mutants appear to be tolerated in highly transmissible viruses seen in the wild. V27A, L26F, and S31N have consistently comprised more than 99% of transmissible M2 mutants; among which S31N is now predominant in 98-100% of the transmissible amantadine-resistant H1N1, H5N1, and H3N2 strains isolated from humans, birds and swine in the last decade.<sup>1</sup> As a result, there is an urgent need to develop second-generation novel M2 inhibitors targeting the most prevalent S31N mutant. However, this mutation presents significant challenges to drug discovery, because the side chain of Asn31 partially fills the amantadine-binding site and also changes its physical properties markedly.

M2 inhibitors generally consist of a hydrophobic portion (adamantane in the drugs amantadine and rimantadine) connected to a polar group, which is generally a positively charged substituent such as an amine in amantadine or an aminoethyl group in rimantadine (Fig. 1). A variety of scaffolds can be substituted for the hydrophobic adamantane group, including fused and spiro-fused multi-cyclic alkanes, acyclic branched alkanes, and silanes.<sup>13, 14</sup> These compounds showed excellent activity for WT, and some were highly active against V27A and L26F.<sup>14, 15</sup> However, none showed significantly better activity vs. S31N than had already been seen for amantadine. The polar group has also been the subject of extensive structure activity relations (SAR). Again, a


variety of substituents were tolerated when the target was WT, but only the simple primary amine found in amantadine gave significant activity against S31N. Thus, after several decades of medicinal chemistry investigations, no progress had been made towards inhibiting this predominant mutant form, suggesting that this variant might be essentially *undruggable*.<sup>16-20</sup>

Our journey to S31N inhibitors discovery started with modification of amantadine (**1**) whose potency against S31N drops approximately 12-fold from 16  $\mu\text{M}$  against WT<sup>21</sup> to 200  $\mu\text{M}$ <sup>22</sup> in a TEVC (two electrode voltage clamp) electrophysiological assay. Amantadine (**1**) is, however, more active than rimantadine (**2**) against S31N (199.9  $\mu\text{M}$  vs > 2 mM)<sup>23</sup> (Fig. **2A**). We envisioned that the S31N inhibitory activity could be restored if we captured additional interactions within the channel. Recent crystallographic and NMR studies have provided major advances in our understanding of the mode of inhibition of M2 by amantadine and related drugs (Fig. **2B** and **2C**).<sup>15,21,24-33</sup>



**Figure 2.** (A) Activity of amantadine and rimantadine against WT (S31) and S31N mutant in the TEVC assay. (B) The M2 channel-lining residues found in transmissible drug-resistant mutants are shown in orange letters. Amantadine (orange bound drug) causes a break in the aqueous path (water molecules, red balls). One helix has been removed for clarity. (C) Snapshot from a simulation of amantadine with WT<sup>15</sup>. Water molecules (red) associate with carbonyl groups (green/red sticks) in a square planar array. This array of water molecules can stabilize the bound ammonium group of amantadine (green and blue bound drug) or a centrally located water molecule (magenta).




The location of the pharmacologically relevant drug binding site in M2 was extensively debated until recently. Our earlier crystallographic structures suggested a pore-binding model,<sup>24</sup> while Chou and co-workers proposed a more peripheral site on the surface of the protein based on NMR studies.<sup>25</sup> However, last year, Chou and co-workers solved a structure, showing that drug binds to the site we had originally identified, ending the controversy, and validating our earlier crystallographic structures.<sup>31</sup> Specifically, the drug binds in a pocket within the channel lumen, and is surrounded by the apolar side chain of Val27, Ala30, the C-beta of Ser31, and the C-alpha of Gly34. The ammonium group projects downward towards the invariant His37 residue, but does not actually interact with this residue. Instead it is stabilized by water molecules, which are in turn stabilized by interactions with backbone carbonyl groups lining the pore.<sup>15</sup> The mutation S31N disrupts the size and polarity of the pore in precisely the region that accommodates the adamantane group of amantadine, explaining the decrease in potency of amantadine from 16  $\mu$ M against WT to 200  $\mu$ M in a TEVC electrophysiological assay. Moreover, NMR studies showed that S31N was more dynamic and possibly also more hydrated in the pore than the corresponding WT protein.<sup>34</sup> This led us to consider strategies in which we either increased the polarity or dimensions of the adamantane core to capture more interactions with the backbone of the channel (Fig. 3, approach I). We synthesized over 300 amantadine analogues and other diverse cores, some of which have been published.<sup>14,15,22,35-38</sup> While we succeeded in inhibiting V27A and L26F, none of these compounds inhibited S31N.<sup>15</sup>



**Figure 3.** Two strategies to overcome drug-resistance

We turned our attention to the strategy of introducing additional groups (warhead groups) to the amine, in order to enhance affinity for S31N as well as WT by introducing direct/water mediated hydrogen binding /electrostatic interactions with backbone or the groups lining the pore of the M2 channel (Fig. 3, approach II). Examination of the structure of M2's channel showed there was sufficient space in the channel to accommodate additional substituents to mediate these interactions. However, this line of modification of amantadine (**1**) was not obvious; primary amines tended to be more potent than secondary amines and many early substitutions of the primary amines decreased activity towards the WT channel.<sup>14</sup> Nevertheless, two reports had shown that aromatic groups could be appended to the amines of 1-adamantanemethylamine (a rimantadine (**2**) derivative)<sup>39</sup> and pinanamine (**4**)<sup>40</sup> and showed activity against M2 channels. We therefore synthesized the most potent inhibitors in these two series, compounds **3**<sup>39</sup> and **4**<sup>40</sup>, and found that they were active against WT (21% and 91% inhibition for **3** and **4**, respectively) (Table 1). However, they showed no activity against S31N (0% inhibition for both) in our electrophysiological TEVC assay. Since amantadine (**1**) is more potent against S31N (35% inhibition) than rimantadine (**2**) (13% inhibition) and pinanamine (0% inhibition<sup>23</sup>) in a TEVC assay, we envisioned that installation of aromatic group to the amine group of amantadine (**5-49**) might improve activity against S31N as well as WT. This led to the discovery of benzyl-substituted amantadine derivatives as dual WT and S31N inhibitors, which is described in the present manuscript. These compounds serve as starting points for design of novel compounds that address the problem of drug-resistant influenza A infections. At the same time of writing this manuscript, a pinanamine bearing imidazole derivative was reported to show weak inhibitory activity against the S31N M2 channel.<sup>41</sup>

Table 1. M2 inhibitors with an appended aromatic group<sup>1</sup>

Structure			
	<b>3</b>	<b>4</b>	<b>5-49</b>
	Rimantadine-based	Pinanamine-based	Amantadine-based (this work)
S31N % inhibition (TEVC)	0%	0%	See the text
WT % inhibition (TEVC)	27 ± 2%	91 ± 2%	

<sup>1</sup> The activity of the inhibitors was measured using the TEVC technique with full length AM2 protein in the *Xenopus* oocytes membrane; average of three measurements was reported (see Experimental Section in the Supporting Information).


A significant obstacle to developing inhibitors of M2 is to devise assays that are specific to inhibition of M2. In establishing an assay for S31N, we found it important to verify that the assay identified true M2 inhibitors, as we found both vesicle proton flux assays and viral inhibition assays are complicated by the fact that many amantadine-like compounds are proton carriers. Proton carriers such as chloroquine are good inhibitors of virus *in vitro*, but lack efficacy in animal models and human trials.<sup>42-44</sup> However, by conducting electrophysiological assays in parallel with antiviral assays it is possible to eliminate many false positives from compounds that acted as proton carriers and delay acidification of the endosome. We report here the first examples of compounds with specificity for both S31N and WT. Such compounds will be particularly important as mechanistic probes as well as serving as starting points for design of novel compounds that address the problem of drug-resistant influenza A infections.



## Chemistry


The synthetic routes of the compounds **5**, **8**, **9**, **11**, **12**, **15**, **17-23**, **29-49** are shown in Scheme 1. A variety of benzyl groups were installed to N-terminal of **1** either through reductive amination (procedure **a** or **b**)<sup>45-47</sup> or amide bond formation followed by lithium aluminium hydride (LAH) reduction (procedure **c**).<sup>48-50</sup> Subsequent reactions included deprotection, oxidation of a thioether,<sup>51</sup> tetrazole formation,<sup>52</sup> or N-methylation, yielding compounds **6**, **10**, **13**, **16**, and **27**, respectively. Due to a low yield in the reductive animation with 2,4-dihydroxybenzaldehyde, compound **44** was synthesized by reductive amination with 2-hydroxy-4-methoxybenzaldehyde followed by demethylation with BBr<sub>3</sub>.<sup>53</sup> Compounds **14**, **24**, **25**, **31**, **33** were prepared via similar routes (Scheme 2). The syntheses of the imine analogue, **26**<sup>40</sup> and the ether analogue, **23**, are shown in Scheme 3.

Scheme 1




Reagents and conditions: (a) NaBH<sub>3</sub>CN, MeOH, RT, overnight. (b) i. Ti(OiPr)<sub>4</sub>; ii. NaBH<sub>4</sub>, MeOH. (c) i. EDCI, HOAT, DIPEA, CH<sub>2</sub>Cl<sub>2</sub>, RT; ii. LAH, THF, Reflux. (d) HCl/dioxane, rt. (e) LAH, THF, Reflux. (f) NaN<sub>3</sub>, CH<sub>3</sub>CN. (g) *m*-CPBA (1.2 equiv), CH<sub>2</sub>Cl<sub>2</sub>, RT. (h) NaH, MeI (1.1 equiv), 0 °C, DMF. (i) BBr<sub>3</sub> (1.2 equiv), CH<sub>2</sub>Cl<sub>2</sub>, -78 to 0 °C.

Scheme 2



Reagents and conditions: (a)  $\text{NaBH}_3\text{CN}$ , MeOH, RT. (b) i. EDCI, HOAT, DIPEA, DMF, RT; ii. LAH, THF, Reflux.

Scheme 3



Reagents and conditions: (a) piperidine (cat.), benzene, Dean-Stark. (b) NaH, PMBCl (1.1 equiv), RT, DMF.

## Structure-Activity Relationship (SAR) Study

The activity of the inhibitors was measured using the TEVC technique with full length M2 protein expressed in *Xenopus* oocytes.<sup>21,54-56</sup> All inhibitors were initially tested at 100  $\mu\text{M}$ ; those that inhibited the AM2 channel activity more than amantadine (Amt) (**1**) (by 35% inhibition) were chosen for measurement of their  $\text{IC}_{50}$ . Inhibitors of M2 enter the channel from the outside of the virus through a narrow opening near Val27. Thus, M2-blockers show slow on/off inhibition, with association occurring on the min. to hour time scale (at 1  $\mu\text{M}$ ) and dissociation hour to day time scale. Due to the limited stability of oocytes at low pH, we measure the percent inhibition after only 2 min, before equilibrium is reached. Due to the *kinetics of inhibition*, this provides an apparent kinetic  $\text{IC}_{50}$  up to 100-fold higher than the true  $K_{\text{diss}}$  measured with much longer times.<sup>57</sup> Nevertheless, the relative potency of the compounds tracks well with their relative potencies measured in other methods. Thus, the low  $\mu\text{M}$   $\text{IC}_{50}$  values of compounds seen in our primary electrophysiologic assays are























1 numerically larger than their antiviral potency in plaque assays. While the TEVC assay underestimates the true  
2 potencies of the compounds, it has the advantage of being remarkably reproducible, allowing us to reliably  
3 interpret even small degrees of inhibition. All measurements were made in triplicate, with a typical standard  
4 error being between 1% and 3%. Moreover, we observed less than 5% error when making repeat measurements  
5 over time and with different batches of oocytes.  
6  
7  
8  
9  
10

11  
12  
13  
14 Since the reported nonamantadine-based aryl-containing M2 inhibitors (Table 1)<sup>39,40</sup> showed essentially no  
15 inhibition against S31N, we started to explore amantadine-based derivatives with an aromatic group appended  
16 to the primary amine group, which led to the discovery of a new series of substituted N-benzyl amantadines as  
17 promising M2 inhibitors against S31N as well as WT. Beginning with the unsubstituted N-benzyl analogue of  
18 amantadine we examined the effects of substitution at various positions in the ring. Because unsubstituted  
19 amantadine is already a relatively hydrophobic drug, we focused the present studies primarily on polar  
20 substituents to capture additional hydrogen bonding or electrostatic interactions as well as to maintain desirable  
21 drug-like properties. Although the unsubstituted derivative **5** was inactive towards S31N (1% inhibition), it did  
22 show significant inhibition against the WT (77% inhibition) at 100  $\mu$ M drug concentration. We therefore  
23 focused on phenyl substituents with hydrogen-bonding donor/acceptor at 4-position (Table 2), and were  
24 gratified to observe that the S31N inhibitory activity did increase gradually for 4-substituted series.  
25 Introduction of a 4-amino substituent (compound **6**) increased S31N activity from 1% to 10% inhibition, but  
26 decreased activity against WT from 77% to 31% inhibition. Interestingly, a Boc derivative of the aniline of **6**  
27 improved S31N activity to 20% inhibition (compound **7**), and acetylation of **6**, further boosted S31N activity to  
28 32% inhibition (compound **8**), which is similar to the activity of **1** (35% inhibition). However, the closely  
29 related benzamide **9** and benzylamine **10** lost considerable potency against both S31N and WT.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Next, heteroaryl substituents at 4-position were examined. Introduction of a pyrazole **11**, or imidazole **12** gave compounds with some activity against S31N (26% and 25% inhibition, respectively), but still less potent than unsubstituted amantadine (**1**). A tetrazole derivative was even less active than the pyrazole and imidazole derivatives (8% inhibition) therefore, we examined other substituents at the 4-position. Replacement of aniline of **6** with its bioisomer, 4-pyridine **14** slightly increased S31N and WT activity. Other substituents at the 4-position such as sulfone **15**, sulfoxide **16**, nitro **17**, cyano **18**, and fluorine **19** showed some improvement in S31N activity compared to the parent compound **5**, but did not reach the same activity as amantadine (**1**).

The 4-phenolic derivative, 4-(adamantan-1-ylaminomethyl)-phenol (**20**) was the first compound that showed slightly higher inhibitory activity against S31N ( $IC_{50} = 166 \mu M$ ) than amantadine (**1**) ( $IC_{50} = 199.9 \mu M$ ). Methylation of the 4-hydroxyl group of **20**, reduced S31N inhibitory activity from 42% to 31% inhibition (compound **21**), which indicated the importance of the 4-hydroxy group. Methyl ketone **23** and methyl ether **21** showed similar S31N activity (31% inhibition). One carbon homologated derivative of **20** (compound **22**), markedly decreased S31N inhibitory activity from 42% to 17% inhibition. Interestingly, 4-cyclohexanol **24**, a saturated version of **20**, dramatically dropped S31N activity from 42% to 19% inhibition, which demonstrated that the aromaticity is crucial for S31N activity. We also found that the 4-hydroxy of **20** could not be replaced by its bioisomer, 3-furan **25**, as this modification led to a marked loss of S31N inhibitory activity from 42% to 8%, respectively.






**Table 2.** SAR of 4-substituted adamantan-1-yl-benzyl-amines<sup>1</sup>

											
R <sub>1</sub> =	-H Amt										
/ ID #	1	5	6	7	8	9	10	11	12	13	14
S31N											
% inh/ IC <sub>50</sub>	35 ± 2/ 199.9 μM	1 ± 1	10 ± 1	20 ± 2	32 ± 2	3 ± 1	8 ± 1	26 ± 3	25 ± 2	8 ± 1	15 ± 1
WT											
% inh/ IC <sub>50</sub>	91 ± 3/ 16.1 μM	77 ± 1	31 ± 2	13 ± 2	3 ± 2	7 ± 2	3 ± 2	4 ± 3	4 ± 2	6 ± 1	6 ± 1
R <sub>1</sub> =											
/ ID #	15	16	17	18	19	20	21	22	23	24	25
S31N											
% inh/ IC <sub>50</sub>	12 ± 1	21 ± 2	14 ± 1	8 ± 2	8 ± 2	42 ± 2/ 166 μM	31 ± 2	17 ± 2	31 ± 2	19 ± 2	8 ± 2
WT											
% inh/ IC <sub>50</sub>	2 ± 2	45 ± 2	49 ± 2	36 ± 2	77 ± 1	67 ± 2	43 ± 1	26 ± 2	2 ± 2	16 ± 1	73 ± 2

<sup>1</sup> The activity of the inhibitors was measured using the TEVC technique with full length AM2 protein in the *Xenopus* oocytes membrane; average of three measurements was reported (see Experimental Section in the Supporting Information).










We also explored the SAR of the linkers (Table 3). A secondary amine is preferred over an imine analogue for S31N activity (42% and 21% inhibition for compounds **20** and **26**, respectively). Secondary amine linker also showed better S31N activity than either a tertiary amine or an ether linkage (31%, 20%, and 13% inhibition for compounds **21**, **27**, and **28**, respectively). Accordingly, in the following SAR studies, we kept the secondary amine linker constant.

Table 3. The SAR of the linkers

Structure					
	<b>20</b>	<b>26</b>	<b>21</b>	<b>27</b>	<b>28</b>
S31N % inh	42 ± 2%	21 ± 1%	31 ± 2%	20 ± 3%	13 ± 1%
WT % inh	67 ± 2%	81 ± 2%	43 ± 1%	34 ± 3%	51 ± 2%












We further examined SAR of 2- and 3-substituted benzyl amine analogues (Table 4). Two phenol regioisomers of **20**, 3-hydroxy **34** and 2-hydroxy **29**, displayed much lower S31N inhibitory activity (6% and 18% inhibition, respectively), but improved WT inhibitory activity (88% and 81% inhibition, respectively). As was the case for 4-methyl ether **21** (Table 2), 2-methyl ether **30** also decreased rather than enhanced the S31N inhibitory activity. Similarly, replacement of the 2-hydroxy in **29** with a 2-furan **31**, 2-nitro-phenyl **32**, or 2-pyridyl **33** group led to the further decrease in activity.

**Table 4.** SAR of 2 and 3-substituted adamantan-1-yl-benzyl-amines

									
R <sub>1</sub> =	-H Amt								
/ ID #	1	5	20	29	30	31	32	33	34
S31N									
% inh/	35 ± 2/	1 ± 1	42 ± 2/	18 ± 2	14 ± 1	6 ± 2	1 ± 1	0%	6 ± 2
IC <sub>50</sub>	199.9 μM		166 μM						
WT									
% inh/	91 ± 3/	77 ± 1	67 ± 2	81 ± 2	22 ± 2	66 ± 3	48 ± 1	12 ± 1	88 ± 1
IC <sub>50</sub>	16.1 μM								

Attention was turned to effects of 3-substituents in the context of the 4-hydroxy **20**, which had significant activity against S31N and WT (Table 5). Both electron withdrawing groups (3-Cl, 3-F, 3-NO<sub>2</sub>) (compounds **37**, **38**, and **39**, respectively) and electron donating groups (3-OH, 3-OMe) (compounds **35** and **36**, respectively) reduced potency, with 3-fluoro derivative **38** showing the smallest decrease in potency (31% inhibition). 3-Methoxy analogue **36** is less effective than 3-hydroxy derivative **35**, which may be due to steric hindrance. This was supported by a more potent inhibitor **41**, in which the methylene at 3-position was constrained into a 5-membered ring. Methylation of the 4-hydroxyl group of **35** (compound **40**) maintained both WT and S31N activity and displayed slightly decreased activity than the more constrained **41**. We also found that 4-hydroxy cannot be replaced with two cyclic bioisomers, **42** and **43**, which displayed much weaker activity against S31N (10% and 12% inhibition, respectively).

Table 5. SAR of 3,4-disubstituted adamantan-1-yl-benzyl-amines










<div></div>											
R <sub>1</sub> =	H										
/ ID #	1	20	35	36	37	38	39	40	41	42	43
S31N											
% inh/	35 ± 2/	42 ± 2/	23 ± 1	3 ± 1	16 ± 1	31 ± 3	13 ± 2	20 ± 1	29 ± 1	10 ± 1	12 ± 1
IC <sub>50</sub>	199.9 μM	166 μM									
WT											
% inh/	91 ± 3/	67 ± 2	44 ± 3	76 ± 2	78 ± 1	65 ± 1	66 ± 1	46 ± 3	54 ± 2	26 ± 1	79 ± 1
IC <sub>50</sub>	16.1 μM										


Because 2- and 4-hydroxyl and methoxy analogues provided better inhibition compared to the 3-hydroxy analogue (Table 4), we further examined whether their beneficial effects were additive. Toward this end, we synthesized 2,4-disubstituted benzyl amine derivatives (Table 6). Pleasingly, 2,4-dihydroxy substituted analog, **44**, markedly enhanced S31N inhibitory activity (67% inhibition at 100 μM, IC<sub>50</sub> = 35.2 μM). Compound **44** was approximately 6-fold more potent against S31N than amantadine (**1**) (IC<sub>50</sub> = 199.9 μM), and only 2-fold less potent than amantadine (**1**) for WT (IC<sub>50</sub> = 16 μM) in the TEVC assay. Compound **44** was also active against the WT with an IC<sub>50</sub> of 59 μM, approximately 3.7-fold less potent than amantadine (**1**). The antiviral activity of **44** was also profiled in viral plaque reduction assays (Fig. 4). The plaque assays showed that the compounds were more active than in the electrophysiological TEVC assays, which underestimated the affinities due to the brief period of inhibition. Compound **44** had an EC<sub>50</sub> of 3.2 μM against S31N and was 7-fold more potent than amantadine (**1**) (EC<sub>50</sub> = 22.5 μM) (Fig. 3A). In the plaque assay, compound **44** was also active against the WT virus with complete inhibition of plaque formation at 1 μM (Fig. 3B).



We also found that methylation of the 4-hydroxy **45** maintained the potency for both WT and S31N with IC<sub>50</sub> values of 79  $\mu$ M and 41.3  $\mu$ M, respectively. The importance of the 2-hydroxy was observed in other 4-substituted analogues. For example, installation of 2-hydroxy to 4-fluoro analogue **19** markedly improved S31N inhibitory activity from 8% to 37% inhibition (compound **46**). However, methylation of 2-hydroxy in **44** and **45** (compounds **47** and **48**, respectively) dramatically reduced both S31N and WT activities, which might be due to the steric hindrance at this position. 2-Chloro analogue **49** also showed much lower S31N activity than **44**, which may be attributed to the fact that chlorine is a weaker hydrogen acceptor than a hydroxyl group. These data might also indicate that a hydrogen bond donating hydroxyl is essential at this position of the ring.

**Table 6.** SAR of 2,4-disubstituted adamantan-1-yl-benzyl-amine of selective compounds

										
R1=	-H Amt									
/ID #	1	20	29	44	45	46	47	48	49	
S31N	% inh	35 $\pm$ 2	42 $\pm$ 2	18 $\pm$ 2	67 $\pm$ 1	66 $\pm$ 1	37 $\pm$ 2	12 $\pm$ 2	11 $\pm$ 1	14 $\pm$ 1
	IC <sub>50</sub> ( $\mu$ M)	199.9 $\mu$ M	166 $\mu$ M		35.2 $\mu$ M	43.1 $\mu$ M	NA	NA	NA	NA
WT	% inh	91 $\pm$ 3	67 $\pm$ 2	81 $\pm$ 2	64 $\pm$ 1	56 $\pm$ 1	87 $\pm$ 2	0	16 $\pm$ 2	78 $\pm$ 1
	IC <sub>50</sub> ( $\mu$ M)	16.1 $\mu$ M	NA		59 $\mu$ M	79 $\mu$ M	NA	NA	NA	NA



**Figure 4.** (A) Plaque reduction assay with A/WSN/33-S31N influenza virus. (B) Plaque reduction assay with A/WSN/33-WT virus. The inhibitory effect of compound **44** and amantadine on influenza A virus replication was evaluated by plaque formation in MDCK cells as described in the Experimental Section.

## Conclusions:

Using a combination of structural information of M2 proton channel and medicinal chemistry approaches, we report the discovery of promising dual M2 inhibitors for S31N and WT M2 channels, demonstrating that M2 S31N is a druggable target. Based on this work, several more potent series of S31N inhibitors have been discovered, one series has already been published<sup>58</sup> and others will be the subject of future reports. Such compounds will be also important as mechanistic probes to address the problem of drug-resistant influenza A infections.

## AUTHORSHIP INFORMATION

### Corresponding Authors:

Jizhou Wang, Tel: (484) 598-2376; Fax: (484) 598-2401; E-mail: jwang@influmeditx.com

William F. DeGrado, Tel: (415) 476-9679; Fax: (415)-502-4690; E-mail: William.DeGrado@ucsf.edu

## ACKNOWLEDGEMENT

We acknowledge support of grant AI74571 from the NIH.

## SUPPORTING INFORMATION

Details of the Two-Electrode Voltage Clamp (TEVC) Assay, Plaque Reduction Assay, and Synthesis of Compounds, **5-49**, are available free of charge via the Internet at <http://pubs.acs.org>.

## LIST OF ABBREVIATIONS

Amantadine (Amt), Serine 31 (S31), Serine 31 Arginine (S31N), Wild type (WT), Compound (Cmpd), Two-Electrode Voltage Clamp (TEVC), % inhibition (% inh), The half maximal inhibitory concentration (IC<sub>50</sub>), Madin-Darby Canine Kidney Cells (MDCK), Lithium aluminium hydride (LAH), 1-Hydroxy-7-azabenzotriazole (HOAT), 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI), Nuclear Magnetic Resonance (NMR), *N,N*-Diisopropylethylamine (DIPEA), Dimethylformamide (DMF), Room temperature (RT), *meta*-Chloroperoxybenzoic acid (*m*-MCPA).

## REFERENCES

- Bright, R. A.; Medina, M. J.; Xu, X. Y.; Perez-Oronoz, G.; Wallis, T. R.; Davis, X. H. M.; Povinelli, L.; Cox, N. J.; Klimov, A. I. Incidence of adamantane resistance among influenza A (H3N2) viruses isolated worldwide from 1994 to 2005: a cause for concern. *Lancet* **2005**, *366*, 1175-1181.

2. Bright, R. A.; Shay, D. K.; Shu, B.; Cox, N. J.; Klimov, A. I. Adamantane resistance among influenza A viruses isolated early during the 2005-2006 influenza season in the United States. *JAMA* **2006**, *295*, 891-894.
3. Cheng, P. K.; Leung, T. W.; Ho, E. C.; Leung, P. C.; Ng, A. Y.; Lai, M. Y.; Lim, W. W. Oseltamivir- and amantadine-resistant influenza viruses A (H1N1). *Emerg. Infect. Dis.* **2009**, *15*, 966-968.
4. Sheu, T. G.; Deyde, V. M.; Okomo-Adhiambo M., e. a. Surveillance for neuraminidase inhibitor resistance among human influenza A and B viruses circulating worldwide from 2004 to 2008. *Antimicrob. Agents Chemother.* **2008**, *52*, 3284-3292.
5. Moscona, A. Global Transmission of Oseltamivir-Resistant Influenza. *N. Engl. J. Med.* **2009**, *360*, 953-956.
6. Moore, C.; Galiano, M.; Lackenby, A.; Abdelrahman, T.; Barnes, R.; Evans, M. R.; Fegan, C.; Froude, S.; Hastings, M.; Knapper, S.; Litt, E.; Price, N.; Salmon, R.; Temple, M.; Davies, E. Evidence of person to person transmission of oseltamivir resistant pandemic influenza A(H1N1) 2009 virus in a hematology unit. *J. Infect. Dis.* **2010**, *203*, 18-24.
7. Hayden, F. G.; deJong, M. D. Emerging Influenza Antiviral Resistance Threats. *J. Infect. Dis.* **2011**, *203*, 6-10.
8. Baz, M. Emergence of Oseltamivir-Resistant Pandemic H1N1 Virus during Prophylaxis. *N. Engl. J. Med.* **2009**, *361*, 2296-2297.
9. Sheu, T. G.; Fry, A. M.; Garten, R. J.; Deyde, V. M.; Shwe, T.; Bullion, L.; Peebles, P. J.; Li, Y.; Klimov, A. I.; Gubareva, L. V. Dual resistance to adamantanes and oseltamivir among seasonal influenza A (H1N1) viruses: 2008-2010. *J. Infect. Dis.* **2011**, *203*, 13-17.
10. Balannik, V.; Carnevale, V.; Fiorin, G.; Levine, B. G.; Lamb, R. A.; Klein, M. L.; DeGrado, W. F.; Pinto, L. H. Functional studies and modeling of pore-lining residue mutants of the influenza A virus M2 ion channel. *Biochemistry* **2010**, *49*, 696-708.

11. Hay, A. J.; Wolstenholme, A. J.; Skehel, J. J.; Smith, M. H. The Molecular-Basis of the Specific Anti-Influenza Action of Amantadine. *Embo. J.* **1985**, *4*, 3021-3024.
12. Grambas, S.; Hay, A. J. Maturation of influenza A virus hemagglutinin-Estimates of the pH encountered during transport and its regulation by the M2 protein. *Virology* **1992**, *190*, 11-18.
13. Hu, W.; Zeng, S.; Li, C.; Jie, Y.; Li, Z.; Chen, L. Identification of hits as matrix-2 protein inhibitors through the focused screening of a small primary amine library. *J. Med. Chem.* **2010**, *53*, 3831-3834.
14. Wang, J.; Ma, C.; Balannik, V.; Pinto, L. H.; Lamb, R. A.; Degrado, W. F. Exploring the Requirements for the Hydrophobic Scaffold and Polar Amine in inhibitors of M2 from Influenza A Virus. *ACS Med. Chem. Lett.* **2011**, *2*, 307-312.
15. Wang, J.; Ma, C.; Fiorin, G.; Carnevale, V.; Wang, T.; Hu, F.; Lamb, R. A.; Pinto, L. H.; Hong, M.; Klein, M. L.; DeGrado, W. F. Molecular dynamics simulation directed rational design of inhibitors targeting drug-resistant mutants of influenza A virus M2. *J. Am. Chem. Soc.* **2011**, *133*, 12834-12841.
16. Kurtz, S.; Luo, G.; Hahnenberger, K. M.; Brooks, C.; Gecha, O.; Ingalls, K.; Numata, K.; Krystal, M. Growth impairment resulting from expression of influenza virus M2 protein in *Saccharomyces cerevisiae*: identification of a novel inhibitor of influenza virus. *Antimicrob. Agents Chemother.* **1995**, *39*, 2204-2209.
17. De Clercq, E. Antiviral agents active against influenza A viruses. *Nat Rev Drug Discov* **2006**, *5*, 1015-1025.
18. Severson, W. E.; McDowell, M.; Ananthan, S.; Chung, D. H.; Rasmussen, L.; Sosa, M. I.; White, E. L.; Noah, J.; Jonsson, C. B. High-Throughput Screening of a 100,000-Compound Library for Inhibitors of Influenza A Virus (H3N2). *J. Biomol. Screen.* **2008**, *13*, 879-887.
19. Lagoja, I. M.; De Clercq, E. Anti-influenza virus agents: Synthesis and mode of action. *Med. Res. Rev.* **2008**, *28*, 1-38.
20. Das, K., Antivirals Targeting Influenza A Virus. *J. Med. Chem.* **2012**, *55*, 6263-6277.

21. Jing, X.; Ma, C.; Ohigashi, Y.; Oliveira, F. A.; Jardetzky, T. S.; Pinto, L. H.; Lamb, R. A. Functional studies indicate amantadine binds to the pore of the influenza A virus M2 proton-selective ion channel. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 10967-10972.
22. Duque, M. D.; Ma, C.; Torres, E.; Wang, J.; Naesens, L.; Juarez-Jimenez, J.; Camps, P.; Luque, F. J.; DeGrado, W. F.; Lamb, R. A.; Pinto, L. H.; Vazquez, S. Exploring the size limit of templates for inhibitors of the M2 ion channel of influenza A virus. *J. Med. Chem.* **2011**, *54*, 2646-2657.
23. Ma, C. *unpublished results*.
24. Stouffer, A. L.; Acharya, R.; Salom, D.; Levine, A. S.; Di Costanzo, L.; Soto, C. S.; Tereshko, V.; Nanda, V.; Stayrook, S.; DeGrado, W. F. Structural basis for the function and inhibition of an influenza virus proton channel. *Nature* **2008**, *451*, 596-599.
25. Schnell, J. R.; Chou, J. J. Structure and mechanism of the M2 proton channel of influenza A virus. *Nature* **2008**, *451*, 591-593.
26. Cady, S. D.; Luo, W.; Hu, F.; Hong, M. Structure and function of the influenza A M2 proton channel. *Biochemistry* **2009**, *48*, 7356-7364.
27. Cady, S. D.; Mishanina, T. V.; Hong, M. Structure of amantadine-bound M2 transmembrane peptide of influenza A in lipid bilayers from magic-angle-spinning solid-state NMR: the role of Ser31 in amantadine binding. *J. Mol. Biol.* **2009**, *385*, 1127-1141.
28. Cady, S. D.; Schmidt-Rohr, K.; Wang, J.; Soto, C. S.; DeGrado, W. F.; Hong, M. Structure of the amantadine binding site of influenza M2 proton channels in lipid bilayers. *Nature* **2010**, *463*, 689-692.
29. Sharma, M.; Yi, M.; Dong, H.; Qin, H.; Peterson, E.; Busath, D. D.; Zhou, H. X.; Cross, T. A. Insight into the mechanism of the influenza a proton channel from a structure in a lipid bilayer. *Science* **2010**, *330*, 509-512.
30. Acharya, R.; Carnevale, V.; Fiorin, G.; Levine, B. G.; Polishchuk, A. L.; Balannik, V.; Samish, I.; Lamb, R. A.; Pinto, L. H.; DeGrado, W. F.; Klein, M. L. Structure and mechanism of proton transport

- through the transmembrane tetrameric M2 protein bundle of the influenza A virus. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 15075-15080.
31. Pielak, R. M.; Oxenoid, K.; Chou, J. J. Structural investigation of rimantadine inhibition of the AM2-BM2 chimera channel of influenza viruses. *Structure* **2011**, *19*, 1655-1663.
32. Cady, S. D.; Wang, J.; Wu, Y.; DeGrado, W. F.; Hong, M. Specific binding of adamantane drugs and direction of their polar amines in the pore of the influenza M2 transmembrane domain in lipid bilayers and dodecylphosphocholine micelles determined by NMR spectroscopy. *J. Am. Chem. Soc.* **2011**, *133*, 4274-4284.
33. Andreas, L. B.; Eddy, M. T.; Chou, J. J.; Griffin, R. G. Magic-Angle-Spinning NMR of the Drug Resistant S31N M2 Proton Transporter from Influenza A. *J. Am. Chem. Soc.* **2012**, *134*, 7215-7218.
34. Pielak, R. M.; Schnell, J. R.; Chou, J. J. Mechanism of drug inhibition and drug resistance of influenza A M2 channel. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 7379-7384.
35. Balannik, V.; Wang, J.; Ohigashi, Y.; Jing, X.; Magavern, E.; Lamb, R. A.; DeGrado, W. F.; Pinto, L. H., Design and pharmacological characterization of inhibitors of amantadine-resistant mutants of the M2 ion channel of influenza A virus. *Biochemistry* **2009**, *48*, 11872-11882.
36. Wang, J.; Fan, X.; Cristian, L. Preparation of adamantane derivatives as antiviral agents. *U.S. Pat. Appl. Publ.* **2011**, US 20110065762 A1.
37. Wang, J.; Fan, X.; Cristian, L. Preparation of adamantane derivatives as antiviral agents. *U.S. Pat. Appl. Publ.* **2011**, US 20110065766 A1.
38. Wang, J.; Ma, C.; Wu, Y.; Lamb, R. A.; Pinto, L. H.; DeGrado, W. F. Exploring organosilane amines as potent inhibitors and structural probes of influenza a virus M2 proton channel. *J. Am. Chem. Soc.* **2011**, *133*, 13844-13847.


39. Zhang, W.; Xu, J.; Liu, F.; Li, C.; Jie, Y.; Chen, S.; Li, Z.; Liu, J.; Chen, Ling; Zhou, G. Heterodimers of histidine and amantadine as inhibitors for wild type and mutant M2 channels of influenza A. *Chin. J. Chem.* **2010**, *28*, 1417-1423.
40. Zhao, X.; Li, C.; Zeng, S.; Hu, W. Discovery of highly potent agents against influenza A virus. *Eur. J. Med. Chem.* **2011**, *46*, 52-57.
41. Zhao, X.; Jie, Y.; Rosenberg, M. R.; Wan, J.; Zeng, S.; Cui, W.; Xiao, Y.; Li, Z.; Tu, Z.; Casarotto, M. G.; Hu, W. Design and synthesis of pinanamine derivatives as anti-influenza A M2 ion channel inhibitors. *Antiviral Res.* **2012**, *96*, 91-99.
42. Di Trani, L.; Savarino, A.; Campitelli, L.; Norelli, S.; Puzelli, S.; D'Ostilio, D.; Vignolo, E.; Donatelli, I.; Cassone, A. Different pH requirements are associated with divergent inhibitory effects of chloroquine on human and avian influenza A viruses. *Viro.l J.* **2007**, *4*, 39.
43. Ooi, E. E.; Chew, J. S.; Loh, J. P.; Chua, R. C. In vitro inhibition of human influenza A virus replication by chloroquine. *Virol. J.* **2006**, *3*, 39.
44. Paton, N. I.; Lee, L.; Xu, Y.; Ooi, E. E.; Cheung, Y. B.; Archuleta, S.; Wong, G.; Wilder-Smith, A. Chloroquine for influenza prevention: a randomised, double-blind, placebo controlled trial. *Lancet Infect. Dis.* **2011**, *11*, 677-6783.
45. Abdel-Magid, A. F.; Carson, K. G.; Harris, B. D.; Maryanoff, C. A.; Shah, R. D. Reductive amination of aldehydes and ketones with sodium triacetoxyborohydride. Studies on direct and indirect reductive amination procedures (1). *J. Org. Chem.* **1996**, *61*, 3849-3862.
46. Dalpathado, D. S.; Jiang, H.; Kater, M. A.; Desaire, H. Reductive amination of carbohydrates using NaBH(OAc)<sub>3</sub>. *Anal. Bioanal. Chem.* **2005**, *381*, 1130-1137.
47. Salmi, C.; Loncle, C.; Letourneux, Y.; Brunel, J. M. Efficient preparation of secondary aminoalcohols through a Ti(IV) reductive amination procedure. Application to the synthesis and antibacterial evaluation of new 3 beta-N-[hydroxyalkyl]aminosteroid derivatives. *Tetrahedron* **2008**, *64*, 4453-4459.




- 1 48. Webb, T. R.; Eigenbrot, C. Conformationally restricted arginine analogues. *J. Org. Chem.* **1991**, *56*,  
2 3009-3016.  
3
- 4 49. Webb, T. H.; Wilcox, C. S. Improved synthesis of symmetrical and unsymmetrical 5,11  
5 methanodibenzo[b,f][1,5]diazocines. Readily available nanoscale structural units. *J. Org. Chem.* **1990**,  
6 363-365.  
7
- 8 50. Kihara, N.; Hashimoto, M.; Takata, T. Redox behavior of ferrocene-containing rotaxane: Transposition  
9 of the rotaxane wheel by redox reaction of a ferrocene moiety tethered at the end of the axle. *Org. Lett.*  
10 **2004**, *6*, 1693-1696.  
11
- 12 51. Zhang, X.; Yu, M.; Yao, J.; Zhang, Y. Palladium-catalyzed regioselective arylation of arene C-H bond  
13 assisted by the removable 2-pyridylsulfinyl group. *Synlett* **2012**, *23*, 463-467.  
14
- 15 52. Finnegan, W. G.; Hennry, R. A.; Lofquist, R. An improved synthesis of 5-substituted tetrazoles. *J. Am.*  
16 *Chem. Soc.* **1958**, *80*, 3908-3911.  
17
- 18 53. Williams, A. B.; Hanson, R. N. Synthesis of substituted asymmetrical biphenyl amino esters as alpha  
19 helix mimetics. *Tetrahedron* **2012**, *68*, 5406-5414.  
20
- 21 54. Balannik, V.; Lamb, R. A.; Pinto, L. H. The oligomeric state of the active BM2 ion channel protein of  
22 influenza B virus. *J. Biol. Chem.* **2008**, *283*, 4895-4904.  
23
- 24 55. Wang, C.; Takeuchi, K.; Pinto, L. H.; Lamb, R. A. Ion channel activity of influenza A virus M2 protein:  
25 characterization of the amantadine block. *J. Virol.* **1993**, *67*, 5585-5594.  
26
- 27 56. Pinto, L. H.; Holsinger, L. J.; Lamb, R. A. Influenza virus M2 protein has ion channel activity. *Cell*  
28 **1992**, *69*, 517-528.  
29
- 30 57. Balannik, V.; Obrdlik, P.; Inayat, S.; Steensen, C.; Wang, J.; Rausch, J. M.; DeGrado, W. F.; Kelety, B.;  
31 Pinto, L. H. Solid-supported membrane technology for the investigation of the influenza A virus M2  
32 channel activity. *Pflugers Arch.* **2010**, *459*, 593-605.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


58. Wang, J.; Wu, Y.; Ma, C.; Fiorin, G.; Wang, J.; Pinto, L. H.; Lamb, R. A.; Klein, M. L.; DeGrado, W. F. Structure and inhibition of the drug-resistant S31N mutant of the M2 ion channel of influenza A virus. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 1315-1320.

Table of Contents Graphic (TOC)






137x146mm (150 x 150 DPI)



211x178mm (150 x 150 DPI)



211x178mm (150 x 150 DPI)