



A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: Enantiodivergent [4+2]-Cycloadditions of Dienolates by Polyfunctional Lewis-Acid / Zwitterion Catalysis

Authors: Vukoslava Miskov-Pajic, Felix Willig, Daniel M. Wanner, Wolfgang Frey, and René Peters

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.202009093

Link to VoR: <https://doi.org/10.1002/anie.202009093>

COMMUNICATION

Enantiodivergent [4+2]-Cycloadditions of Dienolates by Polyfunctional Lewis-Acid / Zwitterion Catalysis

Vukoslava Miskov-Pajic,^[a] Felix Willig,^[a] Daniel M. Wanner,^[a] Wolfgang Frey,^[a] and René Peters*^[a]

Abstract: Diels-Alder reactions are established as one of the most effective ways to prepare stereochemically complex 6-membered rings. Different catalysis concepts have been reported including dienophile activation by Lewis acids or H-bond donors and diene activation by bases. Here we report a new concept, in which an acidic prodiene is acidified by a Lewis acid to facilitate deprotonation by an imidazolium-aryloxide entity within a polyfunctional catalyst. A metal dienolate is thus formed, while imidazolium-ArOH moieties probably form hydrogen bonds with the dienophile. The catalyst type which is readily prepared in few steps with high overall yields, was applied to 3-hydroxy-2-pyrone and -pyridone as well as cyclopentenone prodiene. Maleimide, maleic anhydride, and nitroolefin dienophiles were employed. Kinetic, spectroscopic and control experiments support a cooperative mode of action. Unprecedented TONs up to 3680 were achieved still attaining high enantioselectivity.

Asymmetric Diels-Alder reactions (DAs) provide a very important tool to enantioselectively construct stereochemically complex 6-membered carbocycles.^[1] Attractive features of this reaction type include stereospecificity regarding diene and dienophile components, wide applicability of different substrate types and the possibility to stereoselectively construct a number of stereogenic centers within the formed cyclohexene ring, thus creating structural complexity.^[1] Various concepts for catalytic asymmetric DAs have been reported^[1] including the use of chiral Lewis acids (LA),^[2] primary or secondary amines/ammonium salts,^[3] Brønsted acids,^[4] hydrogen bond donors,^[5] and organic Brønsted bases.^[4] With the traditional LA catalysts the dienophile's LUMO energy is decreased (Scheme 1).^[2] More rarely, metal centers have also served as tethers to allow for quasi-intramolecular reaction pathways.^[6] In contrast, bases were described to increase the HOMO energy of acidic (pro)diene.^[1,3,4]


In this article, we report the use of polyfunctional catalysts, in which the cooperation of a LA^[7] and an aprotic imidazolium aryloxide moiety is utilized to generate the reactive diene.^[8] The prodiene is acidified by coordination to the LA and then deprotonated by the aryloxide moiety.^[9] The LA should stabilize the generated dienolate. Besides, hydrogen bond activation is crucial in this catalyst system, as control experiments show, most likely for activation of the dienophile.^[10,11]

For our initial investigations, 3-hydroxy-2-pyrones **1** were studied as diene component to form densely functionalized bicyclic

lactones. Similar lactones have been described as valuable building blocks for the synthesis of a number of structurally complex, biologically active natural products such as taxol,^[6b] and others.^[12,13] Pyrones **1** are usually quite reluctant to undergo DAs,^[14] unless the electron density is increased by formation of a dienolate.^[15] Consequently, chiral Brønsted bases—in particular cinchona alkaloid derivatives—have been reported for highly enantioselective activation.^[16] Maleimides **2** were initially investigated as dienophiles, because succinimides not only represent an important structural motif in bioactive compounds,^[17] but also because the previously reported methods using a combination of **1** and **2** suffered from moderate enantioselectivity and very limited scope.^[16a-c]

Previous work:

Common catalytic concepts for Diels-Alder reactions:




1) **decreased LUMO energy** by Lewis acids, H-bond donors or secondary amines for dienophile activation

2) **increased HOMO energy** by bases for diene activation

This work:

Lewis acid / zwitterion catalyzed Diels-Alder reactions:



Scheme 1. Previous Diels-Alder approaches compared to presented one.

Cu(II)-imidazolium-/naphthoxide complex **C1a** (5 mol%)^[13] was found to catalyze the reaction of **1a** and **2A** at room temperature with complete *endo*-, yet low enantioselectivity (entry 1). At -40 °C, the *ee* could be significantly increased to 88%, while an excellent yield was maintained (entry 2). Lower reaction temperatures gave a decreased enantioselectivity (not shown). Formally replacing the *N*-Me substituent by the synthetically useful *N*-benzyl protecting group caused a slightly improved enantioselectivity


[a] M.Sc. V. Miskov-Pajic, M.Sc. F. Willig, M.Sc. D. M. Wanner, Dr. W. Frey, Prof. Dr. R. Peters
Universität Stuttgart, Institut für Organische Chemie
Pfaffenwaldring 55, D-70569 Stuttgart, Germany
E-mail: rene.peters@oc.uni-stuttgart.de

Supporting information for this article is given via a link at the end of the document.

COMMUNICATION

employing the same conditions (entry 3). Product **3aB** has been described as valuable precursor to antagonists of neurotransmitter SP, but was previously only accessible with moderate enantioselectivity.^[18] A change in configuration of the iminosulfonamide moiety from (1*S*,2*S*) to (1*R*,2*R*) resulted in the formation of the opposite product enantiomer (*ee* = 91%, entry 4). This shows that the configurational outcome is dominated by the iminosulfonamide moiety, rather than the axially chiral part.

Table 1. Development and optimization of the title reaction.



Entry	C1 (config.)	X	2	T/°C	yield ^[a]	<i>d</i> ^[b]	<i>ee</i> ^[c]
1	C1a (1 <i>S</i> ,2 <i>S</i>)	5	2A	25	99	>98:2	30
2	C1a (1 <i>S</i> ,2 <i>S</i>)	5	2A	-40	97	>98:2	88
3	C1a (1 <i>S</i> ,2 <i>S</i>)	5	2B	-40	97	>98:2	90
4	C1a (1 <i>R</i> ,2 <i>R</i>)	5	2B	-40	95	>98:2	-91
5	C1b (1 <i>S</i> ,2 <i>S</i>)	5	2B	-40	98	>98:2	-96
6	C1b (1 <i>R</i> ,2 <i>R</i>)	5	2B	-40	96	>98:2	-40
7 ^[d]	C1b (1 <i>S</i> ,2 <i>S</i>)	2.5	2B	-40	98	>98:2	-96
8 ^[d]	C1b (1 <i>S</i> ,2 <i>S</i>)	1	2B	-40	95	>98:2	-90
9 ^[e]	C1b (1 <i>S</i> ,2 <i>S</i>)	5	2B	-40	98	>98:2	-98

[a] Yield of isolated product. [b] *Endo/exo* ratios determined by ¹H NMR using the crude product. [c] The enantiomeric excess of the *endo*-isomer was determined by ¹H NMR using saturated CDCl₃ solutions of (*R*)-(+)-binaphthol.^[16a] A minus sign indicates that the antipode of the enantiomer depicted was generated in excess. [d] Reaction time 18 h. [e] **1a** was added by syringe pump.


It was found that the choice of the sulfonamide unit heavily influences the enantioselectivity. Most residues R¹ (Me, *p*-tolyl, C₆F₅, 2- and 4-nitrophenyl, 2-naphthyl) resulted in poor to moderate enantioselectivity (for details see the SI). Nevertheless, with R¹ = 1-naphthyl in **C1b**, which is accessible in an overall yield of 84% starting from (*R*)-BINAM (see SI), the enantiomeric excess increased to 96% (entry 5). Adding **1a** by syringe pump further improved the *ee* to 98% (entry 9).

To our surprise, with **C1b** the optical antipode of **3** was formed, although the (1*S*,2*S*) iminosulfonamide configuration was used (compare entries 3 & 5). In contrast, the (1*R*,2*R*) configuration for **C1b** resulted in no switch of the preferred enantiomer, but a

significantly decreased *ee* value (entry 6). We suggest that the coordination of **1** at Cu might result in different coordination geometries depending on the sulfonamide residues. Lower loadings still resulted in good results (entries 7-8).

Different pyrones **1** were examined as prodiene under conditions of Table 1, entry 9, using catalysts **C1a** and **C1b** (Table 2). Next to **1a**, also the 4-Me and 4-halo substituted substrates were effective and only *endo* products were found (entries 4-6).

Table 2. Investigation of various 3-hydroxy-2-pyrones as well as -pyridones.



#	C1	R	1 or 4	3B or 5B	yield ^[a]	<i>d</i> ^[b]	<i>ee</i> ^[c]
1	C1a	H	1a	3aB	97	>98:2	90
2	C1b	H	1a	<i>ent</i> - 3aB	98	>98:2	-98
3 ^[d]	C1b	H	1a	<i>ent</i> - 3aB	96	>98:2	-97
4	C1a	Me	1b	3bB	93	>98:2	93
5	C1a	Cl	1c	3cB	92	>98:2	95
6	C1a	Br	1d	3dB	94	>98:2	93
7 ^[e]	C1b	H	4a	<i>ent</i> - 5aB	97	>98:2	-94
8 ^[e]	C1b	allyl	4b	<i>ent</i> - 5bB	98	>98:2	-96
9 ^[e]	C1a	allyl	4b	5bB	95	>98:2	95
10 ^[e]	C1b	Cl	4c	<i>ent</i> - 5cB	97	>98:2	-79
11 ^[e,f]	C1b	H	4a	<i>ent</i> - 5aC ^[f]	94	>98:2	-94
12 ^[e,g]	C1b	H	4a	<i>ent</i> - 5aF ^[f]	94	>98:2	-95

[a] Yield of isolated product. [b] *Endo/exo* ratios determined by ¹H NMR using the crude product. [c] The enantiomeric excess was determined by ¹H NMR using saturated CDCl₃ solutions of (*R*)-(+)-binaphthol.^[16a] A minus sign indicates that the antipode of the enantiomer depicted was generated in excess. [d] Reaction was performed on 5 mmol scale. [e] Reaction was performed at 25 °C. [f] 4-O₂N-C₆H₄ was used for Bn in **2**. [g] H was used for Bn in **2**.

To showcase the practicality of this method, it was performed on a 5 mmol scale and provided excellent results (entry 3). In addition, catalyst recovery of **C1b** was successful by silica gel filtration.^[6] Employing the recovered catalyst provided *ent*-**3aB** in diastereopure form in 92% yield and with 98% *ee* for the second run and 92% yield and with 96% *ee* for the third run.


Besides pyrones **1**, we also examined pyridones **4** (entries 7-12) which are valuable to prepare bioactive compounds like the antiviral drug Oseltamivir.^[19] For catalytic asymmetric DAs of pyridones with **2**, there is a single method available providing high enantioselectivity, which is currently limited though to *N*-mesitylsulfonyl substituted substrates.^[20] In our study 2-nosyl (Ns) protected substrates **4** worked fine as well (entries 7-12). Again, only *endo*-isomers were found and different enantiomers were available with catalysts **C1a** and **C1b**.^[21] The 2-Ns protecting group could be readily removed as shown for *ent*-**5aB** by PhSH / DBU at 25 °C in THF (yield: 94%, *ee* 94%).

Different maleimides **2** were also well tolerated (Table 3). Next to unbranched and branched N-alkyl (entries 1-2), a Boc group

COMMUNICATION

(entry 3) was accepted which was removable from **3aD** in 92% yield (*ee* = 92%) using TFA at 25 °C. Moreover, product formation was also achieved with unprotected maleimide (entry 4 and Table 2, entry 12). In addition, aromatic substituents with various electronic properties were tolerated (entry 5 and SI). The reaction could also be extended to maleic anhydride **6** providing product **7**.^[16j] All products were formed in high yields in diastereopure form with high enantioselectivity.^[21]


Table 3. Investigation of maleimide dienophiles **2** and maleic anhydride **6**.



#	2 or 6	R-N / O	(<i>ent</i> -) 3a or 7	yield ^[a]	<i>dr</i> ^[b]	<i>ee</i> ^[c]
1	2A	Me-N	<i>ent</i> - 3aA	93	>98:2	93
2	2D	cyc-Hex-N	<i>ent</i> - 3aD	92	>98:2	96
3 ^[d]	2E	Boc-N	3aE	90	>98:2	-92
4 ^[e]	2F	H-N	<i>ent</i> - 3aF	89	>98:2	90
5	2G	Ph-N	<i>ent</i> - 3aG	94	>98:2	91
6 ^[d,f]	6	O	7	92	>98:2	-84

[a] Yield of isolated product. [b] *Endo/exo* ratios determined by ¹H NMR using the crude product. [c] The enantiomeric excess was determined by ¹H NMR using saturated CDCl₃ solutions of (*R*)-(+)-binaphthol.^[16a] A minus sign indicates that the antipode of the enantiomer depicted was generated in excess. [d] **C1a** (1*S*, 2*S*) was used as catalyst. [e] The reaction was performed at -20 °C. [f] The reaction was performed at 0 °C.

Other types of C-H acidic dienophiles can be applied as well as demonstrated for enone **8** (Scheme 2). With **2B** catalyst **C1a** produced bicycloheptane **9** exclusively as *exo*-diastereomer featuring four contiguous stereocenters (98% *ee*).^[22] Surprisingly, this catalytic asymmetric DA is unknown, although compounds closely related to **9** have been reported as modulators of nuclear hormone receptor function.^[23] The unexpected *exo*-selectivity might possibly point to a stepwise process in that case.




Scheme 2. Extension to different substrate types (**C1a** with (1*S*, 2*S*)-configuration).

In addition, with nitroolefin **10** the cycloaddition proceeded very efficiently (TON 3680, 92% yield) with just 0.025 mol% **C1a**. The previously not available *endo*-product was obtained with an *ee* of 98%. Prior to this study, only the *exo*-isomer of **11** was accessible with moderate *dr* by chiral amine catalysis with TONs <20.^[24,25] To learn more about the role of the aprotic betaine, experiments with several control catalysts were conducted (Table 4). Cs₂CO₃

(2.5 mol%) in the absence of a Cu complex formed racemic product **3aB** in 95% yield as an *endo* / *exo* mixture (84:16).

Table 4. Experiments with control catalyst systems.



[a] Yield of isolated product. [b] *Endo/exo* ratios determined by ¹H NMR using the crude product. [c] The enantiomeric excess was determined by ¹H NMR using saturated CDCl₃ solutions of (*R*)-(+)-binaphthol.^[16a] [d] In the absence of Cs₂CO₃.

With control catalysts **C2-C4** (5 mol%) bearing either an alkyl moiety R, a simple imidazolium or one with an aliphatic alcohol moiety, the reaction outcome was almost identical as with only Cs₂CO₃. The experiments with **C2** and **C4** were also repeated in the absence of base providing yields of 5% (*ee* n.d.) and 10% (racemic), respectively. In contrast, using Cs₂CO₃ and **C5** equipped with a neutral axially chiral residue bearing an aromatic OH group stereoselectivity was considerably higher. However, nearly complete diastereoselectivity was only attained with control catalyst **C6** containing a zwitterionic imidazolium/aryloxide moiety.^[26] In agreement with the observation that the configuration of the axially chiral binaphthol unit has a minor effect on stereocontrol, enantioselectivity was high with **C6**. Overall, these results demonstrate that the zwitterionic moiety is necessary to attain high control.^[27]

Next to the control experiments, kinetic and spectroscopic experiments were performed (for details see SI). Blackmond's reaction progress kinetic analysis (RPKA) was done using ¹H-NMR spectroscopy.^[28] By the "same excess" protocol, catalyst robustness and a possible product inhibition was assessed.^[27] These experiments show that no significant catalyst decomposition and product inhibition occur in the model reaction. The empirical rate law was determined by the variable time normalization analysis (VTNA) method introduced by Burés.^[29] Again, the model reaction of **1a** and **2B** was examined at -20 °C in THF-d₈ using catalyst **C1b** (1*S*, 2*S*). Four reactions with different initial concentrations of **C1b**, **1a** and **2B** were used monitoring product *ent*-**3aB**. The best fit for the normalization of the time scale axis was achieved for the following rate law:

$$r = k * [\text{C1b}]^{1.95} * [\text{1a}]^{0.07} * [\text{2B}]^{1.00}$$

The reaction rate thus follows a second order kinetic dependence for **C1b**, a first order for **2B**, and zero order for **1a**. The latter might be explained by substrate saturation for **1a**. The second order dependence for **C1b** indicates that two catalyst molecules are

COMMUNICATION

involved in the turnover-limiting step.^[30] There is also a significant positive non-linear effect for the model reaction of **1a** and **2B** catalyzed by **C1b** (5 mol%) in THF at $-40\text{ }^{\circ}\text{C}$. We suggest that a dimeric catalyst might be the active species, because a dimeric structure was also found for **C6** by X-ray analysis^[31] (details are in the SI).

UV-Vis titrations were performed in which **C1b** was treated with **1a**. Having added 1.0 equiv. of **1a**, the resulting spectrum is very similar to the one of the precatalyst, in which a neutral naphthol unit is present. With larger amounts of up to 10 equiv. of **1a**, the spectra did not significantly change anymore. This suggests that **1a** is deprotonated by the zwitterionic moiety. In combination with the zero order kinetic dependence for **1a**, we suggest that **1a** binds to the Cu center where it gets deprotonated resulting in substrate saturation. A Cu-dienolate could indeed be found by ESI HRMS (disodium salt: $m/z = 1128.2563$, calculated: 1128.2564). The interpretation was validated by $^1\text{H-NMR}$ titrations, which were done in THF- d_8 at $-20\text{ }^{\circ}\text{C}$. Because the catalyst is paramagnetic, no well resolved signals were detected. By addition of 1 equiv. of **1a**, its signals almost disappeared caused by the paramagnetism. However, with an excess of **1a** the signals were appearing again, thus suggesting that the catalyst is already saturated with 1 equiv. In conclusion, we have reported an efficient way to catalyze [4+2]-cycloadditions with acidic dienophiles. They are activated by coordination to a LA and deprotonation by a zwitterionic moiety forming a metal dienolate. The acidic OH group generated during this proton transfer and the imidazolium unit are crucial for excellent diastereo- and enantioselectivity. In combination with the first order kinetic dependence for maleimide **2B**, it is likely that the cycloaddition step is usually turnover-limiting. The reaction was found to be applicable to different substrate types and TONs of up to 3680 were achieved. Moreover, the catalyst can be prepared in very high overall yield, is stable during catalysis and can be readily recycled by a simple filtration protocol.

Acknowledgements

This work was financially supported by the Deutsche Forschungsgemeinschaft (DFG, project 310990893). VMP thanks the Landesgraduiertenförderung BW for a Ph.D. scholarship.

Keywords: asymmetric catalysis • Diels-Alder • lactones • lactams • polyfunctional catalyst

- [1] a) K. Ishihara, A. Sakakura, in *Science of Synthesis, Stereoselective Synthesis*, Eds.: J. G. De Vries, G. A. Molander, P. A. Evans, **2011**, Vol. 3, pp 67-123; b) H. Du, K. Ding, in *Handbook of Cyclization Reactions*, Ed.: S. Ma, **2010**, Vol. 1, pp 1-57.
- [2] K. Mikami, K. Aikawa, in *Catalytic Asymmetric Synthesis (3rd Edition)*, Ed.: I. Ojima, **2010**, 683-737.
- [3] a) J.-L. Li, T.-Y. Liu, Y.-C. Chen, *Acc. Chem. Res.* **2012**, *45*, 1491-1500; b) Klier, L.; Tur, F.; Poulsen, P. H.; Jørgensen, K. A., *Chem. Soc. Rev.* **2017**, *46*, 1080.
- [4] J. Shen, C.-H. Tan, *Org. Biomol. Chem.* **2008**, *6*, 3229-3236; c) V. Marcos, J. Aleman, *Chem. Soc. Rev.* **2016**, *45*, 6812-6832.
- [5] V. H. Rawal, A. N. Thadani, in *Asymmetric Synthesis (2nd Edition)*, Ed.: M. Christmann, S. Bräse, **2008**, pp 144-148.
- [6] a) Achiral stoichiometric LA: K. Narasaka, S. Shimada, K. Osoda, N. Iwasawa, *Synthesis* **1991**, 1171-1172; b) K. C. Nicolau, J. J. Liu, C.-K. Hwang, W.-M. Dai, R. K. Guy, *J. Chem. Soc., Chem. Commun.* **1992**, 1118-1119; c) chiral stoichiometric LA: D. E. Ward, M. S. Souweha, *Org. Lett.* **2005**, *7*, 3533-3536; d) catalytic asymmetric method: J. Ishihara, S. Nakadachi, Y. Watanabe, S. Hatakeyama, *J. Org. Chem.* **2015**, *80*, 2037-2041.
- [7] Selected reviews on catalytic asymmetric DAs: a) S. Reymond, J. Cossy, *Chem. Rev.* **2008**, *108*, 5359-5406; b) X. Liu, H. Zheng, Y. Xia, L. Lin, X. Feng, *Acc. Chem. Res.* **2017**, *50*, 2621-2631; c) H. Du, K. Ding, in *Comprehensive Enantioselective Organocatalysis*, Ed.: P. I. Dalko, Wiley-VCH, Weinheim, **2013**, Vol. 3, pp 1131-1162.
- [8] Michael additions: F. Willig, J. Lang, A. C. Hans, M. R. Ringenberg, D. Pfeffer, W. Frey, R. Peters, *J. Am. Chem. Soc.* **2019**, *141*, 12029-12043.
- [9] Lewis acid / Brønsted base catalysis: M. Shibasaki, N. Kumagai, in *Cooperative Catalysis – Designing Efficient Catalysts for Synthesis*; R. Peters (Ed.), Wiley-VCH, Weinheim, **2015**.
- [10] For more cooperative polyfunctional LA / azolium / H bond catalysts developed by our group, see: a) M. Mechler, R. Peters, *Angew. Chem. Int. Ed.* **2015**, *54*, 10303-10307; b) J. Schmid, T. Junge, J. Lang, W. Frey, R. Peters, *Angew. Chem. Int. Ed.* **2019**, *58*, 5447-5451.
- [11] Related cooperative bi-/polyfunctional LA catalysts bearing cationic side arms developed by our group: a) D. Brodbeck, S. Álvarez-Barcia, J. Meisner, F. Broghammer, J. Klepp, D. Garnier, W. Frey, J. Kästner, R. Peters, *Chem. Eur. J.*, **2019**, *25*, 1515-1524; b) D. Brodbeck, F. Broghammer, J. Meisner, J. Klepp, D. Garnier, W. Frey, J. Kästner, R. Peters, *Angew. Chem. Int. Ed.* **2017**, *56*, 4056-4060; c) F. Broghammer, D. Brodbeck, T. Junge, R. Peters, *Chem. Commun.* **2017**, *53*, 1156-1159; d) P. Meier, F. Broghammer, K. Latendorf, G. Rauhut, R. Peters, *Molecules* **2012**, *17*, 7121-7150; e) T. Kull, J. Cabrera, R. Peters, *Chem. Eur. J.* **2010**, *16*, 9132-9139; f) T. Kull, R. Peters, *Angew. Chem. Int. Ed.* **2008**, *47*, 5461-5464; for related polymetallic catalysts, see e.g.: g) J. Schmid, W. Frey, R. Peters, *Organometallics* **2017**, *36*, 4313-4324; h) M. Mechler, W. Frey, R. Peters, *Organometallics* **2014**, *33*, 5492-5508; i) M. Mechler, K. Latendorf, W. Frey, R. Peters, *Organometallics* **2013**, *32*, 112-130.
- [12] T. Suzuki, S. Watanabe, S. Kobayashi, K. Tanino, *Org. Lett.* **2017**, *19*, 922-925.
- [13] H. Shimizu, H. Okamura, N. Yamashita, T. Iwagawa, M. Nakatani, *Tetrahedron Lett.* **2001**, *42*, 8649-8651.
- [14] K. Afarinkia, V. Vinader, T. D. Nelson, G. H. Posner, *Tetrahedron* **1992**, *48*, 9111-9171.
- [15] H. Okamura, T. Iwagawa, M. Nakatani, *Tetrahedron Lett.* **1995**, *36*, 5939-5942.
- [16] a) With maleimide: H. Okamura, Y. Nakamura, T. Iwagawa, M. Nakatani, *Chem. Lett.* **1996**, 193-194; b) L. Liu, Y. Cotellet, A.-B. Bornhof, C. Besnard, N. Sakai, S. Matile, *Angew. Chem. Int. Ed.* **2017**, *56*, 13066-13069; c) J. Y.-T. Soh, C.-H. Tan, *J. Am. Chem. Soc.* **2009**, *131*, 6904-6905; d) use of chiral N-acryloyloxazolidinones in combination with cinchona alkaloids: see ref. 13 and H. Okamura, K. Morishige, T. Iwagawa, M. Nakatani, *Tetrahedron Lett.* **1998**, *39*, 1211-1214; e) use of an α -chloroacrylate: see ref. 12, f) use of enones and α,β -unsaturated nitriles: Y. Wang, H. Li, Y.-Q. Wang, Y. Liu, B. M. Foxman, L. Deng, *J. Am. Chem. Soc.* **2007**, *129*, 6364-6365; g) R. P. Singh, K. Bartelson, Y. Wang, H. Su, X. Lu, L. Deng, *J. Am. Chem. Soc.* **2008**, *130*, 2422-2423; h) L.-M. Shi, W.-W. Dong, H.-Y. Tao, X.-Q. Dong, C.-J. Wang, *Org. Lett.* **2017**, *19*, 4532-4535; i) use of nitroolefins: K. J. Bartelson, R. P. Singh, B. M. Foxman, L. Deng, *Chem. Sci.* **2011**, *2*, 1940-1944; j) for a non-enantioselective method with maleic anhydride, see: E. J. Corey, A. P. Kozikowski, *Tetrahedron Lett.* **1975**, *16*, 2389-2392.
- [17] Review: P. Chauhan, J. Kaur, S. S. Chimni, *Chem. Asian J.* **2013**, *8*, 328
- [18] H. Okamura, H. Shimizu, Y. Nakamura, T. Iwagawa, M. Nakatani, *Tetrahedron Lett.* **2000**, *51*, 4147-4150.
- [19] N. Tienabe Kipassa, H. Okamura, K. Kina, T. Hamada, T. Iwagawa, *Org. Lett.* **2008**, *10*, 815-816.
- [20] a) see [16c]; for early studies: b) H. Okamura, H. Nagaike, T. Iwagawa, M. Nakatani, *Tetrahedron Lett.* **2000**, *41*, 8317-8321; additional studies: c) T. Takahashi, U. V. S. Reddy, Y. Kohari, C. Seki, T. Furuyama, N.

COMMUNICATION


- Kobayashi, Y. Okuyama, E. Kwon, K. Uwai, M. Tokiwa, M. Takeshita, H. Nakano, *Tetrahedron Lett.* **2016**, *57*, 5771–5776; d) M. Chennapuram, U. V. S. Reddy, C. Seki, Y. Okuyama, E. Kwon, K. Uwai, M. Tokiwa, M. Takeshita, H. Nakano, *Eur. J. Org. Chem.* **2017**, 4633–4641.
- [21] The absolute configuration for *ent*-**5aC** (Table 2, entry 11, CCDC 2003855) and *ent*-**3aC** (Table 3, entry 10, CCDC 1995951) were determined by X-ray crystal structure analysis, see www.ccdc.cam.ac.uk/data_request/cif.
- [22] Absolute configuration of **9** (CCDC 2002396) and **11a** (after reduction of the keto group (see SI), CCDC 1998669) were determined by X-ray analysis, see www.ccdc.cam.ac.uk/data_request/cif.
- [23] M. E. Salvati, R. M. Attar, M. M. Gottardis, J. A. Balog, D. A. Pickering, R. L. Martinez, C. Sun, PCT Int. Appl. 2002, WO 2002067939 A1.
- [24] J.-G. Fu, Y.-F. Shan, W.-B. Sun, G.-Q. Lin, B.-F. Sun, *Org. Biomol. Chem.* **2016**, *14*, 5229–5232.
- [25] For the parent cyclopentenone, the *endo*-isomer could be formed: R. Mose, M. E. Jensen, G. Preegel, K. A. Jørgensen, *Angew. Chem. Int. Ed.* **2015**, *54*, 13630–13634.
- [26] Ammonium organocatalysts: D. Uraguchi, T. Ooi, *J. Synth. Org. Chem. Jpn.* **2018**, *76*, 1144–1153.
- [27] In experiments using Cu(II) complexes with common enantiopure ligands only poor to low enantioselectivity was attained (see SI).
- [28] a) R. D. Baxter, D. Sale, K. M. Engle, J.-Q. Yu, D. G. Blackmond, *J. Am. Chem. Soc.* **2012**, *134*, 4600–4606; b) D. G. Blackmond, *J. Am. Chem. Soc.* **2015**, *137*, 10852–10866.
- [29] a) J. Burés, *Angew. Chem. Int. Ed.* **2016**, *55*, 16084–16087; b) C. D.-T. Nielsen, J. Burés, *Chem. Sci.* **2019**, *10*, 348–353.
- [30] For cooperative bimetallic enantioselective catalysts, see e.g.: a) M. Shibasaki, N. Yoshikawa, *Chem. Rev.* **2002**, *102*, 2187–2210; b) M. Shibasaki, S. Matsunaga, *Chem. Soc. Rev.* **2006**, *35*, 269–279; c) M. Shibasaki, M. Kanai, S. Matsunaga, N. Kumagai, *Acc. Chem. Res.* **2009**, *42*, 1117–1127; d) S. Handa, V. Gnanadesikan, S. Matsunaga, M. Shibasaki, *J. Am. Chem. Soc.* **2007**, *129*, 4900–4901; e) J. Park, S. Hong, *Chem. Soc. Rev.* **2012**, *41*, 6931–6943; f) S. Mouri, Z. Chen, H. Mitsunuma, M. Furutachi, S. Matsunaga, M. Shibasaki, *J. Am. Chem. Soc.* **2010**, *132*, 1255–1257.
- [31] CCDC 1995950, www.ccdc.cam.ac.uk/data_request/cif.

COMMUNICATION

Entry for the Table of Contents

COMMUNICATION

A very efficient way for catalytic asymmetric [4+2] cycloadditions is reported in which acidic prodienees are deprotonated by a betaine moiety to form a Cu dienolate, whereas the dienophile is activated by hydrogen bond formation.



Vukoslava Miskov-Pajic, Felix Willig,
Daniel M. Wanner, Wolfgang Frey, and
René Peters*

Page No. – Page No.

Enantiodivergent [4+2]-
Cycloadditions of Dienolates by
Polyfunctional Lewis-Acid /
Zwitterion Catalysts