

Communication

Installation of Minimal Tetrazines Through Silver-mediated Liebeskind-Srogl Coupling with Arylboronic Acids

William Lambert, Yinzhi Fang, Subham Mahapatra, Zhen Huang, Christopher W. am Ende, and Joseph M. Fox

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.9b08677 • Publication Date (Web): 11 Oct 2019

Downloaded from pubs.acs.org on October 12, 2019

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Installation of Minimal Tetrazines Through Silver-mediated Liebeskind-Srogl Coupling with Arylboronic Acids

William D. Lambert,[†] Yinzhi Fang,[†] Subham Mahapatra,[‡] Zhen Huang, [§] Christopher W. am Ende,^{*, ‡} and Joseph M. Fox^{*,†}

[†] Department of Chemistry and Biochemistry, University of Delaware, Newark, DE 19716, United States

* Pfizer Worldwide Research and Development, 445 Eastern Point Road, Groton, CT 06340, United States

[§] Pfizer Worldwide Research and Development, 1 Portland St, Cambridge, MA 02139, United States

Supporting Information Placeholder

ABSTRACT: Described is a general method for the installation of a minimal 6-methyltetrazin-3-yl group via the first example of a Ag-mediated Liebeskind-Srogl cross-coupling. The attachment of bioorthogonal tetrazines on complex molecules typically relies on linkers that can negatively impact the physiochemical properties of conjugates. Cross-coupling with arylboronic acids and a new reagent, 3-((p-biphenyl-4-ylmethyl)thio)-6-methyltetrazine (b-Tz), proceeds under mild, PdCl₂(dppf)-catalyzed conditions to introduce minimal, linkerfree tetrazine functionality. Safety considerations guided our design of b-Tz which can be prepared on decagram scale without handling hydrazine and without forming volatile, high-nitrogen tetrazine byproducts. Replacing conventional Cu(I) salts used in Liebeskind-Srogl cross-coupling with a Ag₂O mediator resulted in higher yields across a broad library of aryl and heteroaryl boronic acids and provides improved access to a fluorogenic tetrazine-BODIPY conjugate. A covalent probe for MAGL incorporating 6-methyltetrazinyl functionality was synthesized in high yield and labeled endogenous MAGL in live cells. This new Ag-mediated cross-coupling method using b-Tz is anticipated to find additional applications for directly introducing the tetrazine subunit to complex substrates.

The bioorthogonal reactions of tetrazines have emerged as important tools for chemical biology over the last decade.¹⁻⁶ Cycloadditions involving a range of dienophiles including *trans*-cyclooctenes^{1, 7-10}, cyclopropenes¹¹⁻¹² and norbornenes¹³ have been developed as tools for a variety of applications including cellular labeling¹⁴⁻¹⁷, *in vivo* imaging¹⁸⁻²⁰, unnatural amino acid mutagenesis^{3, 21-22}, targeted drug delivery²³⁻²⁵, proteomics²⁶, as





well as in the fabrication and patterning of biomaterials²⁷. Tetrazines themselves have also found applications in explosives technology²⁸, in metal-organic frameworks²⁹ and in natural product synthesis.³⁰

Conjugates of tetrazines are frequently prepared by amide bond forming reactions as represented in Figure 1A. A major limitation of this approach is that large and hydrophobic linkers can negatively impact the physiochemical properties of an attached ligand.^{6,17} Complementary new methods for the introduction of minimal tetrazines to small molecules may further advance their potential as bioorthogonal probes and chemical reporters. The replacement of bulky derivatives with smaller tetrazines has resulted in fluorophores with improved fluorogenic and cellular wash-out properties³¹⁻³³, better substrates for enzyme-catalyzed protein modification,^{17, 34} and probes for ¹⁸F-PET imaging³⁵. However, there are currently few methods



Figure 2: (A) Tetrazine synthesis based on condensation of nitriles or Pinner reagents with hydrazine (B) Cross-couplings of tetrazine electrophiles with arylboronic acids have been limited to *N*-substituted tetrazines, which are deactivated for bioorthogonal chemistry applications. Stille coupling has been used to couple 3-bromo-6-methyl-tetrazine to fluorophores.

25%

момо

ACS Paragon Plus Environment

for the direct attachment of 'minimal' tetrazine fragments to target molecules.³⁶ Additionally, many approaches to tetrazine synthesis produce high-nitrogen byproducts and involve harsh reaction conditions that can limit scalability and scope. Herein, we describe the decagram synthesis and thermal stability of 3-((*p*-biphenyl-4-ylmethyl)thio)-6-methyltetrazine, (b-Tz, **1a**) and a method to directly introduce the 6-methyltetrazin-3-yl group to arylboronic acids through the first example of a Agmediated Liebeskind-Srogl reaction (Fig 1B).

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

Classical tetrazine synthesis involves the condensation of Pinner salts or nitriles with excess hydrazine followed by oxidation.³⁷⁻³⁸ Catalytic nitrile condensation with neat anhydrous hydrazine, most notably with Zn(OTf)₂ and Ni(OTf)₂, has expanded access to unsymmetrical tetrazines.³⁹ Further, thiol catalysis has been shown to promote tetrazine synthesis from nitriles using hydrazine-hydrate.40 The most practiced procedures utilize excess acetonitrile or formamidine acetate and produce volatile tetrazine byproducts with high-nitrogen content (Fig 2A). Recently, a sulfur-catalyzed reaction of nitriles with hydrazine hydrate and dichloromethane has been described for 3-aryltetrazine synthesis.41 A safety consideration for all of these procedures is the direct addition of an oxidant to a reaction mixture containing hydrazine. While these methods for preparing tetrazines have been transformative to the field of bioorthogonal chemistry, there is a continuing need for safer alternatives with complementary functional group compatibility.

Tetrazines have been used in a limited number of metal catalyzed CH activations, 42-43 cross-couplings, 32-33, 44-51 and in Heck reactions with in situ generated 3-vinyl-6-methyltetrazine³². Recently, 3-amino-6-chlorotetrazines have been cross-coupled under Suzuki conditions (Fig 2B).45 Liebeskind-Srogl crosscouplings have also been reported with 3-amino-6-thiomethyltetrazines at 200 °C (Fig 2B).⁴⁶ The 3-aminotetrazine products of these methods are valuable in medicinal chemistry, but their utility in bioorthogonal chemistry is attenuated by the deactivating amino substituent.^{21, 32} The tetrazines most useful to bioorthogonal chemistry are also sensitive to basic conditions, making them incompatible with many conditions commonly associated with cross-coupling chemistry. Currently, there is a single method of cross-coupling to introduce a 3-methyltetrazine fragment via Stille coupling with 3-bromo-6-methyltetrazine, which is prepared from 3-hydrazino-6-methyltetrazine (Fig 2B).33

We considered that 3-thioalkyl-6-methyltetrazines might serve as useful reagents for the preparation of 3-aryl-6-methyltetrazines, which are attractive bioorthogonal reagents due to their balance of rapid kinetics toward dienophiles and high stability in the cellular environment.^{17, 21, 52} By modifying a method for the synthesis of 3-thiomethyl-6-methyltetrazine,⁵³ we prepared compounds **1a-g** with the rationale that a sacrificial *S*-benzylic substituent could serve to tune cross-coupling efficiency and improve the safety profile of the tetrazine. As



Figure 3: Decagram synthesis and thermal stability of b-Tz (**1a**).

shown in Figure 3, the 4-phenylbenzyl derivative b-Tz (**1a**) was prepared on large scale via alkylation of commercially available thiocarbohydrazide⁵⁴ with 4-bromomethylbiphenyl followed by one-pot condensation with triethylorthoacetate and a novel Cu(OAc)₂-catalyzed air-oxidation of the dihydrotetrazine intermediate. b-Tz was isolated on 27 gram scale with a 47% overall



Figure 4: (A) Rapid decomposition of b-Tz in CuTC. (B) Proposed Liebeskind-Srogl transmetallation mechanism. (C) Optimized Pd-catalyzed cross-coupling of tetrazines b-Tz and **1b** with various nucleophiles (yields determined by GC w/dodecane as a standard). *Conditions*: (a) Pd₂dba₃ (12.5 mol%), Cs₂CO₃ (3.0 eq.), dioxane, 70 °C, 90 min. (b) [Pd(allyl)Cl]₂ (10 mol%), THF, 70 °C, 2 h. (c) Pd(OAc)₂ (10 mol%), TBAF (1.0 eq.), dioxane, 70 °C, 20 h. (d) Pd₂dba₃ (15 mol%), DMF, 60 °C, 20 h. (e) DMF, 60 °C, 20 h. (D) Screening of silver(I) and copper(I) additives for condition e. (E) Screening of tetrazines **1a-g** under condition e. 1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24 25

26

27

28

29

30

31

32

33

34 35

36 37

42 43

44

45

46 47 48

49

50

51

52

53

54

55

56

57

58

59

60

yield after simple silica plug filtration and is a bench-stable crystalline solid (m.p. 141°C). The differential scanning calorimetry (DSC) profile of b-Tz has an onset temperature of 170 °C and a transition enthalpy of 900 J/g and is not flagged as potentially shock sensitive or explosive by a modified Yoshida correlation (Fig S-11).⁵⁵

After extensive screening (Fig S-3 thru S-8), we found copper(I)-mediated Liebeskind-Srogl conditions⁵⁶⁻⁵⁷ could promote cross-coupling of benzylic thioether tetrazines with PhB(OH)₂, PhSnBu₃, and PhSi(OMe)₃ (Fig 4C entries 1-3). Under Cu-mediated conditions tetrazine **1b** was the best substrate; however, the generality under these conditions was modest. The rapid consumption of tetrazine starting materials during the reaction led us to test if Cu(I) was causing decomposition of the reagent. Indeed, heating b-Tz with Cu(I)-thiophene carboxylate (CuTC) at 70 °C resulted in rapid decomposition and produced 4-phenylbenzaldehyde as the only identifiable side product (Fig 4A).

Copper has been proposed to promote the Liebeskind-Srogl reaction by facilitating transmetallation as shown in Figure 4B.⁵⁷⁻ ⁵⁸ We hypothesized that silver(I) salts might be similarly capable as promotors, whereby transmetallation would be promoted in a dual role by the thiophilic capture of benzylic thiolate by silver and the borophilic capture by oxygen. Ag(I) additives have been shown to promote Rh-catalyzed coupling of arylboronic acids with arylmethylsulfides bearing *ortho*-

directing groups⁵⁹⁻⁶⁰, and the Cu-catalyzed coupling of arylboronic acids with aromatic thioesters⁶¹. To our knowledge, a Ag-mediated variation of the Liebeskind-Srogl reaction has not been reported. After extensive optimization (Fig 4C entries 4-7, S-1 and S-2), PdCl₂(dppf) (15 mol%) was found to be especially effective for cross-coupling of 3-thioalkyl-6-methyltetrazines with arylboronic acids in polar, aprotic solvents (DMF, DMSO) at 60 °C. A screening of silver(I) additives revealed Ag₂O as the most general promotor, although Ag₂CO₃ was also effective (Fig 4D). Substitution of Ag₂O by Cu₂O gave only trace product formation. Arylboronic acids are particularly effective nucleophiles, whereas PhBF₃K and PhBPin were both less effective under identical reaction conditions (Fig 4C entries 8-9). Further, a series of 3-arylmethyl-6-methyltetrazines **1a-g** were evaluated as coupling partners (Fig 4E). Of these, the 4-phenylbenzyl derivative b-Tz (1a) was identified as the substrate with both the best cross-coupling yield as well as most favorable thermal stabiliy. We also note that the cost of Ag₂O (currently <\$3/g) is similar to the common promotor CuTC, and is minor in the context of bioorthogonal chemistry reagents which are typically required only in small amounts.

The scope of the Ag-mediated, Pd-catalyzed coupling of b-Tz with arylboronic acids is summarized in Figure 5A. Successful reactions were observed for arylboronic acids containing chloro-, fluoro-, secondary and tertiary amino-, alcohol, ether, nitro, sulfonyl, thioether, nitrile, aldehyde, ester, ketone, carbamate, and styryl groups. Heterocyclic functionality tolerated on



Figure 5: Reaction scope of b-Tz (5A) and **3** (5B). Typical conditions: thioalkyl tetrazine b-Tz or **3** (1.0 equiv.), RB(OH)₂ (1.9 equiv.), PdCl₂(dppf) (15 mol%), Ag₂O (2.5 equiv.), DMF (0.1M), 60°C, 19-21h, average isolated yields of duplicate synthesis (± 5%). ^a 3.0 equiv. of RB(OH)₂. ^b 3.0 equiv. of RB(OH)₂ did not significantly improve yield (< 5%), 1.9 equiv. of RB(OH)₂ was used.

the boronic acid component included quinoline, indole, pyridine, triazole, *N*-methylimidazole, furan and thiophene groups. The protected amino acid **2ae** coupled with b-Tz in 96% yield. Estrone-tetrazine **2ag** was also synthesized in 61%. In general, couplings were carried out using 1.9 equiv. of boronic acid, but 3.0 equiv. were utilized in reactions where homocoupling of the boronic acid was pronounced. *ortho*-Substituted heteroatoms had a deleterious impact with a relatively low yield observed for *ortho*-methoxy tetrazine **2k** and only trace product with *N*-Boc-2-aminophenylboronic acid and 2-hydroxyphenylboronic acid. While protected thiol and amine functionality was well tolerated (Fig 5), additives with free thiol or primary alkyl amine groups were not (Fig S-19). Also unsuccessful were 2pyridyl- and 4-pyridylboronic acids which are regarded as problematic across other cross-coupling reactions.⁶²

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31 32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

This cross-coupling method is not limited to *S*-benzylic thioethers or methyl-substituted tetrazines. 3-(Methylthio)-6-phenyl-tetrazine (**3**) was prepared from triethyl orthobenzoate and evaluated as a reagent in the synthesis of diaryltetrazines (Fig 5B). Successful reactions were observed for arylboronic acids bearing chloro-, alcohol, carbamate, ester, indole and ether groups with yields comparable to b-Tz. Included is an improved synthesis of 3-(4-hydroxymethylphenyl)-6-phenyltetrazine (**4c**), which is used to create cell-contact guiding microfibrous materials for tissue-culture applications.²⁷

We sought to demonstrate the application of b-Tz for the construction of fluorophore-tetrazine conjugates—compounds that have utility in live cell imaging.¹⁵ BODIPY-dye **6** with a directly attached tetrazine has been developed as 'superbright' bioorthogonal probe for fluorogenic labeling in live cells.³¹ The condensation of nitriles with hydrazine produces **6** in 8% yield.³¹ As shown in Figure 6, compound **6** can be accessed in 78% yield through the Ag-mediated cross-coupling of boronic acid **5** with b-Tz.

To demonstrate the utility of b-Tz in synthesizing chemical probes for studying endogenous levels of a protein in a native biological system, we constructed a tetrazine probe for monoacylglycerol lipase (MAGL). MAGL is a serine hydrolase in the endocannabinoid signaling pathway, and has attracted increasing interest as a target for neurological and metabolic disorders.⁶³ We designed a MAGL probe (9) by appending a 6-methyltetrazine moiety to a pyrazolylpiperidine scaffold with an electrophilic hexafluoroisopropyl (HFIP) carbamate warhead for covalently labeling the active site serine (Fig 7A).64 Synthesis was accomplished by cross-coupling of b-Tz with boronic acid 7 resulting in a 77% yield of 8. The reactive HFIP carbamate was installed by Boc deprotection followed by in situ addition to a triphosgene and hexafluoroisopropanol mixture, giving the MAGL reactive probe 9 in 78% yield. The reaction rate of 9 toward trans-cyclooctene is similar to that of 3-methyl-6-(4-aminomethyl)tetrazine (krel 1.1, Fig S-20).65 Probe 9 inhibited MAGL activity with 31 nM IC50 in an in vitro assay.66



Figure 6: Synthesis of 3-BODIPY-6-methyltetrazine 6.



Figure 7: (A) Synthesis of MAGL reactive probe **9**. (B) Live cells were treated with probe **9** for 1 h, followed by $2 \mu M$ TCO-TAMRA for 30 min, cell lysis, and analysis by in-gel fluorescence (C) In-gel fluorescence signals for a dose response of probe **9**. Probe **9** (32 nM, 1 h) was competed by pre-treatment with MAGL inhibitor KML29 (300 nM, 1 h). (D) KML29 also incorporates a HFIP carbamate warhead. (E) Dose response fitting of the fluorescence signals of MAGL normalized by the total protein amount indicated by Coomassie staining. Data are reported as mean ± SEM (n = 2). See Fig S-21 for Coomassie staining.

To test the labeling of endogenous MAGL in live cells, human brain vascular pericytes were treated with probe **9** for 1 h, followed by labeling with 2 μ M of TCO-TAMRA for 30 min in live cells (Fig 7B). After cell lysis, MAGL labeling was assessed with a gel-based activity-based protein profiling (ABPP) analysis (Fig 7C-E).⁶⁷ Strong fluorescence signals were observed for MAGL with minimal non-specific labeling from TCO-TAMRA. The labeling by probe **9** was dose responsive with a cellular IC₅₀ of 8 nM, and was competed by a MAGL inhibitor, KML29.⁶⁴ The HFIP warhead also labeled an additional protein at ~35 kDa, which is consistent with its reactivity with α/β -hydrolase domain 6 (ABHD6), and other off-targets at higher concentrations.^{64, 67}

1

2

3

4

5

6

7

8

9

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

58 59

60

In summary, a method has been described for installing a minimal 6-methyltetrazinyl-3-yl group through the first Ag-mediated Liebeskind-Srogl cross-coupling. A combination of PdCl₂(dppf) catalyst and Ag₂O mediator was found to be uniquely effective for coupling 3-thioalkyl-6-methyltetrazines with arylboronic acids. Safety-testing guided our design of the reactive substrate b-Tz (1a), which can be synthesized from commercially available materials on decagram scale in 47% overall yield. Cross-coupling of b-Tz with boronic acids proceeds under mild conditions with broad functional group tolerance. Alternatively, 3-(Methylthio)-6-phenyl-tetrazine (3) undergoes cross-coupling with arylboronic acids to give 3,6-dia-10 ryltetrazines. Application to the synthesis of chemical biology 11 tools was demonstrated. A BODIPY-tetrazine conjugate was 12 synthesized in 78% yield—substantially higher than what is 13 possible using traditional hydrazine-based synthesis. Finally, a 14 tetrazine-functionalized probe for MAGL was synthesized in 15 high yield and was shown to covalently label endogenous 16 MAGL with good selectivity in live cells. We anticipate that this method for introducing minimal tetrazines to chemical probes 17 will serve as an important tool for studying protein targets at 18 endogenous levels in their native environment. 19

ASSOCIATED CONTENT

Supporting Information

Optimization tables, DSC data, experimental procedures, kinetics, inhibition assays, and ¹H and ¹³C NMR data are displayed. The Supporting Information is available free of charge on the ACS Publications website at DOI: xxxxxxxxxxxx

AUTHOR INFORMATION

Corresponding Author

*imfox@udel.edu, *christopher.amEnde@pfizer.com

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

We would like to thank Jeffery Sperry for helpful discussions, Remzi Duzguner for DSC analysis, and Lucy Stevens for the in vitro potency data on probe 9. This work was supported by NIH GM132460, NSF 1809612 and Pfizer. Instrumentation was supported by NIH P20GM104316, P30GM110758, S10RR026962, S100D016267 and NSF CHE-0840401, CHE-1229234, and CHE-1048367.

REFERENCES

- (1) Selvaraj, R.; Fox, J. M., trans-Cyclooctene--a stable, voracious dienophile for bioorthogonal labeling. Curr. Opin. Chem. Biol. 2013, 17, 753-760.
- (2) Patterson, D. M.; Nazarova, L. A.; Prescher, J. A., Finding the right (bioorthogonal) chemistry. ACS Chem. Biol. 2014, 9, 592-605.
- (3) Lang, K.; Chin, J. W., Bioorthogonal reactions for labeling proteins. ACS Chem. Biol. 2014, 9, 16-20.
- (4) Oliveira, B. L.; Guo, Z.; Bernardes, G. J. L., Inverse electron demand Diels-Alder reactions in chemical biology. Chem. Soc. Rev. 2017, 46, 4895-4950.
- 53 (5) Wu, H.; Devaraj, N. K., Advances in Tetrazine Bioorthogonal Chemistry Driven by the Synthesis of Novel Tetrazines and 54 Dienophiles. Acc. Chem. Res. 2018, 51, 1249-1259. 55
- (6) Cañeque, T.; Müller, S.; Rodriguez, R., Visualizing biologically active 56 small molecules in cells using click chemistry. Nat. Rev. Chem. 2018, 2, 57 202-215.

(7) Blackman, M. L.; Royzen, M.; Fox, J. M., Tetrazine ligation: fast bioconjugation based on inverse-electron-demand Diels-Alder reactivity. J. Am. Chem. Soc. 2008, 130, 13518-13519.

(8) Taylor, M. T.; Blackman, M. L.; Dmitrenko, O.; Fox, J. M., Design and synthesis of highly reactive dienophiles for the tetrazine-transcyclooctene ligation. J. Am. Chem. Soc. 2011, 133, 9646-9649.

(9) Darko, A.; Wallace, S.; Dmitrenko, O.; Machovina, M. M.; Mehl, R. A.; Chin, J. W.; Fox, J. M., Conformationally Strained trans-Cyclooctene with Improved Stability and Excellent Reactivity in Tetrazine Ligation. Chem. Sci. 2014, 5, 3770-3776.

(10) Lambert, W. D.; Scinto, S. L.; Dmitrenko, O.; Boyd, S. J.; Magboo, R.; Mehl, R. A.; Chin, J. W.; Fox, J. M.; Wallace, S., Computationally guided discovery of a reactive, hydrophilic trans-5-oxocene dienophile for bioorthogonal labeling. Org. Biomol. Chem. 2017, 15, 6640-6644.

(11) Patterson, D. M.; Nazarova, L. A.; Xie, B.; Kamber, D. N.; Prescher, J. A., Functionalized cyclopropenes as bioorthogonal chemical reporters. J. Am. Chem. Soc. 2012, 134, 18638-18643.

(12) Yang, J.; Seckute, J.; Cole, C. M.; Devaraj, N. K., Live-cell imaging of cyclopropene tags with fluorogenic tetrazine cycloadditions. Angew. Chem. Int. Ed. 2012, 51, 7476-7479.

(13) Devaraj, N. K.; Weissleder, R.; Hilderbrand, S. A., Tetrazine-based cycloadditions: application to pretargeted live cell imaging. Bioconjug. Chem. 2008, 19, 2297-2299.

(14) Devaraj, N. K.; Upadhyay, R.; Haun, J. B.; Hilderbrand, S. A.; Weissleder, R., Fast and sensitive pretargeted labeling of cancer cells through a tetrazine/trans-cyclooctene cycloaddition. Angew. Chem. Int. Ed. 2009, 48, 7013-7016.

(15) Devaraj, N. K.; Hilderbrand, S.; Upadhyay, R.; Mazitschek, R.; Weissleder, R., Bioorthogonal turn-on probes for imaging small molecules inside living cells. Angew. Chem. Int. Ed. 2010, 49, 2869-2872.

(16) Murrey, H. E.; Judkins, J. C.; am Ende, C. W.; Ballard, T. E.; Fang, Y.; Riccardi, K.; Di, L.; Guilmette, E. R.; Schwartz, J. W.; Fox, J. M.; Johnson, D. S., Systematic Evaluation of Bioorthogonal Reactions in Live Cells with Clickable HaloTag Ligands: Implications for Intracellular Imaging. J. Am. Chem. Soc. 2015, 137, 11461-11475.

(17) Baalmann, M.; Ziegler, M. J.; Werther, P.; Wilhelm, J.; Wombacher, R., Enzymatic and Site-Specific Ligation of Minimal-Size Tetrazines and Triazines to Proteins for Bioconjugation and Live-Cell Imaging. Bioconjug. Chem. 2019, 30, 1405-1414.

(18) Rossin, R.; Verkerk, P. R.; van den Bosch, S. M.; Vulders, R. C.; Verel, I.; Lub, J.; Robillard, M. S., In vivo chemistry for pretargeted tumor imaging in live mice. Angew. Chem. Int. Ed. 2010, 49, 3375-3378.

(19) Zeglis, B. M.; Sevak, K. K.; Reiner, T.; Mohindra, P.; Carlin, S. D.; Zanzonico, P.; Weissleder, R.; Lewis, J. S., A pretargeted PET imaging strategy based on bioorthogonal Diels-Alder click chemistry. J. Nucl. Med. 2013, 54, 1389-1396.

(20) Wang, M.; Vannam, R.; Lambert, W. D.; Xie, Y.; Wang, H.; Giglio, B.; Ma, X.; Wu, Z.; Fox, J.; Li, Z., Hydrophilic (18)F-labeled trans-5-oxocene (oxoTCO) for efficient construction of PET agents with improved tumor-to-background ratios in neurotensin receptor (NTR) imaging. Chem. Commun. 2019, 55, 2485-2488.

(21) Blizzard, R. J.; Backus, D. R.; Brown, W.; Bazewicz, C. G.; Li, Y.; Mehl, R. A., Ideal Bioorthogonal Reactions Using A Site-Specifically Encoded Tetrazine Amino Acid. J. Am. Chem. Soc. 2015, 137, 10044-10047.

(22) Hoffmann, J. E.; Plass, T.; Nikic, I.; Aramburu, I. V.; Koehler, C.; Gillandt, H.; Lemke, E. A.; Schultz, C., Highly Stable trans-Cyclooctene Amino Acids for Live-Cell Labeling. Chemistry 2015, 21, 12266-12270. (23) Li, J.; Chen, P. R., Development and application of bond cleavage reactions in bioorthogonal chemistry. Nat. Chem. Biol. 2016, 12, 129-137.

(24) Versteegen, R. M.; Rossin, R.; ten Hoeve, W.; Janssen, H. M.; Robillard, M. S., Click to release: instantaneous doxorubicin elimination upon tetrazine ligation. Angew. Chem. Int. Ed. 2013, 52, 14112-14116.

(25) Mejia Oneto, J. M.; Khan, I.; Seebald, L.; Royzen, M., In Vivo Bioorthogonal Chemistry Enables Local Hydrogel and Systemic Pro-Drug To Treat Soft Tissue Sarcoma. ACS Cent. Sci. 2016, 2, 476-482.

(26) Kang, K.; Park, J.; Kim, E., Tetrazine ligation for chemical proteomics. Proteome. Sci. 2016, 15, 1-13.

(27) Liu, S.; Moore, A. C.; Zerdoum, A. B.; Zhang, H.; Scinto, S. L.; Zhang, H.; Gong, L.; Burris, D. L.; Rajasekaran, A. K.; Fox, J. M.; Jia, X., Cellular interactions with hydrogel microfibers synthesized via interfacial tetrazine ligation. Biomaterials 2018, 180, 24-35.

(28) DeLuca, L. T.; Shimada, T.; Sinditskii, V. P.; Calabro, M., *Chemical Rocket Propulsion: A Comprehensive Survey of Energetic Materials*. Springer International Publishing: 2017.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

21

22

23

24

25

26

60

- (29) Xue, M.; Ma, S.; Jin, Z.; Schaffino, R. M.; Zhu, G. S.; Lobkovsky, E. B.; Qiu, S. L.; Chen, B., Robust metal-organic framework enforced by tripleframework interpenetration exhibiting high H2 storage density. *Inorg. Chem.* **2008**, *47*, 6825-6828.
- (30) Hamasaki, A.; Zimpleman, J. M.; Hwang, I.; Boger, D. L., Total synthesis of ningalin D. *J. Am. Chem. Soc.* **2005**, *127*, 10767-10770.
- (31) Carlson, J. C.; Meimetis, L. G.; Hilderbrand, S. A.; Weissleder, R., BODIPY-tetrazine derivatives as superbright bioorthogonal turn-on probes. *Angew. Chem. Int. Ed.* **2013**, *52*, 6917-6920.
- (32) Wu, H.; Yang, J.; Seckute, J.; Devaraj, N. K., In situ synthesis of alkenyl tetrazines for highly fluorogenic bioorthogonal live-cell imaging probes. *Angew. Chem. Int. Ed.* **2014**, *53*, 5805-5809.
- (33) Wieczorek, A.; Werther, P.; Euchner, J.; Wombacher, R., Green- to far-red-emitting fluorogenic tetrazine probes - synthetic access and nowash protein imaging inside living cells. *Chem. Sci.* **2017**, *8*, 1506-1510.
- (34) Best, M.; Degen, A.; Baalmann, M.; Schmidt, T. T.; Wombacher, R., Two-step protein labeling by using lipoic acid ligase with norbornene substrates and subsequent inverse-electron demand Diels-Alder reaction. *Chembiochem.* **2015**, *16*, 1158-1162.
- 17 (35) Denk, C.; Svatunek, D.; Filip, T.; Wanek, T.; Lumpi, D.; Frohlich, J.;
 18 Kuntner, C.; Mikula, H., Development of a (18) F-labeled tetrazine with
 19 favorable pharmacokinetics for bioorthogonal PET imaging. *Angew.*20 *Chem. Int. Ed.* 2014, *53*, 9655-9659.
 - (36) For the introduction of bioorthogonal triazines via Suzuki coupling, see: Kamber, D. N.; Liang, Y.; Blizzard, R. J.; Liu, F.; Mehl, R. A.; Houk, K. N.; Prescher, J. A., 1,2,4-Triazines Are Versatile Bioorthogonal Reagents. *J. Am. Chem. Soc.* **2015**, 137, 8388-91.
 - (37) Pinner, A., Ueber die Einwirkung von Hydrazin auf Imidoäther. *Ber. Dtsch. Chem. Ges.* **1893**, *26*, 2126.
 - (38) Tolshchina, S. G.; Rusinov, G. L.; Charushin, V. N., 1,2,4,5-Tetrazines and Azolo[1,2,4,5]tetrazines: Synthesis and Reactions with Nucleophiles. *Chem. Heterocycl. Compd.* **2013**, *49*, 66-91.
- Nucleophiles. Chem. Heterocycl. Compd. 2013, 49, 66-91.
 (39) Yang, J.; Karver, M. R.; Li, W.; Sahu, S.; Devaraj, N. K., Metalcatalyzed one-pot synthesis of tetrazines directly from aliphatic nitriles and hydrazine. Angew. Chem. Int. Ed. 2012, 51, 5222-5225.
- 30 (40) Mao, W.; Shi, W.; Li, J.; Su, D.; Wang, X.; Zhang, L.; Pan, L.; Wu, X.;
 31 Wu, H., Organocatalytic and Scalable Syntheses of Unsymmetrical 1,2,4,5-Tetrazines by Thiol-Containing Promotors. *Angew. Chem. Int. Ed.* 2019, *58*, 1106-1109.
- (41) Qu, Y.; Sauvage, F. X.; Clavier, G.; Miomandre, F.; Audebert, P.,
 Metal-Free Synthetic Approach to 3-Monosubstituted Unsymmetrical
 1,2,4,5-Tetrazines Useful for Bioorthogonal Reactions. *Angew. Chem. Int. Ed.* 2018, *57*, 12057-12061.
- (42) Testa, C.; Gigot, E.; Genc, S.; Decreau, R.; Roger, J.; Hierso, J. C.,
 Ortho-Functionalized Aryltetrazines by Direct Palladium-Catalyzed CHalogenation: Application to Fast Electrophilic Fluorination Reactions. Angew. Chem. Int. Ed. 2016, 55, 5555-5559.
- (43) Mboyi, C. D.; Testa, C.; Reeb, S.; Genc, S.; Cattey, H.; FleuratLessard, P.; Roger, J.; Hierso, J.-C., Building Diversity in orthoSubstituted s-Aryltetrazines By Tuning N-Directed Palladium C-H
 Halogenation: Unsymmetrical Polyhalogenated and Biphenyl sAryltetrazines. ACS Catalysis 2017, 7, 8493-8501.
- (44) Novak, Z.; Kotschy, A., First Cross-Coupling Reactions on
 Tetrazines. Org. Lett. 2003, 5, 3495-3497.
- (45) Bender, A. M.; Chopko, T. C.; Bridges, T. M.; Lindsley, C. W., Preparation of Unsymmetrical 1,2,4,5-Tetrazines via a Mild Suzuki Cross-Coupling Reaction. *Org. Lett.* 2017, *19*, 5693-5696.
- 47 (46) Suzenet, F.; Leconte, N.; Keromnes-Wuillaume, A.; Guillaumet, G.,
 48 Efficient Palladium-Catalyzed Synthesis of Unsymmetrical (Het)aryl-tetrazines. *Synlett* 2007, 2007, 204-210.
 49 (47) Pellegetti L.; Vedeprop. E.; Legger, M.; Lewy, C.; Peutier, S. New,
- (47) Pellegatti, L.; Vedrenne, E.; Leger, J.-M.; Jarry, C.; Routier, S., New
 heteroaromatic aminations on 5-aryl-1,2,4-triazines and 1,2,4,5tetrazines by palladium catalysis. *Tetrahedron* 2010, 66, 4383-4389.
- 52 (48) Quinton, C.; Alain-Rizzo, V.; Dumas-Verdes, C.; Clavier, G.; Vignau,
 53 L.; Audebert, P., Triphenylamine/tetrazine based π-conjugated systems as molecular donors for organic solar cells. *New J. Chem.* 2015,
 54 39, 9700-9713.
 55 (49) Wieczorek, A.; Buckup, T.; Wombacher, R., Rigid tetrazine
 - (49) Wieczorek, A.; Buckup, T.; Wombacher, R., Rigid tetrazine fluorophore conjugates with fluorogenic properties in the inverse

electron demand Diels-Alder reaction. Org. Biomol. Chem. 2014, 12, 4177-4185.

(50) Sołoducho, J.; Doskocz, J.; Cabaj, J.; Roszak, S., Practical synthesis of bis-substituted tetrazines with two pendant 2-pyrrolyl or 2-thienyl groups, precursors of new conjugated polymers. *Tetrahedron* **2003**, *59*, 4761-4766.

(51) Li, Z.; Ding, J.; Song, N.; Lu, J.; Tao, Y., Development of a New s-Tetrazine-Based Copolymer for Efficient Solar Cells. J. Am. Chem. Soc. **2010**, *132*, 13160-13161.

(52) Karver, M. R.; Weissleder, R.; Hilderbrand, S. A., Synthesis and evaluation of a series of 1,2,4,5-tetrazines for bioorthogonal conjugation. *Bioconjug. Chem.* **2011**, *22*, 2263-2270.

(53) Fields, S. C.; Parker, M. H.; Erickson, W. R., A Simple Route to Unsymmetrically Substituted 1,2,4,5-Tetrazines. *J. Org. Chem* **1994**, *59*, 8284-8287.

(54) DSC data for commercially available Thiocarbohydrazide: onset temp. of 172 °C, transition enthalpy of 1634 J/g. As a precaution, all experiments involving thiocarbohydrazide were carried out below 72 °C (100 °C below the DSC transition onset).

(55) Sperry, J. B.; Minteer, C. J.; Tao, J.; Johnson, R.; Duzguner, R.; Hawksworth, M.; Oke, S.; Richardson, P. F.; Barnhart, R.; Bill, D. R.; Giusto, R. A.; Weaver, J. D., Thermal Stability Assessment of Peptide Coupling Reagents Commonly Used in Pharmaceutical Manufacturing. *Org. Process Res. Dev* **2018**, *22*, 1262-1275.

(56) Cheng, H.-G.; Chen, H.; Liu, Y.; Zhou, Q., The Liebeskind-Srogl Cross-Coupling Reaction and its Synthetic Applications. *Asian J. Org. Chem.* **2018**, *7*, 490-508.

(57) Liebeskind, L. S.; Srogl, J., Heteroaromatic Thioether-Boronic Acid Cross-Coupling under Neutral Reaction Conditions. *Org. Lett.* **2002**, *4*, 979-981.

(58) Musaev, D. G.; Liebeskind, L. S., On the Mechanism of Pd(0)-Catalyzed, Cu(I) Carboxylate-Mediated Thioorganic-Boronic Acid Desulfitative Coupling. A Non-innocent Role for Carboxylate Ligand. *Organometallics* **2009**, *28*, 4639-4642.

(59) Hooper, J. F.; Young, R. D.; Pernik, I.; Weller, A. S.; Willis, M. C., Carbon–carbon bond construction using boronic acids and aryl methyl sulfides: orthogonal reactivity in Suzuki-type couplings. *Chem. Sci.* **2013**, *4*, 1568-1572.

(60) Pan, F.; Wang, H.; Shen, P.-X.; Zhao, J.; Shi, Z.-J., Cross coupling of thioethers with aryl boroxines to construct biaryls via Rh catalyzed C–S activation. *Chem. Sci.* **2013**, *4*, 1573-1577.

(61) Ghosh, P.; Ganguly, B.; Perl, E.; Das, S., A synthesis of biaryl ketones via the C-S bond cleavage of thiol ester by a Cu/Ag salt. *Tetrahedron Lett.* **2017**, *58*, 2751-2756.

(62) Markovic, T.; Rocke, B. N.; Blakemore, D. C.; Mascitti, V.; Willis, M. C., Pyridine sulfinates as general nucleophilic coupling partners in palladium-catalyzed cross-coupling reactions with aryl halides. *Chem. Sci.* **2017**, *8*, 4437-4442.

(63) Blankman, J. L.; Cravatt, B. F., Chemical probes of endocannabinoid metabolism. *Pharmacol. Rev.* **2013**, *65*, 849-871.

(64) Chang, J. W.; Niphakis, M. J.; Lum, K. M.; Cognetta, A. B., 3rd; Wang, C.; Matthews, M. L.; Niessen, S.; Buczynski, M. W.; Parsons, L. H.; Cravatt, B. F., Highly selective inhibitors of monoacylglycerol lipase bearing a reactive group that is bioisosteric with endocannabinoid substrates. *Chem. Biol.* **2012**, *19*, 579-588.

(65) Electron withdrawing aromatic substitutents have an accelerating effect on Diels-Alder rate constants with tetrazines, see ref 9 and ref 52.

(66) Butler, C. R.; Beck, E. M.; Harris, A.; Huang, Z.; McAllister, L. A.; Am Ende, C. W.; Fennell, K.; Foley, T. L.; Fonseca, K.; Hawrylik, S. J.; Johnson, D. S.; Knafels, J. D.; Mente, S.; Noell, G. S.; Pandit, J.; Phillips, T. B.; Piro, J. B.; Rogers, B. N.; Samad, T. A.; Wang, L.; W. C. D.

R.; Rogers, B. N.; Samad, T. A.; Wang, J.; Wan, S.; Brodney, M. A., Azetidine and Piperidine Carbamates as Efficient, Covalent Inhibitors of Monoacylglycerol Lipase. *J. Med. Chem.* **2017**, *60*, 9860-9873.

(67) Chang, J. W.; Cognetta, A. B., 3rd; Niphakis, M. J.; Cravatt, B. F., Proteome-wide reactivity profiling identifies diverse carbamate chemotypes tuned for serine hydrolase inhibition. *ACS Chem. Biol.* **2013**, *8*, 1590-1599.

-	For Table of Contents Only
-	b-Tz $\stackrel{CH_3}{\stackrel{N}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{r}{\underset{h}{\underset{h}{\underset{r}{\underset{h}}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\underset{h}{\atop_{h}}{\underset{h}{\underset{h}}{\underset{h}{\underset{h}}}}}}}}}$
-	