## Asymmetric Synthesis

## A Mild and Efficient CH<sub>2</sub>-Extrusion Reaction for the Enantiospecific Synthesis of Highly Configurationally Stable Tröger Bases\*\*

Sandip A. Pujari, Céline Besnard, Thomas Bürgi, and Jérôme Lacour\*

Dedicated to Professor Volker Schurig on the occasion of his 75th birthday

**Abstract:** A novel  $CH_2$ -extrusion reaction leading to the transformation of ethano-Tröger bases into disubstituted methano derivatives is reported (yields up to 93%). Under mild and metal-free oxidative conditions, a loss of  $CH_2$  and a ring contraction are provoked. Despite two bond cleavages at stereogenic nitrogen and carbon centers and a temporary rupture of the bicyclic structure, a very high enantiospecificity (es  $\geq$  98%) is observed for this unusual reaction.

**L**xtrusion reactions, which are formally the reverse of insertion processes, are characterized by the removal of an atom or a group of atoms from a given combination of functional groups and by the reformation of a bond between the atoms to which the departing moiety was attached.<sup>[1]</sup> Such processes are relatively common in organometallic and chalcogen chemistry.<sup>[2,3]</sup> With more classical organic substrates, only few CH<sub>2</sub> extrusions are reported using metal-free conditions and activated benzylamines and a-hydroxy carbonyl compounds as substrates.<sup>[4]</sup> Herein, in the context of nitrogen stereochemistry and Tröger base (TB) derivatives in particular, we report a novel methylene extrusion which sees the enantiospecific transformation of the ethano-TB 1 into the disubstituted methano-TB 2 (Scheme 1, top). Under mild oxidative conditions, the loss of CH<sub>2</sub> occurs with yields of up to 93%. Importantly, a very high enantiospecificity ( $es \ge$ 98%) is obtained in this reaction which proceeds with retention of configuration through a ring opening and two bond cleavages at stereogenic nitrogen and carbon centers.

Recently, it was shown that Tröger bases of type **3** (Scheme 1, bottom),<sup>[5]</sup> which are classical chiral molecules as

Dr. C. Besnard

Laboratory of Crystallography, University of Geneva Quai Ernest Ansermet 24, 1211 Geneva 4 (Switzerland) Prof. T. Bürgi

Department of Physical Chemistry, University of Geneva Quai Ernest Ansermet 30, 1211 Geneva 4 (Switzerland)

[\*\*] We thank the University of Geneva and the Swiss National Science Foundation for financial support. We are also grateful to Dr. Alain de Mesmaeker (Syngenta) and Dr. Théo Berclaz (Geneva) for fruitful discussions. We also acknowledge the contributions of the Sciences Mass Spectrometry (SMS) platform at the Faculty of Sciences, University of Geneva.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201500435.



**Scheme 1.** Extrusion of  $CH_2$  group by concomitant C–C and C–N bond-cleavage reactions (top). Insertion of carbene into a C–N bond (bottom). DDQ = 2,3-dichloro-5,6-dicyano-1,4-benzoquinone.

a result of the presence of stereogenic nitrogen atoms, react in a single step with metal carbenes to afford ethano-Tröger bases of type  $1^{[6]}$  With donor-acceptor diazo reagents as precursors, the enantiospecificity of the ring expansion is excellent (*es* 97–99%).<sup>[7]</sup> The compounds  $1^{[8]}$  which are 1,2diamines, do not racemize under Brønsted- (pH < 1) or Lewis-acidic conditions, which are known to promote the loss of enantiomeric purity of  $3^{[9]}$  These ethano-adducts 1 are essentially unknown. We wondered about their global reactivity and decided to use them as substrates in various reactions, including nitrene insertion reactions.<sup>[10]</sup>

The compound **1a** ( $\mathbf{R} = \mathbf{R}^1 = \mathbf{M}\mathbf{e}$ ,  $\mathbf{A}\mathbf{r} = \mathbf{P}\mathbf{h}$ ) was thus treated with PhI=NTs under rhodium(II) and copper(II) catalysis. To our surprise, instead of obtaining a product with a higher molecular weight of 169, the derivative 2a with a loss of a mass of 14 was isolated in good yield (Table 1, entries 1 and 2). This data indicated a probable loss of a CH<sub>2</sub> group, and was confirmed by <sup>1</sup>H and <sup>13</sup>C spectroscopic analyses. Only an interesting methano-Tröger base structure with two substituents on the aminal carbon atom was consistent with the data, and the motif was ascertained later by X-ray diffraction analysis (see Table 2).<sup>[11]</sup> Starting with (+)-1a (ee 99%), 2a was furthermore obtained with an interesting enantiomeric purity of 47 and 69% (Table 1, entries 1 and 2), thus denoting a rather effective transfer of chirality in this unusual extrusion reaction.<sup>[12]</sup> Trying to obtain some clues on the reaction mechanism and improve the results, (+)-1a was treated separately with an excess of Cu(OTf)<sub>2</sub> and PhI=NTs. Importantly, ring-contracted (+)-2a was isolated again (entries 3 and 4) and this indicated an oxidative nature for the transformation.<sup>[13]</sup> Manganese(III) and manganese(IV) complexes, in combination or without Cu(OTf)<sub>2</sub>, were tested

<sup>[\*]</sup> Dr. S. A. Pujari, Prof. J. Lacour Department of Organic Chemistry, University of Geneva Quai Ernest Ansermet 30, 1211 Geneva 4 (Switzerland) E-mail: jerome.lacour@unige.ch

Table 1: Optimization.[a]

Entry	Oxidant	Solvent	t [h]	<b>2 a</b> Yield [%] <sup>[e]</sup>	ee [%] <sup>[f]</sup>
1	PhINTs/[Rh2(esp)2]	$CH_2Cl_2$	24	62	47
2	PhINTs/Cu(OTf) <sub>2</sub>	CH₃CN	24	75	69
3	Cu(OTf) <sub>2</sub>	CH₃CN	4	30	83
4	PhINTs	CH₃CN	48	79	46
5	$Mn(OAc)_3 \cdot 2H_2O/Cu(OTf)_2$	CH₃CN	6	56	70
6	Mn(OAc)₃·2 H₂O	CH₃CN	48	trace	-
7 <sup>[b]</sup>	MnO <sub>2</sub>	$CH_2Cl_2$	48	trace	-
8 <sup>[c]</sup>	Iodine	THF	16	77	70
<b>9</b> <sup>[c]</sup>	NBS	CH₃CN	6	84	83
10 <sup>[c]</sup>	DDQ	CH₃CN	30	77	98
11 <sup>[c]</sup>	DDQ	$CH_3NO_2$	24	81	98
12 <sup>[c]</sup>	DDQ	CH <sub>3</sub> NO <sub>2</sub> <sup>[d]</sup>	8	85	98

[a] Reaction conditions: the ethano Tröger base (+)-1a (1 equiv), oxidant (2 equiv), solvent (0.05 m). [b] 10 equiv of MnO<sub>2</sub> was used. [c] 1.2 equiv of oxidant was used. [d] Nitromethane containing 1.3% of H<sub>2</sub>O. [e] Yield of isolated product. [f] Determined by HPLC using a chiral stationary phase. esp =  $\alpha$ , $\alpha'$ , $\alpha'$ -tetramethyl-1,3-benzenedipropionate, NBS = *N*-bromosuccinimide, Tf = trifluoromethanesulfonyl, THF = tetrahydrofuran, Ts = 4-toluenesulfonyl.

(entries 5–7).<sup>[14]</sup> Reactions either did not occur or proceeded with only moderate enantioselectivity (70% *ee*). Metal-free conditions were attempted. While iodine and NBS provided (+)-**2a** in good yields and interesting levels of chirality transfer (*ee* 70 and 83% respectively), DDQ (entries 10–12) offered the best results, particularly in wet nitromethane (1.3% H<sub>2</sub>O).<sup>[15]</sup> Under these reaction conditions, at 20°C, (+)-**2a** was obtained in 85% yield and with 98% *ee*. These optimized reaction conditions were used throughout the study.

The generality of the extrusion reaction was established by using a series of ethano-TB derivatives (1a-q). Electronrich and electron-poor substituents were introduced para to the bridgehead nitrogen atoms (Me, OMe and CO<sub>2</sub>Et). Care was also taken to prepare substrates with electron-donating and electron-withdrawing substituents on the aromatic group carried by the stereogenic carbon center. Most of these compounds were synthesized in both racemic and enantioenriched forms (96 to > 99% ee; see Table S1 in the Supporting Information).<sup>[6a]</sup> To our satisfaction, in the 2,8-dimethyl series (1a-i) the methylene extrusion proceeded well. Not too surprisingly, reactions were slightly faster with electron-rich substituents on the aromatic ring (e.g., OMe; 1c). Sterics did not significantly influence the reactivity and stereoselectivity (e.g., **2 f** and **2 g**).<sup>[16]</sup> Changing the ester functional group, from methyl to benzyl and allyl, did not affect the outcome of the reaction either. Very similar results were obtained with 2,8dimethoxy-substituted 1j-p. The reactions were faster thanks to the electron-donating p-MeO substituents (max. 4 h). On the contrary, a seven-day reaction was necessary with 1q to afford 2q in moderate yield (51%).<sup>[17]</sup> For mechanistic purposes, the benzoyl 1r and its reduced alcohol derivative 1s were also tested.<sup>[8c]</sup> While a slow but effective extrusion occurred with the first derivative, no trace of ring contraction could be detected with 1s. The necessity of an electron-





[a] Reaction conditions: the substrate 1 (1 equiv), DDQ (1.2 equiv), CH<sub>3</sub>NO<sub>2</sub>/H<sub>2</sub>O (1.3%) (0.05 M), 20 °C. Yield is that for product isolated after chromatography. The *ee* value was determined by CSP-HPLC. Used 4 equiv of DDQ in the case of 2q and 2r. Structures of substrates 1 are indicated in the Supporting Information.

withdrawing group on the ethano-bridge was confirmed by using  $\mathbf{1t}$ ,<sup>[8a]</sup> which was found to be unreactive.

Importantly, in all examples performed with enantioenriched substrates, an efficient chirality transfer was observed as enantiospecificity values higher than or equal to 98% were constantly obtained. To our surprise, the enantiospecificity was high even under Lewis-acidic conditions which are known



to promote the racemization of **3** (e.g. Cu(OTf)<sub>2</sub>, es 84%; Table 1, entry 3). The configurational stability of the 13,13'disubstituted derivatives was thus tested. Solutions in DMF (N,N-dimethylformamide) of (+)- and (-)-2a were treated with camphorsulfonic acid (2 equiv) for 2 hours at 100 °C. To our satisfaction, these samples retained their enantiomeric purity (97-98% ee)<sup>[18]</sup> while, under the same reaction conditions, a complete racemization was observed with the Tröger base (+)-3 (R = Me, Scheme 1). Also, for these reactions that proceed through bond cleavages at stereogenic nitrogen and carbon atoms, care was taken to determine the absolute configuration of the products 2 with certainty.<sup>[19]</sup> This configuration was established by vibrational circular dichroism (VCD) in view of the rigidity of the compounds 2.<sup>[20]</sup> IR absorption and VCD spectra were measured for solutions  $(CCl_4)$  of both (+)- and (-)-2a and compared to the averaged spectrum calculated for  $(5S_N, 11S_N)$ -2a (Figure 1). Overall,



**Figure 1.** Calculated spectrum of  $(5S_N, 11S_N)$ -**2a** (green). Experimental VCD spectra (CCl<sub>4</sub>, 298 K) of (+)-**2a** (red) and (-)-**2a** (blue). The calculated spectrum represents a weighted average of the VCD spectra of two conformers (see the Supporting Information for details).

a good agreement between the experimental and theoretical spectra was observed, thus allowing the assignment of (+)-**2***a* as  $(5S_N, 11S_N)$ . These results clearly indicate that the methylene extrusion occurs with retention of configuration.<sup>[21]</sup>

At that stage, the fate of the extruded  $CH_2$  group was determined. Careful <sup>1</sup>H NMR monitoring of the oxidation process was performed with a catalytic amount of DDQ (10 mol%) and an excess of  $MnO_2$  (10 equiv).<sup>[22]</sup> It revealed a significant amount of formaldehyde in the crude reaction mixture [up to 15 mol%, Eq. (1)].<sup>[23]</sup> With that information in hand, several mechanistic proposals can be considered, the most likely being that detailed in Scheme 2.

First, compounds of type **1** react in one-electron oxidations to generate nitrogen-centered radical cations of type **A**. These intermediates are formed from the more-electron-rich and sterically accessible bridgehead nitrogen atom. Then, an intramolecular electron-transfer reaction<sup>[24]</sup> occurs to generate the intermediate **B** which undergoes a homolytic C–C cleavage<sup>[25]</sup> favored by the alignment of the nitrogen orbitals



Scheme 2. Proposed reaction mechanism.

and the C-C bond.<sup>[26]</sup> This through-bond-promoted fragmentation creates the intermediate C, which is characterized by the presence of an iminium ion and a radical stabilized by the adjacent nitrogen lone pair (two-center three-electron "halfbond").<sup>[26]</sup> Then, rapid oxidation of the  $\alpha$ -aminomethyl radical forms the cationic intermediate **D** which is trapped by a water molecule  $(\rightarrow E)$ . Intramolecular nucleophilic attack of the tertiary nitrogen atom onto the iminium ion leads to the ring closure  $(\mathbf{E} \rightarrow \mathbf{F})$ . Hydrolysis of the carbinolammonium group leading to the release of the methylene group as formaldehyde and to the formation of the ringcontracted products 2. Finally, it is essential to stress the importance of the aryl and ester-substituted iminium moiety  $(\mathbf{C} \rightarrow \mathbf{E})$  which is conformationally locked and forbids a relaxation of the azocine ring. This restraint leads to the high level of enantiospecificity and retention of configuration.

In conclusion, a highly enantiospecific methylene extrusion reaction is reported and requires only mild oxidative conditions. Interestingly, the products of ring contraction are significantly more configurationally stable than regular Tröger bases. Applications of these derivatives in asymmetric synthesis and catalysis are looked for.

CCDC 1040107 (**2b**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data\_request/cif.

**Keywords:** configuration determination  $\cdot$  enantioselectivity  $\cdot$  heterocycles  $\cdot$  ring contraction  $\cdot$  stereochemistry

How to cite: Angew. Chem. Int. Ed. 2015, 54, 7520–7523 Angew. Chem. 2015, 127, 7630–7633

a) H. N. C. Wong, T. K. Ng, T. Y. Wong, *Heterocycles* **1983**, *20*, 1815–1840; b) L. Henriksen, *Insertion and extrusion reactions*, *Vol. 2*, Wiley, Hoboken, **1987**, pp. 393–420; c) F. S. Guziec, Jr., L. J. Sanfilippo, *Tetrahedron* **1988**, *44*, 6241–6285; d) C. R.



Williams, D. N. Harpp, *Sulfur Rep.* **1990**, *10*, 103–182; e) H. Kato, T. Kobayashi, *Yuki Gosei Kagaku Kyokaishi* **1990**, *48*, 672–680; f) M. Lambla, *NATO ASI Ser. Ser. E* **1995**, *302*, 437–454; g) F. S. Guziec, Jr., L. J. Guziec, *Selenium extrusion reactions*, Oxford University Press, **1999**, pp. 193–206; h) L. J. Guziec, F. S. Guziec, Jr., *Org. React.* **2012**, *78*, 411–549.

- [2] F. S. Guziec, L. James Guziec in PATAI'S Chemistry of Functional Groups, Wiley, Hoboken, 2009.
- [3] Cheletropic cycloreversions are sometime classified as extrusion reactions. See: a) T.-S. Chou, H.-H. Tso, *Org. Prep. Proced. Int.* **1989**, *21*, 257–296; b) M. A. Ogliaruso, M. G. Romanelli, E. I. Becker, *Chem. Rev.* **1965**, *65*, 261–367.
- [4] a) H. Liu, C. Dong, Z. Zhang, P. Wu, X. Jiang, Angew. Chem. Int. Ed. 2012, 51, 12570-12574; Angew. Chem. 2012, 124, 12738-12742; b) D. Liang, Y. He, L. Liu, Q. Zhu, Org. Lett. 2013, 15, 3476-3479; c) J. K. Laha, K. S. S. Tummalapalli, A. Gupta, Org. Lett. 2014, 16, 4392-4395.
- [5] a) J. Tröger, J. Prakt. Chem. 1887, 36, 225-245; b) V. Prelog, P. Wieland, Helv. Chim. Acta 1944, 27, 1127-1134; c) S. H. Wilen, J. Z. Qi, P. G. Williard, J. Org. Chem. 1991, 56, 485-487; d) B. Dolenský, J. Elguero, V. Kral, C. Pardo, M. Valik, Adv. Heterocycl. Chem. 2007, 93, 1-56; e) S. Sergeyev, Helv. Chim. Acta 2009, 92, 415-444; f) A. Révész, D. Schroder, T. A. Rokob, M. Havlik, B. Dolenský, Phys. Chem. Chem. Phys. 2012, 14, 6987-6995; g) X. Gao, C. S. Hampton, M. Harmata, Eur. J. Org. Chem. 2012, 7053-7056; h) J. Artacho, E. Ascic, T. Rantanen, J. Karlsson, C.-J. Wallentin, R. Wang, O. F. Wendt, M. Harmata, V. Snieckus, K. Wärnmark, Chem. Eur. J. 2012, 18, 1038-1042; i) Ö. V. Rúnarsson, J. Artacho, K. Wärnmark, Eur. J. Org. Chem. 2012, 7015-7041; j) P. Ondrisek, R. Schwenk, J. Cvengros, Chem. Commun. 2014, 50, 9168-9171.
- [6] a) A. Sharma, L. Guénée, J.-V. Naubron, J. Lacour, Angew. Chem. Int. Ed. 2011, 50, 3677–3680; Angew. Chem. 2011, 123, 3761–3764; b) A. Sharma, C. Besnard, L. Guenee, J. Lacour, Org. Biomol. Chem. 2012, 10, 966–969.
- [7] High enantiospecificity values (es 97–99%) are obtained with a large variety of electron-donating and electron-withdrawing substituents. Only in the case of p-NO<sub>2</sub> was the selectivity lower (es 64%).
- [8] a) Y. Hamada, S. Mukai, *Tetrahedron: Asymmetry* 1996, 7, 2671–2674; b) D. A. Lenev, D. G. Golovanov, K. A. Lyssenko, R. G. Kostyanovsky, *Tetrahedron: Asymmetry* 2006, 17, 2191–2194; c) C. Michon, A. Sharma, G. Bernardinelli, E. Francotte, J. Lacour, *Chem. Commun.* 2010, 46, 2206–2208.
- [9] O. Trapp, G. Trapp, J. W. Kong, U. Hahn, F. Vögtle, V. Schurig, *Chem. Eur. J.* 2002, *8*, 3629–3634.
- [10] S. A. Pujari, L. Guénée, J. Lacour, Org. Lett. 2013, 15, 3930– 3933.
- [11] There are only few examples of 13,13'-disubstituted derivatives of type 2: a) F. C. Cooper, M. W. Partridge, J. Chem. Soc. 1957, 2888-2893; b) A. Greenberg, N. Molinaro, M. Lang, J. Org. Chem. 1984, 49, 1127-1130; c) D. A. Lenev, I. I. Chervin, K. A. Lyssenko, R. G. Kostyanovsky, Tetrahedron Lett. 2007, 48, 3363-3366; d) M. Periasamy, S. Suresh, S. Satishkumar, Tetrahedron: Asymmetry 2012, 23, 108-116.
- [12] Lowering the reaction temperature to either -30 or 0 °C did not improve the enantiospecificity, but just slowed the reaction progress.

- [13] Tetraalkyl-substituted 1,2-diamines undergo C-C bond cleavage under oxidative photocatalysis: S. Y. Cai, X. Y. Zhao, X. B. Wang, Q. S. Liu, Z. G. Li, D. Z. G. Wang, Angew. Chem. Int. Ed. 2012, 51, 8050-8053; Angew. Chem. 2012, 124, 8174-8177. Under the same [Ru(bpy)<sub>3</sub>Cl<sub>2</sub>]-catalyzed reaction conditions, intramolecular cyclizations of benzylic 1,2 diamines are observed: J. Xuan, Y. Cheng, J. An, L. Q. Lu, X. X. Zhang, W. J. Xiao, Chem. Commun. 2011, 47, 8337-8339.
- [14] a) A. W. J. Logan, S. J. Sprague, R. W. Foster, L. B. Marx, V. Garzya, M. S. Hallside, A. L. Thompson, J. W. Burton, *Org. Lett.* 2014, *16*, 4078–4081; b) A. W. J. Logan, J. S. Parker, M. S. Hallside, J. W. Burton, *Org. Lett.* 2012, *14*, 2940–2943; c) D. G. Hulcoop, H. M. Sheldrake, J. W. Burton, *Org. Biomol. Chem.* 2004, *2*, 965–967; d) A. Toyao, S. Chikaoka, Y. Takeda, O. Tamura, O. Muraoka, G. Tanabe, H. Ishibashi, *Tetrahedron Lett.* 2001, *42*, 1729–1732.
- [15] Water content was determined using a Karl Fischer instrument.
- [16] It is worth mentioning that the presence of water in the reaction medium is beneficial for the stereospecific ring contraction of (+)-1f; performing the reaction in anhydrous nitromethane affords the product (+)-2f in 54% yield and in 51% ee only.
- [17] In this particular case, reaction is particularly sluggish as, after seven days with 4 equivalents of DDQ, traces of unreacted starting material remained.
- [18] The enantiomeric purity of (-)- and (+)-2a was only slightly modified from 99 and 98% to 97 and 98%, respectively.
- [19] A. Aamouche, F. J. Devlin, P. J. Stephens, J. Am. Chem. Soc. 2000, 122, 2346-2354.
- [20] a) T. B. Freedman, X. Cao, R. K. Dukor, L. A. Nafie, *Chirality* 2003, *15*, 743–758; b) G. Holzwarth, E. C. Hsu, H. S. Mosher, T. R. Faulkner, A. Moscowitz, *J. Am. Chem. Soc.* 1974, *96*, 251–252; c) L. A. Nafie, T. A. Keiderling, P. J. Stephens, *J. Am. Chem. Soc.* 1976, *98*, 2715–2723.
- [21] The absolute configuration of (+)-**1a**, that is  $(5S_{\rm N}, 11R_{\rm N}, 14R_{\rm C})$ , was established unambiguously in Ref. [6a].
- [22] MnO<sub>2</sub> does not promote the CH<sub>2</sub> extrusion. This stoichiometric oxidant is used to oxidize DDQH<sub>2</sub> to DDQ under mild conditions: L. Liu, P. E. Floreancig, Org. Lett. 2010, 12, 4686– 4689.
- [23] Reactions were performed in closed vials. Formaldehyde was identified by its  ${}^{1}$ H spectrum and also by spiking the crude samples with aqueous CH<sub>2</sub>O for comparison.
- [24] T. M. Wilson, T. A. Zeidan, M. Hariharan, F. D. Lewis, M. R. Wasielewski, Angew. Chem. Int. Ed. 2010, 49, 2385–2388; Angew. Chem. 2010, 122, 2435–2438.
- [25] a) M. A. Kellett, D. G. Whitten, J. Am. Chem. Soc. 1989, 111, 2314–2316; b) J. W. Leon, D. G. Whitten, J. Am. Chem. Soc. 1993, 115, 8038–8043; c) Y. Wang, K. S. Schanze, J. Phys. Chem. 1995, 99, 6876–6888.
- [26] S. Mondal, B. Gold, R. K. Mohamed, I. V. Alabugin, *Chem. Eur. J.* 2014, 20, 8664–8669.

Received: January 16, 2015 Revised: March 11, 2015 Published online: May 8, 2015