## **RSC Advances**

## PAPER



Cite this: RSC Adv., 2014, 4, 45603

Received 30th July 2014 Accepted 1st September 2014

DOI: 10.1039/c4ra07808j www.rsc.org/advances

## Introduction

2,3-Dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) is a known, powerful oxidizing agent used in organic synthesis; it is mainly used for dehydrogenation of hydroaromatic compounds.1 In addition, DDQ has been used to cause the dehydrogenative C-C coupling leading to interesting products,<sup>2</sup> photocatalytic oxygenation of benzene to phenol,3 and for preparing expanded porphyrins<sup>4</sup> and other  $\pi$ -conjugated molecules.<sup>5</sup> Dolphin and co-workers have reported DDQ oxidation of tripyrrane6 and dipyrromethane<sup>7</sup> to give unexpected, interesting products that have extended  $\pi$ -conjugation.<sup>8</sup> As part of our ongoing activity in developing pyrrole-based anion receptors and ligand systems, we were interested in oxidizing 2,5-bis(3,5-dimethylpyrazolylmethyl)pyrrole<sup>9</sup> 1 with DDQ to give conjugated molecules because 1 possesses two methylene groups. Herein, we report the reactive, fluorescent and  $\pi$ -conjugated quinoidal form of a simple pyrrole, which forms a donor-acceptor complex and reacts with oxygen only under sunlight, and the other

## Unsubstituted quinoidal pyrrole and its reaction with oxygen, charge transfer and palladium(II) complexes *via* DDQ oxidation<sup>†</sup>

Debasish Ghorai and Ganesan Mani\*

The dehydrogenation reaction of 2,5-bis(3,5-dimethylpyrazolylmethyl)pyrrole with 2,3-dichloro-5,6dicyano-1,4-benzoguinone (DDQ) followed by treatment with sodium bicarbonate afforded two new products containing oxygen atoms, 2,5-bis(3,5-dimethylpyrazolylcarbonyl)pyrrole (2) and 5-{bis-(3,5dimethylpyrazolyl)methylpyrrole-2-carbaldehyde (3). Interestingly, the products are different, when the reaction mixture is not treated with base; they are 2,2-tri(3,5-dimethylpyrazolylmethyl)-5-(3,5dimethylpyrazolylcarbonyl)pyrrole (4) and the adduct 5, the 2,3-dichloro-5,6-dicyanohydroquinone (DDQH<sub>2</sub>) adduct of 2,5-bis{di(3,5-dimethylpyrazolyl)methene}-2,5-dihydropyrrole (6) isolated in 22% yield, in which 6 acts as an electron acceptor and  $DDQH_2$  as an electron donor. The treatment of 5 with NaBH<sub>4</sub> or basic alumina afforded the free base 6 in good yield, representing a discrete unsubstituted quinoidal structure of pyrrole. 6 is fluorescent, reactive and could have a biradical resonance structure, which is demonstrated by its reaction with dioxygen only under the irradiation of sunlight, not reacting in darkness, giving the oxygen containing products: 2, bis-(3,5-dimethylpyrazolyl)methanone (7) and 2-(3,5dimethylpyrazolylcarbonyl)-5-(3,5-dimethylpyrazolyl)pyrrole (8). Moreover, the quinoidal molecule loses its fluorescence property on coordinating with palladium metal atom and gives a binuclear complex,  $[Pd_2Cl_4[\mu-C_4H_3N-2,5-(C(Me_2pz)_2)_2-N,N,N,N]]$ , in good yield. The structures of most of the compounds, including this palladium complex, were determined by single crystal X-ray diffraction studies.

> rearranged oxidation products along with a binuclear palladium complex; their structures were determined by X-ray diffraction.

## Results and discussion

As shown in Scheme 1, the oxidation reaction of 2,5-bis(3,5dimethylpyrazolylmethyl)pyrrole 1 with three equivalents of DDQ followed by treatment with a base (NaHCO<sub>3</sub>) afforded the two new products 2 and 3 in 11% and 20% yields, respectively, after silica gel column chromatography separation. Whereas, the same reaction without the treatment with a base afforded 2,3-dichloro-5,6-dicyanohydroquinone (DDQH<sub>2</sub>) adduct 5 in 22% yield, in addition to 2 formed in 11% yield and 4 in very low yield as yellow crystals after silica gel column chromatographic separation. Presumably, in the presence of sodium bicarbonate, the adduct 5 disintegrates into its components and the liberated quinoidal pyrrole molecule, 2,5-bis{di(3,5-dimethylpyrazolyl) methene}-2,5-dihydropyrrole 6, then undergoes further reactions with air, probably during the work-up procedure, to give 2–4 containing oxygen atoms. Therefore, it is likely that in both the reactions, the free quinoidal pyrrole molecule 6 (see below) might have initially formed and then reacted with the byproduct of the reaction, DDQH<sub>2</sub>, to give an adduct 5, where 6 can act as an electron acceptor and DDQH<sub>2</sub> as an electron



View Article Online

View Journal | View Issue

Department of Chemistry, Indian Institute of Technology, Kharagpur 721 302, India. E-mail: gmani@chem.iitkgp.ernet.in

<sup>†</sup> Electronic supplementary information (ESI) available. CCDC 998866–998870, 1011058. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4ra07808j



Scheme 1 The DDQ oxidation products of 2,5-bis(3,5-dimethylpyrazolylmethyl)pyrrole 1.

donor. However, this reaction did not give the expected molecule, that is, a molecule containing one pyrrole ring flanked by two pyrazolylmethylene groups with  $\pi$  conjugation, probably because of the reaction conditions resulting from the presence of three equivalents of DDQ and the considerably reactive nature of the free base **6** with molecular oxygen (see below).

Consistent with the reaction products obtained with sodium bicarbonate, the treatment of the  $DDQH_2$  adduct 5 with another base, NaBH<sub>4</sub>, afforded the free base 6 in 67% yield. In addition, 6 was also obtained after passing the adduct 5 through a basic alumina column (Scheme 2). All these oxidized products are soluble in common organic solvents such as diethyl ether, toluene and chloroform, and their structures were confirmed by X-ray diffraction studies.

The X-ray structures of **2–4** (Fig. 1, 2 and 3, respectively) revealed the presence of carbonyl group along with the



Scheme 2 The generation of the free quinoidal pyrrole molecule 6 from the adduct 5.

rearranged pyrazolate groups attached to the  $\alpha$ -positions of the pyrrole ring. The presence of carbonyl group indicates that these products could have formed by the reaction of the initially formed quinoidal pyrrole molecule **6** with air. A complete planar conformation was observed for **2**, whereas in the cases of **3** and **4**, the planarity is maintained in the carbonyl group side



Fig. 1 Molecular structure of 2 (30% thermal ellipsoids); dotted lines indicate hydrogen bonds. Selected bond lengths (Å) and angles (deg): N1–N2 1.376(5), N2–C6 1.407(5), C6–O1 1.220(5), C6–C7 1.458(6), N3–C7 1.361(5), C7–C8 1.388(6), C8–C9 1.383(7), C9–C10 1.376(6), N3–C10 1.356(5), C10–C11 1.470(6), C11–O2 1.212(5), N4–C11 1.410(5), N4–N5 1.370(5), O1–C6–N2 119.0(4), O1–C6–C7 120.2(4), O2–C11–N4 120.1(4), O2–C11–C10 120.4(4), N1–N2–C6 120.8(3), C7–N3–C10 109.0(4), N5–N4–C11 121.3(3), N2–C6–C7 120.7(4), N3–C10–C11 128.3(4), C10–C11–N4 119.5(4), N3…N1 2.738(5), H1…N1 2.03(5), N3–H1…N1 118(3), N3…N5 2.706(5), H1…N5 2.11(4), N3–H1… N5 110(3), C5…O1 2.824(6), H5a…O1 2.09, C5–H5a…O1 131.8, C12… O2 2.819(6), H12a…O2 2.09, C12–H12a…O2 131.6.



Fig. 2 Molecular structure of **3** (30% thermal ellipsoids); dotted lines indicate hydrogen bonds. Selected bond lengths (Å) and angles (deg): O1-C17 1.2207(19), C16-C17 1.445(2), C14-C16 1.381(2), C13-C14 1.398(2), C12-C13 1.380(2), N5-C16 1.378(2), N5-C12 1.358(2), C1-C12 1.500(2), N1-C1 1.442(2), N3-C1 1.466(2), O1-C17-C16 123.7(2), N5-C16-C17 121.8(2), C12-N5-C16 109.0(1), N5-C12-C1 125.4(2), N1-C1-C12 115.3(1), N3-C1-C12 111.4(1), N5…N2 2.921(2), H5…N2 2.49(2), N5-H5…N2 111(1).



Fig. 3 Molecular structure of 4; dotted lines indicate hydrogen bonds. Selected bond lengths (Å) and angles (deg): C6–O1 1.204(7), C6–N2 1.415(8), C6–C7 1.464(8), C7–C8 1.365(8), C7–N3 1.369(6), C8–C9 1.385(7), C9–C10 1.382(7), C10–N3 1.359(6), C10–C11 1.505(7), C11–N4 1.462(6), C11–N6 1.471(6), C11–N8 1.472(6), O1–C6–N2 119.7(6), O1–C6–C7 121.2(7), N2–C6–C7 119.1(6), N3–C7–C6 127.5(6), N3–C10–C11 123.3(4), N4–C11–N6 107.5(4), N4–C11–N8 107.1(4), N6–C11–C10 111.0(4), N3···N1 2.682(7), H2···N1 2.13(5), N3–H2···N1 118(4), N3···N5 2.739(6), H2···N5 2.18(5), N3–H2···N5 119(4).

of the molecule which consists of the pyrrole ring and the aldehyde (3) or 3,5-dimethylpyrazolylcarbonyl group (4). This planarity indicates the presence of extended  $\pi$ -conjugation among these groups. Moreover, in the structures of 2 and 4, the acidic pyrrolic NH group forms a three centre hydrogen bond<sup>10</sup> with the pyrazolyl nitrogen atoms. Probably because of these hydrogen bonds, the two carbonyl groups in 2 are oriented in one direction. This is in contrast to the thiophene based molecule, which exhibits its carbonyl groups oriented opposite to each other.<sup>11</sup> The <sup>1</sup>H NMR spectra of 2 and 4 in CDCl<sub>3</sub> showed resonances in a fairly deshielded region at  $\delta$  13.00 ppm and 12.04 ppm, respectively, for their pyrrolic NH protons, probably because of the three centre hydrogen bonding interactions in these molecules. Conversely, the structure of 3 showed one intramolecular hydrogen bond between the pyrrolic NH and the adjacent pyrazolyl nitrogen atom; the NH group resonates at 10.74 ppm in the <sup>1</sup>H NMR spectrum.

The adduct 5 is a fluorescent yellow solid that gives absorption maxima at 421 nm and 442 nm and the emission maximum at 490 nm upon excitation carried out at 420 nm in DMSO (see ESI, Fig. S25a†). Its fluorescence property originated from the  $\pi$ -electrons involved in conjugation. The <sup>13</sup>C NMR spectrum of 5 in CDCl<sub>3</sub> shows 17 peaks including the resonances due to DDQH<sub>2</sub>, which is expected for this complex. These chemical shift values shifted only slightly to both the deshielded and shielded regions with respect to those of the free base 6 (see below). Whereas, the <sup>1</sup>H NMR spectrum of 5 shows noticeable changes for the resonances due to the methyl and the pyrrolic NH protons, suggesting that the complex is not dissociated and that the solid state structure of 5 (see below) is preserved in CDCl<sub>3</sub> solution. While, the <sup>1</sup>H NMR of 5 in DMSO $d_6$  (see ESI, Fig. S10†) shows a spectral pattern very close to that

of the free base 6 in DMSO- $d_6$  (see ESI, Fig. S12<sup>†</sup>). This suggests that the charge transfer complex 5 dissociates into its component molecules in the Lewis base solvent DMSO. This is also confimed by the UV-vis spectrum of a dilute solution of 5 in DMSO, which appears to be very close to that of the free base 6. Conversely, the same charge transfer complex in toluene (see ESI, Fig. S26<sup>†</sup>) displays a discrete absorption band  $\lambda_{\text{max}}$  at 353 nm, in addition to the absorbances at 424 nm and 444 nm. The absorption at 353 nm (80.1 kcal  $mol^{-1}$ ) can be regarded as the charge transfer band for this complex, which is in the order of shorter wavelength charge transfer band values reported for pyridinium iodides.12 A dichloromethane solution of 5 also gives a similar UV-vis spectrum. The observation of the charge transfer band in toluene or dichloromethane indicates that the complex 5 is not dissociated in these solvents. In addition, the solid state UV-vis spectrum of 5 (see ESI, Fig. S28<sup>+</sup>) exhibits rather a broad band in the same region, where the absorptions are observed in the solution state spectra. Furthermore, to support the presence of the benzenoid system of DDQH<sub>2</sub> in 5, the FTIR spectrum contains bands at 3222  $\text{cm}^{-1}$  and 2251  $\text{cm}^{-1}$ due to the  $\nu(OH)$  and  $\nu(CN)$  stretching frequencies, respectively, which is confirmed by X-ray structure.

Complex 5 crystallizes in the triclinic space group  $P\overline{1}$ . The free base 6 and one molecule of DDQH<sub>2</sub> constitute the asymmetric unit representing the adduct formation as shown in Fig. 4. The DDQH<sub>2</sub> C–C (1.381(6) to 1.420(6) Å) and C–O (1.339(5) Å and 1.345(5) Å) bond lengths suggest the presence of partial double bond character and a benzenoid system, not quinoid. These distances are similar to the benzenoid systems in other reported charge transfer complexes.<sup>13</sup> Thus, the free base 6 acts as an electron acceptor, and DDQH<sub>2</sub> acts as an electron donor in complex 5.

In contrast to the free base structure 6, which has all pyrazolate rings twisted with respect to the pyrrole ring plane (see below), 5 has two of the pyrazole rings, one from each side almost coplanar to the pyrrole ring plane, which is parallel to that formed by DDQH<sub>2</sub>. This parallel arrangement and the inter planar (C···C) distances indicate that there can be some  $\pi - \pi$ interaction. Whereas, the nitrogen atoms of the other two perpendicularly oriented pyrazole rings with the dihedral angle N1–N2–C16–C15 =  $93.5(5)^{\circ}$  are hydrogen-bonded to DDQH<sub>2</sub>. However, all the pyrazolate nitrogen-methylene carbon bond distances are almost the same and range from 1.409(5) to 1.416(5) Å. Further, these distances are shorter than those single bond distances [1.473(6) Å and 1.466(5) Å] found in the structure of 2,5-bis(3,5-dimethylpyrazolylmethyl)pyrrole 1, indicating the involvement of pyrazolate groups in the extended  $\pi$ -conjugation. The bond distances, C14-C15 = 1.448(5) Å and C12-C13 = 1.448(6) Å, refer to single bonds that are longer than the alternatively located double bond distances (C11–C12 = 1.353(6) Å, C13-C14 = 1.339(6) Å and C15-C16 = 1.345(5) Å), supporting the quinoidal pyrrole structure. On comparing to the free base 6, a noticeable change is observed for the C-C double bond distance in the pyrrole ring (1.339(6) Å (5) and 1.363(10) Å (6)). These metric parameter changes are attributed to the interaction between the quinoidal pyrrole and DDQH<sub>2</sub>. The pyrrolic NH proton forms three-centred hydrogen bonds with the nitrogen atoms of the pyrazolate rings, located in front of it.



Fig. 4 Molecular structure of 5 (30% thermal ellipsoids). Most of the hydrogen atoms are omitted for clarity. Dotted lines indicate hydrogen bonds. Selected bond lengths (Å) and angles (deg): N6-N7 1.373(4), C11-N7 1.409(5), C11-N8 1.411(5), N8-N9 1.379(4), C11-C12 1.353(6), C12-N5 1.387(5), C12-C13 1.448(6), C13-C14 1.339(6), C14-C15 1.448(5), C15-N5 1.380(5), C15-C16 1.345(5), C16-N2 1.416(5), C16-N3 1.411(5), N1-N2 1.379(5), N3-N4 1.373(5), C31-O1 1.345(5), C28-O2 1.339(5), C31-C32 1.407(6), C30-C31 1.383(6), C27-C32 1.381(6), C27-C28 1.394(6), C28-C29 1.389(5), C29-C30 1.420(6), C32-Cl1 1.717(4), C27-Cl2 1.723(4), C29-C34 1.440(6), C34-N11 1.143(5), C30-C33 1.447(6). C33-N10 1.132(5), N7-C11-N8 116.4(4), C12-C11-N7 124.8(4), C12-C11-N8 118.4(4), C11-C12-N5 128.1(4), C15-N5-C12 110.9(3), C16-C15-N5 128.6(4), C15-C16-N3 126.4(4), C15-C16-N2 117.8(4), N5…N6 2.730(5), H5n…N6 2.21(4), N5-H5n…N6 120(4), N5… N4 2.801(5), H5n···N4 2.28(4), N5-H5n···N4 121(4), O1···N9 2.635(5), H10...N9 1.65(5), O1-H10...N9 168(4), O2...N1 2.688(5), H20...N1 1.77(5), O2-H2o···N1 173(4), C12···C31 = 3.5930(2), C13···C30 = 3.4644(2), C14···C29 = 3.4721(2), C15···C28 = 3.5923(2)

Compound **6** is a yellow coloured solid that is stable in air, but decomposes in solution under sunlight. Its emission (490 nm) and absorption (421 nm and 443 nm) spectra are very close to those of **5** in DMSO with a quantum yield of 0.023 (Fig. 5). The <sup>1</sup>H NMR spectrum of **6** in CDCl<sub>3</sub> features a pattern of signals resembling that of **5**. However, the characteristic signal is the resonance due to the pyrrolic NH proton at  $\delta$  9.42 ppm, which is in the slightly shielded region in comparison to that of the adduct **5** ( $\delta$  9.64 ppm), among other chemical shift changes. The <sup>13</sup>C NMR spectrum shows only thirteen peaks, which is in agreement with the X-ray structure.

Compound **6** crystallizes in the orthorhombic space group  $Pna2_1$  and two molecules of **6** constitute the asymmetric unit. These two independent molecules have almost identical metric parameters and an ORTEP view of one of the molecules is shown in Fig. 6. In contrast to the structure of **5**, the X-ray structure of the free base **6** adopts a different conformation that contains none of the pyrazole rings coplanar to the pyrrole ring plane, however the pyrazolate nitrogen atoms (N2, N3, N6 and N8) bonded to the methylene carbons are in the plane defined by the pyrrole ring along with the methylene carbon atoms (C16 and C11). This suggests that these pyrazolate nitrogen atoms are involved in the  $\pi$ -conjugation, similar to the case in **5**. The other structural features are close to those observed in the



Fig. 5 UV-vis and emission (490 nm) spectra of the quinoidal pyrrole molecule 6 in DMSO ( $10^{-5}$  M).

adduct structure 5, and the structure represents a novel discrete quinoidal form of pyrrole with the NH proton retained, in contrast to *N*-substituted<sup>14</sup> and fused quinoidal pyrrole ring<sup>15</sup> molecules.

Molecule **6** is analogous to thiophene-based quinoidal molecules such as 2,5-dimethylene-2,5-dihydrothiophene and its furan and *N*-methylpyrrole derivatives, which have  $\pi$ -conjugations.<sup>16</sup> Interestingly, quinoidal thiophene molecules,<sup>17</sup> Thiele's *p*-xylylene and Chichibabin's hydrocarbons<sup>18</sup> and several other conjugated molecules<sup>19</sup> have been reported to form charge transfer complexes, which have been shown to exhibit biradical structures and react with molecular oxygen. Having mentioned these, **6** could have the biradical canonical



Fig. 6 Molecular structure of 6 (30% thermal ellipsoids); most of the hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (deg): C11–C12 1.335(10), C11–N3 1.413(9), C11–N2 1.425(9), C12–N5 1.392(9), C12–C13 1.451(10), C13–C14 1.363(10), C14–C15 1.426(9), C15–C16 1.362(10), C15–N5 1.387(8), C16–N6 1.414(9), C16–N8 1.407(9), C12–C11–N3 122.6(7), C12–C11–N2 121.5(7), N3–C11–N2 115.9(7), C11–C12–N5 125.0(7), C11–C12–C13 129.6(7), C16–C15–N5 125.3(6), C16–C15–C14 128.5(7), C15–C16–N6 120.2(6), C15–C16–N8 122.3(7), N6–C16–N8 117.4(7), N5…N9 2.885(5), H2…N9 2.30(7), N5–H2…N9 121(5), N14…N15 2.873(8), H1…N15 2.33(7), N14–H1…N15 123(6).

structure, as depicted in Chart 1, contributing to the ground state structure. However, the alkenyl-pyrrole carbon distances (C11-C12 = 1.335(10) Å and C15-C16 = 1.362(10) Å) are not considerably elongated; elongation is expected if this biradical canonical structure contributes to the ground state. However, the biradical form of 6 could be transient and very reactive. Accordingly, a toluene solution of 6 readily reacts with molecular oxygen  $O_2$  under sunlight irradiation (Scheme 3) to give 7 and 8 along with the dicarbonyl compound 2, which were separated by silica gel column chromatography. Interestingly, no reaction was observed when the toluene solution was exposed to sunlight (visible light) for hours under nitrogen atmosphere, suggesting that the observed products 2-4, 7 and 8 are the results of the reaction of 6 with air. Importantly, the reaction between 6 and  $O_2$  did not proceed to completion in the absence of exposure to visible light, that is, in darkness. These observations led us to propose that probably the transient and reactive biradical species or the excited state structure of 6 could have formed in situ upon exposure to visible light, which then reacted with O<sub>2</sub> or air and cleaved the double bond of O<sub>2</sub> to obtain an oxygen containing products as shown in Schemes 1 and 3. All compounds, except 7, are new compounds. The structure of 8 is based on the spectroscopic data, including HRMS (ESI+), which showed the molecular ion  $[M + H^+]$  peak at m/z 284.1581 (calc. 284.1511). Its <sup>1</sup>H NMR spectrum showed a broad resonance at  $\delta$  8.85 ppm for the pyrrolic NH protons. Each of the  $\beta$ -CH protons of the pyrrole ring give one doublet of doublets and two different pyrazole CH protons give two separate resonances, which are consistent with the structure. The presence of a carbonyl group is confirmed by the IR spectrum, which displays the  $\nu$ (CO) stretching frequency at 1700 cm<sup>-1</sup>.

In addition, a preliminary study of the coordination behaviour of **6** was carried out with palladium metal. The reaction of the quinoidal pyrrole **6** with [Pd(PhCN)<sub>2</sub>Cl<sub>2</sub>] gave the bis-chelate complex **9** in 72% yield (Scheme 3). Interestingly, upon complexation the fluorescence of **6** is quenched, suggesting that the pyrazole  $\pi$ -electrons are involved in the origin of the fluorescence. The UV-vis absorption spectrum of **9** in acetonitrile showed a blue shifted  $\lambda_{max}$  at 371 nm ( $\varepsilon = 32400 \text{ M}^{-1} \text{ cm}^{-1}$ ) compared to that of **6** (Fig. S32†).

The X-ray structure of **9** is given in Fig. 7 along with the selected bond lengths and angles; its refinement data are given in Table 1. Complex **9** crystallizes in the monoclinic  $P2_1/n$  space group and the asymmetric unit contains one molecule of **9**, one molecule of acetonitrile, and one and a half molecules of water



Chart 1 The proposed biradical resonance structure for the quinoidal pyrrole molecule 6.



Scheme 3 The reaction of the quinoidal pyrrole molecule 6 with dioxygen under sunlight irradiation and the synthesis of its binuclear palladium(II) complex.

in the form of the solvents of crystallization, formulated as **9**  $CH_3CN \cdot 1.5H_2O$ . The X-ray structure revealed that the pyrazolyl nitrogen atoms on both the sides of the molecule are chelated to the palladium metal atom, thus forming two six-membered palladacycles adopting a boat conformation. This type of boat conformation has been reported for several other palladium complexes containing pyrazolyl groups.<sup>20</sup> The molecule can also be regarded to be a compartmental complex, which contains the two distorted square planar palladium( $\pi$ ) atoms located above the plane formed by the pyrrole ring and the methylene carbons and pyrazolyl nitrogen atoms are facing each other, which can be regarded to be the *cis* orientation. This is in contrast to the structure of the palladium( $\pi$ ) complex of 2,3,4,5-tetrakis(3,5-



Fig. 7 Molecular structure of 9 (30% probability ellipsoids); most of the hydrogen atoms, acetonitrile and water molecules are omitted for clarity. Selected bond lengths (Å) and angles (deg): N1-Pd1 2.020(11), N3-Pd1 2.031(13), Cl1-Pd1 2.284(5), Cl2-Pd1 2.281(4), N6-Pd2 2.039(11), N8-Pd2 2.025(12), Cl3-Pd2 2.273(4), Cl4-Pd2 2.285(4), C11-C12 1.360(18), C11-N2 1.410(17), C11-N4 1.418(18), C12-N5 1.382(17), C12-C13 1.455(18), C13-C14 1.317(19), C14-C15 1.446(18), C15-C16 1.297(18), C15-N5 1.404(16), C16-N9 1.425(16), C16-N7 1.431(16), N1-Pd1-N3 86.5(5), N1-Pd1-Cl2 177.8(4), N3-Pd1-Cl2 91.6(3), N1-Pd1-Cl1 91.9(4), N3-Pd1-Cl1 177.5(4), Cl2-Pd1-Cl1 89.99(18), N8-Pd2-N6 86.9(5), N8-Pd2-Cl3 177.0(4), N6-Pd2-Cl3 91.9(4), N8-Pd2-Cl4 91.3(3), N6-Pd2-Cl4 174.4(4), Cl3-Pd2-Cl4 89.72(16), C12-C11-N2 124.0(13), C12-C11-N4 119.9(13), N2-C11-N4 116.0(11), C11-C12-N5 126.0(13), C11-C12-C13 128.1(14), C16-C15-C14 128.5(13), N5-C15-C14 105.6(12), C15-C16-N9 122.2(13), C15-C16-N7 125.1(13), N9-C16-N7 112.8(12).

#### Table 1 Crystallographic data for compounds 2-6 and 9

	2	3	4	5	6	9
Empirical	$C_{16}H_{17}N_5O_2$	$C_{16}H_{19}N_5O$	C <sub>26</sub> H <sub>31</sub> N <sub>9</sub> O	$C_{34}H_{33}Cl_2N_{11}O_2$	$C_{26}H_{31}N_9$	$C_{28}H_{37}Cl_4N_{10}O_{1.5}Pd_2$
formula						
Formula	311.35	297.36	485.60	698.61	469.60	892.28
weight						
Wavelength (Å)	0.71073	1.54178	0.71073	0.71073	0.71073	0.71073
Temperature (K)	293(2)	110(2)	293(2)	293(2)	293(2)	293(2)
Crystal system	Monoclinic	Triclinic	Monoclinic	Triclinic	Orthorhombic	Monoclinic
Space group	Сс	ΡĪ	$P2_1/c$	ΡĪ	$Pna2_1$	$P2_1/n$
a/Å	19.167(6)	8.6780(9)	19.389(3)	8.1453(7)	23.991(4)	13.058(3)
b/Å	4.5557(15)	10.1309(12)	13.8717(17)	12.1246(10)	8.3841(13)	15.125(3)
c/Å	18.226(6)	10.3932(11)	9.8796(12)	18.4077(16)	25.825(4)	20.479(4)
α/degree	90.00	104.171(6)	90.00	106.647(3)	90.00	90.00
β/degree	100.720(10)	108.391(6)	104.655(4)	99.820(3)	90.00	106.625(7)
γ/degree	90.00	110.259(6)	90.00	94.826(3)	90.00	90.00
Volume ( $Å^3$ )	1563.7(9)	746.32(14)	2570.7(6)	1699.1(3)	5194.6(13)	3875.6(13)
Ζ	4	2	4	2	8	4
$D_{\rm calcd}$ , g cm <sup>-3</sup>	1.322	1.323	1.255	1.366	1.201	1.529
$\mu/\mathrm{mm}^{-1}$	0.091	0.702	0.082	0.241	0.076	1.241
F(000)	656	316	1032	728	2000	1788
$\theta$ range	2.16 to 25.38	4.89 to 59.08	1.09 to 25.34	1.18 to 25.00	1.58 to 25.00	1.66 to 25.00
(degree)						
Limiting	$-22 \le h \le 22,$	$-9 \le h \le 9$ ,	$-23 \le h \le 23,$	$-9 \le h \le 9$ ,	$-28 \le h \le 28$ ,	$-14 \le h \le 15$ ,
indices	$-5 \le k \le 5$ ,	$-11 \leq k \leq 10$ ,	$-16 \le k \le 16,$	$-14 \le k \le 14,$	$-9 \le k \le 9$ ,	$-17 \le k \le 17$ ,
	$-21 \leq l \leq 21$	$-11 \leq l \leq 11$	$-10 \leq l \leq 11$	$-21 \leq l \leq 21$	$-29 \le l \le 30$	$-24 \le l \le 24$
Total/unique	8729/2449	8494/2031	30 038/4528	20 611/5978	59 470/8778	44 801/6824
no. of reflns						
R <sub>int</sub>	0.0404	0.0985	0.1460	0.0825	0.2162	0.2343
Data/restr./	2449/2/212	2031/0/207	4528/0/328	5978/0/451	8778/1/638	6824/0/415
params.						
$\operatorname{GOF}\left(F^{2}\right)$	1.007	0.942	1.005	1.002	1.106	1.002
R1, wR2	0.0553, 0.1353	0.0362, 0.0886	0.0726, 0.1659	0.0625, 0.1393	0.0945, 0.1616	0.0924, 0.2214
R1, wR2 (all data)	0.0829, 0.1532	0.0428, 0.0909	0.1961, 0.2341	0.1418, 0.1806	0.2114, 0.2085	0.2083, 0.2836
Peak and hole (e Å <sup>-3</sup> )	0.197 and -0.170	0.159 and -0.236	0.296 and -0.283	0.346 and -0.329	0.200 and -0.281	1.685 and -0.603

dimethylpyrazol-1-ylmethyl)pyrrole that contains two palladium atoms above and below the pyrrole ring plane.<sup>21</sup> Compared to the free molecule **6**, upon coordination a significant decrease in the bond distance is observed in **9** for the pyrrole ring  $\beta$  C–C distance: C13–C14 = 1.317(19) Å against 1.363(10) Å in **6**. Further, the Pd–N(pz) and Pd–Cl bond distances fall within the range reported for the pyrazole nitrogen coordinated palladium complexes.<sup>22</sup> The molecule appears to be rigid, which is shown by the <sup>1</sup>H NMR spectrum displaying one separate singlet for each pyrazolyl CH proton and methyl group, probably because of the conjugation to be maintained. The pyrrolic NH group is hydrogen-bonded to one of the co-crystallized water molecules and gives a broad singlet at 8.52 ppm in the <sup>1</sup>H NMR spectrum.

## Conclusions

In conclusion, a new class of fluorescent quinoidal pyrrole with four high electron affinity nitrogen atoms attached to the methylene carbons causing it to act as an electron acceptor was developed through the dehydrogenation reaction of **1** with DDQ; this represents the first structurally characterised molecule of its type. The cleavage of  $O_2$  under sunlight demonstrates the considerable reactive nature of this pyrrole and gives insight into its possible biradical resonance structure. The isolated DDQ oxidation products are observed to correlate with the isolated products of the reaction of the quinoidal pyrrole with oxygen. The structures for most of the compounds, including a binuclear palladium(n) complex, are unambiguously established by X-ray diffraction methods. Further reactions for adduct and metal complex synthesis are in progress.

## **Experimental section**

#### Experimental section, general

All reactions were carried out under a nitrogen atmosphere using standard Schlenk-line techniques. Work-up procedures including NaHCO<sub>3</sub> treatment were carried out under aerobic conditions. Petroleum ether (bp 40–60 °C) and other solvents were distilled according to the standard procedures. Other chemicals were obtained from commercial sources and used without further purification. <sup>1</sup>H NMR (200

#### Paper

MHz and 400 MHz) and <sup>13</sup>C NMR (50.3 MHz and 100.6 MHz) spectra were recorded on a Bruker ACF200 spectrometer. Chemical shifts are referenced with respect to the chemical shift of the residual protons present in the deuterated solvents. FTIR spectra were recorded using Perkin Elmer Spectrum Rx. High Resolution Mass Spectra (ESI) were recorded using the Xevo G2 Tof mass spectrometer (Waters). Elemental analyses were carried out using a Perkin Elmer 2400 CHN analyzer. Melting points were determined in open capillaries and are corrected using benzophenone as a reference. UV-vis spectra were recorded using a Varian Cary 5000 UV-vis-NIR spectrophotometer.

## Synthesis of 2,5-bis(3,5-dimethylpyrazolylcarbonyl)pyrrole, 2, and 5-{bis(3,5-dimethylpyrazolyl)methyl}pyrrole-2-carbaldehyde, 3

To a dichloromethane (~50 mL) solution of 1 (1.0 g, 3.13 mmol) at 0 °C was added solid DDQ (2.13 g, 9.4 mmol). The colour of the solution immediately changed to dark green. After stirring at room temperature for 2 h, a saturated aqueous NaHCO<sub>3</sub> solution (2  $\times$  50 mL) was added to the solution and stirred under aerobic conditions. The organic layer was separated and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The solvent was removed from the organic layer and the residue was loaded onto a silica gel column. Elution using ethyl acetate-petroleum ether (1:3, v/v) afforded the first fraction from which the solvent was removed to give 2 as a colourless solid (0.11 g, 0.35 mmol, 11%). Further elution using the same solvent gave the second fraction from which the solvent was removed under vacuum to obtain compound 3 as a colourless solid (0.19 g, 0.64 mmol, 20%). Single crystals of 2 were obtained by the slow evaporation of a solution of 2 in methanol. 3 was crystallized by layering petroleum ether upon a solution of 3 in CH<sub>2</sub>Cl<sub>2</sub>.

For 2: mp 134 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz, 25 °C):  $\delta$  = 13.00 (br s, 1H, NH), 7.38 (d, 2H, pyrrole CH), 6.06 (s, 2H, pyrazole CH), 2.65 (s, 6H, CH<sub>3</sub>), 2.38 (s, 6H, CH<sub>3</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 51.3 MHz, 25 °C):  $\delta$  = 157.7, 152.9, 146.1, 129.2, 120.2, 111.0, 15.0, 14.3. FT-IR (KBr, cm<sup>-1</sup>):  $\nu$  = 3437 (br, s), 3310 (s), 3112 (w), 2976 (w), 2923 (w), 1676 (vs), 1585 (m), 1518 (m), 1486 (m), 1411 (m), 1365 (s), 1340 (s), 1312 (m), 1275 (m), 1195 (m), 1148 (w), 1106 (m), 1028 (m), 968 (w), 870 (m), 816 (m), 768 (m), 731 (w), 627 (w). HRMS (+ESI): calcd *m*/*z* for [M + H<sup>+</sup>] C<sub>16</sub>H<sub>18</sub>N<sub>5</sub>O<sub>2</sub>: 312.1455, found: 312.1458.

For 3: mp 146 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz, 25 °C):  $\delta$  = 10.74 (br s, 1H, NH), 9.54 (s, 1H, CHO), 7.60 (s, 1H, CH), 6.93 (s, 1H, pyrrole CH), 6.25 (s, 1H, pyrrole CH), 5.80 (s, 2H, pyrazole CH), 2.22 (s, 12H, CH<sub>3</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 51.3 MHz, 25 °C):  $\delta$  = 179.5, 149.2, 141.1, 134.2, 132.7, 121.1, 111.5, 107.5, 67.9, 13.8, 11.5. IR (KBr, cm<sup>-1</sup>):  $\nu$  = 3299 (s), 2923 (w), 2865 (w), 1666 (s), 1555 (m), 1468 (m), 1418 (s), 1380 (m), 1316 (m), 1268 (w), 1223 (m), 1168 (s), 1116 (w), 1035 (m), 976 (w), 875 (w), 786 (s), 757 (m), 719 (m), 650 (w), 516 (w). HRMS (+ESI): calcd *m*/*z* for [M + Na]<sup>+</sup> C<sub>16</sub>H<sub>19</sub>N<sub>5</sub>ONa: 320.1482, found: 320.1485.

# Synthesis of 2-tri(3,5-dimethylpyrazolylmethyl)-5-(3,5-dimethylpyrazolylcarbonyl)pyrrole, 4, and the DDQH<sub>2</sub> adduct of 2,5-bis{di(3,5-dimethylpyrazolyl)methene}-2,5-dihydropyrrole, 5

To a dichloromethane ( $\sim$ 50 mL) solution of 1 (1.0 g, 3.13 mmol) at 0 °C was added solid DDQ (2.13 g, 9.40 mmol). The colour of the solution immediately changed to dark green and stirring was continued for an additional 2 h at room temperature. The solution was concentrated and loaded onto a silica gel column. Elution using ethyl acetate-petroleum ether (1 : 3, v/v) afforded the first fraction, from which the solvent was removed to obtain 2 as a colourless solid (0.11 g, 0.35 mmol, 11%). Removal of the solvent from the second orange coloured fraction obtained compound 5 as a yellow solid (0.5 g, 0.72 mmol, 22%). Slow evaporation of the reaction mixture in acetonitrile gave a few single crystals of 4. Suitable single crystals of 5 were obtained by the slow evaporation of a solution of 5 in toluene at room temperature.

For 4: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz, 25 °C):  $\delta$  = 12.04 (br s, 1H, NH), 5.99 (s, 1H, pyrazole CH), 5.91 (s, 3H, pyrazole CH), 5.79 (dd, J = 4.0, 2.4 Hz, 2H, pyrrole CH), 2.62 (s, 3H, CH<sub>3</sub>), 2.27 (s, 3H, CH<sub>3</sub>), 2.17 (s, 9H, CH<sub>3</sub>), 1.69 (s, 9H, CH<sub>3</sub>). Further characterization could not be carried out because of the very low yield.

For 5: mp 140 °C (decomposed). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz, 25 °C):  $\delta$  = 9.64 (br s, 1H, NH), 6.00 (s, 2H, pyrrole CH), 5.88 (s, 2H, pyrazole CH), 5.79 (s, 2H, pyrazole CH), 2.44 (s, 6H, CH<sub>3</sub>), 2.34 (s, 6H, CH<sub>3</sub>), 1.91 (s, 6H, CH<sub>3</sub>), 1.75 (s, 6H, CH<sub>3</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 51.3 MHz, 25 °C):  $\delta$  = 151.8, 149.8, 149.7, 144.0, 141.0, 137.5, 133.4, 126.0, 113.6, 109.0, 107.6, 106.8, 106.2, 14.0, 13.5, 11.0, 10.6. IR (KBr, cm<sup>-1</sup>):  $\nu$  = 3222 (br, s), 2925 (w), 2526 (w), 2364 (w), 2251 (m), 1661 (w), 1564 (m), 1450 (s), 1414 (m), 1355 (s), 1274 (m), 1193 (s), 1078 (w), 1039 (w), 979 (w), 891 (m), 778 (w), 692 (w), 623 (w). HRMS (+ESI): calcd *m*/*z* for [M + Na]<sup>+</sup> C<sub>34</sub>H<sub>33</sub>Cl<sub>2</sub>N<sub>11</sub>NaO<sub>2</sub>: 720.2093, found: 720.2114. UV-vis (DMSO):  $\lambda_{max}$ , nm ( $\varepsilon$ , M<sup>-1</sup> cm<sup>-1</sup>) = 421 (35 000), 442 (32 200).

#### Synthesis of 2,5-bis{di(3,5-dimethylpyrazolyl)methene}-2,5dihydropyrrole 6 from 5

Method A. To a solution of 5 (1.0 g, 1.43 mmol) in dry MeOH ( $\sim$ 50 mL) at 0 °C was slowly added solid NaBH<sub>4</sub> (2.7 g, 71.50 mmol). The solution was stirred at room temperature for 16 h. After adding water ( $\sim$ 150 mL) to the solution, the solution was extracted for three times with dichloromethane (50 mL). The dichloromethane solution was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and filtered. The filtrate was concentrated and loaded onto a silica gel column. Elution using ethyl acetate-petroleum ether (1:2, v/v) gave the first fraction, from which the solvent was removed to obtain 6 as a yellow solid (0.45 g, 0.96 mmol, 67%). mp 147 °C. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz, 25 °C):  $\delta = 9.42$  (br s, 1H, NH), 6.22 (d, 2H, pyrrole CH), 5.90 (s, 2H, pyrazole CH), 5.82 (s, 2H, pyrazole CH), 2.28 (s, 6H, CH<sub>3</sub>), 2.25 (s, 6H, CH<sub>3</sub>), 2.00 (s, 6H, CH<sub>3</sub>), 1.69 (s, 6H, CH<sub>3</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 51.3 MHz, 25 °C): δ = 149.7, 149.4, 143.2, 140.9, 137.8, 126.7, 109.5, 107.2, 106.4,14.0, 13.8, 11.2, 10.5. IR (KBr,  $cm^{-1}$ ):  $\nu = 3357$  (m), 2956 (w), 2925 (w), 2861 (w), 1676 (w), 1647 (w), 1562 (m), 1454 (m), 1415

(s), 1358 (s), 1204 (w), 1117 (w), 1027 (w), 974 (w), 909 (w), 838 (w), 792 (m), 724 (w), 656 (w), 629 (w), 581 (w). HRMS (+ESI): calcd *m*/*z* for  $[M + H]^+$  C<sub>26</sub>H<sub>32</sub>N<sub>9</sub>: 470.2775, found: 470.2797. UV-vis (DMSO):  $\lambda_{max}$ , nm ( $\varepsilon$ , M<sup>-1</sup> cm<sup>-1</sup>) = 420 (23 700), 443 (21 900) and 259 (16 500).

**Method B.** A solution of 5 in ethyl acetate-petroleum ether (1/1, v/v) was passed through a basic alumina column. The first fraction was collected and the solvent was removed under vacuum to obtain an oily residue, which was washed with hexane to obtain 6 in the form of a yellow solid. The <sup>1</sup>H and <sup>13</sup>C NMR spectra of 6 obtained by this method were identical to the spectra obtained from the above mentioned procedure, Method A.

#### The sunlight irradiation of 6

A solution of **6** (0.10 g, 0.21 mmol) in toluene (20 mL) was connected to a bladder containing oxygen gas and stirred under sunlight irradiation for 6 h at room temperature. The solvent was removed under vacuum and the residue was loaded onto a silica gel column. Elution using ethyl acetate–petroleum ether (1:5 v/v) gave the first fraction. Elution using ethyl acetate–petroleum ether (1:1 v/v) gave the second and the third fractions. Removal of solvents from these fractions under vacuum gave compound **2** as a colourless solid (0.01 g, 0.032 mmol, 15%), compound **7** as a colourless solid (0.03 g, 0.11 mmol, 50%), respectively. The identity of **2** obtained by this method was confirmed after comparing the NMR and IR data with those of **2** obtained from the above-described reaction of **1** with DDQ.

For bis-(3,5-dimethylpyrazolyl)methanone.<sup>23</sup> 7 <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz, 25 °C):  $\delta = 5.85$  (s, 2H, pyrazole CH), 2.28 (s, 12H, CH<sub>3</sub>). IR (KBr, cm<sup>-1</sup>):  $\nu = 3440$  (s), 2925 (m), 2852 (m), 1740 (m), 1699 (s), 1650 (s), 1560 (m), 1541 (m), 1513 (w), 1460 (m), 1418 (w), 1393 (w), 1261 (w), 1189 (w), 1105 (s), 1029 (w), 800 (w), 697 (w), 629 (w). HRMS (+ESI): calcd *m*/*z* for [M + H]<sup>+</sup> C<sub>11</sub>H<sub>15</sub>N<sub>4</sub>O: 219.1246, found: 219.1292.

For 2-(3,5-dimethylpyrazolylcarbonyl)-5-(3,5-dimethylpyrazolyl)pyrrole 8. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz, 25 °C):  $\delta$  = 8.85 (br s, 1H, NH), 6.84 (dd, *J* = 5.6, 1.6 Hz, 1H, pyrrole CH), 6.13 (dd, *J* = 5.6, 1.6 Hz, 1H, pyrrole CH), 5.90 (s, 1H, pyrazole CH), 2.27 (s, 3H, CH<sub>3</sub>), 2.26 (s, 3H, CH<sub>3</sub>) 1.97 (s, 3H, CH<sub>3</sub>), 1.60 (s, 3H, CH<sub>3</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 51.3 MHz, 25 °C):  $\delta$  = 171.3, 152.4, 151.1, 143.6, 142.4, 136.3, 129.4, 124.8, 116.8, 109.3, 107.9, 13.9, 13.7, 11.0, 10.3. IR (KBr, cm<sup>-1</sup>):  $\nu$  = 3384 (m), 3250 (m), 2925 (s), 2856 (m), 1700 (vs), 1567 (s), 1450 (s), 1393 (s), 1352 (s), 1235 (w), 1192 (m), 1119 (m), 1073 (m), 1033 (m), 975 (m), 915 (m), 807 (s), 732 (s), 799 (m), 632 (w), 590 (w), 513 (w). HRMS (+ESI): calcd *m*/*z* for [M + H]<sup>+</sup> C<sub>15</sub>H<sub>18</sub>N<sub>5</sub>O: 284.1506, found: 284.1581.

#### Synthesis of [Pd<sub>2</sub>Cl<sub>4</sub>{µ-C<sub>4</sub>H<sub>3</sub>N-2,5-(C(Me<sub>2</sub>pz)<sub>2</sub>)<sub>2</sub>-N,N,N,N}], 9

To a solution of  $[Pd(PhCN)_2Cl_2]$  (0.045 g, 0.12 mmol) in toluene (10 mL) was added **6** (0.028 g, 0.059 mmol). The solution was stirred at room temperature for 16 h to obtain a brown precipitation of **9**. The solution was filtered and the precipitate was washed with diethyl ether (3 × 10 mL) and then dried under

vacuum. Yield: 72% (0.035 g, 0.042 mmol). A suitable single crystal of **9** was grown from an acetonitrile solution of **9** by slow evaporation. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz, 25 °C):  $\delta$  = 8.52 (br s, 1H, NH), 6.90 (s, 2H, pyrrole CH), 6.03 (s, 2H, pyrazole CH), 5.98 (s, 2H, pyrazole CH), 2.58 (s, 6H, CH<sub>3</sub>), 2.56 (s, 6H, CH<sub>3</sub>), 2.42 (s, 6H, CH<sub>3</sub>), 2.38 (s, 6H, CH<sub>3</sub>). IR (KBr, cm<sup>-1</sup>):  $\nu$  = 3454 (br), 3126 (w), 2961 (w), 2930 (w), 1656 (m), 1625 (m), 1561 (m), 1464 (m), 1416 (m), 1393 (m), 1261 (m), 1098 (s), 803 (m), 654 (w), 471 (w). HRMS (+ESI): calcd *m*/*z* for [M – 2Cl + Na]<sup>+</sup> C<sub>26</sub>H<sub>31</sub>Cl<sub>2</sub>N<sub>9</sub>NaPd<sub>2</sub>: 774.0041, found: 774.0657. UV-vis (acetonitrile):  $\lambda_{max}$  ( $\varepsilon$ ) = 371 nm (32 400 M<sup>-1</sup> cm<sup>-1</sup>).

#### X-ray crystallography

Suitable single crystals of 2-6 and 9 were obtained from solvents mentioned in their respective synthetic procedures. Single crystal X-ray diffraction data collections for these crystals were performed using a Bruker-APEX-II CCD diffractometer with graphite monochromated molybdenum K $\alpha$  radiation ( $\lambda$  = 0.71073 Å). The structures were solved by SIR-92 (ref. 24) available in the WinGX program which successfully located most of the non-hydrogen atoms. Subsequently, least-squares refinements were carried out on F2 using SHELXL-97 (ref. 25) (WinGX version) to locate the remaining non-hydrogen atoms. Typically for all the structures, hydrogen atoms attached to carbon atoms were fixed in calculated positions. The pyrrolic NH or OH hydrogen atoms were located from the difference Fourier map and freely refined isotropically with their thermal parameters set as equivalent to 1.2 times that of their parent atoms. In case of the structure of 9, water hydrogens did not appear and, thus are not located; the pyrrolic NH hydrogens are fixed. Crystallographic refinement data are given in Table 1.

## Acknowledgements

We thank the CSIR and DST for support and for the X-ray and NMR facilities. We also thank Dr Joseph H. Reibenspies, Department of Chemistry, Texas A&M University for one data collection and structure refinement.

## Notes and references

- (*a*) D. Walker and J. D. Hiebert, *Chem. Rev.*, 1967, **67**, 153; (*b*)
  L. Yu, K. Muthukumaran, I. V. Sazanovich, C. Kirmaier,
  E. Hindin, J. R. Diers, P. D. Boyle, D. F. Bocian, D. Holten and J. S. Lindsey, *Inorg. Chem.*, 2003, **42**, 6629; (*c*)
  S. B. Bharate, *Synlett*, 2006, 496.
- 2 (a) H. Liu, L. Cao, J. Sun, J. S. Fossey and W.-P. Deng, *Chem. Commun.*, 2012, 48, 2674; (b) S. Bhunia, S. Ghosh, D. Dey and A. Bisai, *Org. Lett.*, 2013, 15, 2426; (c) Z.-L. Wang, H.-L. Li, L.-S. Ge, X.-L. An, Z.-G. Zhang, X. Luo, J. S. Fossey and W.-P. Deng, *J. Org. Chem.*, 2014, 79, 1156.
- 3 K. Ohkubo, A. Fujimoto and S. Fukuzumi, *J. Am. Chem. Soc.*, 2013, **135**, 5368.
- 4 (a) J.-Y. Shin, H. Furuta, K. Yoza, S. Igarashi and A. Osuka, J. Am. Chem. Soc., 2001, 123, 7190; (b) S. Ramakrishnan,

K. S. Anju, A. P. Thomas, E. Suresh and A. Srinivasan, *Chem. Commun.*, 2010, **46**, 4746.

- 5 (a) J. L. Sessler, R. S. Zimmerman, C. Bucher, V. Král and B. Andrioletti, *Pure Appl. Chem.*, 2001, 73, 1041; (b) T. E. Wood and A. Thompson, *Chem. Rev.*, 2007, 107, 1831; (c) T. Guchhait, V. K. Jha and G. Mani, *Org. Biomol. Chem.*, 2013, 11, 2818.
- 6 J.-Y. Shin, S. S. Hepperle, B. O. Patrick and D. Dolphin, *Chem. Commun.*, 2009, 2323.
- 7 J.-Y. Shin, B. O. Patrick and D. Dolphin, *Org. Biomol. Chem.*, 2009, 7, 2032.
- 8 J.-Y. Shinab and D. Dolphin, New J. Chem., 2011, 35, 2483.
- 9 D. Ghorai, S. Kumar and G. Mani, *Dalton Trans.*, 2012, 41, 9503.
- 10 I. Rozas, I. Alkorta and J. Elguero, *J. Phys. Chem. A*, 1998, **102**, 9925.
- 11 I. A. Guzei, L. C. Spencer, M. G. Tshivashe and J. Darkwa, *Acta Crystallogr.*, 2009, **E65**, 02743.
- 12 E. J. Poziomek, R. A. Mackay and T. J. Novak, *Mikrochim. Acta*, 1974, **62**, 281.
- 13 (a) E. Gebert Jr, A. H. Reis, J. S. Miller, H. Rommelmann and A. J. Epstein, *J. Am. Chem. Soc.*, 1982, **104**, 4403; (b) A. O. Patil, D. Y. Curtin and I. C. Paul, *J. Am. Chem. Soc.*, 1984, **106**, 4010; (c) M. Bouvet, B. Malezieux and P. Herson, *Chem. Commun.*, 2006, 1751; (d) C. Yi, S.-X. Liu, A. Neels, P. Renaud and S. Decurtins, *Org. Lett.*, 2009, **11**, 5530; (e) X. Guegano, J. Hauser, S.-X. Liu and S. Decurtins, *Acta Crystallogr.*, 2011, **E67**, 02967.
- 14 Y. Qiao, J. Zhang, W. Xu and D. Zhu, J. Mater. Chem., 2012, 22, 5706.
- 15 (a) T. M. Pappenfus, B. J. Hermanson, T. J. Helland, G. G. W. Lee, S. M. Drew, K. R. Mann, K. A. McGee and S. C. Rasmussen, *Org. Lett.*, 2008, **10**, 1553; (b) Y. A. Getmanenko, T. A. Purcell, D. K. Hwang, B. Kippelen and S. R. Marder, *J. Org. Chem.*, 2012, 77, 10931.
- 16 (a) S. Gronowitz and B. Uppstrom, Acta Chem. Scand., Ser. B, 1974, 28, 981; (b) A. Ishii, Y. Horikawa, I. Takaki, J. Shibata, J. Nakayama and M. Hoshino, Tetrahedron Lett., 1991, 32, 4313.
- 17 (a) K. Takahashi, T. Suzuki, K. Akiyama, Y. Ikegami and
  Y. Fukazawat, J. Am. Chem. Soc., 1991, 113, 4576; (b)

- T. Takahashi, K.-I. Matsuoka, K. Takimiya, T. Otsubo and Y. Aso, J. Am. Chem. Soc., 2005, **127**, 8928; (c) S. D. Motta, F. Negri, D. Fazzi, C. Castiglioni and E. V. Canesi, J. Phys. Chem. Lett., 2010, **1**, 3334; (d) E. V. Canesi, D. Fazzi, L. Colella, C. Bertarelli and C. Castiglioni, J. Am. Chem. Soc., 2012, **134**, 19070; (e) J. Casado, R. P. Ortiz and J. T. L. Navarrete, Chem. Soc. Rev., 2012, **41**, 5672; (f) T. M. Pappenfus, K. B. Schliep, A. Dissanayake, T. Ludden, B. Nieto-Ortega, J. T. L. Navarrete, M. C. R. Delgado and J. Casado, J. Chem. Educ., 2012, **89**, 1461.
- 18 L. K. Montgomery, J. C. Huffman, E. A. Jurczak and M. P. Grendze, J. Am. Chem. Soc., 1986, 108, 6004.
- 19 (a) A. Kikuchi, F. Iwahori and J. Abe, *J. Am. Chem. Soc.*, 2004, 126, 6526; (b) M. Abe, *Chem. Rev.*, 2013, 113, 7011.
- 20 (a) M. C. Carrion, G. Dura, F. A. Jalo, B. R. Manzano and A. M. Rodríguez, *Cryst. Growth Des.*, 2012, 12, 1952; (b) A. Sánchez-Méndez, E. de Jesús, J. C. Flores and P. Gómez-Sal, *Eur. J. Inorg. Chem.*, 2010, 1881; (c) S. Tsuji, D. C. Swenson and R. F. Jordan, *Organometallics*, 1999, 18, 4758.
- 21 D. Ghorai and G. Mani, Inorg. Chem., 2014, 53, 4117.
- 22 (a) G. Minghetti, M. A. Cinellu, A. L. Bandini, G. Banditelli, F. Demartin and M. Manassero, J. Organomet. Chem., 1986, 315, 387; (b) N. Arroyo, F. Gomez-de la Torre, F. A. Jalon, B. R. Manzano, B. Moreno-Lara and A. M. Rodríguez, J. Organomet. Chem., 2000, 603, 174; (c) J. Garcia-Anton, J. Pons, X. Solans, M. Font-Bardia and J. Ros, Eur. J. Inorg. Chem., 2002, 3319; (d) M. P. Conley, C. T. Bruns and R. F. Jordan, Organometallics, 2007, 26, 6750; (e) A. Sánchez-Méndez, E. de Jesús, J. C. Flores and P. Gómez-Sal, Inorg. Chem., 2007, 46, 4793; (f) A. John, M. M. Shaikh, R. J. Butcher and P. Ghosh, Dalton Trans., 2010, 39, 7353; (g) A. Sánchez-Méndez, E. de Jesús, J. C. Flores and P. Gómez-Sal, Eur. J. Inorg. Chem., 2010, 141; (h) A. J. Hallett, C. J. Adams, K. M. Anderson, R. A. Baber, N. G. Connelly and C. J. Prime, Dalton Trans., 2010, 39, 5899.
- 23 T. C. Higgs and C. J. Carrano, Inorg. Chem., 1997, 6, 291.
- 24 A. Altomare, G. Cascarano, C. Giacovazzo and A. Guagliardi, J. Appl. Crystallogr., 1993, 26, 343.
- 25 G. M. Sheldrick, A short history of SHELX, *Acta Crystallogr.*, 2008, **A64**, 112.