

Accepted Manuscript

Triacetic Acid Lactone as a Common Intermediate for the Synthesis of 4-Hydroxy-2-pyridones and 4-Amino-2-pyridones

George A. Kraus, Umayangani K. Wanninayake, Jashaun Bottoms

PII: S0040-4039(16)30151-4
DOI: <http://dx.doi.org/10.1016/j.tetlet.2016.02.043>
Reference: TETL 47316

To appear in: *Tetrahedron Letters*

Received Date: 22 December 2015
Revised Date: 8 February 2016
Accepted Date: 11 February 2016

Please cite this article as: Kraus, G.A., Wanninayake, U.K., Bottoms, J., Triacetic Acid Lactone as a Common Intermediate for the Synthesis of 4-Hydroxy-2-pyridones and 4-Amino-2-pyridones, *Tetrahedron Letters* (2016), doi: <http://dx.doi.org/10.1016/j.tetlet.2016.02.043>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.





Tetrahedron Letters
journal homepage: www.elsevier.com

Triacetic Acid Lactone as a Common Intermediate for the Synthesis of 4-Hydroxy-2-pyridones and 4-Amino-2-pyrones

George A. Kraus*, Umayangani K. Wanninayake and Jashaun Bottoms

Department of Chemistry, Iowa State University, Ames, IA 50011, United States
Iowa State University Center for Biorenewable Chemicals

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

Keywords:

2-pyrone

Triacetic acid lactone

Amine

4-amino-2-pyrones

4-hydroxy-2-pyridones

ABSTRACT

At ambient temperature, triacetic acid lactone reacts with amines to produce 4-amino-2-pyrones. If the temperature is raised to 100 °C, 4-hydroxy-2-pyridones are generated.

© 2015 Elsevier Ltd. All rights reserved.

Triacetic acid lactone (**1**) is readily available either through the acid catalyzed deacetylation of dehydroacetic acid or through microbial transformation of glucose.¹ It is a useful intermediate for the synthesis of penstyrylpyrone (**3**)² and pogostone (**4**)³, shown in Figure 1. A related synthetic compound PH797804 (**5**)⁴ is a potent p38 MAPK inhibitor. As part of a program to expand the potential of **1** as a platform chemical⁵, we studied the reaction of **1** and its tosylate **2** with amines. A number of groups have reported limited studies of **1** with primary amines and with glycine.⁶⁻¹⁰





Figure 1. Triacetic acid lactone derivatives

Reaction of 1.1 equivalents of a primary amine with **1** at 100 °C in water afforded 2-pyridones **6**, as shown in Scheme 1. The structure assignment of **6a** was supported by a shift in the NMR resonance of the methyl group at C-6 and by a strong NOE interaction between the methyl group at C-6 and the methylene of the ethyl group.



Scheme 1. Reaction with amines

The products of primary amines with **1** are shown in Figure 2. Both aliphatic and aromatic amines react with **1**. The pyridones **6a-6g** were polar solids whose insolubility made them difficult to purify by silica gel chromatography.¹⁹ Fortunately, the pyridones were readily separable from **1** by differential solubility in ethyl acetate.

*Corresponding author. Tel.: +1 515 294 7794; fax: +1 515 294 0105.
E-mail address: gakraus@iastate.edu (G.A. Kraus).




Figure 2. Pyridone derivatives

In contrast to the reactivity of **1**, tosylate **2**¹⁸ reacted with amines at C-4 (Method A). Adducts **7a-7d** were not as polar as the pyridones.²⁰ A slight shift of the chemical shifts of the hydrogens at C-3 and C-5 plus the strong NOE interaction between the hydrogens at C-3 and C-5 with the methylene in **7a** supported the structure assignments.






Figure 3. 4-amino-2-pyrone derivatives

We subsequently found that **1** reacted with amines at ambient temperature in water to form **7** (Method B). We attribute the remarkable chemoselectivity of this addition to the rapid reaction of the amine with the keto tautomer at C-4. When **7a** was boiled in water, **6a**, likely the thermodynamic product, was isolated in 35% yield.



Scheme 2. Reaction with amines at ambient temperature and 100 °C

We also examined the reactions with diamines. As shown in Scheme 3, the reaction of tosylate **2** with piperazine afforded the 2:1 adduct **8** in 21% yield.²¹ Later, we found that simply mixing two equivalents of **1** with piperazine at ambient temperature in water afforded a cleaner product.



Scheme 3. Reaction with diamines




Figure 4. Products with diamine and tetramine

Triacetic acid lactone constitutes a useful platform for the direct introduction of nitrogen functionality. The reactions proceed in good yields and are operationally convenient. The extension to the reactions of **1** or **2** with diamines and tetramines leads rapidly to new materials.²²

Acknowledgments

We thank the NSF Engineering Center for Biorenewable Chemicals (CBiRC) which was awarded NSF grant EEC-0813570 for support of this research. Jashaun Bottoms, an undergraduate at Tuskegee University, was a 2015 CBiRC REU participant.

References

- Moreno-Manas, M.; Pleixats, R. *Adv. Heterocyclic Chem.* **1992**, *53*, 1-84.
- Kraus, G. A.; Wanninayake, U. K. *Tetrahedron Lett.* **2015**, *56*, 7112-7114.
- Wanninayake, U. K.; Kraus, G. A. Abstracts of Papers, 250th ACS National Meeting, Boston, MA, **2015**, IEC-88.
- Selness, S. R.; Devraj, R. V.; Devadas, B.; Walker, J. K.; Boehm, T. L.; Durley, R. C.; Shieh, H.; Xing, Li; Rucker, P. V.; Jerome, K. D.; *Bioorg. Med. Chem. Lett.* **2011**, *21*(13), 4066-4071.
- Kraus, G. A.; Basemann, K.; Guney, T. *Tetrahedron Lett.* **2015**, *56*(23), 3494-96.
- Castillo, Simone; Ouadahi, Hamid; Herault, Valentin. *Bull. Soc. Chim. France.* **1982**, 7-8, Pt. 2, 257-61.
- Kong, X.; Chen, G.; Wang, J.; Zheng, L.; Huang, Y. *Faming Zhuanli Shenqing* **2013**, CN 103145666 A 20130612.
- Kherfi, H.-N.; Hamdi, M.; Speziale, V. *J. Heterocyclic Chem.* **1990**, *27*, 1401.
- Stoyanov, E. V.; Ivanov, I. C. *Molecules*, **2004**, *9*, 627.
- Stoncius, S.; Orentas, E.; Butkus, E.; Ohrstrom, L.; Wendt, O. F.; Wammack, K. *J. Am. Chem. Soc.* **2006**, *128*, 8272.
- Hansen, A.L.; Skrydstrup, T. *Org. Lett.*, **2005**, *7*, 5585-5587.
- Haydon, David John.; Czaplowski, Lloyd George. **2009**, WO 2009074812 A1 20090618.
- Morone, M.; Sato, A.; Azuma, Y. *Heterocycles*. **2003**, *60*, 2241 - 2249.
- Patel, B. H.; Mason, A. M.; Barrett, A. G. M. *Org. Lett.* **2011**, *13*(19), 5156-5159.
- Rana, S.; Hong, H.-S.; Barrigan, L.; Jin, L.-W.; Hua, D. H. *Bioorganic & Medicinal Chemistry Letters*. **2009**, *19*, 670-674.

16. Miura, T.; Tagashira, J.; Sekimoto, R.; Ishida, R.; Aoki, H.; Ohgiya, T. **2012**, US 2012/0225896 A1.
17. Defant, A.; Mancini, I. *Tetrahedron*, **2013**, *69*, 4586-4590.
18. Experimental procedure for the synthesis of 6-methyl-2-oxo-2H-pyran-4-yl 4-methylbenzenesulfonate (**2**) (96%) colorless solid: 4-Hydroxy-6-methyl-2H-pyran-2-one (triacetic acid lactone, **1**) (0.63 g, 5 mmol) and tosyl chloride (0.955 g, 5 mmol) was dissolved in CH₂Cl₂ (37.5 mL). Triethylamine (2.05 mL, 15 mmol) was added and the reaction mixture was left stirring at rt. overnight (23 h). CH₂Cl₂ (50 mL) was added and the organic phase was washed with water (25 mL) and brine (25 mL). The organic phase was dried over MgSO₄. After concentration in vacuo the crude product was purified by column chromatography using hexane/ethyl acetate (4:1) as eluent to afford the desired product.¹¹ Rf = 0.18 (silica gel, hexanes/EtOAc 3:1); ¹H NMR (300 MHz, Chloroform-*d*) δ = 7.82 (d, 2H), 7.37 (d, 2H), 6.00 (s, 1H), 5.80 (s, 1H), 2.45 (s, 3H), 2.23 (s, 3H) ppm.
19. Representative procedure for the preparation of pyridones **6a-6g**: A mixture of triacetic acid lactone (**1**) (0.5 g, 3.96 mmol, 1 eq) and primary amine (4.35 mmol, 1.1 eq) in water (2.5 mL) was heated to 100 °C overnight. After completion of the reaction (TLC monitoring), the reaction mixture was cooled down and the precipitate was filtered and washed with ethyl acetate and dried under vacuum to obtain the desired product.¹²
- 1-ethyl-4-hydroxy-6-methylpyridin-2(1H)-one (6a)* (79%) Light brown solid: mp 240-242 °C (Lit. 247 °C)⁶; Rf = 0.16 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 5.92 (d, 1H), 5.72 (d, 1H), 4.05 (q, 2H), 2.39 (s, 3H), 1.24 (t, 3H) ppm; ¹³C NMR (126 MHz, Methanol-*d*₄) δ = 165.74, 164.21, 145.70, 100.64 (2C), 37.09, 17.14, 11.18 ppm; LRMS (ESI-QTOF) calcd for C₈H₁₂NO₂ [M + H]⁺ 154.0868, found 154.0872.
- 4-hydroxy-1-isobutyl-6-methylpyridin-2(1H)-one (6b)* (56%) Light brown solid: mp 173-175 °C; Rf = 0.35 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 5.92 (d, 1H), 5.73 (d, 1H), 3.85 (d, 2H), 2.37 (s, 3H), 2.26-2.05 (m, 1H), 1.04-0.80 (m, 6H) ppm; ¹³C NMR (126 MHz, Methanol-*d*₄) δ = 165.41, 164.67, 146.18, 100.54, 94.73, 48.51, 26.18, 17.90, 17.20 (2C) ppm; LRMS (ESI-QTOF) calcd for C₁₀H₁₆NO₂ [M + H]⁺ 182.1181, found 182.1191.
- 1-benzyl-4-hydroxy-6-methylpyridin-2(1H)-one (6c)* (20%) White solid: mp 219-220 °C (Lit. 217 °C)⁶; Rf = 0.30 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 7.33 (m, 2H), 7.27 (m, 1H), 7.11 (m, 2H), 5.97 (d, 1H), 5.82 (d, 1H), 5.32 (s, 2H), 2.25 (s, 3H) ppm; LRMS (ESI-QTOF) calcd for C₁₃H₁₄NO₂ [M + H]⁺ 216.1025, found 216.1040. ¹³C NMR data agreed with the literature.¹⁴
- 1-(benzo[1,3]dioxol-5-ylmethyl)-4-hydroxy-6-methylpyridin-2(1H)-one (6d)* (80%): mp 206-209 °C; Rf = 0.30 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Chloroform-*d*) δ = 6.76 (s, 1H), 6.72 (d, 1H), 6.69 (d, 1H), 5.94 (s, 2H), 5.92 (d, 1H), 5.68 (d, 1H), 5.13 (s, 2H), 2.15 (s, 3H) ppm; ¹³C NMR (126 MHz, DMSO-*d*₆) δ = 163.85, 155.28, 147.80, 146.57, 145.65, 133.25, 120.72, 108.65, 108.58, 101.35, 101.30, 99.40, 45.55, 20.37 ppm; LRMS (ESI-QTOF) calcd for C₁₄H₁₄NO₄ [M + H]⁺ 260.0923, found 260.0932.
- 4-hydroxy-1-(4-methoxybenzyl)-6-methylpyridin-2(1H)-one (6e)* (48%) White solid: mp 221-225 °C; Rf = 0.24 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 7.27 (d, 2H), 7.07 (d, 2H), 5.86 (d, 1H), 5.50 (d, 1H), 5.19 (s, 2H), 3.77 (s, 3H), 2.20 (s, 3H) ppm; LRMS (ESI-QTOF) calcd for C₁₄H₁₆NO₃ [M + H]⁺ 246.1130, found 246.1142. ¹³C NMR data agreed with the literature.¹⁵
- 4-hydroxy-6-methyl-1-(3,4,5-trimethoxybenzyl)pyridin-2(1H)-one (6f)* (69%) Light yellow solid: mp 251-255 °C; Rf = 0.11 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 6.43 (s, 2H), 5.97 (d, 1H), 5.82 (d, 1H), 5.26 (s, 2H), 3.80 (s, 6H), 3.76 (s, 3H), 2.29 (s, 3H) ppm; ¹³C NMR (126 MHz, Methanol-*d*₄) δ = 163.37, 159.92, 151.91, 132.02 (2C), 131.43, 115.89, 101.54 (2C), 100.73, 94.46, 58.12, 53.55 (2C), 45.61, 17.52 ppm; LRMS (ESI-QTOF) calcd for C₁₆H₂₀NO₅ [M + H]⁺ 306.1341, found 306.1347.
- 4-hydroxy-6-methyl-1-phenethylpyridin-2(1H)-one (6g)* (29%) White solid: mp 244-247 °C (Lit. 252 °C)⁶; Rf = 0.34 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 7.35-7.14 (m, 5H), 5.85 (d, 1H), 5.77 (d, 1H), 4.17 (t, 2H), 2.97 (t, 2H), 2.12 (s, 3H) ppm; LRMS (ESI-QTOF) calcd for C₁₄H₁₆NO₂ [M + H]⁺ 230.1181, found 230.1173. NMR data agreed with the literature.¹⁶
20. Representative procedure for the preparation of 4-aminopyrones **7a-7f**: **Method A**: A mixture of 6-methyl-2-oxo-2H-pyran-4-yl 4-methylbenzenesulfonate (**2**) (0.08 g, 0.3 mmol, 1 eq) and amine (0.66 mmol, 2.2 eq) in ethanol (4 mL) was stirred at rt. overnight. After completion of the reaction (TLC monitoring), solvent was evaporated. The crude compound was purified by preparative thin layer chromatography (EtOAc/dichloromethane) to afford the desired product. **Method B**: A mixture of triacetic acid lactone (**1**) (0.17 g, 1.4 mmol, 1 eq) and amine (1.54 mmol, 1.1 eq) in water (1.5-2 mL) was stirred at rt. overnight. After completion of the reaction (TLC monitoring), solvent was evaporated. The crude compound was purified by preparative thin layer chromatography (EtOAc/dichloromethane) to afford the desired product.
- 4-(ethylamino)-6-methyl-2H-pyran-2-one (7a)* (Method A: 51%, Method B: 83%) Colorless liquid: Rf = 0.11 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 5.92 (s, 1H), 5.72 (s, 1H), 3.32 (m, 2H), 2.39 (s, 3H), 1.24 (t, 3H) ppm; ¹³C NMR (126 MHz, Methanol-*d*₄) δ = 163.68, 157.62, 152.78, 97.77, 89.99, 35.12, 27.46, 16.60 ppm; LRMS (ESI-QTOF) calcd for C₈H₁₂NO₂ [M + H]⁺ 154.0868, found 154.0871.
- 4-(butylamino)-6-methyl-2H-pyran-2-one (7b)* (Method A: 82%, Method B: 44%) White solid: mp >260 °C; Rf = 0.24 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 5.75 (s, 1H), 4.86 (s, 1H), 2.93 (m, 2H), 1.64 (m, 2H), 1.41 (m, 2H), 0.93 (t, 3H) ppm; ¹³C NMR (126 MHz, Methanol-*d*₄) δ = 160.55, 160.06, 154.39, 102.35, 85.08, 42.10, 37.52, 18.34, 18.01, 11.06 ppm; LRMS (ESI-QTOF) calcd for C₁₀H₁₆NO₂ [M + H]⁺ 182.1181, found 181.1176.
- 4-(isopropylamino)-6-methyl-2H-pyran-2-one (7c)* (Method A: 65%, Method B: 69%) White solid: mp >260 °C; Rf = 0.13 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 5.76 (s, 1H), 5.21 (s, 1H), 3.63 (m, 1H), 2.13 (s, 3H), 1.20 (d, 6H) ppm; ¹³C NMR (126 MHz, Methanol-*d*₄) δ = 165.87, 158.99, 156.69, 97.95, 76.32, 41.94, 19.01 (2C), 16.63 ppm; LRMS (ESI-QTOF) calcd for C₉H₁₄NO₂ [M + H]⁺ 168.1025, found 168.1030.
- 4-(benzylamino)-6-methyl-2H-pyran-2-one (7d)* (Method A: 32%) White solid: mp 126-130 °C; Rf = 0.43 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 7.41 - 7.17 (m, 5H), 5.90 (s, 1H), 4.94 (s, 1H), 4.33 (d, 2H), 2.15 (s, 3H) ppm; LRMS (ESI-QTOF) calcd for C₁₃H₁₄NO₂ [M + H]⁺ 216.1025, found 216.1034. ¹³C NMR data agreed with the literature.¹⁷
- 4-(diethylamino)-6-methyl-2H-pyran-2-one (7e)* (Method A: 95%, Method B: 47%) Yellow solid: mp 96-98 °C; Rf = 0.30 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 5.80 (d, 1H), 4.99 (d, 1H), 3.41 (q, 4H), 2.21 (s, 3H), 1.19 (t, 6H) ppm; ¹³C NMR (126 MHz, Methanol-*d*₄) δ = 165.56, 160.24, 156.37, 94.74, 77.67, 42.67 (2C), 17.09 (2C), 16.63 ppm; LRMS (ESI-QTOF) calcd for C₁₀H₁₆NO₂ [M + H]⁺ 182.1181, found 182.1189.
- 6-methyl-4-(phenylamino)-2H-pyran-2-one (7f)* (Method A: 55%) White solid: mp 192-195 °C (Lit. 195-196 °C)¹⁵; Rf = 0.43 (silica gel, CH₂Cl₂/EtOAc 1:1); ¹H NMR (300 MHz, Acetone-*d*₆) δ = 7.48 - 7.36 (m, 2H), 7.32 - 7.14 (m, 3H), 5.95 (d, 1H), 5.27 (d, 1H), 2.14 (s, 3H) ppm; ¹³C NMR (126 MHz, Methanol-*d*₄) δ = 165.54, 160.18, 156.21, 141.31, 127.65 (2C), 123.91, 121.73 (2C), 97.47, 79.47, 16.84 ppm; LRMS (ESI-QTOF) calcd for C₁₂H₁₂NO₂ [M + H]⁺ 202.0868, found 202.0879.
21. Experimental procedure for the preparation of 4,4'-(piperazine-1,4-diyl)bis(6-methyl-2H-pyran-2-one) (**8**) (Method A: 21%, Method B: 54%) white solid: **Method A**: A mixture of 6-methyl-2-oxo-2H-pyran-4-yl 4-methylbenzenesulfonate (**2**) (0.1 g, 0.36 mmol, 2 eq) and piperazine (0.18 mmol, 1 eq) in ethanol (8 mL) was stirred at rt. overnight. After completion of the reaction (TLC monitoring), solvent was evaporated. The crude compound was purified by preparative thin layer chromatography (EtOAc/dichloromethane) to afford the desired product. **Method B**: A mixture of triacetic acid lactone (**1**) (0.15 g, 1.2 mmol, 2 eq) and piperazine (0.6 mmol, 1 eq) in water (1.5 mL) was stirred at rt. overnight. After completion of the reaction (TLC monitoring), solvent was evaporated. The crude compound was purified by preparative thin layer chromatography (EtOAc/dichloromethane) to afford the desired product: Rf = 0.13 (silica gel, CH₂Cl₂/EtOAc 7:1); ¹H NMR (300 MHz, Methanol-*d*₄) δ = 5.77 (s, 2H), 4.86 (s, 2H), 3.11 (s, 8H), 2.14 (s, 6H) ppm; ¹³C NMR (400 MHz, Methanol-*d*₄) δ = 179.72 (2C), 169.40 (2C), 161.39 (2C), 105.51 (2C), 87.39 (2C), 42.94 (2C), 18.22 (2C) ppm; LRMS (ESI-QTOF) calcd for C₁₆H₁₉N₂O₄ [M + H]⁺ 303.1345, found 303.1347.
22. Experimental procedure for the preparation of 4,4',4'',4'''-(1,4,7,10-tetraazacyclododecane-1,4,7,10-tetrayl)tetrakis(6-

Tetrahedron

methyl-2H-pyran-2-one (**7g**) (51%) White solid: A mixture of triacetic acid lactone (**1**) (0.14 g, 1.1 mmol, 4 eq) and cyclen (0.047 g, 0.27 mmol, 1 eq) in water (2 mL) was stirred at rt. overnight. After completion of the reaction (TLC monitoring), solvent was evaporated. The crude compound was purified by preparative thin layer chromatography (EtOAc/dichloromethane)

to afford the desired product: Rf = 0.14 (silica gel, CH₂Cl₂/EtOAc 7:1); ¹H NMR (400 MHz, Methanol-*d*₄) δ = 5.67 (s, 4H), 5.03 (s, 4H), 2.92 (s, 16H), 2.05 (s, 12H) ppm; ¹³C NMR (400 MHz, Methanol-*d*₄) δ = 179.97 (4C), 169.46 (4C), 161.35 (4C), 105.68 (4C), 87.34 (4C), 43.16 (8C), 18.22 (4C) ppm.

ACCEPTED MANUSCRIPT

Graphical Abstract

Leave this area blank for abstract info.

Triacetic Acid Lactone as a Common Intermediate for the Synthesis of 4-Hydroxy-2-pyridones and 4-Amino-2-pyrones

George A. Kraus*, Umayangani K. Wanninayake and Jashaun Bottoms

