

Ferrat Mélodie (Orcid ID: 0000-0001-7004-3565)

Dahl Kenneth (Orcid ID: 0000-0003-2948-2042)

"In-loop" carbonylation – a simplified method for carbon-11 labeling of drugs and radioligands

Mélodie Ferrat<sup>a\*</sup>, Kenneth Dahl<sup>b</sup>, Christer Halldin<sup>a</sup> and Magnus Schou<sup>a,c</sup>

<sup>a</sup>Department of Clinical Neuroscience, Center for Psychiatry Research, Karolinska Institutet and Stockholm County Council, SE-171 76 Stockholm, Sweden

<sup>b</sup> CAMH & University of Toronto, Toronto, ON M5T-1R8, Canada

<sup>c</sup> AstraZeneca PET Science Centre, Department of Clinical Neuroscience, Karolinska Institutet, S-171 76 Stockholm, Sweden

\*Author to whom correspondence should be addressed: Email: melodie.ferrat@ki.se

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/jlcr.3805

This article is protected by copyright. All rights reserved.

### Abstract:

Transition-metal mediated carbonylation with <sup>11</sup>C-labelled carbon monoxide ([<sup>11</sup>C]CO) is a versatile method for introducing <sup>11</sup>C ( $t_{1/2} = 20.3$  min) into drugs and radioligands for subsequent use in positron emission tomography (PET). The aim of the current study was to perform the <sup>11</sup>C-carbonylation reaction on the interior surface of a stainless-steel loop used for high performance liquid chromatography (HPLC). In the experimental setup, cyclotron produced <sup>11</sup>C-labelled carbon dioxide ( $[^{11}C]CO_2$ ) was converted to  $[^{11}C]CO$  by reduction over heated Molybdenum and swept into an HPLC loop pre-charged with the appropriate reaction mixture. Following a 5 min reaction, the radiochemical purity (RCP) and the trapping efficiency (TE) of the reaction mixture was determined. After optimization, [<sup>11</sup>C]N-Benzylbenzamide was obtained in quantitative radiochemical yield (RCY) following a 5 min reaction at room temperature. The methodology was further applied to label [<sup>11</sup>C]benzoic acid (RCP≥99%, TE>91%), [<sup>11</sup>C]methyl benzoate (RCP≥99%, TE>93%) and [<sup>11</sup>C]phtalide (RCP>99%, TE>88%). A set of pharmaceuticals was finally radiolabelled using nonoptimized conditions. Excellent yields were obtained for the histamine-3 receptor radioligand  $[^{11}C]AZ13198083$ , the oncology drug  $[^{11}C]olaparib$  and the dopamine D2 receptor radioligand [<sup>11</sup>C]raclopride, whereas a moderate yield was observed for the high-affinity dopamine D2 receptor radioligand [<sup>11</sup>C]FLB457. The presented "in-loop" process proved efficient for diverse <sup>11</sup>C-carbonylations, providing [<sup>11</sup>C]amides, [<sup>11</sup>C]esters and <sup>[11</sup>C]carboxylic acids in moderate to excellent RCYs. Based on the advantages associated with performing the radiolabelling step as an integrated part of the purification system, this methodology may become a valuable addition to the toolbox of methodologies used for <sup>11</sup>Ccarbonylation of drugs and radioligands for PET.

**Keywords:** Loop, <sup>11</sup>C-carbonylation, Carbon-11, Radioligands, Radiopharmaceutical chemistry

Accept

#### 1. INTRODUCTION

Positron emission tomography (PET) is a nuclear medicine imaging technique frequently used in translational research and drug development. After injection of a radiolabelled compound, PET produces a series of three-dimensional images of the tracer tissue distribution inside the living organism. A fundamental pre-requisite for PET is that the tracer molecule is labelled with a positron emitting radionuclide, such as <sup>11</sup>C ( $t_{1/2} = 20.3$  min). An important advantage with <sup>11</sup>C is that it can be introduced in the place of <sup>12</sup>C in bio-organic compounds without any structural modification of the target compound.

Due to the short half-life of carbon-11, its radiochemistry needs to be rapid, efficient and preferably automated to minimize radiation exposure to the radiochemist. The radionuclide is commonly produced by a medical cyclotron in the form of [<sup>11</sup>C]methane or [<sup>11</sup>C]carbon dioxide. Because of the convenient conversion of these species into [<sup>11</sup>C]methyl iodide,<sup>1,2</sup> the radionuclide is most commonly introduced into tracer molecules via <sup>11</sup>C-methylation. Although methyl groups are common in bioactive molecules, the majority lack this structural motif, which makes other labelling methods necessary to avoid structural modification of the tracer molecule.

Carbonyl groups are present in most biologically active molecules and are featured in a large number of drug molecules. Significant efforts have thus been put into the development of methods that use [<sup>11</sup>C]carbon monoxide ([<sup>11</sup>C]CO), available in one step from [<sup>11</sup>C]CO<sub>2</sub>, to introduce the <sup>11</sup>C-carbonyl motif into tracer molecules. Radiochemistry with [<sup>11</sup>C]CO is far from trivial, however, because of its low reactivity, poor solubility in organic solvents, high dilution in inert gas and the low concentration of isotopically labelled [<sup>11</sup>C]CO.<sup>3</sup> Many of these challenges have been successfully addressed via recirculating [<sup>11</sup>C]CO through the reaction media,<sup>4</sup> by performing the <sup>11</sup>C-carbonylation in a micro-autoclave (at 350 bar)<sup>3,5</sup> or inside a microtube reactor.<sup>6</sup> Other fruitful methods for improving the yield for these reactions include chemical complexation of [<sup>11</sup>C]CO with diborane (BH<sub>3</sub>-[<sup>11</sup>C]CO),<sup>7</sup> copper scorpionates (Cu-Sc-[<sup>11</sup>C]CO),<sup>8</sup> or introducing [<sup>11</sup>C]CO into the reaction media in a stream of Xenon carrier gas.<sup>9</sup> These developments have resulted in numerous PET studies with tracer molecules labelled using palladium mediated <sup>11</sup>C-carbonylation chemistry.<sup>10,11,12,13,14,15</sup>

In an effort to simplify <sup>11</sup>C-carbonylation reactions from a technical perspective, we were inspired by previous work that employed captive solvents and reagents confined within plastic or stainless-steel tubes for the radiochemical synthesis.<sup>16,17</sup> It was considered that such an approach, which maximizes the surface area between the gas and liquid phase, may be ideally suited for [<sup>11</sup>C]CO radiochemistry because of the increased opportunity for the radioactive gas to equilibrate with solution. In addition, an approach in which the HPLC loop is the reaction vessel for the <sup>11</sup>C-carbonylation was considered ideal, since transfer losses in such case are minimized and process automation simplified according to Wilson et al.<sup>17</sup> Thus, the aim of this study was to develop a novel "in-loop" method for PET tracer synthesis via

<sup>11</sup>C-carbonylation using [<sup>11</sup>C]CO (Fig. 1).

#### **RESULTS AND DISCUSSION**

An initial requirement for most "in-loop" <sup>11</sup>C-labeling processes is that the product must be formed in a one-pot, one-step procedure. We thus started our investigation on the "in-loop" <sup>11</sup>C-aminocarbonylation of iodobenzene with benzylamine, forming the product  $[^{11}C]N$ benzylbenzamide, a reaction that has been well characterized previously.<sup>18</sup> DMF was first tested as a solvent for the reaction because of its captive properties and compatibility with "in-loop" <sup>11</sup>C-methylation.<sup>17</sup> Likewise, the Pd-xantphos catalytic system was used as a starting point for the optimization, since it has been shown to be efficient in promoting <sup>11</sup>Caminocarbonylation reactions.<sup>18,19</sup> To our delight, a 5 min reaction at room temperature with Pd-xantphos in DMF furnished  $[^{11}C]N$ -benzylbenzamide at high trapping efficiency (TE), albeit at a moderate radiochemical purity (RCP) (Table 1, entry 1). Other solvents, such as DMSO and toluene, also provided good TEs in the "in-loop" <sup>11</sup>C-carbonylation (Table 1, entries 2 and 3), but the best yields were obtained in 1,4-dioxane and THF with near quantitative radiochemical yields (RCYs) (Table 1, entries 4 and 5). It is worth noting that an autoradiography experiment revealed that the majority of the radioactivity was trapped in the first quarter (~0.5 mL) of the stainless steel loop (Fig. 2). In comparison, a control reaction with  $Pd(PPh_3)_4$  provided  $[^{11}C]N$ -benzylbenzamide in 70% radiochemical purity, but with a poor TE (0.2%) and thus low RCY (0.14%).

With the improved conditions in hand, the methodology was next applied to radiolabel functional groups other than amides. Thus, a lactone, an ester and a carboxylic acid, were each radiolabelled with only minor modifications to those previously used in the radiosynthesis of [<sup>11</sup>C]*N*-benzylbenzamide (Scheme 1). Modifications to the original procedure include an increase in aryl halide concentration to improve trapping efficiency for [<sup>11</sup>C]*phtalide*, and an increased temperature to improve the RCP of [<sup>11</sup>C]*methyl b*enzoate. The observed yields from these reactions are on par with those observed for reactions performed in either vial or in micro-reactor setups, thus demonstrating that the utility of the developed "in-loop" <sup>11</sup>C-carbonylation method extends beyond aminocarbonylation.<sup>18,20</sup>

The "in-loop" carbonylation was finally tested on a set of druglike molecules (Scheme 2). Thus, carbon-11 radiolabelling of the histamine type-3 receptor radioligand AZ13198083, the oncology drug olaparib and the dopamine D<sub>2</sub> receptor radioligands raclopride<sup>14</sup> and FLB457 were performed. Again, some modifications to the developed procedure were important to achieve good RCYs for these compounds. Most importantly, heating was required for all substrates to obtain the target tracers at high RCP. In addition, as was previously demonstrated by Skrydstrup et al., [<sup>11</sup>C]olaparib cannot be produced with the Pd-xantphos system because of a phenyl scrambling from xantphos onto the aryl palladium complex.<sup>21</sup> Instead, the Pd-<sup>1</sup>Bu<sub>3</sub> system was used to produce the target compound, thus demonstrating that other catalytic systems than the Pd-xantphos system are also compatible with the "inloop" <sup>11</sup>C-carbonylation methodology. It is noteworthy that [<sup>11</sup>C]FLB457 could only be produced at a moderate TE under the employed conditions, but this may well be improved following further optimization. Finally, after integrating the loop system into a commercially available radiochemistry system (GE tracerlab FX-C, Uppsala, Sweden), [<sup>11</sup>C]AZ13198083

was obtained in a 38% decay-corrected RCY. The isolated radiochemical yield was lower than expected because of losses during HPLC isolation, formulation and sterile filtration. However, the RCP was >99% and the molar activity was 110 GBq/ $\mu$ mol.

# CONCLUSION

The presented "in-loop" process proved to be useful for diverse <sup>11</sup>C-carbonylation, providing both simple reference compounds (*e.g.* [<sup>11</sup>C]amides, [<sup>11</sup>C]esters and [<sup>11</sup>C]carboxylic acids) as well as PET tracer molecules (*e.g.* [<sup>11</sup>C]raclopride, [<sup>11</sup>C]olaparib) in moderate to excellent RCYs. Given its simplicity and efficiency, the methodology has the potential to be widely implemented for carbonylations in PET tracer synthesis.

# MATERIALS AND METHODS

### **General experimental information**

All chemical and solvents were obtained from Sigma-Aldrich (Sweden) and used without further purification, unless specified otherwise.

High performance liquid chromatography analysis (HPLC) was performed using a Hitachi L-6200 gradient pump and a Hitachi L-4000 variable wavelength UV-detector in a series with a Bioscan  $\beta$ +-flow detector. Analytical HPLC analysis was performed using a reverse phase column (XBridge, C18, 5 µm, 4.6 x 150 mm):

- a- For [<sup>11</sup>C]*N*-benzylbenzamide, eluted with a gradient between acetonitrile (A) and 0.1% TFA (B). The gradient was linear between 10 90% over 5 minutes and isocratic in between 5-8 min (ACN: 0.1% TFA, 90:10), at a flow rate of 3 ml/min.
- b- [<sup>11</sup>C]phtalide, eluted with a gradient between acetonitrile (A) and 0.1 M NH<sub>4</sub>HCO<sub>2</sub>
  (B). The gradient was linear between 10 90% over 9 minutes and isocratic in between 9-10 min (ACN: 0.1 M NH<sub>4</sub>HCO<sub>2</sub>, 90:10), at a flow rate of 2 ml/min.
- c- [<sup>11</sup>C]methyl benzoate, eluted with a gradient between acetonitrile (A) and 0.1% TFA (B). The gradient was linear between 10 90% over 5 minutes and isocratic in between 5-10 min (ACN: 0.1% TFA, 90:10), at a flow rate of 3 ml/min.
- d- [<sup>11</sup>C]benzoic acid, eluted with a gradient between acetonitrile (A) and 0.1% TFA (B).
  The gradient was linear between 10 90% over 7 minutes and isocratic in between 7-10 min (ACN: 0.1% TFA, 90:10), at a flow rate of 3 ml/min.
- e- For [<sup>11</sup>C]AZ13198083, eluted with a gradient between acetonitrile (A) and 0.1 M NH<sub>4</sub>HCO<sub>2</sub> (B). The gradient was linear between 10 90% over 9 minutes and isocratic in between 9-10 min (ACN: 0.1 M NH<sub>4</sub>HCO<sub>2</sub>, 90:10), at a flow rate of 2 ml/min.
- f- For [<sup>11</sup>C]olaparib, eluted with a gradient between acetonitrile (A) and 0.01 M  $NH_4HCO_2$  (B). The gradient was linear between 30 70% over 5 minutes and

isocratic in between 5-6 min (ACN: 0.1 M NH<sub>4</sub>HCO<sub>2</sub>, 90:10), at a flow rate of 3 ml/min.

g- For [<sup>11</sup>C]raclopride, eluted with a gradient between acetonitrile (A) and 0.1% TFA (B). The gradient was linear between 10 - 90% over 9 minutes and isocratic in between 9-10 min (ACN: 0.1% TFA, 90:10), at a flow rate of 2 ml/min.

h- For [<sup>11</sup>C]FLB457, eluted with a gradient between acetonitrile (A) and 0.1 M NH<sub>4</sub>HCO<sub>2</sub> (B). The gradient was linear between 30 - 70% over 9 minutes and isocratic in between 9-10 min (ACN: 0.1 M NH<sub>4</sub>HCO<sub>2</sub>, 90:10), at a flow rate of 2 ml/min.

The identity of all radioactive products was confirmed by co-elution with the corresponding non-radioactive compound.

#### Radiochemistry

All experiments were performed in accordance with local and national rules and laws that govern the work with open sources of radiation. The study was carried out at the PET Centre, Karolinska Institutet, (Stockholm, Sweden).

<sup>[11</sup>C]CO<sub>2</sub> was produced by a GEMS PETtrace cyclotron (General Electrics Medical Systems, Uppsala, Sweden). The <sup>14</sup>N(p, $\alpha$ )<sup>11</sup>C nuclear reaction, performed by proton bombardment of a pressurized gas target containing nitrogen (AGA, Nitrogen 6.0) and 1% oxygen (AGA, Oxygen 4.8), was used to produce [<sup>11</sup>C]CO<sub>2</sub>. At the end of bombardment (EOB), [<sup>11</sup>C]CO<sub>2</sub> was transferred from the target to a [<sup>11</sup>C]CO-synthesizer prototype (General Electrics Medical Systems, Uppsala, Sweden).  $[^{11}C]CO_2$  was trapped on a molecular sieve column (0.6 mg packed in a <sup>1</sup>/<sub>4</sub>" tube, mesh 80/100, GRACE, USA) at room temperature. The accumulated  $[^{11}C]CO_2$  was released into a controlled stream of helium (10 ml/min) while heating the molecular sieve trap to 360°C. The purified  $[^{11}C]CO_2$  was pre-concentrated by a stainless steel loop trap immersed in liquid nitrogen (-196°C) prior to on-line reduction over a heated Molybdenum column (850°C). The effluent gas was purified through an ascarite<sup>®</sup> (sodium hydroxide-coated silica, 20-30 mesh, Sigma-Aldrich, Sweden) column prior to concentrating the  $[^{11}C]CO$  on a silica gel trap immersed in liquid nitrogen. Following completed entrapment, [<sup>11</sup>C]CO was released in a stream of Helium into a stainless steel HPLC loop for the <sup>11</sup>C-carbonylation reaction. Untrapped [<sup>11</sup>C]CO was collected in a gas-tight bag (Tedlar<sup>®</sup> gas sampling bag push lock valve 1L, Sigma-Aldrich, Sweden).

#### In-loop <sup>11</sup>C-carbonylation setup

A commercial HPLC injection valve (VICI, #C52-12061, Switzerland) was equipped with a stainless-steel loop (2 mL, VICI, #SL2KCUW, Switzerland) and an injection port (for 1/16" fill port, VICI #VISF-2, Switzerland). Prior to start of synthesis, the injection valve was set to LOAD position (Fig. 1) and a solution of aryl halide, palladium-source, supporting ligand and nucleophile in anhydrous solvent, was injected using a 1 mL syringe capped with a filter

(Acrodisc 13 mm minispike with 0,45  $\mu$ m GHP Membrane, Sigma Aldrich, Sweden) through the injection port of the loop. The isolated and concentrated [<sup>11</sup>C]CO was then swept into the loop at 5 mL/min, and after a 5 min reaction, the contents of the loop were transferred to a sealed test tube by purging the loop with THF (~2 mL). After transfer to the test tube, the radioactivity in the test tube was measured before and after it was purged with N<sub>2</sub> (to assess TE), followed by a measurement of the gas waste bag that was connected on the outlet of the HPLC loop. The radioactivity in the test tube was also analyzed using radio-HPLC to assess the RCP that gives information about reaction selectivity. Moreover, as [<sup>11</sup>C]CO is a radioactive gas, the parameter TE needs to be taken into account. TE is the total amount of [<sup>11</sup>C]CO trapped in the reaction solution that is converted to non-volatile entities at the end of the reaction. RCY was calculated from the RCP and TE (*i.e.* RCY = RCP x TE).

#### **Application to GE Tracerlab FX-C**

A Tracerlab FX-C system (GE PETtrace, Uppsala, Sweden) was used for semi-preparative HPLC purification, in parallel with the [<sup>11</sup>C]CO-synthesizer prototype (General Electrics Medical Systems, Uppsala, Sweden) to produce [<sup>11</sup>C]**7**. In short, <sup>11</sup>C-carbon monoxide was transferred from the [<sup>11</sup>C]CO-system to the stainless-steel loop in the Tracerlab FX-C system where the reaction was taking place. The crude product was obtained in the HPLC loop and then purified using a semi-preparative HPLC column (ACE C-18, 5µm, 10 × 250 mm; Advanced Chromatography Technologies Ltd; Aberdeen, UK) eluted with an isocratic mobile phase consisting of 30% MeCN in HCO<sub>2</sub>NH<sub>4</sub> (0.1M). The eluate was monitored for absorbance ( $\Box \Box = 254$  nm) and for radioactivity. [<sup>11</sup>C]**7** ( $t_R = 8-10$  min) was collected and diluted with water (~50 mL) and then concentrated on a C-18 SepPak (Waters, Sweden). The SepPak was then washed with water (~8 mL) prior to elution with ethanol (99%, 1.5 mL) into saline (10 mL). The formulation was finally filtered through a yellow sterile filter (0.22 µm, Millipore, Millex@GV) to yield the final product in a sterile injection vial.

#### Synthesis of [<sup>11</sup>C]*N*,*N*-benzylbenzamide

An oven-dried disposable 4 mL vial was equipped with a screw thread cap, a rubber septum and cooled under Nitrogen (N<sub>2</sub>) atmosphere. Pd<sub>2</sub>[ $\pi$ -cinnamyl]Cl<sub>2</sub> (2.3 mg; 1eq), xantphos (5,0 mg; 2 eq), iodobenzene (1.4 µL, 1.42 eq) were dissolved in anhydrous THF (700 µL) and the resulting mixture was flushed with N<sub>2</sub> until the THF was fully evaporated (20 min). 1,4-Dioxane (150 µL) and Benzylamine (10 µL) were added and the reaction mixture was introduced into the HPLC loop approximately 2 min prior to start of synthesis. The resulting mixture was allowed to react for 5 min at r.t.



### Synthesis of [<sup>11</sup>C]Phtalide

Pd<sub>2</sub>[ $\pi$ -cinnamyl]Cl<sub>2</sub> (6.7 mg; 3 eq), xantphos (7.5 mg; 3 eq) and iodobenzylalcohol (3.0  $\mu$ L, 3 eq) were dissolved in anhydrous THF (700  $\mu$ L) kept at r.t. for 15 min and then heated at 75°C during 5 min. THF (150  $\mu$ L) was added and the resulting mixture was introduced into the HPLC loop approximately 2 min prior to start of synthesis. The resulting mixture was allowed to react for 5 min at 75°C.

## Synthesis of [<sup>11</sup>C]Methyl benzoate

 $Pd_2[\pi\text{-cinnamyl}]Cl_2$  (2.3 mg; 1eq), xantphos (5.0 mg; 2eq) and iodobenzene (2.2 µL, 2.9 eq) were dissolved in anhydrous THF (700 µL) and the resulting mixture was flushed with N<sub>2</sub> until the THF was fully evaporated (20 min). THF (150 µL) and a solution of lithium methoxide in MeOH (1M, 40 µL) were added and the reaction mixture was introduced into the HPLC loop approximately 2 min prior to start of synthesis. The resulting mixture was allowed to react for 5 min at 120°C.

## Synthesis of [<sup>11</sup>C]Benzoic acid

 $Pd_2[\pi$ -cinnamyl]Cl<sub>2</sub> (2.2 mg), xantphos (5.0 mg) and iodobenzene (1.4 µL) were dissolved in anhydrous THF (700 µL) and the resulting mixture was flushed with N<sub>2</sub> until the THF was fully evaporated (20 min). 1,4-Dioxane (150 µL) and a solution of NaOH (0.35 M, 3.5 µL) were added and the reaction mixture was introduced into the HPLC loop approximately 2 min prior to start of synthesis. [<sup>11</sup>C]CO was swept at r.t to the loop in which the coupling reagents were already added. The resulting mixture was allowed to react for 5 min at r.t.

## Synthesis of [<sup>11</sup>C]AZ13198083

Pd<sub>2</sub>[ $\pi$ -cinnamyl]Cl<sub>2</sub> (2.2 mg; 1 eq), xantphos (5.0 mg; 2 eq), ((2S)-2-(4-bromophenyl)cyclopropyl)(4-cyclobutylpiperazin-1-yl)methanone (3.3 µL) were dissolved in anhydrous THF (700 µL) and the resulting mixture was flushed with N<sub>2</sub> until the THF was fully evaporated (20 min). A solution of ethylamine in THF (2M, 600 µL) was added and the reaction mixture was introduced into the HPLC loop approximately 2 min prior to start of synthesis. The resulting mixture was allowed to react for 5 min at 100°C.

# Synthesis of [<sup>11</sup>C]Olaparib

Pd(dba)<sub>2</sub> (11.5 mg), P<sup>t</sup>Bu<sub>3</sub> (4.9  $\mu$ L) and 4-(4-fluoro-3-iodobenzyl)phthalazin-1(2H)-one (7.6 mg) were dissolved in anhydrous THF (700  $\mu$ L) and the resulting mixture was flushed with N<sub>2</sub> until the THF was fully evaporated (20 min). THF (150  $\mu$ L) and cyclopropyl(piperazin-1-yl)methanone (10.1  $\mu$ L) were added and the reaction mixture was introduced into the HPLC

loop approximately 2 min prior to start of synthesis. The resulting mixture was allowed to react for 5 min at 100°C.

## Synthesis of [<sup>11</sup>C]Raclopride

 $Pd_2[\pi$ -cinnamyl]Cl<sub>2</sub> (6.4 mg; 1 eq), xantphos (7.1 mg; 1eq) and 4,6-dichloro-2-iodo-3methoxyphenol (5.9 mg, 1.5 eq) were dissolved in anhydrous THF (1 mL) and the resulting mixture was flushed with N<sub>2</sub> until the THF was fully evaporated (20 min). THF (150 µL) and (S)-(-)-2-Aminomethyl-1-ethylpyrrolidine (30 µL) were added and the reaction mixture was introduced into the HPLC loop approximately 2 min prior to start of synthesis. The resulting mixture was allowed to react for 5 min at 75°C.



Synthesis of [ C]FLD457

Pd<sub>2</sub>[ $\pi$ -cinnamyl]Cl<sub>2</sub> (2.2 mg; 1 eq), Xantphos (5.0 mg; 2 eq), 5-bromo-1-iodo-2,3dimethoxybenzene (5.1 mg) were dissolved in anhydrous THF (700 µL) and the resulting mixture was flushed with N<sub>2</sub> until the THF was fully evaporated (20 min). 1,4-Dioxane (150 µL) and (S)-(-)-2-Aminomethyl-1-ethylpyrrolidine (27 µL) were added and the reaction mixture was introduced into the HPLC loop approximately 2 min prior to start of synthesis. The resulting mixture was allowed to react for 5 min at 75°C.

# ACKNOWLEDGMENTS

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement N°675071.

The authors would like to thank members of PET group at Karolinska Institutet.

Acce

### REFERENCES

- 1. Långström B, Lundqvist H. The preparation of <sup>11</sup>C-methyl iodide and its use in the synthesis of <sup>11</sup>C-methyl-L-methionine. Int J Appl Radiat Isot 1976;27(7):357-363.
- 2. Larsen P, Ulin J, Dahlström K, Jensen M. Synthesis of [<sup>11</sup>C]Iodomethane by Iodination of [<sup>11</sup>C]Methane. Appl Radiat Isot 1997;48:153-157.
- 3. Långström B, Itsenko O, Rahman O. [<sup>11</sup>C]Carbon monoxide, a versatile and useful precursor in labelling chemistry for PET-ligand development. J Label Compd Radiopharm 2007;50(9-10):794-810.
- 4. Lidström P, Kihlberg T, Långström B. [<sup>11</sup>C]Carbon monoxide in the palladium-mediated synthesis of <sup>11</sup>C-labelled ketones. J Chem Soc Perkin Trans 1 1997;(18):2701-2706.
  - 5. Kihlberg T, Långströms B. Biologically active C-11labeled amides using palladiummediated reactions with aryl halides and [C-11]carbon monoxide. J Org Chem 1999;64(25):9201-9205.
  - Miller PW, Long NJ, De Mello AJ, Vilar R, Audrain H, Bender D, Passchier J, Gee A. Rapid multiphase carbonylation reactions by using a microtube reactor: applications in positron emission tomography 11 C-radiolabeling. Angew Chem Int Ed 2007;46(16):2875-2878.
  - 7. Audrain H, Martarello L, Gee A, Bender D. Utilisation of <sup>11</sup>C-labelled boron carbonyl complexes in palladium carbonylation reaction. Chem Commun 2004;(5):558-559.
- 8. Kealey S, Miller PW, Long NJ, Plisson C, Martarello L, Gee AD. Copper(I) scorpionate complexes and their application in palladium-mediated [<sup>11</sup>C]carbonylation reactions. Chem Commun 2009;(25):3696-3698.
- Eriksson J, Hoek JVD, Windhorst AD. Transition metal mediated synthesis using [<sup>11</sup>C]CO at low pressure a simplified method for <sup>11</sup>C-carbonylation. J Label Compd Radiopharm 2012;55(6):223-228.
- Lindhe Ö, Almqvist P, Kågedal M, Gustafsson S-Å, Bergström M, Nilsson D, Antoni G, Lindhe Ö, Almqvist P, Kågedal M, Gustafsson S-Å, Bergström M, Nilsson D, Antoni G. Autoradiographic mapping of 5-HT(1B/1D) binding sites in the rhesus monkey brain using [carbonyl-C]zolmitriptan. Int J Mol Imaging 2011;2011(2011):6.
- Wall A, Kågedal M, Bergström M, Jacobsson E, Nilsson D, Antoni G, Frändberg P, Gustavsson SA, Långström B, Yates R. Distribution of Zolmitriptan into the CNS in Healthy Volunteers. Drugs in R&D 2005;6(3):139-47.
- 12. Lu S, Hong J, Itoh T, Fujita M, Inoue O, Innis RB, Pike VW. [carbonyl-11C]Benzyl acetate: Automated radiosynthesis via Pd-mediated [11C]carbon monoxide chemistry and PET measurement of brain uptake in monkey. J Label Compd Radiopharm 2010;53(8):548-551.
- 13. Takashima-Hirano M, Ishii H, Suzuki M. Synthesis of (11)CAm80 via novel Pd(0)mediated rapid (11)CCarbonylation using arylboronate and (11)CCarbon Monoxide. ACS Med Chem Lett 2012; 3(10):804-807.
- 14. Rahman O, Takano A, Amini N, Dahl K, Kanegawa N, Långströms B, Farde L, Halldin C. Synthesis of ([<sup>11</sup>C]carbonyl)raclopride and a comparison with ([<sup>11</sup>C]methyl)raclopride in a monkey PET study. Nucl Med Biol 2015;42(11):893-898.

- 15. Dahl K, Nakao R, Amini N, Moein MM, Finnema S, Malmquist J, Varnäs K, Schou M. Development of [Carbonyl-<sup>11</sup>C]AZ13198083, a novel histamine type-3 receptor radioligand with favorable kinetics. *ACS Chem Neurosci* **2018**;*9*(5):906-911.
- McCarron JA, Turton DR, Pike VW, Poole K.G. Remotely-controlled production of the 5-HT1A receptor radioligand, [carbonyl-<sup>11</sup>C]WAY-100635, via <sup>11</sup>C-carboxylation of an immobilized Grignard reagent. J Label Compd Radiopharm 1996;38(10):941-953.
- 17. Wilson AA, Garcia A, Jin L, Houle S. Radiotracer synthesis from [C-11]-iodomethane: A remarkably simple captive solvent method. Nucl Med Biol 2000;27(6):529-532.
- 18. Dahl K, Schou M, Amini N, Halldin C. Palladium-mediated [<sup>11</sup>C]carbonylation at atmospheric pressure: A general method using xantphos as supporting ligand. Eur J Org Chem 2013;(7):1228-1231.
- 19. Martinelli JR, Watson DA, Freckmann DMM, Barder TE, Buchwald SL. Palladiumcatalyzed carbonylation reactions of aryl bromides at atmospheric pressure: A general system based on xantphos. J Org Chem 2008;73(18):7102-7107.
- 20. Dahl K, Schou M, Ulin J, Sjöberg CO, Farde L, Halldin C. <sup>11</sup>C-carbonylation reactions using gas–liquid segmented microfluidics. RSC Adv 2015;5(108):88886-88889.
- 21. Andersen TL, Friis SD, Audrain H, Nordeman P, Antoni G, Skydstrup T. Efficient <sup>11</sup>Ccarbonylation of isolated aryl palladium complexes for PET: Application to challenging radiopharmaceutical synthesis. J Am Chem Soc 2015;137(4):1548-1555.

Accepted

Table 1. Investigation of solvent effects on the radiochemical yield of  $[^{11}C]N$ -benzylbenzamide.



Reaction conditions:  $Pd_2(\pi$ -cinnamyl)Cl<sub>2</sub> (1 eq), Xantphos (2eq), Iodobenzene(1,4  $\mu$ L), THF(700  $\mu$ L) in appropriate solvent (150  $\mu$ L) and Benzylamine (10  $\mu$ L)

 $^a$  Oxidative addition takes place during evaporation of THF (700  $\mu L)$ 

<sup>b</sup> n = 3 (triplicate)

Acceptec



Figure 1. Schematic representation of the HPLC loop applied in the <sup>11</sup>C-carbonylation procedure. The yellow arrows indicate the flow direction for [<sup>11</sup>C]CO.

Accepted



Figure 2. Picture (left) and autoradiographical image (right) of the HPLC loop following entrapment of [<sup>11</sup>C]CO in the synthesis of [<sup>11</sup>C]*N*-benzylbenzamide. The arrows indicate the entry and exit points of the He carrier gas through the HPLC loop.





[<sup>11</sup>C]methylbenzoate

RCC ≥ 99% ; TE = 93%



[<sup>11</sup>C]benzoic acid RCC  $\ge$  99% ; TE = 91%

Scheme 1. Synthesis of [<sup>11</sup>C]phtalide, [<sup>11</sup>C]methyl benzoate and [<sup>11</sup>C]benzoic acid, via "inloop" <sup>11</sup>C-carbonylation.





#### **Graphical Abstract**



A new "in-loop" methodology was developed to simplify palladium-mediated carbonylations with [<sup>11</sup>C]carbon monoxide. The methodology proved to be useful in generating <sup>11</sup>C-labelled amides, esters and carboxylic acids. Given its simplicity, efficiency and compatibility with automation, this methodology may become a valuable addition to the toolbox of methodologies used for <sup>11</sup>C-carbonylation of drugs and radioligands for PET.