

Design and Synthesis of 4-(4-Benzoylaminophenoxy)phenol Derivatives As Androgen Receptor Antagonists


Ayumi Yamada, Shinya Fujii,* Shuichi Mori, and Hiroyuki Kagechika

Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, 2-3-10 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-0062, Japan

Supporting Information

ABSTRACT: We report the design and synthesis of novel 4-(4-benzoylaminophenoxy)phenol derivatives that bind to the androgen receptor (AR) ligand-binding domain and exhibit potent androgen-antagonistic activity. Compound 22 is one of the most potent of these derivatives, inhibiting the dihydrotestosterone-promoted growth of SC-3 cell line bearing wild-type AR (IC_{50} 0.75 μ M), LNCaP cell line bearing T877A-mutated AR (IC_{50} 0.043 μ M), and 22Rv1 cell line bearing H874Y-mutated AR (IC_{50} 0.22 μ M). Structure–activity relationship studies confirmed that the pharmacophore of these novel AR antagonists is distinct from the nitro- or cyano-substituted anilide substructure of other nonsteroidal AR antagonists. This novel pharmacophore is expected to provide a basis for designing new antiprostata cancer agents.

KEYWORDS: Androgen receptor (AR), AR antagonist, prostate cancer, pharmacophore, 4-(4-benzoylaminophenoxy)phenol



SC-3 cells (wild-type AR) : IC_{50} 0.75 μ M
LNCaP cells (T877A AR) : IC_{50} 0.043 μ M
22Rv1 cells (H874Y AR) : IC_{50} 0.22 μ M

Androgen receptor (AR) is a member of the nuclear receptor superfamily of ligand-dependent transcriptional factors.^{1,2} AR is activated by binding of endogenous androgens, such as testosterone and its activated form, dihydrotestosterone (DHT), and regulates numerous physiological processes, including growth, maintenance of the male reproductive system, and homeostasis of bone and muscle.^{3,4} Because AR plays an important role in progression of prostate cancer, AR antagonists are used clinically for treatment of prostate cancer. However, chronic administration of AR antagonists often leads to development of resistance, so-called castration-resistant prostate cancer (CRPC),^{5,6} and mutation of AR is thought to be the main cause.^{7–9} For example, a representative nonsteroidal AR antagonist, flutamide (1), and its activated metabolite, hydroxyflutamide (2), act as agonists of AR bearing T877A, the most common mutation of AR in CRPC and consequently exacerbate the cancer.¹⁰

All the nonsteroidal AR antagonists currently in clinical use are derivatives of flutamide bearing a nitrophenyl or a cyanophenyl group as the pharmacophore (Figure 1),¹¹ and these antagonists are often ineffective for mutated ARs.¹² Intensive studies based on flutamide derivatives have been investigated, and recently, novel flutamide derivative MDV3100 (5) was developed as a potent AR antagonist effective for CRPC.¹³ Another possible approach to overcome CRPC would be to develop AR antagonists bearing a novel pharmacophore, different from those of flutamide analogues. For example, nicotinamide derivative DIMN (6) was developed as novel type of nonsteroidal AR antagonists.¹⁴

With the aim of developing novel AR antagonists, we focused on the structure of the natural pigment curcumin (7, Figure 2). Several curcumin analogues possess antiandrogenic activity,^{15,16} and 7 does not possess the cyanophenyl or nitrophenyl group




Figure 1. Structures of nonsteroidal AR antagonists.

that is the key structure of potent nonsteroidal AR antagonists so far developed. Structurally, curcumin (7) consists of two phenolic groups linked by an unsaturated 1,3-diketone (or its enol form), which would be chemically and metabolically reactive.¹⁷ Therefore, we assumed that at least one phenolic group of 7 was essential for the AR antagonistic activity and that the overall structural framework of 7 could be replaced by stable, drug-like structures. Our previous studies on retinoids suggested that benzanilide is a versatile structure for replacing conjugated polyene substructure.^{18,19} Recent studies by us and others also suggested that a Ph–X–Ph skeleton, such as diphenylmethane (X = carbon),²⁰ diphenyl ether (X =

Received: May 10, 2013

Accepted: August 18, 2013




Figure 2. Design of novel AR antagonists bearing phenol substructure.


oxygen),²¹ diphenylamine (X = nitrogen),²² or diphenylsilane (X = silane),²³ is a suitable hydrophobic substructure for nuclear receptor ligands.

On the basis of those findings, we designed the phenoxybenzanilide core structure as a candidate scaffold of AR antagonists, and we initially synthesized the parent compound **8** (Figure 2). Biological evaluation revealed that **8** exhibited significant AR antagonistic activity, being a potent inhibitor of androgen-dependent SC-3 cell²⁴ proliferation (IC₅₀ = 9.4 μM). Here, we report the synthesis and biological activity of **8** and its derivatives (general formula shown in Figure 2) as candidate new-generation AR antagonists bearing a novel pharmacophore distinct from those of conventional flutamide analogues.

Compound **8** and 4-phenoxyphenols **12a–p** were synthesized as shown in Scheme 1.²⁵ S_NAr reaction using 4-fluoronitrobenzene (**9**) and hydroquinone gave diphenyl ether **10**, and then reduction of the nitro group afforded primary amine **11**. Amide bond formation of **11** with benzoic acid or benzoyl chloride derivatives gave compounds **8** and **12a–p**. Compounds **12c**, **12f**, and **12h** were synthesized from the corresponding *O*-monomethyl (for **12c**) or *O*-monoacetyl (for **12f** and **12h**) derivatives by removal of the *O*-substituent. Diphenyl ether **15** without a phenolic hydroxyl group was also prepared for investigation of the significance of the phenolic hydroxyl group. S_NAr reaction using **7** and phenol gave diphenyl ether **13**, and then reduction of the nitro group afforded primary amine **14**. Amide bond formation of **14** using benzoyl chloride gave compound **15** (Scheme 1).

Scheme 2 shows the synthesis of compounds **22** and **23** bearing methyl group(s) at the central benzene ring and the synthesis of heterocyclic analogues **30** and **31**. S_NAr reaction using hydroquinone and fluoronitrobenzene **16** or **17** gave diphenyl ethers **18** and **19**, respectively. Reduction of the nitro group gave amine **20** and **21**, and following amide formation afforded the target compounds **22** and **23**. Compounds bearing pyridine or pyrimidine as the central aromatic ring were also similarly synthesized. Reaction of 2-chloro-5-nitropyridine **24** or 2-chloro-5-nitropyrimidine **25** with 4-benzyloxyphenol gave diaryl ethers **26** and **27**, respectively. Reduction of the nitro


Scheme 1. Synthesis of **8**, **12a–p**, and **15**^a



^aReagents and conditions: (a) hydroquinone, NaOH, DMSO–H₂O, 50 °C, 56%; (b) Pd–C, H₂, MeOH, rt, 92–98%; (c) method A, aryl chloride, THF, rt, 55–86%; method B, aryl chloride, pyridine, rt, 50–60%; method C, (i) (COCl)₂, DMF, CH₂Cl₂, rt; (ii) **11**, THF, rt, 39–84%; method D, (i) (COCl)₂, DMF, CH₂Cl₂, rt; (ii) **11**, pyridine, THF, rt, 46%; (d) BBr₃, CH₂Cl₂, 40 °C, 50%; (e) K₂CO₃, MeOH–H₂O, rt, 95%; (f) NaOEt, EtOH, rt, 30%; (g) phenol, K₂CO₃, DMF, 40 °C, 90%.

group and deprotection of the benzyl group by catalytic hydrogenation afforded amines **28** and **29**, and then, amide condensation gave the target compounds **30** and **31**.

The antiandrogenic activity of the synthesized compounds was initially evaluated by assay of growth-inhibitory activity toward SC-3.²⁴ SC-3 cells have wild-type AR and proliferate androgen-dependently. None of the synthesized compounds promoted growth of SC-3 cells. Table 1 shows the inhibitory potency of the test compounds toward DHT-promoted cell growth of SC-3. Compound **12b** exhibited the most potent antiandrogenic activity among the synthesized compounds, and **12l** also exhibited potent antiandrogenic activity exceeding that of the lead compound **8**. Deletion of the phenolic hydroxyl group (compound **15**) resulted in significant loss of potency (20% inhibition at the concentration of 10 μM). Ohtsu et al. reported that the phenolic hydroxyl group was not essential for its AR antagonistic activity,¹⁵ and therefore, the importance of this hydroxyl group is intriguing. Interestingly, compounds **12i**, **12j**, and **12k** bearing a nitrophenyl group did not exhibit significant antiandrogen activity. The nitrophenyl group is a common pharmacophore of conventional nonsteroidal AR antagonists, and this result suggested that the 4-(4-benzoylamino-phenoxy)phenol derivatives are a novel class of AR antagonists bearing a unique pharmacophore, different from those of AR antagonists so far known. Bisphenol compound **12f** and compounds bearing a pyridine ring **12n** and **12o** exhibited moderate activity similar to that of **8**. These results indicate that the 3-methoxybenzoyl group of **12b** was the most suitable as the terminal functionality.

Scheme 2. Synthesis of 22, 23, 30, and 31^a

^aReagents and conditions: (a) hydroquinone, NaOH, DMSO–H₂O, 50 °C to rt, 66% (18), 55% (19); (b) for 20, Pd–C, H₂, MeOH, rt, 94%; for 21, Pd–C, H₂, THF, rt, 90%; (c) for 22, *m*-anisoyl chloride, THF, rt, 63%; for 23, *m*-anisoyl chloride, pyridine, THF, rt, 67%; (d) for 26, 4-benzyloxyphenol, *t*-BuOK, DMF, rt, 46%; for 27, *p*-benzyloxyphenol, K₂CO₃, DMF, rt, 38%; (e) Pd–C, H₂, THF, rt, quant (28), 80% (29); (f) *m*-anisoyl chloride, THF, rt, 76% (30), 79% (31).

Table 1. AR Antagonistic Activity of Synthesized Compounds by Means of Inhibition of DHT-Promoted SC-3 Cell Proliferation^a

| compd | R | X | IC ₅₀ (μM) |
|-------|------------------------|----|-----------------------|
| 2 | | | 0.37 |
| 8 | Ph | OH | 9.4 |
| 12a | 2-MeO–Ph | OH | >10 |
| 12b | 3-MeO–Ph | OH | 2.0 |
| 12d | 4-MeO–Ph | OH | >10 |
| 12f | 2-HO–Ph | OH | 8.8 |
| 12c | 3-HO–Ph | OH | n.d. ^b |
| 12h | 4-HO–Ph | OH | n.d. ^b |
| 12i | 2-NO ₂ –Ph | OH | >10 |
| 12j | 3-NO ₂ –Ph | OH | >10 |
| 12k | 4-NO ₂ –Ph | OH | >10 |
| 12l | 3-NMe ₂ –Ph | OH | 3.0 |
| 12m | 4-NMe ₂ –Ph | OH | 6.7 |
| 12n | 2-pyridyl | OH | 8.7 |
| 12o | 3-pyridyl | OH | 9.7 |
| 12p | 4-pyridyl | OH | >10 |
| 15 | Ph | H | >10 |

^aSC-3 cell growth was promoted by 1 nM DHT. ^bInhibition was not detected at the concentration of 10 μM.

For further structural development of these novel AR antagonists, the structure–activity relationship of the central aromatic ring was investigated. Compounds bearing phenol and methoxybenzoyl groups as terminal aromatic rings and different center rings were synthesized, and their activities were compared by means of antiandrogen activity assay with SC-3 cells, competitive AR binding assay, and luciferase assay (Table 2). Compound 22 bearing a methyl group at the center ring exhibited the most potent activity toward SC-3 cells, and the potency (IC₅₀ = 0.75 μM) of this compound was of the same order as that of hydroxyflutamide (2). Dimethylated derivative 23 also exhibited similar potency to 12b, whereas heterocyclic compounds 30 and 31 exhibited lower potency than 12b in SC-3 assay. Compounds 22 and 23 exhibited strong binding affinity for wild-type AR, whereas heterocyclic derivatives 30 and 31 did not. The results of luciferase assay using MDA-Kb2 cell

Table 2. Androgen Antagonistic Activity of Synthesized Phenol Derivatives by Means of SC-3 Assay, Binding Affinity, and Reporter Assay

| compd | Z ₁ | Z ₂ | SC-3 assay ^a | binding affinity ^b | luciferase assay ^c |
|-------|----------------|----------------|-------------------------|-------------------------------|-------------------------------|
| | | | IC ₅₀ (μM) | IC ₅₀ (μM) | % inhibition |
| 2 | | | 0.37 | 7.1 | 60 |
| 12b | CH | CH | 2.0 | 20 | 6 |
| 22 | CMe | CH | 0.75 | 9.2 | 66 |
| 23 | CMe | CMe | 2.9 | 5.4 | 69 |
| 30 | N | CH | 7.2 | >10 | n.d. ^d |
| 31 | N | N | 5.9 | >10 | n.d. ^d |

^aCell growth was promoted by 1 nM DHT. ^bCompetitive binding assay using hAR-LBD and ³H-labeled DHT (4 nM). ^cInhibition (% of control) of luciferase activity by 10 μM test compound using MDA-Kb2 cell line. The concentration of DHT was 0.1 nM. ^dInhibition was not detected at the concentration of 10 μM.

line²⁶ also supported the idea that the AR antagonistic activities of compounds 22 and 23 are due to their AR-modulating activity.

Then, we investigated the antiandrogenic activity of the selected compounds toward human prostate cancer cell lines LNCaP and 22Rv1, which have mutation in the AR ligand-binding domain. The LNCaP cell line expresses T877A AR, which is the most frequently detected mutation of AR in castration-resistant prostate cancer.^{27,28} Hydroxyflutamide (2) and some other AR antagonists act as agonists toward T877A AR,¹⁰ and therefore, we use 4 as positive control of this assay. The 22Rv1 cell line expresses H874Y AR.²⁹ None of the compounds examined promoted growth of these prostate cancer cell lines, and all of them exhibited significant inhibitory activity toward DHT-induced proliferation of these cell lines (Table 3). Among them, compound 22, which was the most potent AR antagonist in SC-3 assay, exhibited the most potent antiandrogenic activity toward LNCaP cell line (IC₅₀ = 0.043 μM). Compound 22 also exhibited potent antiandrogenic activity toward 22Rv1 cell line (IC₅₀ = 0.22 μM). The antiandrogenic activities of compound 22 are more potent than those of bicalutamide (4), one of the most effective

Table 3. Antiandrogenic Activity and Cytotoxicity of Selected Phenol Derivatives toward Three Human Prostate Cancer Cell Lines

| compd | LNCaP (T877A-AR) | 22Rv1 (H874Y-AR) | PC-3 |
|-------|------------------------------------|------------------------------------|------------------------------------|
| | IC ₅₀ (μM) ^a | IC ₅₀ (μM) ^a | IC ₅₀ (μM) ^b |
| 4 | 0.94 | 4.6 | >10 |
| 12b | 0.29 | 1.9 | 5.6 |
| 22 | 0.043 | 0.22 | 5.7 |
| 23 | 2.4 | 1.3 | >10 |

^aCell growth was promoted by 10 nM DHT. ^bIC₅₀ values for androgen-independent cell proliferation.

nonsteroidal AR antagonists. These results suggest that compound **22** is a potent AR antagonist active toward both wild-type and mutant AR. We also examined the effect of compounds toward androgen-independent prostate cancer cell line PC-3. Compound **12b** and **22** exhibited cell-growth inhibiting potency at micromolar concentrations. This result indicates that these compounds possess anticancer or cytotoxic activity independent to AR antagonistic activity, though the effective concentrations are comparatively higher than that of antiandrogenic activities. It was reported that curcumin and its derivatives have various physiological activities and antiproliferation activity toward prostate cancer by interfering with many transcription pathways.^{30–33} The compounds developed in this study certainly bind to AR ligand-binding domain and exhibit AR antagonistic activity; however, these compounds probably possess another biological activity other than AR modulation like curcumin.

In summary, we have developed a new class of nonsteroidal AR antagonists without the nitrophenyl or cyanophenyl substructure of known AR antagonists. We designed and synthesized a series of 4-phenoxyphenol derivatives as AR antagonist candidates. We found that the 4-(4-benzoylamino-phenoxy)phenol skeleton serves as a novel core structure of AR antagonists. Compound **22** is one of the most potent of these non-nitrophenyl and non-cyanophenyl type nonsteroidal AR antagonists and exhibited antiandrogenic activity toward several prostate cancer cell lines, including two with mutated ARs. This novel non-flutamide derivative type AR antagonist is a promising candidate for antiandrogen therapy of prostate cancer.

■ ASSOCIATED CONTENT

Supporting Information

Analytical data and experimental procedures for synthetic preparation and biological evaluation of compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*(S.F.) E-mail: fujis.chem@tmd.ac.jp.

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the manuscript.

Funding

This study was partially supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology, Japan (Grant No. 21790128 to S.F.).

Notes

The authors declare no competing financial interest.

■ REFERENCES

- (1) Evans, R. M. The steroid and thyroid hormone receptor superfamily. *Science* **1988**, *240*, 889–894.
- (2) McEwan, I. J. Gene regulation through chromatin remodeling by members of the nuclear receptor superfamily. *Biochem. Soc. Trans.* **2000**, *28*, 369–373.
- (3) Mooradian, A. D.; Morley, J. E.; Korenman, S. G. Biological actions of androgens. *Endocr. Rev.* **1987**, *8*, 1–28.
- (4) Bagatell, C. J.; Bremner, W. J. Androgens in men; uses and abuses. *N. Engl. J. Med.* **1996**, *334*, 707–714.
- (5) Taplin, M.-E. Androgen receptor: Role and novel therapeutic prospects in prostate cancer. *Expert Rev. Anticancer Ther.* **2008**, *8*, 1495–1508.
- (6) Ross, R. K.; Pike, M. C.; Coetzee, G. A.; Reinhardt, J. K. V.; Yu, M. C.; Feigelson, H.; Stanczyk, F. Z.; Kolonel, L. N.; Henderson, B. E. Androgen metabolism and prostate cancer: Establishing a model of genetic susceptibility. *Cancer Res.* **1998**, *58*, 4497–4504.
- (7) Scher, H. I.; Steineck, G.; Kelly, W. K. Hormone-refractory (D3) prostate cancer: refining the concept. *Urology* **1995**, *46*, 142–148.
- (8) Miyamoto, H.; Rahman, M. M.; Chang, C. Molecular basis for the antiandrogen withdrawal syndrome. *J. Cell. Biochem.* **2003**, *91*, 3–12.
- (9) Bohl, C. E.; Gao, W.; Miller, D. D.; Bell, C. E.; Dalton, J. T. Structural basis for antagonism and resistance of bicalutamide in prostate cancer. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102*, 6201–6206.
- (10) Steketee, K.; Timmerman, L.; Ziel-van der Made, A. C.; Doesburg, P.; Brinkmann, A. O.; Trapman, J. Broadened ligand responsiveness of androgen receptor mutants obtained by random amino acid substitution of H874 and mutation hot spot T877 in prostate cancer. *Int. J. Cancer* **2002**, *100*, 309–317.
- (11) Gao, W.; Bohl, C. E.; Dalton, J. T. Chemistry and structural biology of androgen receptor. *Chem. Rev.* **2005**, *105*, 3352–3370.
- (12) Jung, M. E.; Ouk, S.; Yoo, D.; Sawyers, C. L.; Chen, C.; Tran, C.; Wongvipat, J. Structure–activity relationship for thiohydantoin androgen receptor antagonists for castration-resistant prostate cancer (CRPC). *J. Med. Chem.* **2010**, *53*, 2779–2796.
- (13) Song, C.-H.; Yang, S. H.; Park, E.; Cho, S. H.; Gong, E.-Y.; Khadka, D. B.; Cho, W.-J.; Lee, K. Structure-based virtual screening and identification of a novel androgen receptor antagonist. *J. Biol. Chem.* **2012**, *287*, 30769–30780.
- (14) Yang, S. H.; Song, C.-H.; Van, H. T. M.; Park, E.; Khadka, D. B.; Gong, E.-Y.; Lee, K.; Cho, W.-J. SAR based design of nicotinamides as a novel class of androgen receptor antagonists for prostate cancer. *J. Med. Chem.* **2013**, *56*, 3414–3418.
- (15) Ohtsu, H.; Xiao, Z. Y.; Ishida, J.; Nagai, M.; Wang, H. K.; Itokawa, H.; Su, C. Y.; Shih, C.; Chiang, T. Y.; Chang, E.; Lee, Y. F.; Tsai, M. Y.; Chang, C. S.; Lee, K. H. Antitumor agents. 217. Curcumin analogues as novel androgen receptor antagonists with potential as anti-prostate cancer agents. *J. Med. Chem.* **2002**, *45*, 5037–5042.
- (16) Zhou, J.; Geng, G.; Shi, Q.; Sauriol, F.; Wu, J. H. Design and synthesis of androgen receptor antagonists with bulky side chains for overcoming antiandrogen resistance. *J. Med. Chem.* **2009**, *52*, 5546–5550.
- (17) Esatbeyoglu, T.; Huebbe, P.; Ernst, I. M. A.; Chin, D.; Wagner, A. E.; Rimbach, G. Curcumin: From molecule to biological function. *Angew. Chem., Int. Ed.* **2012**, *51*, 5308–5332.
- (18) Kagechika, H.; Kawachi, E.; Hashimoto, Y.; Himi, T.; Shudo, K. Retinobenzoic acids. 1. Structure–activity relationships of aromatic amides with retinoidal activity. *J. Med. Chem.* **1988**, *31*, 2182–2192.
- (19) Kagechika, H.; Shudo, T. Synthetic retinoids: Recent developments concerning structure and clinical utility. *J. Med. Chem.* **2005**, *48*, 5875–5883.
- (20) Boehm, M. F.; Fitzgerald, P.; Zou, A.; Elgort, M. G.; Bischoff, E. D.; Mere, L.; Mais, D. E.; Bissonnette, R. P.; Heyman, R. A.; Nadzan, A. M.; Reichman, M.; Allegretto, E. A. Novel nonsecosteroidal vitamin D mimics exert VDR-modulating activities with less calcium

mobilization than 1,25-dihydroxyvitamin D₃. *Chem. Biol.* **1999**, *6*, 265–275.

(21) Mitchell, L. H.; Hub, L.-Y.; Nguyen, M.; Fakhoury, S.; Smith, Y.; Iula, D.; Kostlan, C.; Carroll, M.; Dettling, D.; Du, D.; Pocalyko, D.; Wade, K.; Lefker, B. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 2176–2178.

(22) Takahashi, B.; Ohta, K.; Kawachi, E.; Fukasawa, H.; Hashimoto, Y.; Kagechika, H. Novel retinoid X receptor antagonists: Specific inhibition of retinoid synergism in RXR–RAR heterodimer actions. *J. Med. Chem.* **2002**, *45*, 3327–3330.

(23) Nakamura, M.; Makishima, M.; Hashimoto, Y. Development of silicon-containing bis-phenol derivatives as androgen receptor antagonists: Selectivity switching by C/Si exchange. *Bioorg. Med. Chem.* **2013**, *21*, 1643–1651.

(24) Hiraoka, D.; Nakamura, N.; Nishizawa, Y.; Uchida, N.; Noguchi, S.; Matsumoto, K.; Sato, B. Inhibitory and stimulatory effects of glucocorticoid on androgen-induced growth of murine shionogi carcinoma 115 in vivo and in cell culture. *Cancer Res.* **1987**, *47*, 6560–6564.

(25) One-pot synthesis of this scaffold was also reported: Chen, W.; Li, J.; Fang, D.; Feng, C.; Zhang, C. Copper-catalysed one-pot multicomponent coupling reaction of phenols, amides, and 4-bromophenyl iodide. *Org. Lett.* **2008**, *10*, 4565–4568.

(26) Wilson, V. S.; Bobseine, K.; Lambright, C. R.; Gray, L. E., Jr. A novel cell line, MDA-kb2, that stably expresses an androgen- and glucocorticoid-responsive reporter for the detection of hormone receptor agonists and antagonists. *Toxicol. Sci.* **2002**, *66*, 69–81.

(27) Veldscholte, J.; Berrevoets, C. A.; Brinkmann, A. O.; Grootegeed, J. A.; Mulder, E. Anti-androgens and the mutated androgen receptor LNCaP cells: Differential effects on binding affinity, heat-shock protein interaction, and transcription activation. *Biochemistry* **1992**, *31*, 2393–2399.

(28) Zhou, J.; Liu, B.; Geng, G.; Wu, J. H. Study of the impact of the T877A mutation on ligand induced helix-12 positioning of the androgen receptor resulted in design and synthesis of novel antiandrogens. *Proteins* **2010**, *78*, 623–637.

(29) Sramkoski, R. M.; Pretlow, T. G., II; Giaconia, J. M.; Pretlow, T. P.; Schwartz, S.; Sy, M.-S.; Marengo, S. R.; Rhim, J. S.; Zhang, D.; Jacobberger, J. W. A new human prostate carcinoma cell line, 22Rv1. *In Vitro Cell. Dev. Biol.-Anim.* **1999**, *35*, 403–409.

(30) Aggarwal, B. B. Prostate cancer and curcumin. *Prostate* **2001**, *47*, 293–303.

(31) Singh, R. P.; Agarwal, R. Mechanism of action of novel agents for prostate cancer chemoprevention. *Endocr.-Relat. Cancer* **2006**, *13*, 751–778.

(32) Fuchs, J. R.; Pandit, B.; Bhasin, D.; Etter, J. P.; Regan, N.; Abdelhamid, D.; Li, C.; Lin, J.; Li, P.-K. Structure–activity relationship studies of curcumin analogues. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 2065–2069.

(33) Shi, Q.; Wada, K.; Ohkoshi, E.; Lin, L.; Huang, R.; Morris-Natschke, S. L.; Goto, M.; Lee, K.-H. Antitumor agents 290. Design, synthesis and biological evaluation of a new LNCaP and PC-3 cytotoxic curcumin analogs conjugated with anti-androgens. *Bioorg. Med. Chem.* **2012**, *20*, 4020–4031.