

View Article Online View Journal

RSC Advances

This article can be cited before page numbers have been issued, to do this please use: V. G. Jadhav, J. M. Bhojane and J. M. Nagarkar, *RSC Adv.*, 2014, DOI: 10.1039/C4RA12827C.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This Accepted Manuscript will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/advances

RSC Advances

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxx

ARTICLE TYPE

Palladium on Manganese ferrite: an efficient catalyst for one pot synthesis of primary amides from iodobenzene.

Vilas Gangadhar Jadhav^a, Jeevan Manohar Bhojane^a and Jayashree Milind Nagarkar^{*a}

Received (in XXX, XXX) Xth XXXXXXXX 20XX, Accepted Xth XXXXXXXX 20XX 5 DOI: 10.1039/b000000x

Amidation of aryl halide in one pot is reported using Pd-MnFe₂O₄ as a catalyst. The catalyst was characterized by various techniques such as XRD, FEG-SEM, EDS, TEM, BET surface area and ICP AES. K₄ [Fe(CN)₆]is used as non toxic cyanation reagent for in situ generation of benzonitrile and hydrolyzed as soon as it formed. The catalyst was found to be efficient and can be used for several cycles ¹⁰ without loss in activity. Moderate to good yield of primary amides was obtained.

1. Introduction.

Published on 12 December 2014. Downloaded by University of Virginia on 14/12/2014 12:50:11.

Primary amides are the most important functionality in natural products, pharmaceuticals, agrochemicals and biologically active molecules¹⁻³. Generally, primary amides are 15 synthesized either by reaction of carboxylic acid or its derivative with ammonia or its equivalent⁴. They are also synthesized by different methodologies such as catalytic rearrangement of aldoximes⁶, reaction of aldehydes with ammonia⁷, oxidation of benzylic alcohols in presence of ammonia⁸ and catalytic 20 oxidation of benzyl amines9. Aminocarbonylation of aryl halide by using carbon monoxide and ammonia or ammonia equivalent is also reported as the efficient method for the synthesis of primary amides. However, this process involves handling of toxic gas and reduces its use at kilogram level ⁵. Aryl halides with

- ²⁵ ammonia in presence of metal carbonyls also have been used for the synthesis of primary amides, but it leads to release of toxic carbon monoxide gas and also the metal carbonyl complexes are volatile and unstable¹⁰. Hydration of nitriles is an atom economical process and numbers of catalytic methods have been
- ³⁰ developed for nitrile synthesis¹¹. Hence preparation of nitriles as an intermediate for amide synthesis is of great importance. Nitriles are prepared by reaction of aryl halides with various nitrile sources such as alkali cyanide, CuCN, TMSCN and K₄[Fe(CN)₆] salt in presence of transition metal catalyst. Among

³⁵ these K₄[Fe(CN)₆] is used as a less toxic and cheaper cyanide source for cyanation of aryl halides¹². Recently oxidation of benzyl amine to primary amide by using

OMS-2 catalyst was reported ^{9c}. However, the reaction requires very high pressure (6 bars) and temperature (160°C) and also ⁴⁰ gives side products, where as we have reported general synthesis

of primary amides at mild reaction conditions using magnetically retrievable catalyst.

Magnetically separable catalyst has attracted much attention in ⁴⁵ the area of heterogeneous catalyst as it is easily separated by external magnet and can be used several times without losing its activity¹³. Super paramagnetic materials, spinnel ferrites MFe₂O₄ (M= Ni, Co, Zn,) are one of the important advance materials used in drug delivery and biomedical applications such as MRI¹⁴. ⁵⁰ Besides this, they are also found to be good support for transition metal catalysts¹⁵. In particular, palladium supported on super paramagnetic material is the most important heterogeneous catalyst as it offers high surface area and easy separation. In continuation of our work with metal nano particles supported on ⁵⁵ super paramagnetic material^{15b}, herein we report palladium nanoparticles (PdNPs) supported on MnFe₂O₄ and its application in one pot synthesis of primary amides



Fig.1 General synthesis of primary amide.

Manganese ferrite (MnFe₂O₄) nano crystals possess higher ⁶⁰ magnetization as compared to magnetite, CoFe₂O₄ and NiFe₂O₄ nanoparticles¹⁶. Here we have carried out the tandem synthesis of primary amide from iodobenzene via in situ generation of benzonitrile in presence of DMSO: H₂O solvent system using Pd nano particles supported on MnFe₂O₄. This is the first report of ⁶⁵ the application of the heterogeneous catalyst for the synthesis of primary amide and that too using nontoxic nitrile source

```
This journal is © The Royal Society of Chemistry [year]
```

 $K_4[Fe(CN)_6]$. Very good yields of the desired products were obtained.

2. Experimental

5 2.1 Catalyst preparation.

Catalyst was prepared by using ultrasound assisted coprecipitation method. FeCl₃. $6H_2O$ (9.5 mmol, 50 mL) and MnCl₂.4H₂O (4.4 mmol, 50 mL) solutions were added in 250 mL round bottom flask followed by addition of 0.5 g KCl and PdCl₂

¹⁰ (60 mg). This mixture was placed in ultrasound bath for 5 min. Then pH of this mixture was adjusted at 12 by adding 3N NaOH followed by heating at 60 °C for 30 min. The obtained catalyst was filtered and washed with distilled water several times and then by ethanol to remove water. The catalyst was dried under ¹⁵ vacuum at 60 °C for 24 h, and subsequently at 200°C for 4h.

2.2 Characterization of material

The as synthesized catalyst was characterized by various techniques such as X-ray diffraction (XRD), Field emission gun Scanning electron microscopy(FEG-SEM),Transmission electron ²⁰ microscopy (TEM), FTIR, ICP-AES analysis, BET surface area

The XRD analysis was performed on Shimadzu XRD 2400 instrument using Cu K α radiation (λ = 1.5406 A°) with scanning rate 2 degree per minute. TEM analysis was performed over PHILIPS 2200 instrument. FEG-SEM analysis was performed on ²⁵ TESCAN MIRA Instrument. The energy dispersive X-ray spectral analysis (EDS) image was recorded with an Oxford instrument at 10 kV. Beam intensity was kept high to get good response by the detector. Inductively coupled plasma Atomic Absorption spectrometry (ICP AES) analysis was performed on ³⁰ ARCOS from M/s. Spectro, Germany. GC analysis is performed over PerkinElmer Clarus 480 instrument. GC-MS Spectra recorded over Shimadzu QP-2010 Instrument. 1H and 13C NMR spectra's were recorded on Agilent 400MHz and 100MHz instrument respectively.



Fig.2 a) XRD and b) EDS analysis of Pd-MnFe₂O₄

3. Results and Discussion

XRD analysis showed peak at $2\theta = 30.1$, 35.14, 43.41, 54.601, 40 57.0671 and 63.6651 which represent the Bragg reflections from the (220), (311), (400), (422), (511) and (440) planes respectively. The crystalline data matches with the JCPDS Card No. (74-2403). Palladium peaks at 40, 46 θ values were not observed because of the low loading of the palladium (Fig 2a). 45 Displacing value from XRD and TEM SEAD pattern is same and

found to be 2A°. (Fig. 3 c and d)

The elemental analysis of Pd-MnFe₂O₄ was confirmed by EDS analysis as shown in (fig.2 b). The distribution of elements in Pd-⁵⁰ MnFe₂O₄ was Mn =11.45%, Fe = 28.15%, O =59.31% and Pd=1.09%.



Fig.3 FEG-SEM image a) fresh and b) reused catalyst after fifth cycle. c) TEM image of catalyst and d) SEAD pattern.

- 55 FEG-SEM analysis shows highly crystalline morphology of the catalyst. (Fig. 3 c)
- The TEM image of Pd-MnFe₂O₄ NPs. (Fig.3 c) clearly indicates that NPs are uniform in size with average particle size less than 45 nm. This is in accordance with the average crystallite size
- ⁶⁰ obtained by using the Debye–Scherrer equation. The image also exhibits that the Pd NPs on MnFe₂O₄ are well dispersed. Pd NPs size ranges from 2-10 nm.
- From the N_2 adsorption-desorption isotherm of Pd-MnFe₂O₄ NPs, the BET surface area of the particles was found to be 65 24.9346m2/g and the adsorption average pore width was 315 A°.
- Single point surface area at P/Po = 0.254238207: 24.5190 m²/g (Fig.2 Supporting Information)

Reaction of iodobenzene with K4[Fe(CN)6] was treated as a

35

model reaction and various reaction parameters were optimized for this reaction. Initially (1 mmol) of iodobenzene was treated with 0.25 mmol of $K_4[Fe(CN)_6]$ in the presence of Pd-MnFe₂O₄ catalyst (30 mg, 1mol% Pd), K_2CO_3 as a base in DMF :H₂O s (1:1) solvent system, at 100°C, which offered 58 % yield of honoramide. (Table 1 entry 1) Series of experiments were

- benzamide (Table 1, entry 1) Series of experiments were performed to optimize the various reaction parameters such as catalyst loading, solvent, base and temperature of the reaction.
- Loading of the catalyst was also the important parameter and it ¹⁰ was found that 1 mmol of the catalyst was enough to achieve the maximum yield of the desired product (Table 1 entry 4). Solvent study was the important part for this transformation. Various solvent systems such as DMF: H₂O, DMSO: H₂O and NMP: H₂O were used out of which



20 Scheme 1. One pot synthesis of amide from iodobenzene

. Table1. Optimization of reaction parameters	.Table1.	Optimization	of reaction	parameters
---	----------	---------------------	-------------	------------

			A		
Entry	Pd- MnFe2O4 (mol %)	Base	Solvent: H2O	Temperature °C	Yield ^b %
1	0.5	K ₂ CO ₃	DMF:H ₂ O(1:1)	100	58
2	0.5	K_2CO_3	$NMP:H_2O(1:1)$	100	64
3	0.5	K_2CO_3	$DMSO:H_2O(1:1)$	100	70
4	1	K ₂ CO ₃	$DMSO:H_2O(1:1)$	100	78
5	1	K_2CO_3	$DMSO:H_2O(2:1)$	100	84
6	1	K_2CO_3	$DMSO:H_2O(2:1)$	110	88
7	2	K_2CO_3	$DMSO:H_2O(2:1)$	110	88
8	3	K_2CO_3	$DMSO:H_2O(2:1)$	110	88
9	1	Cs ₂ CO ₃	DMSO: $H_2O(2:1)$	110	74
10	1	TEA	$DMSO:H_2O(2:1)$	110	trace
11	1	DBU	DMSO:H ₂ O(2:1)	110	trace
12	1	KOH	$DMSO:H_2O(2:1)$	110	20
13	1	K_2CO_3	DMSO:H ₂ O(2:1)	90	19
14	1	K ₂ CO ₃	DMSO: $H_2O(2:1)$	120	91
15	1	K_2CO_3	$DMSO:H_2O(2:1)$	140	90
16	1	K_2CO_3	H_2O	110	-
17	1	K_2CO_3	DMSO	110	_*
18	1	K_2CO_3	DMSO:H ₂ O(2:1)	110	90 °
19	1	K_2CO_3	DMSO:H ₂ O(2:1)	110	90 ^d

^aReaction conditions: Aryl halide (1mmole), K_4 [Fe(CN)]₆ (0.25 mmole), Time 12h, base (1.2 mmole), ^bisolated yield., ^c time 18 h. ^e24 h,

25 *benzonitrile 92%

DMSO:H₂O (2:1) was found to be the most suitable solvent system for this transformation(Tablel entry 5). When reaction was carried out in DMSO solvent only, benzonontrile was formed ³⁰ (Table1 entry 17). Iodobenzene was insitu converted to benzonitrile by the cyanating reagent K₄[Fe(CN)₆] which immediately gets hydrated to benzamide in the presence of water.

The reaction was also carried out with various inorganic and ³⁵ organic bases such as TEA, DBU, K_2CO_3 , KOH and Cs_2CO_3 , out of which K_2CO_3 was found to be the most efficient base giving maximum yield of the benzamide (Table 1 entry 8). The reaction was also carried out at various temperatures under optimized

This journal is © The Royal Society of Chemistry [year]

reaction conditions. The product yield increased dramatically 40 from 19% to 90% when temperature increased from 90°C to 110°C whereas beyond 110°C no significant increase in the product yield with increase in temperature was observed (Table 1 entry 13-15)

We also studied the effect of concentration of K₄ [Fe (CN)₆] salt ⁴⁵ on the product yield by increasing the amount from 0.1 mmol to 0.25 mmol. Conversion of aryl halide to benzonitrile was 100% at 0.25 mmol concentration of K₄[Fe(CN)₆]. The reaction was also carried out at various time intervals and 100% conversion was obtained in 18 h. For optimized reaction TON and [‡]TOF for ⁵⁰ amidation of iodobenzene was found to be 90 and 5 h⁻¹.

The applicability of the protocol was examined for various substituted aryl iodides (Table 2). It was found that electron rich substrate gave very good yield of the desired product.

⁵⁵ **Table 2**. One pot synthesis of primary amides from aryl iodides^a.



Page 3 of 5





Fig 4. Recycle study of the catalyst.



Fig.5 Plausible Mechanism of the reaction.

³⁵ Plausible mechanism suggests that iodobenzene is oxidatively added to the catalyst and then in presence of K_4 [Fe(CN)₆] salt gives benzonitrile which subsequently get hydrolyzed in presence of water and catalyst to give primary amide.



40 Fig.6 Magnetic separation of the catalyst a) before and b) after applying external magnet.



^aReaction conditions: Aryl halide (1mmole), K₄[Fe(CN)₆] (0.25 mmole), 12h, base (1.2 mmole), Pd-MnFe₂O₄ (30 mg, 1mol % Pd) ^bisolated yield.

In case of p-NO₂ iodobenzene electron withdrawing effect was ⁵ more prominent givinng moderate yield of the desired products. However, in m-Nitro iodobenzene the yield was increased due to minimum electron withdrawing effect. Results are summarized in (Table 2. entry 6 and 12).

Donating substituents at ortho positions were well tolerated ¹⁰ giving good yield of the corresponding primary amides. We further studied amidation of bromobenzene as well as chlorobenzene and observed that bromobenzene gave only 20% yield of the desired product where as cholrobenzene was unreactive with optimized reaction conditions(Table 2 entry 2 and ¹⁵ 3). Hehteroaryl iodide also gave good yield of corresponding

amide (Table 2 entry 16).

Heterogeneous test for the catalyst

We also examined the heterogeneous nature of the catalyst by hot

²⁰ filtration test. The catalyst was separated from reaction mass after half of the reaction time and the filtrate was stirred for required time under same reaction conditions. The reaction did not

Conclusions

In summary, we have developed the simple and practically applicable protocol for the synthesis of primary amides from aryl iodides. Pd-MnFe2O4 is the first heterogeneous catalyst for such

5 transformation. Catalyst was found to be efficient for the transformation of iodobenzenes to respective amides. $K_4[Fe(CN)_6]$ was used as the non toxic cyanating reagent and in situ generated benzonitrile was hydrated as soon as it forms. The catalyst can be retrieved using external magnet and reused up to 10 five consecutive cycles without much loss in its activity.

Acknowledgements

The authors are very thankful to the UGC-SAP, New Delhi, India for awarding the fellowship.

Notes and references

- 15 ^a Department of Chemistry, Institute of Chemical Technology, Matunga, Mumbai – 400019, India. Fax: +91 22 33611020; Tel.: +91 22 33611111/2222; E-mail: jm.nagarkar@ictmumbai.edu.in
 - † Electronic Supplementary Information (ESI) available:
- † Electronic Supplementary Information (ESI) available: [Experimental 20 procedure, Characterization of the catalyst, Mass and 1H NMR spectrum of representative compounds are given]. See DOI: 10.1039/b000000x/ ‡ TOF = [mol of product /(mol catalyst X h)].
- Arthur. The Amide Linkage: Selected Structural Aspects in 1. G. Chemistry, Biochemistry, and Materials Science; Wiley-Interscience, 25 2000
 - 2. S. Budavari, The Merck Index, Merck, Rahway, USA, 11th edn, 1989
 - 3. V. R. Pattabiraman, J. W. Bode, Nature., 480, 2011,471.
- 4. (a) G. G. Arzoumanidis, F. C. Rauch, J. Org. Chem., 1981 46,
- 3930-3932(b) C. L Allen.;, J. M. Williams, J. Chem. Soc. Rev., 2011, 30 40, 3405-3415; (c) E. Valeur, M. Bradley, Chem. Soc. Rev. 2009, 38, 606-631. (d) H. Lundberg, F. Tinnis, N. Selander and H. Adolfsson, Chem. Soc. Rev., 2014, 43, 2714
- 5. (a) P. G. Alsabeh, M. Stradiotto, H. Neumann, M., Beller, Adv. Synth. Catal., 2012, 354, 3065-3070; (b) X.-F. Wu, H. Neumann, 35 M.Beller, Chem. Asian J., 2010, 5, 2168-2172; (c) Wu, X.-F.; Neumann, H.; Beller, M. Chem. Eur. J., 2010,16, 9750-9753; (d) Wu, X.-F.; Neumann, H.; Beller, M. Chem. Eur. J., 2012, 18,419-422; (e) X.-F. Wu, J. Schranck, H. Neumann, M. Beller, ChemCatChem., 2012,
- 4, 69-71. (f) T. Xu, H. Alper, Tetrahedron Lett., 2013, 54, 5496-5499. 40 (g) S. T. Gadge, , B.M. Bhanage, Synlett., 2014, 25, 85 - 88
- A. Martinez-Asencio, M. Yus, D. J 6. (a) Ramon Tetrahedron., 2012, 68, 3948 - 3951. (b) F. Li, P. Qu, J. Ma; X. Zou, C. Sun, ChemCatChem., 2013, 5, 2178 - 2182. (c) N. A. Owston,.; A.
- J. Parker, J. M. Williams, Org. Lett., 2007, 9, 3599 3601. (d) C. Sun, P. Qu, F. Li, Cat. Sci. and Tech., 2014, 4, 988 - 996
- 7. (a) U. B. Patil, A. S. Singh, J. M. Nagarkar, RSC Advances., 2014, 4, 1102 - 1106. (b) Md. A. Ali, T. Punniyamurthy, Adv. Synth. Catal., 2010, 352, 288-292.
- 50 8. (a) K. Yamaguchi, H. Kobayashi, T. Oishi and N. Mizuno, Angew. Chem., Int. Ed., 2012, 51, 544-547; (b) K. Yamaguchi, H. Kobayashi, Y. Wang, T. Oishi, Y. Ogasawara and N. Mizuno, Catal. Sci. Technol., 2013, 3, 318-327; (c) R. Nie, J. Shi, S. Xia, L. Shen, P. Chen, Z. Hou and F. Xiao, J. Mater. Chem., 2012, 22, 18115-18115; (d) R. Ohmura,
- M. Takahata and H. Togo, Tetrahedron Lett., 2010, 51, 4378-4381; (e) 55 R. Das and D. Chakraborty, Catal. Commun., 2012, 26, 48-53; (f) X. -Q. Li, W. -K. Wang, Y. -X. Han and C. Zhang, Adv. Synth. Catal., 2010. 352. 2588-2598
- 9. (a) X. -F. Wu, C. B. Bheeter, H. Neumann, P. H. Dixneuf and M.
- Beller, Chem. Commun., 2012, 48, 12237-12239; (b) J. W. Kim, K. 60 Yamaguchi and N. Mizuno, Angew. Chem., Int. Ed., 2008, 47, 9249-9251; (c) Y. Wang, H. Kobayashi, K. Yamaguchi and N. Mizuno, Chem. Commun., 2012, 48, 2642-2644
- 10. (a) W. Ren, M. Yamane, J. Org. Chem., 2010, 75, 8410 8415. (b) W. Ren, M. Yamane, J. Org. Chem., 2010, 75, 3017 - 3020.

- 11. (a) J. Mauger, T. Nagasawa, H. Yamada Tetrahedron., 1989, 45, 1347 - 1354 (b) A. Goto, K. Endo, S. Saito, Anewg. Chemie - Int. Ed., 2008, 47, 3607 - 3609. (c) R.S. Ramon;, N. Marion, S. P Nolan, Chem. - A Eur. J., 2009, 15, 8695 - 8697 (d) M. Muranaka,;
- I. Hyodo, , W. Okumura, T. Oshiki, Cat. Today., 2011, 164, 552 -555 (e) E. Bolyog-Nagy, A. Udvardy, F. Joo,; A. Katho, Tetrahedron Lett., 2014, . 55, 3615 - 3617 (f) S. Kumar, P. Das, New J. Chem., 2013, 37, 2987 - 2990. (g) V. Polshettiwar, R.S. Varma, Chem. - A Eur. J., 2009, 15, 1582 - 1586 h) T. Hirano,
- K.Uehara,; K. Kamata, N. Mizuno, J. Am. Chem. Soc., 2012, 134, 6425 – 6433. (i) Nippon Shokubai Co. Ltd Patent: JP2005/170821 A, 2005. (j) Y. Wang,; F. Wang,; Q. Song, Q. Xin,; S. Xu, J.Xu, J. Am. Chem. Soc., 2013, 135, 1506 - 1515.(k) K-T. Liu, M.-H. Shih, H.-W. Huang,; C.-J. Hu, Synthesis., 1988, 9, 715-717.
- 80 12. (a) T. Schareina, A. Zapf, M. Beller, Chem. Commun., 2004, 40, 1388 (b) T. Schareina, R. Jackstell, T. Schulz, A. Zapf, A. Cotte, M. Gotta, M. Beller, Adv. Synth .Catal., 2009, 351, 643 (c) J. Huiliang , J. Jianzhong, W. Huixian, C. Chun, Catal. Lett., 2013 143, 1195-1199.
- 13. (a) W. L. Choon, In.S. Lee Nano Today., 2010 5, 412-434. (b) V. Polshettiwar, R. Luque, A. Fihri, H. Zhu, M. Bouhrara and J-M. 85 Basset Chem. Rev., 2011, 111, 3036-3075. (c) A. H. Latham and M. E. Williams, Acc. Chem. Res., 2008, 41, 411-420. (d) S.A Sarode, J.M. J. M. Nagarkar, Tetrahedron Lett., 2014 Bhojane, doi:10.1016/j.tetlet.2014.11.065.(e) R.S. Shelkar, S. H. Gund, J. M. Nagarkar. RSC Advances., 2014 4 53387-53396. 90
- 14. J. Lu, S.Ma, J. Sun, C. Xia , C. Liu, Z. Wang, X.Zhao, F. Gao, Q. Gong , B. Song , X. Shuai , H. Ai, Z. Gua Biomaterials., 2009, 30, 2919-2928.
- 15. (a)K. K.Senapati, S. Roy, C. Borgohain, P. Phukan J. Mol.Cat. A: Chem., 2012, 352, 128-134. (b) A.S. Singh, U.B. Patil, J.M. Nagarkar, Cat. Comm. 2013 35 11-16. (c) S. R. Borhade, S. B. Waghmode; Ind. J. Chem., 2008, 47, 1549-1554. (d) D. Wang., D. Astruc Molecules., 2014, 19, 4635-4653. (e) N. Baig, R. S. Verma. Chem. Commun., 2013, 49, 752-770. (f) V. Polshettiwar, R. S. Verma. Org. Bimol. Chem., 2009, 7, 37-40. 100
 - 16. (a) U.I. Tromsdorf, N.C. Bigall, M.G. Kaul, O.T. Bruns, M.S. Nikolic, B. Mollwitz Nano Lett., 2007, 7, 2422. (b) J.H. Lee, Y.M. Huh, Y. Jun, J. Seo, J. Jang, HT Song, Nat. Med., 2007,13, 95.

This journal is © The Royal Society of Chemistry [year]