

Cobalt-Catalyzed Cross-Dehydrogenative Coupling Reaction in Water by Visible Light

Cheng-Juan Wu, Jian-Ji Zhong, Qing-Yuan Meng, Tao Lei, Xue-Wang Gao, Chen-Ho Tung, and Li-Zhu Wu*

Key Laboratory of Photochemical Conversion and Optoelectronic Materials, Technical Institute of Physics and Chemistry & University of Chinese Academy of Sciences, the Chinese Academy of Sciences, Beijing 100190, P. R. China

Supporting Information

ABSTRACT: By using catalytic amount of $CoCl_2$ with dmgH (dimethylglyoxime) as ligand to form a photosensitizer in situ, a highly selective, efficient, and environmentally benign visible light $C_{sp3} - H + \begin{cases} H - C_{sp} \\ H - C_{sp2} \\ H - C_{sp3} \end{cases} + \begin{cases} Visible light \\ CoCl_2, dmgH \\ air, water \end{cases}$ ABSTRACT: By using catalytic amount of CoCl₂ with dmgH mediated cross-dehydrogenative coupling (CDC) reaction has been

-C_{sp2} C_{sp3}-

developed in aqueous medium. The desired cross-coupling C-C bonds that involve C_{sp3} with C_{sp}, C_{sp2}, and C_{sp3}, respectively, were achieved exclusively in high yields without formation of any other byproduct.

onstruction of C–C bonds has always been a research focus of organic chemists, and transition-metal-catalyzed activation of C-H bonds is well-known to provide atomeconomic and efficient ways for such transformation.¹ As one of the most powerful C-H activation processes, cross-dehydrogenative-coupling (CDC) reaction² constructs C-C bonds directly from two different C-H bonds under oxidative conditions. The oxidation protocol involves metal catalysts together with oxygen or organic oxidants. This straightforward reaction avoids the prefunctionalization and defunctionalization of substrates, which has been part of traditional synthetic design, and has made considerable progress in the past decades.³ In particular, the visible light⁴ catalyzed CDC reaction⁵ has appeared at the forefront owing to its mild, clean, and environmentally benign characteristics. Various noble metal complexes based on Ir(III),^{5a} Ru(II),^{5b} Pt(II),^{5h} Au(III),^{5f,i} and Pd(II)^{5j} have been employed as photocatalysts to initiate the cross-coupling reactions efficiently. However, these catalysts were expensive and suffered complicated synthesis. Developing inexpensive and easily available metal complexes would be a superior choice and would meet the requirements of green chemistry.

On the other hand, it is noteworthy that most of the crosscoupling reactions have been performed in organic solvents. The use of water as a solvent for organic transformations has attracted considerable attention because it not only offers several "green chemistry" benefits but also accelerates the rate and increases the selectivity of the reaction. To realize this type of transformation in water featured with greenness, nontoxicity, and low cost has yet to be explored. To date, only limited examples utilizing water as the reaction medium have been reported.⁶ Li et al. made use of copper bromide to catalyze a cross-coupling reaction between two sp³ C-H bonds in water.^{6b} Our group³¹ realized an efficient aerobic CDC reaction in water catalyzed by recyclable graphene-supported RuO2 nanoparticles. Despite advances, to the best of our knowledge,

there are no reports about visible light catalyzed CDC reactions performed in water.

As an extension of our continuous research efforts to explore greener and milder CDC reactions, we developed a new system for the activation of C-H bonds mediated by visible light in water in ambient conditions. During this reaction, one cobaloxime complex generated in situ from earth-abundant and cost-effective transition metal cobalt salts⁷ and dmgH (dimethylglyoxime) was used as a photocatalyst instead of traditional light-harnessing photocatalysts. Good to excellent yields of the desired cross-coupling products of N-phenyl-1,2,3,4-tetrahydroisoquinolines and indoles were obtained in aqueous solution under visible light irradiation (Scheme 1).





Noticeably, the developed simple system herein could compete with photocatalyzed CDC reactions based on traditional noble metal complexes in yield as well as selectivity. On the other hand, compared with traditional thermal CDC reactions that were carried out in organic solvents, the established system still maintained the efficiency in water, which satisfied the need of modern organic chemistry.

Our preliminary studies focused on the cross-coupling of Nphenyl-1,2,3,4-tetrahydroisoquinoline (1a) with indole (2a) under the irradiation of blue LEDs (λ_{max} = 450 nm) at room temperature. To our delight, when the mixture containing

Received: December 29, 2014

catlytic amount of $CoCl_2$ with dmgH (dimethylglyoxime) as ligand and the substrates in CH_3CN was irradiated under visible light, the formation of the desired cross-coupling product was observed in 40% yield (Table 1, entry 1). To

Table 1. Optimization of the Reaction Conditions^a

$ \begin{array}{c} & & \\ & & $				
	1a	2a	3а	
entry	CoCl ₂ (mol %)	dmgH (mol %)	solvent	yield ^b (%)
1	16	32	CH ₃ CN	40
2	16	32	DMF	none
3	16	32	CH_2Cl_2	trace
4	16	32	CH ₃ OH	43
5 ^c	16	32	H_2O/CH_3CN	58
6^d	16	32	H_2O/CH_3CN	66
7	16	32	H_2O	83
8	8	16	H ₂ O	84
9	8	8	H ₂ O	73
10	8	24	H ₂ O	75
11^e	8	16	H ₂ O	76
12^{f}	8	16	H ₂ O	95
13^g	8	16	H ₂ O	96

^{*a*}Reaction conditions: 0.1 mmol of **1a**, 0.3 mmol of **2a**, corresponding CoCl₂, and dmgH were added in 1 mL of solvent, the solution was irradiated under blue LEDs ($\lambda = 450$ nm) in air for 24 h. ^{*b*}Yields determined by ¹H NMR using diphenylacetonitrile as an internal standard. ^{*c*}Volume ratio of H₂O/CH₃CN was 1:1. ^{*d*}Volume ratio of H₂O/CH₃CN was 1:0.5. ^{*c*}0.2 mmol of **2a** was used. ^{*f*}0.4 mmol of **2a** was used.

enhance the reaction efficiency, we tried to optimize the reaction conditions. When the reaction was conducted in DMF or CH₂Cl₂, negligible amount of the desired cross-coupling product could be observed (Table 1, entries 2 and 3). Compared with the reaction in CH₃CN, a similar yield of 43% was obtained in CH₂OH (Table 1, entry 4). Furthermore, a different performance was shown when the reaction was performed in different volume ratio of H₂O/CH₃CN mixture. The increased ratio of H₂O/CH₃CN resulted in an improved yield of the cross-coupling product (Table 1, entries 5-6). This finding suggested the possibility of H₂O as the green solvent. Indeed, when the reaction was performed in H₂O, the yield achieved 83% (Table 1, entry 7). Afterward, we found that 8 mol % of CoCl₂ and 16 mol % of dmgH were the optimal amounts (Table 1, entries 8-10). Increasing the amount of nucleophilic indole to 4 equiv resulted in the best yield of 95% in the reaction (Table 1, entries 8 and 11-13). As a result, the visible light mediated CDC reaction was able to efficiently proceed in the optimized conditions, i.e., 8 mol % of CoCl₂ with 16 mol % of dmgH in H2O, irradiated by blue LEDs for 24 h in aerobic conditions at room temperature. Control experiments indicated that each component of the reaction is critical to the reaction efficiency (Supporting Information, Table S1, entries 1-4). Significantly, during the process, only the desired cross-coupling product could be generated, and no byproduct was detected.

Molecular oxygen played an important role in this system. None of the desired cross-coupling product was observed when the reaction was carried out in N_2 atmosphere (Supporting

Information, Table S1, entry 4). Moreover, the yield would achieve 95% if the reaction underwent under oxygen atmosphere (Table S1, entry 5). In order to determine the function of the molecular oxygen in the process, we studied the active species of oxygen in the system by electron-spin resonance $(ESR)^8$ spectroscopy as shown in Figure 1. 2,2,6,6-

a) provided the second of the

D) Whyther when the propher was a start of the second of the second s



Figure 1. ESR measurements of (a) a solution in H_2O/CH_3CN of $CoCl_2$, dmgH without 1a in the presence of TEMP under the irradiation of blue LEDs; (b) a solution in H_2O/CH_3CN of $CoCl_2$, dmgH with 1a in the presence of TEMP under the irradiation of blue LEDs; (c) a solution in H_2O/CH_3CN of $CoCl_2$, dmgH without 1a in the presence of DMPO under the irradiation of blue LEDs; (d) a solution in H_2O/CH_3CN of $CoCl_2$, dmgH with 1a in the presence of DMPO under the irradiation of blue LEDs; (d) a solution in H_2O/CH_3CN of $CoCl_2$, dmgH with 1a in the presence of DMPO under the irradiation of blue LEDs; (d) a solution in H_2O/CH_3CN of $CoCl_2$, dmgH with 1a in the presence of DMPO under the irradiation of blue LEDs.

Tetramethylpiperidine (TEMP) was employed as a probe to trap ${}^{1}O_{2}$, and 5,5-dimethyl-1-pyrroline *N*-oxide (DMPO) was used to capture $O_{2}^{\bullet-}$. No any signal was detected when the mixture containing TEMP, CoCl₂, and dmgH was irradiated by blue LEDs, even if *N*-phenyl-1,2,3,4-tetrahydroisoquinoline (1a) was present. There was no signal when DMPO was added into the H₂O/CH₃CN solution containing CoCl₂ and dmgH. However, a characteristic signal of $O_{2}^{\bullet-}$ captured by DMPO was clearly observed after addition of 1a. This finding indicated that the addition of 1a resulted in photoinduced electron transfer to generate $O_{2}^{\bullet-}$, which was a crucial intermediate in the reaction.

To understand the interaction, UV-vis absorption spectra of the subcomponents and their mixture were monitored (Figure 2). After the mixture was stirred for 2 h in air, the absorption spectra of the mixture of CoCl₂ and dmgH in H₂O/CH₃CN became similar to that of $Co^{III}(dmgH)_2Cl_2$, and the highresolution mass spectra have also proved the generation of Co^{III}(dmgH)₂Cl₂ (Figure S1, Supporting Information). Based on above results, we speculated that Co^{III}(dmgH)₂Cl₂ generated in situ was the catalyst in essence. To verify the speculation, Co^{III}(dmgH)₂Cl₂ was used instead of CoCl₂ and dmgH in the reaction under the same condition and a yield of 98% could be achieved after 24 h (Supporting Information, Table S1, entry 6). In addition, CoCl₂ was found to be unable to interact with dmgH to form Co^{III}(dmgH)₂Cl₂ in CH₂Cl₂ and DMF (Figure S2, Supporting Information), leading to poor reaction performance (Table 1, entries 2 and 3).

On the basis of the above results, we proposed a plausible mechanism shown in Scheme 2. $CoCl_2$ together with dmgH in

Organic Letters



Figure 2. UV–vis absorption spectra of system: (a) $CoCl_2$ (1.0×10^{-4} M) in H₂O/CH₃CN 1:1; (b) dmgH (2.0×10^{-4} M) in H₂O/CH₃CN 1:1; (c) mixture of CoCl₂ (1.0×10^{-4} M) and dmgH (2.0×10^{-4} M) in H₂O/CH₃CN 1:1; (d) mixture of CoCl₂ (1.0×10^{-4} M) and dmgH (2.0×10^{-4} M) in H₂O/CH₃CN 1:1 after irradiation under blue LEDs in air for 2 h; (e) mixture of CoCl₂ (1.0×10^{-4} M) and dmgH (2.0×10^{-4} M) in H₂O/CH₃CN 1:1 after irradiation under blue LEDs in air for 2 h; (f) Co^{III}(dmgH)₂Cl₂ (1.0×10^{-4} M) in H₂O/CH₃CN 1:1 after irradiation under blue LEDs in air for 24 h; (f) Co^{III}(dmgH)₂Cl₂ (1.0×10^{-4} M) in H₂O/CH₃CN 1:1.





air conditions resulted in the formation of $Co^{III}(dmgH)_2Cl_2$. Upon irradiation by visible light, $Co^{III}(dmgH)_2Cl_2$ reached its exicited state and then abstracted one electron from **1a** to generate **1a**^{•+} and Co (II) intermediate.⁹ Subsequently, Co(II) intermediate reacted with O₂ to produce the superoxide radical anion (O₂^{•-}) and regenerate Co^{III}(dmgH)₂Cl₂. O₂^{•-} abstracted one proton from **1a**^{•+} to create the HOO[•] radical, which continued to capture a hydrogen atom to form H₂O₂.¹⁰ The generated **1a**[•] further lose one electron to afford iminium ion **2**, followed by nucleophile addition to give rise to the desired product **3a**.

With the understanding of the reaction mechanism, we attempted to investigate the universality of the visible light mediated cobalt-catalyzed CDC reaction in water. First, the scope with respect to tetrahydroisoquinolines was examined (Scheme 3, 3a-g,r). For both electron-withdrawing and electron-donating groups, the desired coupling products could be obtained in good to excellent yields. When tetrahydroisoquinoline was linked with electron-donating groups, the yield of the product decreased slightly which may be due to the electronic effect. The strong electron-withdrawing nitro group would reduce the yield greatly (Scheme 3, 3r). Meanwhile, it



"Reaction conditions: 0.1 mmol of 1, 0.4 mmol of 2, 0.008 mmol of CoCl₂, and 0.016 mmol of dmgH were added in 1 mL of water; the solution was irradiated under blue LEDs ($\lambda = 450$ nm) in air for 24 h. ^bYields determined by ¹H NMR using diphenylacetonitrile as an internal standard; isolated yields are given in parentheses.

can be seen that steric hindrance has no effect on the outcome of this transformation. Furthermore, good to excellent yields were also obtained by using a variety of substituted nucleophilic indoles (Scheme 3, 3h-q). Note that indoles containing electron-donating groups proceeded better than those with electron-withdrawing groups which can be explained by the fact that the electron-donating substituents could enhance the nucleophilicity of the indoles. Significantly, the desired products with halogen substituent on the rings were potential intermediates for further functionalization. Other nucleophilic substrates such as sp^3 C–H of nitroalkane and dimethyl malonate even sp C–H of phenylacetylene are also effective for such transformation (Scheme S1, Supporting Information).

In summary, we have developed a highly selective, efficient, and environmentally benign method for the visible light

Organic Letters

mediated cobalt-catalyzed CDC reaction in aqueous medium. By using earth-abundant and simple cobalt salt with dmgH as a ligand to form a photosensitizer in situ, good to excellent yields of the desired cross-coupling products could be achieved in aerobic conditions at room temperature under visible light irradiation. The catalytic process has no other byproduct showing exclusive selectivity for the cross-coupling C–C bonds formation of sp³ C–H with sp³ C–H, sp² C–H, and sp C–H. Moreover, the reaction is performed in water which is environmentally benign. Such an approach enriches low cost metal catalysis and represents a lower cost, milder, and greener process for organic transformation.

ASSOCIATED CONTENT

Supporting Information

Experimental details and characterization of all compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

*E-mail: lzwu@mail.ipc.ac.cn.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Financial support for this research from the Ministry of Science and Technology of China (2013CB834804, 2014CB239402 and 2013CB834505), the National Natural Science Foundation of China (21390404, 91427303 and 21402217), the Key Research Programme of the Chinese Academy of Sciences (KGZD-EW-T05), and the Chinese Academy of Sciences is gratefully acknowledged.

REFERENCES

(1) (a) Arndtsen, B. A.; Bergman, R. G.; Mobley, T. A.; Peterson, T. H. Acc. Chem. Res. 1995, 28, 154. (b) Ritleng, V.; Sirlin, C.; Pfeffer, M. Chem. Rev. 2002, 102, 1731. (c) Liu, C.; Zhang, H.; Shi, W.; Lei, A. Chem. Rev. 2011, 111, 1780. (d) Sun, C.-L.; Li, B.-J.; Shi, Z.-J. Chem. Rev. 2011, 111, 1293. (e) Ackermann, L. Acc. Chem. Res. 2013, 47, 281. (f) Tang, X.; Huang, L.; Xu, Y.; Yang, J.; Wu, W.; Jiang, H. Angew. Chem., Int. Ed. 2014, 53, 4010. (g) Yamamoto, Y. Chem. Soc. Rev. 2014, 43, 1575.

(2) (a) Li, C.-J. Acc. Chem. Res. 2009, 42, 335. (b) Yeung, C. S.; Dong, V. M. Chem. Rev. 2011, 111, 1215. (c) Girard, S. A.; Knauber, T.; Li, C.-J. Angew. Chem., Int. Ed. 2014, 53, 74.

(3) (a) Murahashi, S.-I.; Komiya, N.; Terai, H.; Nakae, T. J. Am. Chem. Soc. 2003, 125, 15312. (b) Li, Z.; Li, C.-J. Org. Lett. 2004, 6, 4997. (c) Li, Z.; Li, C.-J. J. Am. Chem. Soc. 2004, 126, 11810. (d) Li, Z.; Li, C.-J. J. Am. Chem. Soc. 2005, 127, 3672. (e) Li, Z.; Li, C.-J. J. Am. Chem. Soc. 2005, 127, 6968. (f) Catino, A. J.; Nichols, J. M.; Nettles, B. J.; Doyle, M. P. J. Am. Chem. Soc. 2006, 128, 5648. (g) Li, Z.; Cao, L.; Li, C.-J. Angew. Chem., Int. Ed. 2007, 46, 6505. (h) Perry, A.; Taylor, R. J. K. Chem. Commun. 2009, 22, 3249. (i) Liu, P.; Zhou, C.-Y.; Xiang, S.; Che, C.-M. Chem. Commun. 2010, 46, 2739. (j) Wang, M.-Z.; Zhou, C.-Y.; Wong, M.-K.; Che, C.-M. Chem.-Eur. J. 2010, 16, 5723. (k) Boess, E.; Schmitz, C.; Klussmann, M. J. Am. Chem. Soc. 2012, 134, 5317. (l) Meng, Q. Y.; Liu, Q.; Zhong, J. J.; Zhang, H. H.; Li, Z. J.; Chen, B.; Tung, C. H.; Wu, L. Z. Org. Lett. 2012, 14, 5992. (m) Ratnikov, M. O.; Doyle, M. P. J. Am. Chem. Soc. 2013, 135, 1549. (n) Ratnikov, M. O.; Xu, X.; Doyle, M. P. J. Am. Chem. Soc. 2013, 135, 9475. (o) Cheng, Y.; Dong, W.; Wang, L.; Parthasarathy, K.; Bolm, C. Org. Lett. 2014, 16, 2000. (p) Wang, F.-F.; Luo, C.-P.; Deng, G.; Yang, L. Green Chem. 2014, 16, 2428. (q) Leskinen, M. V.; Madarász, Á.; Yip,

K.-T.; Vuorinen, A.; Pápai, I.; Neuvonen, A. J.; Pihko, P. M. J. Am. Chem. Soc. 2014, 136, 6453.

(4) (a) Narayanam, J. M. R.; Stephenson, C. R. J. Chem. Soc. Rev. 2011, 40, 102. (b) Prier, C. K.; Rankic, D. A.; Macmillan, D. W. Chem. Rev. 2013, 113, 5322. (c) Schultz, D. M.; Yoon, T. P. Science 2014, 343, 985.

(5) (a) Condie, A. G.; Gonzalez-Gomez, J. C.; Stephenson, C. R. J. J. Am. Chem. Soc. 2010, 132, 1464. (b) Rueping, M.; Vila, C.; Koenigs, R. M.; Poscharny, K.; Fabry, D. C. Chem. Commun. 2011, 47, 2360. (c) Hari, D. P.; König, B. Org. Lett. 2011, 13, 3852. (d) Pan, Y.; Kee, C. W.; Chen, L.; Tan, C.-H. Green Chem. 2011, 13, 2682. (e) Möhlmann, L.; Baar, M.; Rieß, J.; Antonietti, M.; Wang, X.; Blechert, S. Adv. Synth. Catal. 2012, 354, 1909. (f) To, W.-P.; Tong, G. S.-M.; Lu, W.; Ma, C.; Liu, J.; Chow, A. L.-F.; Che, C.-M. Angew. Chem., Int. Ed. 2012, 51, 2654. (g) Liu, Q.; Li, Y.-N.; Zhang, H.-H.; Chen, B.; Tung, C.-H.; Wu, L.-Z. Chem.—Eur. J. 2012, 18, 620. (h) Zhong, J.-J.; Meng, Q.-Y.; Wang, G.-X.; Liu, Q.; Chen, B.; Feng, K.; Tung, C.-H.; Wu, L.-Z. Chem.—Eur. J. 2013, 19, 6443. (i) Xue, Q.; Xie, J.; Jin, H.; Cheng, Y.; Zhu, C. Org. Biomol. Chem. 2013, 11, 1606. (j) To, W.-P.; Liu, Y.; Lau, T.-C.; Che, C.-M. Chem.—Eur. J. 2013, 19, 5654.

(6) (a) Zhang, H.-B.; Liu, L.; Chen, Y.-J.; Wang, D.; Li, C.-J. Eur. J. Org. Chem. 2006, 4, 869. (b) Basle, O.; Li, C.-J. Green Chem. 2007, 9, 1047. (c) Nishikata, T.; Lipshutz, B. H. Org. Lett. 2010, 12, 1972. (d) Alagiri, K.; Kumara, G. S. R.; Prabhu, K. R. Chem. Commun. 2011, 47, 11787. (e) Willis, N. J.; Smith, J. M. RSC Adv. 2014, 4, 11059.

(7) (a) Ikeda, Y.; Nakamura, T.; Yorimitsu, H.; Oshima, K. J. Am. Chem. Soc. 2002, 124, 6514. (b) Gandon, V.; Aubert, C.; Malacria, M. Chem. Commun. 2006, 2209. (c) Teo, Y. C.; Chua, G. L. Chem.—Eur. J. 2009, 15, 3072. (d) Cahiez, G.; Moyeux, A. Chem. Rev. 2010, 110, 1435. (e) Toma, G.; Yamaguchi, R. Eur. J. Org. Chem. 2010, 6404. (f) Gao, K.; Yoshikai, N. Acc. Chem. Res. 2014, 47, 1208.

(8) (a) Lion, Y.; Delmelle, M.; Van De Vorst, A. Nature 1976, 263, 442. (b) Ma, J.; Zhao, J.; Jiang, L. Photochem. Photobiol. 2001, 74, 143.
(c) Ou, Z.-Z.; Chen, J.-R.; Wang, X.-S.; Zhang, B.-W.; Cao, Y. New J. Chem. 2002, 26, 1130.

(9) (a) Weiss, M. E.; Kreis, L. M.; Lauber, A.; Carreira, E. M. Angew. Chem., Int. Ed. 2011, 50, 11125. (b) Kreis, L. M.; Krautwald, S.; Pfeiffer, N.; Martin, R. E.; Carreira, E. M. Org. Lett. 2013, 15, 1634.

(10) The generation of H_2O_2 was evidenced by the iodometric titration experiments. For the detailed procedures, see: Mair, R. D.; Graupner, A. J. Anal. Chem. **1964**, 36, 194.