



Contents lists available at ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl

New salicylamide and sulfonamide derivatives of quinoxaline 1,4-di-N-oxide with antileishmanial and antimalarial activities

Carlos Barea^a, Adriana Pabón^b, Denis Castillo^c, Mirko Zimic^d, Miguel Quiliano^d, Silvia Galiano^a, Silvia Pérez-Silanes^a, Antonio Monge^a, Eric Deharo^{e,f}, Ignacio Aldana^{a,*}

^aUnidad en Investigación y Desarrollo de Medicamentos, Centro de Investigación en Farmacobiología Aplicada (CIFA), University of Navarra, Pamplona, Spain

^bGrupo Malaria, Universidad de Antioquia, Medellín, Colombia

^cUnidad de Parasitología Celular, Laboratorios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Lima, Peru

^dBioinformatics Unit – Drug Design Group, Laboratorios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Lima, Peru

^eUniversité de Toulouse; UPS; UMR 152 (PHARMA-DEV, Pharmacochimie et Pharmacologie pour le Développement), 118, rte de Narbonne, F-31062 Toulouse cedex 9, France

^fInstitut de Recherche pour le Développement; UMR-152; Mission IRD Casilla 18-1209 Lima, Peru

ARTICLE INFO

Article history:

Received 12 May 2011

Revised 30 May 2011

Accepted 31 May 2011

Available online 12 June 2011

Keywords:

Quinoxaline

N-Oxides

Malaria

Leishmaniasis

ABSTRACT

Continuing with our efforts to identify new active compounds against malaria and leishmaniasis, 14 new 3-amino-1,4-di-N-oxide quinoxaline-2-carbonitrile derivatives were synthesized and evaluated for their in vitro antimalarial and antileishmanial activity against *Plasmodium falciparum* Colombian FCR-3 strain and *Leishmania amazonensis* strain MHOM/BR/76/LTB-012A. Further computational studies were carried out in order to analyze graphic SAR and ADME properties. The results obtained indicate that compounds with one halogenous group substituted in position 6 and 7 provide an efficient approach for further development of antimalarial and antileishmanial agents. In addition, interesting ADME properties were found.

© 2011 Elsevier Ltd. All rights reserved.


Malaria is by far the world's most important tropical parasitic disease and one of the oldest diseases known to mankind. For nearly half a century, chloroquine (CQ) has been the primary therapy of choice. However, CQ-resistant *Plasmodium falciparum* is now observed in nearly all of the malaria-endemic regions and causes the most deadly form of malaria.¹ Mortality, currently estimated at over a million people per year, has risen in recent years, probably due to increasing resistance. Therefore, it is necessary to develop cheaper and more effective drugs against the parasite.^{2,3} Leishmaniasis is generally recognized for its cutaneous form which causes non-fatal disfiguring lesions, although epidemics of the potentially fatal visceral form cause thousands of deaths. Today, this disease threatens about 350 million people, and 12 million people are believed to be currently infected, with about 1–2 million estimated new cases occurring every year. Most available drugs against Leishmaniasis are costly, require long treatment regimes and are becoming more and more ineffective, requiring the discovery of new drugs.⁴ Moreover, since the introduction of miltefosine at the beginning of this century, no new antileishmanial compounds have been approved for human treatment.⁵

Many classes of organic compounds have been tested, with special attention being paid to nitrogen heterocycles, five- and six-membered rings. Quinoxalines display diverse pharmacological activities as antibacterial, antiviral, anticancer and antiparasitic agents and more specifically, their 1,4-di-N-oxides, which increase the biological properties enormously.⁶ Our research project led to the publication of several papers, in which both synthesis and biological activity assessments have been described for a large number of quinoxaline and quinoxaline 1,4-di-N-oxide derivatives with a variety of substituents in positions 2, 3, 6 and 7 of the quinoxaline ring.^{7–10} In an attempt to exalt antiparasitic activity of quinoxaline derivatives, our group has synthesized different series with promising results, one of them being the introduction of a carbonitrile group in position 2, which increases the antiparasitic activity, and an amine group in position 3, with the aim of linking together new molecules with interesting activities. We synthesized and evaluated in vitro 14 new 3-amino-1,4-di-N-oxide quinoxaline-2-carbonitrile analogues against *P. falciparum* Colombian FCR-3 strain and *Leishmania amazonensis* MHOM/BR/76/LTB-012A strain. The ADME properties of all molecules were calculated in silico.


We have prepared 14 new quinoxaline 1,4-di-N-oxide derivatives. The benzofuroxane starting compounds (BFX, **I**, Schemes 1 and 2) have been prepared using previously described methods.^{11,12} The 3-amino-1,4-di-N-oxide quinoxaline-2-carbonitrile derivatives (cyanoamines, **II**) were obtained by the Beirut reaction from the

* Corresponding author. Tel.: +34 948 425653; fax: +34 948 425652.

E-mail address: ialdana@unav.es (I. Aldana).



Scheme 1. General synthesis of sulfonamides. Reagents and conditions: (i) malononitrile, DMF, triethylamine, 15–90%; (ii) pyridine, 0° C, 13–15%.



Scheme 2. General synthesis of acetoxybenzamides. Reagents and conditions: (i) malononitrile, DMF, triethylamine, 15–90%; (ii) dry THF, 25%.

corresponding BFX with malononitrile using *N,N*-dimethylformamide (DMF) as solvent and triethylamine as catalyst.¹³

With regard to **Scheme 1**, the next stage consists of reacting 3-amino-1,4-di-*N*-oxide quinoxaline-2-carbonitrile derivatives with aromatic sulfonyl chloride derivatives in order to synthesize sulfonamides which have shown interesting antileishmanial activity.¹⁴

These quinoxaline derivatives were tested for their activity against two parasites: *P. falciparum* (responsible for malaria) and *L. amazonensis* (responsible for cutaneous New World leishmaniasis), and against non-tumorigenic cell line (Murine Peritoneal Macrophages: MPM). Biological activities and graphical representation of the structure–activity relationship for series 1 and 2 are presented in **Table 1**, **Table 2** and **Figure 1(a,b)**. None of the sulfonamides (compounds **1–7**) showed interesting activity against malaria parasite, being 60 to >80 times less active than chloroquine. Similarly, most of acetoxybenzamides were scarcely active, being 90 to >100 times less active than chloroquine. Only compounds **9** and **14** showed some activity, with an IC_{50} of approximately 10 μ M (although they were almost 30 times less active than CQ). Interestingly, a Cl (**9**) or a F (**14**) in R^7 position increased the activity 4–7 times when compared to a H or a CH_3 in the same position. As shown in **Figure 1(b)**, the position of one electronegative atom (F, Cl or O), represented as a blue field, is clearly important in order for the atoms to act as possible hydrogen bond acceptors. This contrasts with the groups CH_3 and H, which projected hydrophobic regions (represented as golden fields).


Compound **9**, with an acetoxybenzamide in position 3, is almost twice as active as compound **4**, the most active compound of the first series. From a spatial point of view, according to **Figure 1(a,b)**, this could mean that a strong concentration of electronegative atoms (potential hydrogen bond acceptors) in one same plane is a good feature for acetoxybenzamides. Therefore, we can say that *o*-nitrophenyl, *p*-nitrophenyl and 2-naphtyl sulfonamides lower the activity. The most active compound against malaria was compound **14** (IC_{50} 7.4 μ M), followed by compound **9** (10.8 μ M).

With regard to *Leishmania*, compounds **4** and **5** inhibited 50% of parasite growth at approximately 3 μ M, being almost 10–15 times less active than the reference drug amphotericin B. As we mentioned before, in order to be active, the R^7 position must be occupied by one electronegative atom Cl, because when missing (**7**), the activity drops dramatically (86 μ M). The paranitrophenyl substituent is also responsible for the activity in association with Cl in R^7 position, because once it is charged by a naphthyl group (**1**), the activity lowers ten times. Interestingly, the presence of withdrawing groups, such as *o*-nitro and *p*-nitro, in the sulfonamide region is correlated to the activity (see **Fig. 1b**). Nevertheless, compounds **4** and **5** showed a low Selectivity Index (2–4), almost 10 times lower than the SI of amphotericin B. Although compound **2** and **6** were almost five times less active than compounds **4** and **5**, we decided to test them against *Leishmania*-infected macrophages, because these products were not toxic against non-infected macrophages (>245 and 89 μ M respectively).

In infected macrophages, compound **2** was 7 times more active than **6** and showed the best Selectivity Index (>15), being very close to that of amphotericin B. In the second series, the most active compound against *Leishmania* (**14**) was 2–3 times less active than compounds **4** and **5**. It appears that in order to be leishmanicidal, the R^7 position must be occupied by a CH_3 or a F. The activity drops when Cl or CH_3O are in R^7 position. Nevertheless, when the active compounds were tested on non-infected macrophage, they showed some toxicity, being almost as toxic on macrophage as on *Leishmania*.

A computational study designed to predict the ADME properties of our salicylamide and sulfonamide derivatives of quinoxaline was performed (results are presented in **Table 3**). Topological polar surface area (TPSA) is a good indicator of drug absorption in the intestines, Caco-2 monolayer penetration, and blood–brain barrier crossing.¹⁵ TPSA was used to calculate the percentage of absorption (%ABS) following the equation: %ABS = $109 - 0.345 \times \text{TPSA}$, as reported.¹⁶ In addition, the number of rotatable bonds (n-ROTB), and Lipinski's rule of five were also calculated.¹⁷ The most active compounds found for antileishmanial and antimalarial bioassays

Table 1
Biological characterization of sulfonamides



Compd	MW	R ⁶	R ⁷	Ar	IC ₅₀ ^a (μM)	CQ ^b index	IC ₅₀ ^c amas (μM)	Anf ^d index	CC ₅₀ ^e Mø (μM)	SI ^f
1	426.6	H	Cl	2-Naphtyl	>23.4	>78	20 ± 1.4	100	88.6 ± 5.6	4.4
2	406.1	H	CH ₃	2-Naphtyl	>24.6	>82	16.3 ± 0.8	81.5	>245.6	>15.1
3	420.1	CH ₃	CH ₃	2-Naphtyl	>23.8	>79	>100	>100	NT	NT
4	421.6	H	Cl	<i>o</i> -Nitrophenyl	17.4 ± 0.5	58	3.1 ± 0.1	15.5	6.8 ± 0.8	2.2
5	421.6	H	Cl	<i>p</i> -nitrophenyl	>23.7	>79	2.1 ± 0.1	10.5	7.9 ± 0.3	3.8
6	456.1	Cl	Cl	<i>p</i> -Nitrophenyl	18.7 ± 1	62	15.9 ± 1.3	–15	89.1 ± 6.4	5.6
7	387.1	H	H	<i>p</i> -Nitrophenyl	20 ± 2	67	86.3 ± 8.5	430	NT	–
CQ	320				0.2 ± 0.1	1	20 ± 1.4	100	88.6 ± 5.6	4.4
Anf B	320						0.2	1	4.4	22

Selectivity index (SI): CC₅₀ drug/IC₅₀ drug.

MW molecular weight; NT: not tested.

^a IC₅₀ against *P. falciparum* FCR-3.

^b CQ Index: IC₅₀ drug/IC₅₀ CQ.


^c IC₅₀ against axenic amastigotes of *L. amazonensis*.

^d Anf Index: IC₅₀ of compounds /IC₅₀ amphotericin B.

^e Cytotoxicity in murine peritoneal macrophages.

^f Selectivity index.

Table 2
Biological characterization of acetoxybenzamides



Compd	MW	R ⁶	R ⁷	IC ₅₀ ^a (μM)	CQ ^b index	IC ₅₀ ^c amas (μM)	Anf index ^d	CC ₅₀ ^e Mø (μM)	SI ^f
8	364	H	H	41.4 ± 1.1	138	111.8 ± 3.2	559	NT	–
9	398.5	H	Cl	10.8 ± 1.8	36	33.6 ± 1.3	168	NT	–
10	378	H	CH ₃	52.8 ± 1.4	176	18.8 ± 0.3	94	21.7 ± 1.9	1.1
11	394	H	CH ₃ O	27 ± 1.7	90	42.4 ± 1.7	212	NT	–
12	400	F	F	36 ± 0.9	120	14.8 ± 0.1	74	10.5 ± 0.7	0.7
13	392	CH ₃	CH ₃	59.4 ± 2.0	198	17.6 ± 0.6	88	12.5 ± 1.1	0.7
14	382	H	F	7.4 ± 0.6	25	7.3 ± 0.1	36.5	13.6 ± 1.6	1.8
CQ	320			0.3 ± 0.01	1				
Anf B						0.2 ± 0.01	1	4.4 ± 0.1	22

Selectivity Index (SI): CC₅₀ drug/ IC₅₀ drug. MW molecular weight; NT: not tested.

^a IC₅₀ against *P. falciparum* FCR-3.

^b CQ Index: IC₅₀ drug/ IC₅₀ CQ.

^c IC₅₀ against axenic amastigotes of *L. amazonensis*.

^d Anf Index: IC₅₀ of compounds /IC₅₀ amphotericin B.

^e Cytotoxicity in murine peritoneal macrophages.

^f Selectivity index.

(compounds **4**, **5**, **9** and **14**) showed medium percentages of intestinal absorption, with mean values of 58%, and an excellent *n*-ROTB, ranging only from 3 to 4.¹⁸ None of these compounds violated any of the Lipinski's parameters, an important characteristic for future drug-development.

On the other hand, correlations between molecular descriptors such as molecular weight, log *P* and activities of acetoxybenzamides

and sulfonamides derivatives were found. Simple Pearson correlation values, which are useful for measuring the association between two variables, were calculated between biological activities reported in Table 1 and Table 2, and physical chemical properties in Table 3. In contrast to previously reported results^{8,9}, whose derivatives only correlate between log *P* and activities, in our case, an important element for the activity was also conferred




Figure 1. Graphical representation of the structure-activity relationship for sulfonamides (a) and acetoxybenzamides (b). Field Points are colored as follows: Blue; negative field points (like to interact with positives/H-bond donors on a protein), red; positive field points (like to interact with negatives/H-bond acceptors on a protein), yellow; van der Waals surface field points (describing possible surface/vdW interactions), gold/orange; Hydrophobic field points (describe regions with high polarizability/hydrophobicity).

by MW of the derivatives, an important parameter that is directly related with the size of the molecule, which is useful to illustrate the influence of the shape and structural features of a molecule. In the second series, the reported values showed a correlation between MW and antimalarial activity and antileishmanial activity respectively, expressed as $\log(1/IC_{50})$, in which higher values exponentially indicate greater potency (the logarithm of inverse of the IC_{50} is classically used in QSAR studies). Compounds

in the second series followed a positive linear relationship ($r=0.47$ for malaria, $r=0.64$ for *Leishmania* and undetermined for cytotoxicity in murine peritoneal macrophages) while the first series only followed a negative linear relationship between $\log P$ and antimalarial activity ($r=-0.56$), also expressed as $\log(1/IC_{50})$.

The seven new sulfonamides presented in this paper were prepared through the synthetic route illustrated in Scheme 1.

Table 3
Physical Chemical Properties of quinoxaline derivatives^a

ID rule	%ABS	TPSA (Å) ≤140	n-RT	Molecular weight <500	mi Log P <5	Lipophilicity descriptor ^b (Σπ)	n-OH/NH donors ≤5	n-NO acceptors ≤10	Lipinski's violations ≤1
1	67.28	120.9	3	426.84	1.19	0.65	1	8	0
2	67.27	120.8	3	406.42	0.96	0.43	1	8	0
3	67.27	120.8	3	420.45	1.34	0.80	1	8	0
4	51.48	166.7	4	421.79	−0.08	0.65	1	11	1
5	51.48	166.7	4	421.79	−0.03	0.65	1	11	1
6	51.48	166.7	4	456.22	0.57	1.25	1	11	1
7	51.48	166.7	4	387.33	−0.68	0	1	11	1
8	69.98	113.1	4	364.32	−0.47	0	1	9	0
9	64.11	130.1	4	398.76	0.18	0.65	1	9	0
10	64.11	130.1	4	378.34	−0.04	0.43	1	9	0
11	60.94	139.3	5	394.34	−0.43	0.04	1	10	0
12	64.11	130.1	4	400.29	−0.23	0.24	1	9	0
13	64.11	130.1	4	392.37	0.33	0.80	1	9	0
14	64.11	130.1	4	396.33	−0.33	0.14	1	9	0

^a %ABS, percentage of absorption, calculated by: %ABS = 109 − (0.345 × TPSA); TPSA, topological polar surface area; n-RT, number of rotatable bonds; mi log P, logarithm of compounds partition coefficient between *n*-octanol and water.

^b Hydrophilic–lipophilic fragments determined for changes in R⁷ and R⁶, calculated by the sum of Σπ values, using compounds 7 and 8, and the fragment constant method as Ref.20.

With regard to Scheme 2, the final synthetic route consists in combining 3-amino-1,4-di-N-oxide quinoxaline-2-carbonitrile derivatives with *o*-acetylsalicyloyl chloride, used for the treatment of infections caused by protozoans, bacteria and viruses.¹⁹

In the last stage of the synthesis of sulfonamides, the temperature is critical and must remain below 0 °C, whereas the last stage of synthesis of acetoxybenzamides is carried out at room temperature. In addition, there is a loss of yield due to the purification by column chromatography, decreasing from 50% to 25% (acetoxybenzamides) and 13–15% (sulfonamides).

A computational study for prediction of ADME properties of all the molecules was performed using Molinspiration online property calculation toolkit (MolinspirationCheminformatics). Parameters such as solubility (mi log P), Topological Polar Surface Area (TPSA)¹⁵, and absorption (%ABS) were calculated using the formula: %ABS = 109 − (0.345 × TPSA).¹⁶ Violations of Lipinski's rule-of-five were evaluated.¹⁷

Analysis of structure–activity relationship was performed using Cresset's field technology (<http://www.cresset-group.com/>) which condenses the molecular fields down to a set of points around the molecule, termed 'Field Points'.²⁵ Field Points are the local extrema of the electrostatic, van der Waals and hydrophobic potentials of the molecule. Throughout FielView[®] software, the Field Points are colored as follows: Blue; negative field points (like to interact with positives/H-bond donors on a protein), red; positive field points (like to interact with negatives/H-bond acceptors on a protein), yellow; van der Waals surface field points (describing possible surface/vdW interactions), or gold/orange; Hydrophobic field points (describe regions with high polarizability/hydrophobicity).

Quinoxaline derivatives harboring a halogenous group in positions 6 and 7 present interesting activity against *Plasmodium* and *Leishmania* parasite and show interesting in silico ADME characteristics. These products should be the starting point for the synthesis of antiprotozoal compounds.

Acknowledgements

The authors gratefully acknowledge the 'Oficina de Cooperación de la Embajada de Bélgica' for funding a Master Scholarship to

Denis Castillo. Carlos Barea is indebted to the University of Navarra (Spain) for PhD scholarship.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.bmcl.2011.05.125.

References and notes

- Vicente, E.; Charnaud, S.; Bongard, E.; Villar, R.; Burguete, A.; Solano, B.; Ancizu, S.; Perez-Silanes, S.; Aldana, I.; Vivas, L.; Monge, A. *Molecules* **2008**, *13*, 69.
- Biot, C.; Chibale, K. *Infect. Disord. Drug Targets* **2006**, *2*, 173.
- WHO: http://www.who.int/malaria/world_malaria_report_2010/en/index.html.
- WHO: <http://www.who.int/leishmaniasis/en/>. Accessed 27 March, 2011.
- Kedzierski, L.; Sakthianandeswaren, A.; Curtis, J. M.; Andrews, P. C.; Junk, P. C.; Kedzierska, K. *Curr. Med. Chem.* **2009**, *16*, 599.
- Carta, A.; Corona, P.; Loriga, M. *Curr. Med. Chem.* **2005**, *12*, 2259.
- Monge, A.; Palop, J. A.; Piñol, A.; Martínez-Crespo, F. J.; Narro, S.; González, M.; Sáinz, Y.; López de Ceráin, A. *J. Heterocycl. Chem.* **1994**, *31*, 1135.
- Burguete, A.; Estevez, Y.; Castillo, D.; González, G.; Villar, R.; Solano, B.; Vicente, E.; Pérez-Silanes, S.; Aldana, I.; Monge, A.; Sauvain, M.; Deharo, E. *Memorias do Oswaldo Cruz* **2008**, *103*, 778.
- Estevez, Y.; Quiliano, M.; Burguete, A.; Zimic, M.; Málaga, E.; Verástegui, M.; Pérez-Silanes, S.; Aldana, I.; Monge, A.; Castillo, D.; Deharo, E. *Exp. Parasitol.* **2011**, *127*, 745.
- Ancizu, S.; Moreno, E.; Torres, E.; Burguete, A.; Perez-Silanes, S.; Benitez, D.; Villar, R.; Solano, B.; Marin, A.; Aldana, I.; Cerecetto, H.; Gonzalez, M.; Monge, A. *Molecules* **2009**, *14*, 2256.
- Ortega, M. A.; Sainz, Y.; Montoya, M. E.; Jaso, A.; Zarranz, B.; Aldana, I.; Monge, A. *Arzneim.-Forsch.* **2002**, *52*, 113.
- González, M.; Cerecetto, H. *Springer* **2007**, *10*, 265.
- Ley, K.; Seng, F. *Synthesis* **1975**, 415.
- Dea-Ayuela, M. A.; Castillo, E.; González-Álvarez, M.; Vega, C.; Rolón, M.; Bolás-Fernández, F.; Borrás, J.; González-Rosende, M. E. *Bioorg. Med. Chem.* **2009**, *17*, 7449.
- Ertl, P.; Rohde, B.; Selzer, P. *J. Med. Chem.* **2000**, *43*, 3714.
- Zhao, Y. H.; Abraham, M. H.; Le, J.; Hersey, A.; Luscombe, C. N.; Beck, G.; Sherborne, B.; Cooper, I. *Pharm. Res.* **2002**, *19*, 1446.
- Lipinski, C. A.; Lombardo, F.; Dominy, B. W.; Feeney, P. J. *Adv. Drug Delivery Rev.* **2001**, *46*, 3.
- Veber, D. F.; Johnson, S. R.; Cheng, H. Y.; Smith, B. R.; Ward, K. W.; Kopple, K. D. *J. Med. Chem.* **2002**, *45*, 2615.
- De Carvalho, L. P. S.; Lin, G.; Jiang, X.; Nathan, C. J. *Med. Chem.* **2009**, *52*, 5789.
- Gordon, L. F. *J. Pharm. Sci.* **1980**, *69*, 1109.
- Cheeseright, T.; Mackey, M.; Rose, S.; Vinter, J. G. *J. Chem. Inf. Model.* **2006**, *46*, 665.