This article was downloaded by: [Universite Laval] On: 25 December 2014, At: 15:13 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Synthesis of Bis(arylmethylidene)thiopyranones and Crystal Structure of the Phenyl Derivative

M. Saeed Abaee^a, Mohammad M. Mojtahedi^a,
M. Mehdi Zahedi^a, A. Wahid Mesbah^a, Nafiseh
Mohaddes Ghandchi^a & Werner Massa^b

^a Chemistry and Chemical Engineering Research Center of Iran , Tehran, Iran

^b Fachbereich Chemie der Philipps-Universitaet Marburg , Marburg, Germany Published online: 03 Dec 2007.

To cite this article: M. Saeed Abaee , Mohammad M. Mojtahedi , M. Mehdi Zahedi , A. Wahid Mesbah , Nafiseh Mohaddes Ghandchi & Werner Massa (2007) Synthesis of Bis(aryImethylidene)thiopyranones and Crystal Structure of the Phenyl Derivative, Phosphorus, Sulfur, and Silicon and the Related Elements, 182:12, 2891-2895, DOI: 10.1080/10426500701542759

To link to this article: http://dx.doi.org/10.1080/10426500701542759

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and

are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Synthesis of Bis(arylmethylidene)thiopyranones and Crystal Structure of the Phenyl Derivative

M. Saeed Abaee Mohammad M. Mojtahedi M. Mehdi Zahedi A. Wahid Mesbah Nafiseh Mohaddes Ghandchi Chemistry and Chemical Engineering Research Center of Iran, Tehran, Iran

Werner Massa Fachbereich Chemie der Philipps-Universitaet Marburg, Marburg, Germany

A convenient method is offered for double aldol condensation of tetrahydrothiopyran-4-one with aromatic aldehydes under LiBr catalysis. An X-ray analysis verifies the proposed structures.

 $\label{eq:keywords} {\bf Keywords} \ {\bf Aldol \ condensation; \ bis (arylmethylidene) thiopyranones; \ lithium \ bromide; \ thiopyranone$

INTRODUCTION

Synthesis of bis(arylmethylidene)cycloalkanones is facilitated via aldol condensation of homocyclic ketones with aromatic aldehydes¹ and several procedures are offered so far for this purpose using Lewis acid catalysis,² solid-supported reactions,³ ionic liquids,⁴ microwave irradiation,^{3a} and ultrasound mediation.^{3b} In contrast, similar chemistry for heterocyclic analogs is much less developed. We recently disclosed the synthesis of several novel bisarylmethylidene derivatives of thiopyranone⁵ and pyranone⁶ structures using very

Received April 2, 2007; accepted June 5, 2007.

Partial financial support by the Ministry of Science, Research, and Technology of Iran is greatly appreciated.

Address correspondence to Mohammad M. Mojtahedi, Chemistry and Chemical Engineering Research Center of Iran, P. O. Box 14335-186, Tehran, Iran. E-mail: mojtahedi@ ccerci.ac.ir facile synthetic methods. The structures were assigned logically based on the spectroscopic characterization of the products. However, to distinguish between the possible isomers (Scheme 1) and verify the proposed structures we decided to determine the crystal structure of a representative compound by X-ray diffraction.



SCHEME 1

In the present article, a convenient and rapid procedure is offered for the synthesis of bis(arylmethylidene)thiopyranones under catalysis of LiBr which has found many applications as a mild Lewis-acid in recent years to ease up various synthetic organic transformations.⁷ The X-ray structure analysis of the phenyl derivative confirms the presence of exocyclic double bonds in the products in Z-Z configuration (**3**).

RESULTS AND DISCUSSION

An equimolar mixture of thiopyranone **4** and an aldehyde (as listed in Table I) was treated with LiBr (10 mol%) and Et_3N in dichloromethane. Complete conversion of the starting materials to the desired product was observed within a few hours as reactions were monitored by TLC.

	S 4	ArCHO LiBr, Et ₃ N	Ar	O Ar S 3a-g
Entry	ArCHO	M.P. (° C)	Product	Yield% ^a
1	benzaldehyde	142–144	3a	95
2	(p-Me)benzaldehyde	186 - 188	3b	91
3	(p-MeO)benzaldehyde	174 - 176	3c	93
4	(p-CI)benzaldehyde	126 - 128	3d	95
5	2-thiophenecarbaldehyde	155 - 157	3e	92
6	furfural	156 - 158	3f	88
7	3-pyridinecarbaldehyde	186 - 188	3g	94

TABLE I Alodol Condensation of 4 with Aldehydes under LiBrCatalysis

^aIsolated yields.



FIGURE 1 Structure of **3a** at 193 K in the crystal. Displacement ellipsoids at 50% probability level. C1–C2 1.495(2), C2–C3 1.498(2), C2–C6 1.344(2), C3–S1 1.802(2), S1–C4 1.799(2), C4–C5 1.499(2), C5–C1 1.503(2), C5–C13 1.331(2) Å.

Control experiments confirmed the combined promoting and catalytic effects of LiBr; an alternative reaction in the absence of the catalyst led to formation of less than 10% of **3a** after 24 h.

The structure of the products was assigned with spectroscopic methods and compared with the literature data.^{5,8} In order to verify this structure, a single crystal of **3a** was prepared and investigated by X-ray diffraction. The result, as depicted in the Figure 1, clearly supports the proposed structure with exocyclic double bonds C2–C6 and C5–C13. The experimental data and crystal parameters are given in the Table II.

The atoms of the central moiety C1, O1, C2, C3, C4, C5, C6, C7, C13, and C14 are almost coplanar (maximum deviation from the best plane 0.065(2) Å). The folding angle of this main plane and the C3, S1, C4 plane is $122.8(6)^{\circ}$. The planes of the phenyl rings are rotated out of the main plane as documented in torsion angles C5-C13-C14-C15 $-143.0(2)^{\circ}$ and C2-C6-C7-C12 $147.3(2)^{\circ}$, resulting in a molecule conformation resembling a landing bird (Fig. 2).

In summary, a LiBr promoted aldol condensation of thiopyran **4** with various aldehydes was carried out within few hours. Reactions proceeded with catalytic amounts of the Lewis acid, and the use of harsh conditions and tedious work up procedures were avoided. High yields of the products and rapid completion of the reactions are among other advantages of this methodology. The proposed structure was confirmed for the first time by single crystal X-ray diffraction analysis.

EXPERIMENTAL

Typical Procedure for the Synthesis of 3a

A mixture of 4 (5.0 mmol), benzaldehyde (5.0 mmol), LiBr (0.5 mmol), and Et_3N (10.0 mmol) in CH_2Cl_2 (15 mL) was stirred in a flask. The course of the reaction was monitored by TLC and complete

Empirical formula	C ₁₉ H ₁₆ O S
Formula weight	292.38
Habitus, color	column, colorless
Crystal size	$0.33 imes 0.15 imes 0.12~{ m mm}^3$
Crystal system	monoclinic
Space group	$P2_1/n, \mathbf{Z} = 4$
Unit cell dimensions	a = 10.035(1) Å
	$\mathrm{b} = 9.540(1)~\mathrm{\AA}~eta = 95.32(1)^{\circ}.$
	c = 15.595(1) Å
Volume	$1486.55(15){ m \AA}^3$
Wavelength	0.71073 Å
Temperature	193(2) K
Density (calculated)	1.306 Mg/m^3
Absorption coefficient	0.213 mm^{-1} , no correction
F(000)	616
Theta range for data collection	$2.32 ext{ to } 25.87^{\circ}.$
Index ranges	$-12{<}{=}h{<}{=}12,-11{<}{=}k{<}{=}11,-19{<}{=}l{<}{=}19$
Reflections collected	14181
Independent reflections	2870 [R(int) = 0.0308]
Completeness to theta $= 25.87^{\circ}$	99.6 %
Refinement method	Full-matrix least-squares on F ²
Treatment of hydrogen atoms	refined with isotropic displacement parameters
Data / restraints / parameters	2870 / 0 / 254
Goodness-of-fit on F ²	0.715
Final R indices [I>2sigma(I)]	R1 = 0.0307, wR2 = 0.0758
R indices (all data)	R1 = 0.0505, wR2 = 0.0828
Largest diff. peak and hole	0.185 and –0.159 e.Å ^{–3}

TABLE II Crystal and Experimental Data for the StructureDetermination of 3a

disappearance of the starting materials was observed within few hours. The mixture was diluted by dichloromethane (20 mL) and washed twice by water (20 mL portions). The organic phase was dried over Na_2SO_4 , the solvent was removed at reduced pressure and the product was precipitated after removal of the volatile portion. The NMR, and IR spectra and melting point of the product were obtained and compared perfectly with those existing in the literature.^{5,8}

(3Z,5Z)-3,5-Dibenzylidene-tetrahydrothiopyran-4-one (3a)

Yellow crystals were obtained in 95% yield. M.p. 142–144°C; IR (KBr, cm⁻¹) 1599, 1444, 1269;¹H NMR (CDCl₃) δ 3.84 (s, 4H), 7.30 (s, 10H), 7.72 (s, 2H);¹³C NMR (CDCl₃) δ 0.0, 128.4, 128.7, 129.8, 133.7, 134.9, 136.6, 188.6; MS (70 eV) m/z (%) 292 (M⁺), 147, 115.



FIGURE 2 Side views of the structure of **3a** in the crystal showing the molecular conformation.

X-Ray Crystal Structure Analysis of 3a

A crystal of **3a**, grown from hexane, was investigated on an IPDS area detector system (Stoe) at -80° C using MoK α -radiation. Crystal and experimental data are given in the Table II. The structure was solved by direct methods and refined using the SHELX97 programs. Crystallographic data (excluding structure factors) for the structure reported in this paper have been deposited with the Cambridge Crystallographic Data Center as supplementary publication no. CCDC-637988. Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK [Fax: int. Code +44(1223)336-033; E-mail: deposit@ccdc.cam.ac.uk or via www.ccdc.cam.ac.uk/conts/retrieving.html]

REFERENCES

- (a) B. A. Hathaway, J. Chem. Educ., 64, 367 (1987); (b) M. Zheng, L. Wang, J. Shao, and Q. Zhong, Synth. Commun., 27, 351 (1997); (c) N. Iranpoor and E. Kazemi, Tetrahedron, 54, 9475 (1998); (d) T. Nakano and T. Migita, Chem. Lett., 2157 (1993).
- [2] (a) L. Wang, J. Sheng, H. Tian, J. Han, Z. Fan, and C. Qian, Synthesis, 3060 (2004);
 (b) G. Sabitha, K. K. Reddy, K. B. Reddy, and J. S. Yadav, Synthesis, 263 (2004); (c) N. Iranpoor, B. Zeynizadeh, and A. Aghapour, J. Chem. Res., 554 (1999); (d) Y. Zhu and Y. Pan, Chem. Lett., 668 (2004); (e) M. S. Abaee, M. M. Mojtahedi, R. Sharifi, M. M. Zahedi, H. Abbasi, and K. Tabar-Heidar, J. Iran Chem. Soc., 3, 293 (2006); (f) M. S. Abaee, M. M. Mojtahedi, M. Mojtahedi, M. M. Zahedi, R. Sharifi, A. W. Mesbah, W. Massa, W. First report on the synthesis and structural elucidation of novel bisarylmethylidenes of cyclic enones. Synth. Commun., 37, 2949 (2007).
- [3] (a) J. Wang, L. Kang, Y. Hu, and B. Wei, Synth. Commun., 32, 1691 (2002); (b) J. Li,
 W. Yang, G. Chen, T. Li, Synth. Commun., 33, 2619 (2003).
- [4] (a) X. Zheng and Y. Zhang, Synth. Commun., **33**, 161 (2003); (b) X. Hu, X. Fan, X. Zhang, and J. Wang, J. Chem. Res., 684 (2004); (c) X. Zhang, X. Fan, H. Niu, and J. Wang, Green Chem., **5**, 267 (2003).
- [5] (a) M. S. Abaee, M. M. Mojtahedi, and M. M. Zahedi, *Synlett*, 2317 (2005); (b) M. S. Abaee, M. M. Mojtahedi, M. M. Zahedi, and R. Sharifi, *Heteroatom Chem.*, 18, 44 (2007).
- [6] M. S. Abaee, M. M. Mojtahedi, M. M. Zahedi, and M. Bolourtchian, Synth. Commun., 36, 199 (2006).
- [7] (a) W. F. Bailey, M. R. Luderer, K. P. Jordan, J. Org. Chem., 71, 2825 (2006); (b)
 L. X. Shao, M. Shi. Synlett, 1269 (2006); (c) S. Rudrawar, Synlett, 1197 (2005) and references cited therein.
- [8] (a) M. S. Puar, G. C. Rovnyak, A. I. Cohen, B. Toeplitz, J. Z. Gougoutas, J. Org. Chem., 44, 2513 (1979); (b) N. J. Leonard, D. Choudhury, J. Am. Chem. Soc., 79, 156 (1957).