

## Correction to 2-Pyridone Synthesis Using 2-(Phenylsulfinyl)acetamide

Masaya Fujii, Takuya Nishimura, Takahiro Koshihara, Satoshi Yokoshima, and Tohru Fukuyama\*

*Org. Lett.* **2013**, *15*(1), 232–234. DOI: 10.1021/ol303320c

### **S** Supporting Information

**P**eaks of solvents and impurities had been removed from the NMR spectra reported for compounds **6b** (S21), **6c** (S23), **6g** (S29), **6h** (S31), **6i** (S33), **6k** (S37), **6l** (S39 and S40), **6m** (S41 and S42), **6n** (S43 and S44), **4m** (S53), **4n** (S55), and **4o** (S57). Original FIDs were located, and the spectra were reprocessed and have been replaced for the above compounds in the revised Supporting Information submitted with this correction. The spectra editing did not affect any of the conclusions of the published paper. The purities calculated on the basis of the revised spectra and corrected yields are as follows: **6b** (99% purity, 67% yield), **6c** (98% purity, 39% yield), **6k** (94% purity, 83% yield), **6l** (90% purity, 74% yield), **6m** (97% purity, 70% yield), **6n** (89% purity, 72% yield), **4m** (99% purity, 79% yield). The samples of **6g**, **6h**, **6i**, **4n**, and **4o** included some unknown impurities; thus, the exact purity could not be determined.

### ■ ASSOCIATED CONTENT

#### **S** Supporting Information

Revised Supporting Information including reprocessed spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Published: February 3, 2014