

Communication

Total Synthesis of Legionaminic Acid as Basis for Serological Studies

Stefan Matthies, Pierre Stallforth, and Peter H Seeberger

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.5b00455 • Publication Date (Web): 10 Feb 2015

Downloaded from <http://pubs.acs.org> on February 18, 2015

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications
High quality. High impact.

Total Synthesis of Legionaminic Acid as Basis for Serological Studies

Stefan Matthies,^{†,‡} Pierre Stallforth,^{†,§} and Peter H. Seeberger^{*,†,‡}

[†]Max Planck Institute of Colloids and Interfaces, Department of Biomolecular Systems, Am Mühlenberg 1, 14476 Potsdam, Germany

[‡]Freie Universität Berlin, Department of Biology, Chemistry, Pharmacy, Arnimallee 22, 14195 Berlin, Germany

Supporting Information Placeholder

ABSTRACT: Legionaminic acid is a 9-carbon diamino monosaccharide that is found coating the surface of various bacterial human pathogens. Its unique structure makes it a valuable biological probe but access *via* isolation is difficult and no practical synthesis has been reported. We describe a stereoselective synthesis that yields a legionaminic acid building block as well as linker-equipped conjugation-ready legionaminic acid, starting from cheap D-threonine. To set the desired amino and hydroxyl group pattern of the target we designed a concise sequence of stereoselective reactions. The key transformations rely on chelation-controlled organometallic additions and a Petasis multi-component reaction. The legionaminic acid was synthesized in a form that enables attachment to surfaces. Glycan microarray containing legionaminic acid revealed that human antibodies bind the synthetic glycoside. The synthetic bacterial monosaccharide is a valuable probe to detect an immune response to bacterial pathogens such as *Legionella pneumophila*, the causative agent of Legionnaires' disease.

The human immune system can identify bacterial pathogens by interacting with specific carbohydrates on the outer membrane of the microbe.¹ In particular, homopolymers built from sialic acid monomers (**1–3**) are key players involved in recognition processes.^{2,3} While *N*-acetylneuraminic acid **1** is ubiquitous in both mammalian and bacterial glycomes, legionaminic acids **2** and **3** are specific to bacteria.^{3,4} Structurally, they differ from their closest mammalian relative, *N*-acetylneuraminic acid **1**, by lacking one oxygen at C9 and by exhibiting an amine in place of an oxygen at C7 (Scheme 1).

A key virulence factor involved in Legionnaire's disease, a devastating form of pneumonia in humans, is the lipopolysaccharide (LPS) of *Legionella pneumophila*.^{3,5} Legionaminic acid **2** is the major component of the LPS, a homopolymer of 5-*N*-acetimidoyl-7-*N*-acetyl legionaminic acids.^{4,6–8}

The presence of legionaminic acids **2** and **3** (LegA) within glycoconjugates of bacterial pathogens^{3,4} makes them ideal targets for pathogen detection and for the development of vaccines. Interestingly, legionaminic acid is also found in other prominent human pathogens such as *Acinetobacter*


*baumannii*⁹, *Enterobacter cloacae*¹⁰ and *Campylobacter jejuni*.^{3,11}

The structural complexity of legionaminic acids that renders them exquisite biological probes has a major drawback: the total synthesis of these compounds is notoriously difficult. So far no synthetic strategy has been described to access fully functionalized legionaminic acid **2** and **3** glycosylating agents^{3,8,12–14} for the construction of oligosaccharides or for the conjugation to slides or carrier proteins.^{15,16}

Here, we provide a versatile synthetic route to legionaminic acid building block **5** and linker-equipped legionaminic acid **4**. We describe a *de novo* strategy^{17–27} starting from inexpensive and commercially available D-threonine **8**.


The retrosynthetic disconnection of linker-LegA **4** is shown in Scheme 1 with hemiketal **5** as universal building block. The latter is accessible by a sequence of an oxidation and an indium-mediated allylation of aminol **6**. Acetamide **6** is derived from α -hydroxy aldehyde **7** via a Petasis-borono-Mannich reaction. Aldehyde **7** in turn originates from D-threonine **8** via an intramolecular inversion and a chelation-controlled organometallic addition reaction.

Scheme 1. A) Structures of *N*-Acetylneuraminic acid **1**, Legionaminic acid **2** and 4-*Epi*-legionaminic acid **3**; B) Retrosynthesis of Legionaminic Acid Building Block **5** as well as Conjugation-Ready LegA **4**.



The total synthesis of legionaminic acid commenced with preparation of aldehyde **9** setting the required stereogenic information at C7 and C8 in the final building block (Scheme 2). Since *D*-*allo*-threonine methyl ester is prohibitively expensive, aldehyde **9** was synthesized from *D*-threonine **8** via a slightly modified literature procedure.²⁸ The necessary C3-epimerization was achieved by anchimeric assistance. Methyl ester formation, *N*-benzoylation, followed by thionyl chloride-induced cyclization, acidic hydrolysis of the intermediate oxazoline and esterification yielded *D*-*allo*-threonine methyl ester in 98% over five steps without chromatographic purification. The latter was transformed to the corresponding *N*-toluenesulfonamide, followed by an acetalization and DIBAL-H reduction of the intermediate ester provided *D*-*allo*-threoninal **9** in 84% overall yield.²⁹

Scheme 2. Synthesis of Hemiketal Donor **5**.^a



^aReagents and conditions: a) SOCl₂, MeOH, reflux, 1 h, quant.; BzCl, NEt₃, MeOH, 0°C, 4 h, quant.; SOCl₂, 0°C, 5 days, 99%; 6N HCl, reflux, 5 h; SOCl₂, MeOH, reflux, 1 h, 99%; TsCl, NEt₃, DCM, 0°C, 4 h, 99%; 2,2-DMP, *p*-TsOH, Toluene, 120°C, 4 h, 98%; DIBAL-H, Toluene, -78°C, 2 h, 87%; b) *n*-BuLi, furan, MgBr₂·OEt₂, DME, -78°C, 4 h, d.r.=5:1 (*syn/anti*), 80% (*syn*); c) O₃, DCM, MeOH, -78°C, 1 h, then PPh₃ and TMSCHN₂, 0°C, 59%; d) LiBH₄, THF, 0°C → rt, 12 h, 92%; e) NaOCl, cat. TEMPO, KBr, DCM, sat. aq. NaHCO₃, 0°C, 10 min, quant.; f) (*E*)-styrylboronic acid, HNR₂, d.r.>19:1 (*anti/syn*), yield and conditions see: Table 1; g) DMBA, Pd(PPh₃)₄, DCM, 35°C, 2 h, then Ac₂O, NaHCO₃, MeOH, rt, 5 h, 99%; h) O₃, PPh₃ (resin, crosslinked), then i) In, methyl 2-(bromomethyl)-acrylate, EtOH, sat. aq. NH₄Cl, rt, sonication, 30 min, 99%, d.r.=2.5:1 (*syn/anti*); j) O₃, MeOH, -78°C, 15 min, then Me₂S, rt, 4 h, 46% (*syn*), 15% (*anti*); k) Ac₂O, pyr, DMAP, DCM, 0°C → rt, 12 h, 82%. *p*-TsOH = *p*-toluenesulfonic acid, Bz = Benzoyl, DMBA = 1,3-dimethylbarbituric acid, DMP = Dimethoxypropane, Ts = Tosyl.

With threoninal **9** in hand, a Cram-chelate organometallic addition reaction of 2-lithiofuran to threoninal **9** resulted in the formation of the desired *syn*-configured alcohol **10** (Scheme 2).²⁹ Crystallization gave diastereomerically pure **10**

in 80% yield. Since the oxidation of the furan moiety failed using ruthenium-based protocols,³⁰ we treated alcohol **10** with ozone followed by methylation to give methyl ester **11** in 59% yield. The conversion of ester **11** to diol **12** employing lithium borohydride in THF³¹ proceeded in 92% yield. X-ray crystallographic analysis of diol **12** confirmed the stereochemical assignment (see SI).³²

α -Hydroxy aldehyde **7** was required as precursor to introduce a protected amine via a Petasis-borono-Mannich reaction.^{33,34} Careful optimization of the chemoselective oxidation of diol **12** to aldehyde **7** was crucial in circumventing the notorious propensity of α -hydroxy aldehydes to decompose or epimerize. Subjecting alcohol **12** to Dess-Martin oxidation³⁵ resulted in low conversion, while TEMPO-mediated oxidation employing trichloroisocyanuric acid as the stoichiometric oxidant³⁶ did not proceed reproducibly. Finally, when using Anelli's modification of the TEMPO oxidation,³⁷ the desired aldehyde **7** was obtained in quantitative yield without loss of stereochemical purity.³⁸

To access protected diamines **13** or **14**, a Petasis-borono-Mannich reaction between aldehyde **7**, (*E*)-styrylboronic acid and an amine component was used (Table 1). The newly formed stereocenter displays *anti*-configuration relative to the directing free hydroxyl group. The high degree of stereocontrol reported by Petasis and Zavialov³⁴ arises from two energetically different conformations of the intermediate ate complexes (not shown, for review see³³).

Table 1. Optimization of the Petasis Multi Component Reaction

Entry	Amine	Solvent	Temp.	Time	Yield
1	NHAI ₂ ^[a]	EtOH	rt	36 h	49%
2	NHAI ₂ ^[b]	DCM	120°C ^[f]	0.5 h	<10%
3	NHAI ₂ ^[b]	DCM/HFIP ^[c]	40°C	24 h	46%
4	NHAI ₂ ^[b]	DCM/HFIP ^[d]	40°C	24 h	57%
5	NHAI ₂ ^[b]	EtOH/water ^[e]	40°C	2 h	23%
6	NHAI ₂ ^[b]	DCM	rt	24 h	0% ^[f]
7	NH ₂ AI ^[a]	EtOH	rt	24 h	76%

^[a]Equivalents (7:(*E*)-styrylboronic acid:amine) 1:1:1.

^[b]Equivalents (7:(*E*)-styrylboronic acid:amine) 1:1.5:1.5, ^[c]3:1,

^[d]9:1, ^[e]4:1, ^[f]microwave irradiation, ^[f]*N,O*-Isopropylidene hydrolyzed. HFIP = 1,1,1,3,3,3-Hexafluoro-2-propanol

Table 1 summarizes the optimization process of the key Petasis-borono-Mannich reaction. Initially, we applied standard reaction conditions using ethanol as the solvent and equimolar amounts of all reactants as well as a reaction time of 36 h (entry 1).³⁴ Despite quantitative conversion of **7**, the isolated yield of aminol **13** was low (49%). Microwave irradiation is known to accelerate and facilitate Petasis reactions.^{38,39} However, in our hands it did not result in acceptable rates of conversion (entry 2). Polar, protic solvents such as hexafluoroisopropanol are known to increase reaction rates and improve yields for this transformation. Unfortunately, the yields were comparable to the initial trials (entries 3 and 4). Using an ethanol-water mixture, as well as dichloromethane as solvent resulted in *N,O*-isopropylidene

cleavage rather than improved yields (entries 5 and 6). Eventually, the use of monoallyl amine provided aminol **14** in 76% yield (entry 7). As desired, the exclusive formation of the *anti*-diastereomers **13** and **14** with a d.r. > 19:1 was observed in all cases.


Conversion of the *N*-allyl group of **14** into the corresponding *N*-acetate was achieved by Pd(o)-catalyzed deallylation followed by chemoselective *N*-acetylation to give amide **6** in 99% yield (Scheme 2).³⁸ Oxidative cleavage of *N*-cinnamylacetamide **6** to give aldehyde **15** required careful optimization of the ozonolytic conditions to prevent epimerization and decomposition of the labile α -aminoaldehyde moiety.^{27,40} Zinc and dimethyl sulfide reduced the ozonide but the transformation required long reaction times.²⁷ The use of resin-bound triphenyl phosphine⁴¹ resulted in complete reduction of the intermediate ozonide without epimerization. Thus, quantitative conversion of hydroxylamine **6** to *N*-acetamide aldehyde **15** was achieved. Indium-mediated allylation of aldehyde **15** to enoate **16** is an established method to introduce a masked pyruvate unit.^{42–44} While other potential methods such as the Cornforth synthesis⁴⁵ are performed using harsh conditions, the In-mediated allylation offers mild reaction conditions and can be performed in aqueous media.⁴³ Quantitative conversion of aldehyde **15** to the desired allylic alcohol **16** was achieved using sonication and a solvent mixture consisting of ethanol and aqueous ammonium chloride. Ammonium chloride was used as alternative to aqueous hydrogen chloride,⁴⁴ which would hydrolyze the *N,O*-isopropylidene acetal. Labile enoate **16** was isolated in high yield (99%) in favor of the *syn*-configuration (d.r. = 2.5:1 (*syn/anti*)). Oxidative cleavage of the double bond on enoate **16**, proved problematic, initially leading to decomposition or low yields. The use of a reductive workup (dimethyl sulfide) after ozonolytic cleavage of the alkene moiety in methanol to provide legionaminic acid **17** and 4-epilegionaminic acid derivative **18** in good yield.

To obtain linker-equipped legionaminic acid **4**, the installation of an anomeric linker via glycosidation of a suitable legionaminic acid building block was required. Screening of protocols for the glycosidation of sialic acid building blocks⁴⁶ (e.g. anomeric chlorides, acetates and phosphites, thioglycosides and *N*-phenyl-trifluoroacetimidate, results not shown) were irreproducible. Finally, we found that a dehydrative glycosylation protocol⁴⁷ furnished the desired linker-conjugate. Thus, diol **17** yet had to be elaborated into C4-OAc hemiketal donor **5** by selective acylation. Careful optimization allowed for selective C4-O-acetylation when using 2 equivalents of acetic anhydride and pyridine in the presence of catalytic amounts of DMAP to yield monoacetylated hemiketal **5** in high yield.⁴⁴ Starting from D-threonine, the overall yield of acetate **5** was 10%. To confirm the stereochemical configuration of diol **17**, the corresponding C2/C4-OAc derivative was characterized (see SI).

Dehydrative glycosylation of 4Ac5NAcLeg donor **5** provided crude thioether **19** (Scheme 3).⁴⁷ Surprisingly, attempts to purify crude **19** all failed due to decomposition of the material. Thus, acetal hydrolysis using aqueous TFA was performed prior to purification to provide glycoside **20** (63% β -anomer). Although the α -anomer was formed, it proved to be inseparable from minor decomposition products by means of silica gel chromatography. A 1D heteronuclear correlation experi-

ment (selective, coupled, 1D ¹H-¹³C-HMBC) was performed to determine the heteronuclear coupling constant. With ³J_{C-1H-3ax} = 1.9 Hz the anomeric configuration proved the β -anomeric center.⁴⁶ Starting from thioether **20**, acetate removal gave alcohol **21** in close to quantitative yield. Hydrolysis of the methyl ester, using 20 equivalents of lithium hydroxide to prevent the formation of elimination by-products, gave carboxylate **22** in quantitative yield.

Scheme 3. Synthesis of Glycoside **4**.^a




^aReagents and conditions: a) Ph₂SO, Tf₂O, HOC₂H₄SBn, 4 Å MS, -78°C → -50°C, 2.5 h; then b) DCM, TFA, H₂O (10:3:1 v/v), 0°C, 5 min 63%; c) NaOMe, MeOH, rt, 12 h, 95%; d) 20 eq. LiOH, MeOH/H₂O (1:1 v/v), rt, 15 h, 99%; e) Na_(s), NH₃(l), -78°C, 45 min, then O₂ atmosphere, 99%; f) NaOH, Ac₂O, H₂O, 0°C, 60 min, then MeOH, 2 h, rt, 98%.

Global deprotection by reductive cleavage of the thioether and the *N*-sulfonamide moiety was achieved by Birch reduction (Scheme 3). Carboxylate **22** was treated with sodium in liquid ammonia without co-solvent to provide the desired amine **23** as the disulfide in quantitative yield. Amide **4** was finally obtained by treatment of amine **23** with sodium hydroxide and acetic anhydride. Starting from D-threonine, the overall yield of *N*-acetamide **4** was 6%.

To evaluate whether a legionaminic acid monosaccharide is a sufficient epitope for antibody recognition, human sera were screened against the synthetic structures using glycan microarrays (Scheme 4).^{48–51} Carbohydrates **4** and **23** were immobilized via their thiol linker on maleimide coated glass slides. Treatment with pooled human sera revealed that legionaminic acids **4** and **23** were recognized, presumably as a result of previous immune responses to bacterial pathogens displaying a legionaminic acid motif. IgG serum antibody levels against acetamide **4** were substantially higher than against amine **23** (Scheme 4). This side chain modification appears to be crucial for the specificity of antibody recognition. Since no bacterial glycans isolated so far display free amino groups on C7, we chose amine **23** as negative control to exclude unspecific binding. We further confirmed the preferential binding of human antibodies to antigen **4** using two other pooled human antisera (CDC1992⁵² and anti-Hib human reference serum⁵³, results not shown). Legionaminic acid-binding antibodies in human sera show that this rare

carbohydrate from bacterial pathogen origin is recognized by the human immune system. Conjugate **4** is a valuable tool for the production of novel pathogen detection systems or vaccines.

Scheme 4. Microarray analysis of synthetic LegA.



A: Printing Pattern. Legionaminic acid structures (**4** and **23**) printed in concentration from 1 mM to 0.0016 mM. B: Representative microarray scan representing IgG antibodies to LegA **4** and **23** after incubation with Human Reference Serum 007sp.

In summary, we have completed the *de novo* synthesis of orthogonally protected legionaminic acid **5**. This building block served as a glycosylating agent in the synthesis of linker-equipped legionaminic acid **4**. Glycan arrays containing synthetic antigen **4** showed that this epitope is recognized by human antibodies present in blood sera. Immunization studies and additional binding studies are currently ongoing.

ASSOCIATED CONTENT

Supporting Information

Experimental details and characterization data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

peter.seeberger@mpikg.mpg.de

Present Addresses

§ Leibniz Institute for Natural Product Research and Infection Biology e.V. (HKI), Department of Chemistry of Microbial Communication, Beutenbergstr. 11, 07745 Jena, Germany

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENT

We thank the Max Planck Society and the Studienstiftung des deutschen Volkes (Fellowship to P.S.) for generous funding. The authors would like to thank Dr. A. Schäfer (FU Berlin) for NMR analysis, Dr. A. Hagenbach (FU Berlin) for X-ray crystallographic analysis and Dr. F. Pfengle for critical comments to the manuscript and A. Geissner for help with microarray experiments.

REFERENCES

(1) Varki, A.; Cummings, R. D.; Esko, J. D.; Freeze, H. H.; Stanley, P.; Bertozzi, C. R.; Hart, G. W.; Etzler, M. E. *Essentials of Glycobiology; Second Edition*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 2009.

- (2) Mühlenhoff, M.; Eckhardt, M.; Gerardy-Schahn, R. *Curr. Opin. Struct. Biol.* **1998**, *8*, 558.
- (3) Zunk, M.; Kiefel, M. J. *RSC Adv.* **2014**, *4*, 3413.
- (4) Knirel, Y. A.; Shashkov, A. S.; Tsvetkov, Y. E.; Jansson, P.-E.; Zähringer, U. In *Adv. Carbohydr. Chem. Biochem.*; Horton, D., Ed.; Academic Press: 2003; Vol. 58, p 371.
- (5) Cazalet, C.; Jarraud, S.; Ghavi-Helm, Y.; Kunst, F.; Glaser, P.; Etienne, J.; Buchrieser, C. *Genome Res.* **2008**, *18*, 431.
- (6) Kooistra, O.; Herfurth, L.; Lüneberg, E.; Frosch, M.; Peters, T.; Zähringer, U. *Eur. J. Biochem.* **2002**, *269*, 573.
- (7) Mas Pons, J.; Dumont, A.; Sautejeau, G.; Fugier, E.; Baron, A.; Dukan, S.; Vauzeilles, B. *Angew. Chem. Int. Ed.* **2014**, *53*, 1275.
- (8) Tsvetkov, Y. E.; Shashkov, A. S.; Knirel, Y. A.; Zähringer, U. *Carbohydr. Res.* **2001**, *331*, 233.
- (9) Vinogradov, E.; MacLean, L.; Xu, H. H.; Chen, W. *Carbohydr. Res.* **2014**, *390*, 42.
- (10) Filatov, A. V.; Wang, M.; Wang, W.; Perepelov, A. V.; Shashkov, A. S.; Wang, L.; Knirel, Y. A. *Carbohydr. Res.* **2014**, *392*, 21.
- (11) MacLean, L. L.; Vinogradov, E.; Pagotto, F.; Perry, M. B. *Carbohydr. Res.* **2011**, *346*, 2589.
- (12) Tsvetkov, Y. E.; Shashkov, A. S.; Knirel, Y. A.; Zähringer, U. *Carbohydr. Res.* **2001**, *335*, 221.
- (13) Glaze, P. A.; Watson, D. C.; Young, N. M.; Tanner, M. E. *Biochemistry* **2008**, *47*, 3272.
- (14) Watson, D. C.; Leclerc, S.; Wakarchuk, W. W.; Young, N. M. *Glycobiology* **2011**, *21*, 99.
- (15) Schumann, B.; Anish, C.; Pereira, C. L.; Seeberger, P. H. In *Biotherapeutics: Recent Developments using Chemical and Molecular Biology*; The Royal Society of Chemistry: 2013; p 68.
- (16) Seeberger, P. H.; Werz, D. B. *Nature* **2007**, *446*, 1046.
- (17) Schmidt, R. R. *Pure Appl. Chem.* **1987**, *59*, 415.
- (18) Vogel, P. In *The Organic Chemistry of Sugars*; Levy, D. E., Fügedi, P., Eds.; CRC Press: Boca Raton, 2005.
- (19) Kirschning, A.; Jesberger, M.; Schöning, K.-U. *Synthesis* **2001**, 507.
- (20) Martin, S. F.; Zinke, P. W. *J. Am. Chem. Soc.* **1989**, *111*, 2311.
- (21) Aljhdali, A. Z.; Shi, P.; Zhong, Y.; O'Doherty, G. A. *Adv. Carbohydr. Chem. Biochem.* **2013**, *69*, 55.
- (22) Guo, H.; O'Doherty, G. A. *J. Org. Chem.* **2008**, *73*, 5211.
- (23) Yu, X.; O'Doherty, G. In *Chemical Glycobiology*; American Chemical Society: 2008; Vol. 990, p 3.
- (24) Brasholz, M.; Reissig, H.-U. *Angew. Chem. Int. Ed.* **2007**, *46*, 1634.
- (25) Leonori, D.; Seeberger, P. H. *Beilstein J. Org. Chem.* **2013**, *9*, 332.
- (26) Leonori, D.; Seeberger, P. H. *Org. Lett.* **2012**, *14*, 4954.
- (27) Stallforth, P.; Matthies, S.; Adibekian, A.; Gillingham, D. G.; Hilvert, D.; Seeberger, P. H. *Chem. Commun.* **2012**, *48*, 11987.
- (28) Liang, X.; Lee, C.-J.; Chen, X.; Chung, H. S.; Zeng, D.; Raetz, C. R. H.; Li, Y.; Zhou, P.; Toone, E. J. *Bioorg. Med. Chem.* **2011**, *19*, 852.
- (29) Szechner, B.; Achmatowicz, O.; Galdecki, Z.; Fruzinski, A. *Tetrahedron* **1994**, *50*, 7611.
- (30) Kasai, M.; Ziffer, H. J. *Org. Chem.* **1983**, *48*, 2346.
- (31) Watanabe, A.; Kiyota, N.; Yamasaki, T.; Tanda, K.; Miyagoe, T.; Sakamoto, M.; Otsuka, M. *J. Heterocycl. Chem.* **2011**, *48*, 1132.
- (32) CCDC 1019693 contains the supplementary crystallographic data for this paper. These data can be obtained from The Cambridge Crystallographic Data Centre at www.ccdc.cam.ac.uk/data_request/cif.
- (33) Candeias, N. R.; Montalbano, F.; Cal, P. M. S. D.; Gois, P. M. P. *Chem. Rev.* **2010**, *110*, 6169.
- (34) Petasis, N. A.; Zavialov, I. A. *J. Am. Chem. Soc.* **1998**, *120*, 11798.
- (35) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155.
- (36) De Luca, L.; Giacomelli, G.; Porcheddu, A. *Org. Lett.* **2001**, *3*, 3041.

(37) Lucio Anelli, P.; Biffi, C.; Montanari, F.; Quici, S. *J. Org. Chem.* **1987**, *52*, 2559.

(38) Soulé, J.-F.; Mathieu, A.; Norsikian, S.; Beau, J.-M. *Org. Lett.* **2010**, *12*, 5322.

(39) McLean, N. J.; Tye, H.; Whittaker, M. *Tetrahedron Lett.* **2004**, *45*, 3.

(40) Ito, A.; Takahashi, R.; Baba, Y. *Chem. Pharm. Bull.* **1975**, *23*, 3081.

(41) Prakash, G. K. S.; Mandal, M.; Schweizer, S.; Petasis, N. A.; Olah, G. A. *Org. Lett.* **2000**, *2*, 3173.

(42) Gordon, D. M.; Whitesides, G. M. *J. Org. Chem.* **1993**, *58*, 7937.

(43) Pae, A. N.; Cho, Y. S. *Curr. Org. Chem.* **2002**, *6*, 715.

(44) Lee, Y. J.; Kubota, A.; Ishiwata, A.; Ito, Y. *Tetrahedron Lett.* **2011**, *52*, 418.

(45) Cornforth, J. W.; Firth, M. E.; Gottschalk, A. *Biochem. J.* **1958**, *68*, 57.

(46) Boons, G.-J.; Demchenko, A. V. *Chem. Rev.* **2000**, *100*, 4539.

(47) Haberman, J. M.; Gin, D. Y. *Org. Lett.* **2003**, *5*, 2539.

(48) de Paz, J. L.; Seeberger, P. H. *QSAR Comb. Sci.* **2006**, *25*, 1027.

(49) Liu, Y.; Palma, A. S.; Feizi, T. *Biol. Chem.* **2009**, *390*, 647.

(50) Park, S.; Gildersleeve, J. C.; Blixt, O.; Shin, I. *Chem. Soc. Rev.* **2013**, *42*, 4310.

(51) Geissner, A.; Anish, C.; Seeberger, P. H. *Curr. Opin. Chem. Biol.* **2014**, *18*, 38.

(52) Holder, P. K.; Maslanka, S. E.; Pais, L. B.; Dykes, J.; Plikaytis, B. D.; Carlone, G. M. *Clin. Diagn. Lab. Immunol.* **1995**, *2*, 132.

(53) Madore, D. V.; Johnson, C. L.; Phipps, D. C.; Myers, M. G.; Eby, R.; Smith, D. H. *Pediatrics* **1990**, *86*, 527.

