

ORIGINAL ARTICLE

Incubation of 2-methylisoborneol synthase with the intermediate analog 2-methylneryl diphosphate

Wayne KW Chou, Colin A Gould and David E Cane

Incubation of synthetic 2-methylneryl diphosphate (2-MeNPP, 10) with 2-methylisoborneol synthase (MIBS) gave a mixture of products that differed significantly from that derived from the natural substrate (*E*)-2-methylgeranyl diphosphate (3, 2-MeGPP). The proportion of (-)-2-methylisoborneol (1) decreased from 89 to 17% while that of 2-methylenebornane (4) increased from 10 to 26%, with the relative yields of the isomeric homo-monoterpenes 2-methyl-2-bornene (5) and 1-methylcamphene (6) remaining essentially unchanged (<1% each), as determined by chiral GC–MS analysis. The majority of the product mixture resulting from the MIBS-catalyzed cyclization of 2-MeNPP (10) consisted of the anomalous monocyclic homo-monoterpenes (\pm)-2-methyllimonene (15, 39%) and 2-methyl- α -terpineol (13, 10%), as well as the acyclic derivatives 2-methylnerol (11, 7%) and 2-methylinalool (14, <1%). The steady-state kinetic parameters of the MIBS-catalyzed reaction, determined using [$1\text{-}^3\text{H}$]-2-methylneryl diphosphate (2-MeNPP), were k_{cat} $0.0046 \pm 0.0003 \text{ s}^{-1}$, K_{m} $18 \pm 6 \mu\text{M}$ and $k_{\text{cat}}/K_{\text{m}}$ $2.55 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$. In comparison, the natural substrate 2-MeGPP had a k_{cat} $0.105 \pm 0.007 \text{ s}^{-1}$, K_{m} $95 \pm 49 \mu\text{M}$ and $k_{\text{cat}}/K_{\text{m}}$ $1.11 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$. Taken together with earlier X-ray crystallographic studies of MIBS, as well as previous investigations of the mechanistically related plant monoterpene cyclase, bornyl diphosphate synthase, these results provide important insights into the binding and cyclization of both native substrates and intermediates and their analogs.

The Journal of Antibiotics advance online publication, 1 March 2017; doi:10.1038/ja.2017.24

INTRODUCTION

Since the discovery of 2-methylisoborneol (2-MIB, 1) by Gerber in 1969, there has been considerable interest in the detection, bioremediation and biosynthesis of this volatile, odiferous homo-monoterpene (Figure 1).^{1–4} Possessing a musty odor and muddy off-taste, as well as an extremely low threshold of detection by humans (<10 ng l⁻¹), 2-MIB is produced by a wide range of Gram-positive actinomycetes, myxobacteria and cyanobacteria, being second in occurrence only to the common odiferous sesquiterpene geosmin.⁵

The elucidation of the biosynthesis of 2-MIB at the molecular genetic and biochemical level was independently reported in 2008 by two research groups (Figure 1).^{6–8} The 2-MIB biosynthetic gene cluster in *Streptomyces coelicolor* was shown to harbor a terpene synthase gene (*sco7700*) translationally coupled to the gene for an *S*-adenosyl-*L*-methionine-dependent C-methyl transferase, (*sco7701*), as well as a third gene (*sco7699*) encoding a protein of still unknown function, annotated only as a nucleotide-binding protein.⁷ This three-gene biosynthetic operon is highly conserved across the genome sequences of more than three dozen bacterial species.^{3,5,6} The *S*-adenosyl-*L*-methionine-dependent geranyl diphosphate C-methyl transferase (GPPMT; *Sco7701*) was shown to catalyze the first step in the biosynthetic pathway in *S. coelicolor*, the electrophilic methylation of the olefinic C-2 position of geranyl diphosphate

(GPP, 2) to form (*E*)-2-methylgeranyl diphosphate (2-MeGPP, 3).⁷ *Sco7700*, 2-MIB synthase (MIBS), catalyzes the Mg²⁺-dependent multistep cyclization of the acyclic 2-MeGPP precursor to the bicyclic alcohol 2-MIB (1) which is accompanied by small amounts of the bicyclic homo-monoterpenes 2-methylenebornane (4), 2-methylbornene (5) and 1-methylcamphene (6). The only other enzyme known to utilize 2-MeGPP as substrate is the closely related 2-methylenebornane synthase (Pfl_1841) of *Pseudomonas fluorescens* PfO-1.⁹ These findings were fully consistent with the results of independent whole-cell precursor incorporation experiments involving the feeding of [*methyl*-¹³C]methionine and samples of deuterated mevalonate to the myxobacterium *Nannocystis exedens* and MS analysis of the resulting 2-MIB in the head-space volatiles.⁴

The proposed 2-MIB synthase mechanism is based on close analogy to the well-documented cyclization mechanisms established for a wide range of monoterpene synthases (Figure 2).¹⁰ MIBS initiates the electrophilic cyclization by ionizing the Mg²⁺-complexed substrate 2-MeGPP (3) to generate the corresponding allylic cation–pyrophosphate ion pair which then collapses to the *transoid* conformer of the allylic isomer (3*R*)-2-methylinalyl diphosphate (2-MeLPP, 7). Rotation about the C-2,3-bond generates the *cisoid* conformer of 7, folded in the *anti-endo*-boat conformation, which then undergoes ionization and cyclization to generate the (4*R*)-2-methyl- α -terpinyl

Department of Chemistry, Box H, Brown University, Providence, RI, USA

Correspondence: Dr DE Cane, Department of Chemistry, Box H, Brown University, Providence, RI 02912-9108, USA.

E-mail: david_cane@brown.edu

This article is dedicated to Professor Satoshi Ōmura, 2015 Nobel Laureate, in Physiology or Medicine, in honor of his profound contributions to basic science and the improvement of human health as well as his long-standing support of the arts, and in deep appreciation for his continuing support and friendship.

Received 20 October 2016; revised 1 December 2016; accepted 27 January 2017




Figure 1 Biosynthesis of 2-methylisoborneol (1).




Figure 2 Mechanism of the 2-methylisoborneol synthase (MIBS)-catalyzed cyclization of 2-methylgeranyl diphosphate (3) to 2-methylisoborneol (1). A full color version of this figure is available at the *The Journal of Antibiotics* journal online.

cation (8). Electrophilic attack on the cyclohexenyl double bond of 8 generates the 2-methyl-2-bornyl cation (9), which is then quenched on the *exo* face by bound water to generate 2-MIB (1). Competing deprotonation of intermediate 9 or its proximal rearrangement product readily accounts for formation of the co-products 4, 5 and 6. The MIBS cyclization mechanism, including the deduced stereochemistry and conformations of the various intermediates, closely resembles that firmly established for (+)-bornyl diphosphate synthase/ α -pinene synthase from *Salvia officinalis* which converts GPP (2) to (+)-bornyl diphosphate (75%) as well as a mixture of the monoterpene hydrocarbons (+)- α -pinene, (+)-camphene, (\pm)-limonene and terpinolene (25% total olefins; Figure 3).^{11–13} There are of course two key differences between the reactions catalyzed by MIBS and by (+)-bornyl diphosphate synthase/ α -pinene synthase: (1) MIBS cyclizes 2-MeGPP instead of GPP. Although MIBS can cyclize GPP to a mixture of monocyclic and bicyclic monoterpenes, the observed k_{cat} is four orders of magnitude lower than that for the natural methylated substrate 2-MeGPP.⁷ (2) The final step in the MIBS-catalyzed cyclization cascade is the quenching of the 2-methyl-2-bornyl cation exclusively on the *exo* face by a bound water (Figure 2), while in the (+)-bornyl diphosphate synthase/ α -pinene synthase-catalyzed reaction, the paired pyrophosphate ion is recombined with the bornyl cation exclusively by *endo* attack (Figure 3).

High-resolution crystal structures have recently been determined for MIBS in both an unliganded state and in complex with a number of




Figure 3 Mechanism of bornyl diphosphate/ α -pinene synthase. A full color version of this figure is available at the *The Journal of Antibiotics* journal online.

substrate or intermediate analogs.^{14,15} The 1.80–1.95 Å structures of MIBS bound to geranyl-S-thiolodiphosphate and 2-fluorogeranyl diphosphate, respectively, show each analog coordinated with two Mg^{2+} ions and bound in extended conformations in which the 6,7-double bond of each analog is twisted away from the parallel relationship expected for the postulated *anti-endo-boat* conformation of the actual cyclizing *cisoid* (3*R*)-2-MeLPP intermediate (Figure 2).¹⁴ Although this deviation may simply be the result of crystallographic trapping of a thermodynamically stable, rather than a kinetically or mechanistically relevant conformer, more interestingly it reflects the fact that the required 180° rotation of the 2-propenyl substituent about the C-2,3 bond of the *anti-endo-boat* conformer of the (3*R*)-2-methylallyl diphosphate intermediate is sterically forbidden by a clash with the 6,7-double bond that is augmented by the presence of the C-2-methyl substituent. It is thus probable that adoption of the *endo-boat* conformation takes place *subsequent* to conversion of the *transoid* to the *cisoid* conformer of 2-MeLPP. Such extended 6,7-double bond conformations of bound substrate analogs have previously been observed in the structures of a number of canonical monoterpene synthases.^{16,17} Also thought-provoking were the unexpected results from co-crystallization of MIBS with racemic




Figure 4 Synthesis of 2-methylnerol diphosphate (**10**, 2-MeNPP) and 2-methyl- α -terpineol (**13**).

2-fluorolinalyl diphosphate (2-FLPP), which had been predicted to be an unreactive analog of the natural (3*R*)-2-methylallyl diphosphate intermediate due to the strongly electron-withdrawing 2-fluoro substituent. The 2.00 Å structure of the resulting complex revealed that the MIBS-bound 2-FLPP had undergone a reverse allylic rearrangement to generate bound 2-fluoroneryl diphosphate in complex with 2 Mg²⁺ ions.¹⁵ Also unexpectedly, it was found that in solution MIBS converted the supposedly unreactive 2-FLPP to camphor, no doubt formed by cyclization of 2-FLPP and spontaneous elimination of hydrogen fluoride from enzymatically generated 2-fluoro-2-isoborneol. The observation of enzyme-generated 2-fluoroneryl diphosphate trapped within the active site of MIBS raises the question whether the *cis* geometric isomer of 2-FGPP is a true intermediate of the MIBS-catalyzed ionization and cyclization of 2-FLPP, the nominally unreactive analog of the natural intermediate 2-MeLPP (7), or merely an anomalous shunt metabolite of the reverse allylic diphosphate rearrangement. We have therefore now examined the incubation of MIBS with 2-MeNPP (**10**), the *cis* isomer of the natural substrate, 2-MeGPP (**3**), and report the results below.

RESULTS

Preparation of 2-methylnerol diphosphate (**10**) and (±)-2-methyl- α -terpineol (**13**)

To prepare 2-MeNPP (**10**), we used a modification of the previously reported synthesis of the *trans* isomer 2-MeGPP (**3**; Figure 4).⁷ Wadsworth–Horner–Emmons reaction of 6-methyl-5-hepten-2-one with triethyl 2-phosphonopropionate followed by LiAlH₄ reduction of the resulting mixture of *Z* and *E* conjugated esters gave a 1:1 mixture of (*Z*)-2-methylnerol (**11**) and (*E*)-2-methylgeraniol (**12**) which were cleanly resolved by flash column chromatography (see Supplementary Information for details). The purified 2-methylnerol (**11**) was reacted with triphenylphosphine and CCl₄ in an Appel reaction and the resulting 2-methylneryl chloride was then directly reacted without further purification with tris(tetrabutylammonium) hydrogen pyrophosphate to give 2-MeNPP (**10**). The corresponding [1-³H]-2-MeNPP ([1-³H]-**10**) was similarly prepared by oxidation of the mixture of 2-methylnerol and 2-methylgeraniol to the derived mixture of aldehydes, followed by reduction with NaBH₃T and chromatographic separation of the resulting mixture of [1-³H]-2-methylnerol and [1-³H]-2-methylgeraniol using AgNO₃-impregnated silica gel.

For the planned enzyme incubations, we also needed an authentic reference sample of the homo-monoterpene alcohol (±)-2-methyl- α -terpineol (**13**),¹⁸ which was readily prepared by biomimetically-modeled, *p*-TsOH-catalyzed solvolysis of 2-methylnerol (**11**) in nitromethane (Figure 4).

In vitro incubation of 2-methylneryl diphosphate with 2-methylisoborneol synthase

Before examining the cyclization of 2-MeNPP by methylisoborneol synthase, we carried out individual control incubations in the absence of enzyme with 2-MeGPP and 2-MeNPP and analyzed the pentane-soluble extracts by chiral GC–MS in order to identify and quantify the formation of Mg²⁺-catalyzed solvolysis products (Figures 5a and b). The control incubation with 2-MeGPP produced only racemic (±)-2-methylallalool (**14**). The exclusive formation of this tertiary allylic alcohol contrasts with the well-known Mg²⁺-catalyzed solvolysis of GPP itself, which typically gives a 3:1 mixture of linalool and geraniol.¹⁸ The preferred formation of 2-methylallalool (**14**) by solvolysis of 2-MeGPP presumably reflects the higher free energy of 2-methylgeraniol due to the increased steric hindrance about the tetra-substituted double bond compared to geraniol. In contrast, the control Mg²⁺-catalyzed solvolysis of 2-MeNPP gave a mixture of three racemic products, consisting of acyclic (±)-2-methylallalool (**14**) as well as the monocyclic (±)-2-methylallimonene (**15**), and (±)-2-methyl- α -terpineol (**13**; Figure 5b). The Mg²⁺-catalyzed formation of both (±)-**13** and (±)-**15** from 2-MeNPP, but not from 2-MeGPP, is consistent with the fact that the (*E*)-allylic diphosphate 2-MeGPP (**3**) is geometrically incapable of direct cyclization. As a positive control, incubation of 2-MeGPP (**3**) with 2-MIB synthase and chiral GC–MS analysis confirmed the formation of (–)-2-MIB (**1**) as the major product (89%) accompanied by minor amounts of enantiomerically pure 2-methylenebornane (**4**; 10%), 1-methylcamphene (**6**; <1%) and 2-methyl-2-bornene (**5**; <1%), as previously observed (Figure 5c and Table 1).^{6,7}

Incubation of 2-MeNPP (**10**) with MIBS and analysis by chiral GC–MS revealed the production of the same group of homochiral bicyclic homo-monoterpenes, (–)-2-MIB (**1**, 17%), 2-methylenebornane (**13**, 26%), 1-methylcamphene (**6**) (<1%) and 2-methyl-2-bornene (**5**) (<1%), but in both substantially different proportions and lower overall yield compared to 2-MeGPP (Figures 5d and 6). Significantly, four additional, previously unobserved products




Figure 5 Chiral GC-MS analysis (total ion current, TIC) of incubations with methylisoborneol synthase (MIBS). (a) Control incubation of 2-MeGPP (**3**) in Mg²⁺-containing buffer without MIBS. (b) Control incubation of 2-MeNPP (**10**) in Mg²⁺-containing buffer without MIBS. (c) Incubation of MIBS with 2-MeGPP (**3**). (d) Incubation of MIBS with 2-MeNPP (**10**). Previously unobserved product peaks are shown in red. Peaks denoted with *are geraniol internal standard. A full color version of this figure is available at the *The Journal of Antibiotics* journal online.


Table 1 Distribution of products from the incubation of 2-MeGPP (3) and 2-MeNPP (10) with 2-MIB synthase

Substrate	Product (%)							
	MIB (1)	4	5	6	13	(±)-15	11	14
2-MeGPP (3)	89	10	<1	<1	ND	ND	ND	ND
2-MeNPP (10)	17	26	<1	<1	10 ^a	39 ^a	7	<1 ^a
					(10/0) ^b	(17/22) ^b		(0/<1) ^b

Abbreviations: MeGPP, methylgeranyl diphosphate; MeNPP, methylneryl diphosphate; MIB, methylisoborneol; ND, not detected.

^aPercent corrected for background Mg²⁺-catalyzed solvolysis.

^bPercent first eluted peak/percent second eluted peak.

**Figure 6** Mechanism of the methylisoborneol synthase (MIBS)-catalyzed transformation of 2-methylneryl diphosphate (2-MeNPP; 10).

constituting >56% of the overall mixture were formed: monocyclic 2-methyl- α -terpineol (13, 10%) along with the acyclic alcohols 2-methylnerol (11, 7%) and 2-methyl- α -terpinene (15, 39%) and 2-methyl- α -terpineol (13, 10%), as well as the acyclic derivatives 2-methylnerol (11, 7%) and 2-methyl- α -terpinene (15, 39%). Interestingly, correction for background Mg²⁺-catalyzed solvolysis of the 2-MeNPP substrate revealed that both 2-methyl- α -terpineol (13) and 2-methyl- α -terpinene (15) were formed as single enantiomers, while 2-methyl- α -terpineol (13) and 2-methyl- α -terpinene (15) were generated as a racemic mixture. The absolute configurations of the enzymatically generated products 13 and 14 were not determined.

Steady-state kinetics

The steady-state kinetic parameters for the MIBS-catalyzed conversion of [1-³H]-2-MeNPP (10) to total pentane-soluble homo-monoterpene products were determined and directly compared with those re-determined for the natural substrate [1-³H]-2-MeGPP (3) under identical conditions (Table 2). While the turnover number, k_{cat} , for 2-MeNPP (0.0045 s⁻¹) was ~20-fold lower than that for 2-MeGPP, the observed k_{cat}/K_m , of 255 M⁻¹ s⁻¹ for 2-MeNPP was only 4-fold lower than that for 2-MeGPP (1105 M⁻¹ s⁻¹), reflecting a partially compensating ~5-fold decrease in the observed K_m for 2-MeNPP compared to 2-MeGPP.⁸

Table 2 Steady-state kinetic parameters for total product formation from incubation with methylisoborneol synthase

Substrate	k_{cat} (s ⁻¹)	K_m (μ M)	k_{cat}/K_m (M ⁻¹ s ⁻¹)
2-MeGPP (3)	0.105 ± 0.007	95 ± 49	1110
2-MeNPP (10)	0.0046 ± 0.0003	18 ± 6	255

DISCUSSION

Although (*Z*)-2-MeNPP (10, 2-MeNPP) shows only a modest fourfold decrease in the specificity constant k_{cat}/K_m compared to the natural substrate, the *trans* geometric isomer (*E*)-2-MeGPP (3), the substantially altered product distributions resulting from incubation of 2-MeNPP with 2-MIB synthase establish that 2-MeNPP is a poor surrogate for 2-MeGPP. Thus, while 2-MeNPP could, in principle, be ionized to the same *cisoid* allylic cation–pyrophosphate ion pair as the presumptive intermediate (3*R*)-2-methylallyl diphosphate (7, 2-MeLPP), in practice incubation of 2-MeNPP with 2-MIB synthase generated a substantially altered array of bicyclic and monocyclic homo-monoterpene products. Thus, although cyclization of 2-MeNPP generated only a single enantiomer of each of the four natural bicyclic natural products, 2-MIB (1), 2-methylenebornane (4), 2-methylbornene (5) and 1-methylcamphene (6), the relative proportion of 1 was reduced more than fivefold from 89% to 17% of the total product mixture, only partially offset by an increase in the fraction of 2-methylenebornane (4) which increased from 10 to 26% (Table 1). The majority of the product mixture resulting from the MIBS-catalyzed cyclization of 2-MeNPP (10) consisted of the anomalous monocyclic homo-monoterpenes (±)-2-methyl- α -terpineol (15, 39%) and a single enantiomer of 2-methyl- α -terpineol (13, 10%), as well as the acyclic derivatives 2-methylnerol (11, 7%) and a single enantiomer of 2-methyl- α -terpinene (15, 39%). Interestingly, while 13, 14 and 15 are not detected among the products of the *in vitro* incubation of 2-MIB synthase with 2-MeGPP, all three of these homo-monoterpenes, of unspecified chiral purity, have been reported as minor components of the volatile head-space extracts of a small number of 2-MIB-producing actinomycetes.^{19,20}

Substantial changes in product distribution have been reported when the *cis* isomer neryl diphosphate is incubated with monoterpene synthases in place of the natural *trans* allylic substrate, geranyl diphosphate. For example, in spite of only minor differences in overall k_{cat} , incubation (+)-BPP/(+)-pinene synthase with NPP instead of GPP resulted in a decrease in the proportion of the bicyclic monoterpenes α -pinene and camphene from 79% to 23%, while the fraction of the monocyclic products limonene and α -terpinolene exhibited a substantial increase from 15 to 71% of total monoterpenes.²¹ The enhanced formation of abortive cyclization products is almost certainly the consequence of aberrant positioning of 2-MeNPP and NPP in the active sites of the respective homo-monoterpene or monoterpene synthases, compared to the binding of the native substrates 2-MeGPP or GPP.

The fact that cyclization of 2-MeNPP by 2-MIB synthase generates the naturally occurring enantiomers of 2-MIB (1), 2-methylenebornane (4), 2-methylbornene (5) and 1-methylcamphene (6) indicates that at least a portion of this anomalous substrate is bound to MIBS in a position and conformation that is compatible with, but certainly not identical to, the precise folding of the natural substrate 2-MeGPP (3) and the derived intermediate (3*R*)-2-MeLPP (7) in the MIBS active site. Although the 2-methyl- α -terpineol (13) and 2-methyl- α -terpinene (15) formed from 2-MeNPP are both formally derivable from a common 2-methyl- α -terpinyl cation, the fact that 15 is generated as a racemic mixture while 13 is produced as

a single enantiomer indicates clearly that they cannot both be derived from a single-monocyclic carbocation intermediate. The surrogate 2-MeNPP substrate therefore must be bound in at least two distinct conformations during MIBS-catalyzed cyclization. The fact that only a single enantiomer of each of the bicyclic, monocyclic and acyclic alcohols **1**, **13** and **14** is produced by incubation of 2-MeNPP with MIBS also suggests that the same bound water molecule is responsible for quenching the corresponding carbocation intermediates.

EXPERIMENTAL PROCEDURE

Materials

Reagents and solvents were purchased from Sigma-Aldrich (St Louis, MO, USA) or Fisher Scientific (Waltham, MA, USA), were of the highest quality available, and were used without further purification. NaBH_3T solution (6 μmol , 80 Ci mmol^{-1} specific activity, 500 mCi total in 1 ml of 0.01 N NaOH) was purchased from American Radiolabeled Chemicals (St Louis, MO, USA). Isopropylthio-D-galactopyranoside was purchased from Invitrogen (Waltham, MA, USA). Ni-NTA affinity resin was from Qiagen (Valencia, CA, USA). Amicon Ultra Centrifugal Filter Units (Amicon Ultra-15 10 000 MWCO) were purchased from Millipore (St Charles, MO, USA). Purified SCO7700 protein was overexpressed and purified as previously described.⁷

Methods

GC – MS analyses were performed using an Agilent 5977A Series GC/MSD instrument (70 eV, electron impact), a 1- μl injection volume, and a 3 min solvent delay. Achiral GC – MS conditions used an HP-5 ms capillary column (0.25 mm ID \times 30 m length \times 0.25 μm film, Agilent Technologies (Santa Clara, CA, USA)) and a temperature program with a 2 min hold at 60 °C, a 20 °C min^{-1} increment to 280 °C, followed by a 2 min hold at 280 °C. Chiral GC – MS separations were performed using a CP-ChiralSil-Dex column (0.32 mm ID \times 25 m length \times 0.25 μm film, Agilent) and a temperature program with a 1 min hold at 50 °C, a 10 °C min^{-1} increment to 200 °C, followed by a 1 min hold at 200 °C. Compounds detected by GC – MS were directly compared to their authentic standards using the MassFinder 4.2.1 program (<http://www.massfinder.com>). Retention indices were measured using C_8 – C_{20} and C_{10} – C_{40} alkane standards.

LC–MS analyses were performed using a Finnigan LXQ LC–MS instrument in negative (electrospray ionization ESI) mode, with a Waters Symmetry C_{18} column (35 μm , 2.1 \times 50 mm), a 10 μl injection volume and 1 min signal delay. The method used was a 1 min isocratic gradient of 95:5 water:acetonitrile, a gradient to 5:95 water:acetonitrile over 5 min, followed by isocratic elution with 5:95 water:acetonitrile for 9 min. NMR spectra were obtained using a Bruker Avance AV400 NMR spectrometer operating at a 400 MHz ^1H frequency or a Bruker Avance AV300 NMR spectrometer operating at a 300 MHz ^1H frequency. Liquid scintillation counting was performed using a Beckman-Coulter LS6500 scintillation counter and Opti-Flour scintillation cocktail from Perkin-Elmer (Waltham, MA, USA).

All proteins were handled at 4 °C unless otherwise stated. Protein concentrations were determined according to the method of Bradford with bovine serum albumin as the standard.²² Protein purity was estimated using SDS PAGE gel electrophoresis and visualized using Coomassie Blue stain according to the method of Laemmli.²³

2-methylneryl diphosphate (10)

The *cis* isomer 2-methylnerol (**11**) was obtained by preparation of a 1:1 mixture of 2-methylnerol (**11**) to 2-methylgeraniol (**12**) and separation of the isomeric alcohols using column chromatography as previously described.⁷ The purified 2-methylnerol was then converted to 2-MeNPP (**10**) using the method developed by Poulter.²⁴ Triphenylphosphine (1.27 g, 4.75 mmol) was added to a stirring solution of 2-methylnerol (**11**; 200 mg, 1.19 mmol) in anhydrous CCl_4 (16 ml) and the mixture was refluxed for 24 h. The solvent was evaporated *in vacuo* and the resulting slurry was triturated with pentane (30 ml) and filtered by suction filtration. Concentration of the pentane filtrate at room temperature provided crude 2-methylneryl chloride, which was used in the subsequent reaction without further purification. Tris(tetrabutylammonium) hydrogen pyrophosphate (1.58 g,

2.38 mmol) was added to the 2-methylneryl chloride in dry CH_3CN (10 ml) under nitrogen and the reaction was stirred at room temperature for 2 days. The solvent was removed under a gentle stream of N_2 to give a viscous, light brown liquid, followed by the addition of 30 ml of 0.05 M KHCO_3 and extraction with ether (5 \times 5 ml) to remove organic impurities. The resulting aqueous phase was applied to a DEAE-Sephadex column that had been equilibrated with 0.05 M KHCO_3 . After loading, the column was washed with 100 ml of 0.05 M KHCO_3 and eluted with a linear gradient of 0.05 M KHCO_3 (300 ml) to 1.0 M KHCO_3 (300 ml) at a flow rate of 2.0 ml min^{-1} . Fractions (35 ml) were collected and lyophilized, and those containing 2-MeNPP (**10**) were dissolved in a minimum amount of dH_2O and applied to a CHP-20P column equilibrated with dH_2O . The column was run with a stepwise gradient of 100 ml of dH_2O , 100 ml of 5% aqueous CH_3CN , 100 ml of 10% aqueous CH_3CN , 100 ml of 20% aqueous CH_3CN in H_2O and 100 ml of 60% aqueous CH_3CN in H_2O . Fractions were lyophilized to yield 105 mg of 2-MeNPP (**10**) as a pale yellow solid (27% yield). $R_f=0.35$ (6:3:1 propanol to NH_4OH to water). ^1H NMR (400 MHz, D_2O) δ 5.08 (t, $J=4.5$ Hz, 1H, H-6), 4.31 (d, $J=4.0$ Hz, 2H, H-1), 2.05 (m, 2H, H-4), 1.98 (m, 2H, H-5), 1.61 (s, 3H, H-11), 1.58 (s, 3H, H-9), 1.55 (s, 3H, H-10), 1.49 (s, 3H, H-8). ^{31}P $\{^1\text{H}\}$ NMR (161 MHz, D_2O) δ -6.6 (d, $J=22.5$ Hz, P_β), -10.4 (d, $J=22.5$ Hz, P_α). ^{13}C $\{^1\text{H}\}$ NMR (75 MHz, D_2O): δ 16.1 (CH_3), 17.0 (CH_3), 18.2 (CH_3), 24.9 (CH_3), 26.6 (CH_2), 33.3 (CH_2), 66.3 (CH_2), 124.1 (CH), 133.8 (C), 135.6 (C), 160.3 (C). LC–MS (neg ion ESIMS) m/z 326.99.

[1- ^3H]-2-Methylneryl diphosphate ([1- ^3H]-10) and [1- ^3H]-2-methylgeranyl diphosphate ([1- ^3H]-3)

A 1.0 ml solution of NaBH_3T (6 μmol , 80 Ci mmol^{-1} , 500 mCi in 1 ml of 0.01 N NaOH) was added to a round bottom flask. The reagent vial was rinsed with 4 \times 1 ml of 0.01 N NaOH which was added to the solution and then placed under argon, cooled in an ice bath. An ice-cold 1:1 mixture of 2-methylnerol to 2-methylgeraniol (425 mg, 2.55 mmol), freshly prepared as previously described,⁷ in dry EtOH (7.5 ml) was added slowly over 10 min. The reaction was stirred at 0 °C for 3 h, and the remaining aldehydes were reduced by the slow addition of unlabeled NaBH_4 powder (75 mg, 2.0 mmol). The resulting solution was warmed to room temperature and stirred overnight. After a total of 22 h, the reaction was quenched by careful transfer of the mixture into ice-cold half-saturated aqueous NH_4Cl (10 ml). The aqueous layer was extracted with Et_2O (5 \times 10 ml) and the combined extracts were dried, filtered and concentrated to give 310 mg of yellowish liquid (72% crude chemical yield, 375 mCi, 75% radiochemical yield). The residue was purified by flash chromatography on 10% (w/w) AgNO_3 -impregnated silica gel (EtOAc /hexanes (1:9); 4 \times 16-cm column) giving 100 mg of [1- ^3H]-2-methylnerol ([1- ^3H]-11, 25% chemical yield, 128 mCi, 26% radiochemical yield) and 210 mg of [1- ^3H]-2-methylgeraniol ([1- ^3H]-12, 48% chemical yield, 247 mCi, 49% radiochemical yield). The purified [1- ^3H]-2-methylnerol ([1- ^3H]-11) and [1- ^3H]-2-methylgeraniol ([1- ^3H]-12) were individually converted to the corresponding diphosphate esters using the same method employed for the synthesis of unlabeled **10** and **3** to yield 98 mg of [1- ^3H]-2-MeNPP ([1- ^3H]-10), 37% chemical yield, 6.2 mCi total radioactivity, 2.2% radiochemical yield, 28 mCi/mmol specific activity) and 200.1 mg of [1- ^3H]-2-MeGPP ([1- ^3H]-3, 36% chemical yield, 105 mCi total radioactivity, 36% radiochemical yield and 232.8 mCi/mmol specific activity). The NMR spectra of ([1- ^3H]-10 and of [1- ^3H]-3 matched those of unlabeled **10** and **3**.⁷

2-Methyl- α -terpineol (13)

p-TsOH (40 mg, 0.20 mmol) was added to a solution of 2-methylnerol (200 mg, 1.19 mmol) in 10 ml of MeNO_2 . The reaction mixture was stirred at rt for 1 h and monitored by TLC (5:1 hexanes/ethyl acetate). The crude product was concentrated *in vacuo* and purified by column chromatography (5:1 hexanes/ethyl acetate) yielding 44 mg of 2-methyl- α -terpineol (**13**, 22% yield). The NMR data matched those previously reported for **13** prepared by an alternative method.¹⁹ ^1H NMR (400 MHz, CDCl_3) δ 1.98, 1.48 (m, 2H, H-6), 1.85 (m, 2H, H-3), 1.72 (m, 1H, H-5), 1.65 (s, 6H, H-10, 11), 1.48 (m, 2H, H-4), 1.22 (s, 6H, H-8, 9). ^{13}C $\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 19.4 (C-9, CH_3), 19.5 (C-8, CH_3), 24.3 (C-3, CH_2), 26.3 (C-6, CH_2), 27.3 (C-4, CH_2), 32.7 (C-11, CH_3), 33.2 (C-10, CH_3), 45.8 (C-5, CH), 72.9 (C-7, C), 125.0 (C-2, C), 125.8 (C-1, C). GC–MS (EI) m/z 168.1.

Incubations with 2-methylisoborneol synthase (SCO7700)

Purified 2-MIB synthase (SCO7700; 10 μ M) was added to a glass test tube containing 3.0 ml of assay buffer (50 mM PIPES, 100 mM NaCl, 15 mM MgCl₂, 5 mM β -mercaptoethanol and 20% glycerol, pH 6.7) and 60 μ M of either 2-MeNPP (10) or 2-MeGPP (3). The enzymatic reaction mixture was overlaid with 3.0 ml of pentane and incubated at 30 °C for 2 h. The enzymatic products were extracted with 3 \times 3.0 ml of pentane and the organic extracts dried over Na₂SO₄, filtered, concentrated *in vacuo* to 100 μ l and analyzed by GC–MS. Control reactions, measuring the background Mg²⁺-catalyzed hydrolysis of each substrate, were conducted similarly as above minus the addition of SCO7700 to the incubation mixture. An internal geraniol standard (25 pmol) was added to the organic extracts of the enzymatic and control incubations after extraction.

Steady-state kinetics

Kinetic analyses were performed in 1 ml of assay buffer (50 mM PIPES, 15 mM MgCl₂, 100 mM NaCl, 5 mM β -mercaptoethanol, 20% glycerol and pH 6.7) with either 2-MeNPP (10, 10–500 μ M) and [1-³H]-2-MeNPP ([1-³H]-10, 8.4 Ci/mol) or 2-MeGPP (3, 10–500 μ M) and [1-³H]2-MeGPP ([1-³H]-3, 16.5 Ci mol⁻¹). The reactions were initiated by the addition of 0.20 μ M of SCO7700 protein, overlaid with 1 ml of pentane and incubated at 30 °C for 17 min. The reactions were quenched by the addition of 75 μ l of 500 mM EDTA (pH 8.0) and vortexing for 30 s. The pentane layer was loaded onto a silica column (2 cm) in a Pasteur pipette and forced through with a stream of nitrogen into a scintillation vial containing 7 ml of Opti-Fluor. The enzymatic reaction mixture was extracted 3 more times with 1 ml of ether, and all organic extracts were passed through the silica column and collected. The combined extracts were assayed by liquid scintillation counting using a Beckman-Coulter LS6500 scintillation counter. Kinetic constants were calculated using the program Kaleidagraph 4.0 and were fit to the Michaelis–Menten equation. Reported standard deviations in the steady-state kinetic parameters represent the calculated statistical errors in the nonlinear, least squares regression analysis.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by NIH grant GM30301 (DEC). We thank Tun-Li Shen for assistance with MS and Russell Hopson for assistance with NMR spectroscopy.

- 1 Gerber, N. N. A volatile metabolite of actinomycetes, 2-methylisoborneol. *J. Antibiot.* **22**, 508–509 (1969).
- 2 Jüttner, F. & Watson, S. B. Biochemical and ecological control of geosmin and 2-methylisoborneol in source waters. *Appl. Environ. Microbiol.* **73**, 4395–4406 (2007).

- 3 Giglio, S., Chou, W. K., Ikeda, H., Cane, D. E. & Monis, P. T. Biosynthesis of 2-methylisoborneol in cyanobacteria. *Environ. Sci. Technol.* **45**, 992–998 (2011).
- 4 Dickschat, J. S. *et al.* Biosynthesis of the off-flavor 2-methylisoborneol by the myxobacterium *Nannocystis excedens*. *Angew. Chem. Int. Ed. Engl.* **46**, 8287–8290 (2007).
- 5 Yamada, Y. *et al.* Terpene synthases are widely distributed in bacteria. *Proc. Natl Acad. Sci. USA* **112**, 857–862 (2015).
- 6 Komatsu, M., Tsuda, M., Omura, S., Oikawa, H. & Ikeda, H. Identification and functional analysis of genes controlling biosynthesis of 2-methylisoborneol. *Proc. Natl Acad. Sci. USA* **105**, 7422–7427 (2008).
- 7 Wang, C. M. & Cane, D. E. Biochemistry and molecular genetics of the biosynthesis of the earthy odorant methylisoborneol in *Streptomyces coelicolor*. *J. Am. Chem. Soc.* **130**, 8908–8909 (2008).
- 8 Wang, C. M. & Cane, D. E. Biochemistry and molecular genetics of the biosynthesis of the earthy odorant methylisoborneol in *Streptomyces coelicolor* (corn to *J. Am. Chem. Soc.* **130**, 8908 (2008)). *J. Am. Chem. Soc.* **132**, 9509 (2010).
- 9 Chou, W. K., Ikeda, H. & Cane, D. E. Cloning and characterization of Pfl₁₈₄₁, a 2-methylenebornane synthase in *Pseudomonas fluorescens* PfO-1. *Tetrahedron* **67**, 6627–6632 (2011).
- 10 Wise, M. L. & Croteau, R. in *Comprehensive Natural Products Chemistry*, Vol. 2 (ed. D. E. Cane) 97–153 (Elsevier, New York, NY, USA, 1999).
- 11 Croteau, R., Satterwhite, D. M., Cane, D. E. & Chang, C. C. Biosynthesis of monoterpenes: enantioselectivity in the enzymatic cyclization of (+)- and (–)-linalyl pyrophosphate to (+)- and (–)-bornyl pyrophosphate. *J. Biol. Chem.* **261**, 13438–13445 (1986).
- 12 Croteau, R., Satterwhite, D. M., Cane, D. E. & Chang, C. C. Enantioselectivity in the enzymatic cyclization of (+)- and (–)-linalyl pyrophosphate to (+)- and (–)-pinene and (+)- and (–)-camphene. *J. Biol. Chem.* **263**, 10063–10071 (1988).
- 13 Wise, M. L., Savage, T. J., Katahira, E. & Croteau, R. Monoterpene synthases from common sage (*Salvia officinalis*). cDNA isolation, characterization, and functional expression of (+)-sabinene synthase, 1,8-cineole synthase, and (+)-bornyl diphosphate synthase. *J. Biol. Chem.* **273**, 14891–14899 (1998).
- 14 Koksai, M., Chou, W. K., Cane, D. E. & Christianson, D. W. Structure of 2-methylisoborneol synthase from *Streptomyces coelicolor* and implications for the cyclization of a noncanonical C-methylated monoterpene substrate. *Biochemistry* **51**, 3011–3020 (2012).
- 15 Koksai, M., Chou, W. K., Cane, D. E. & Christianson, D. W. Unexpected reactivity of 2-fluorolinalyl diphosphate in the active site of crystalline 2-methylisoborneol synthase. *Biochemistry* **52**, 5247–5255 (2013).
- 16 Whittington, D. A. *et al.* Bornyl diphosphate synthase: structure and strategy for carbocation manipulation by a terpenoid cyclase. *Proc. Natl Acad. Sci. USA* **99**, 15375–15380 (2002).
- 17 Hyatt, D. C. *et al.* Structure of limonene synthase, a simple model for terpenoid cyclase catalysis. *Proc. Natl Acad. Sci. USA* **104**, 5360–5365 (2007).
- 18 George-Nascimento, C., Pont-Lezicka, R. & Cori, O. Nonenzymic formation of nerolidol from farnesyl pyrophosphate in the presence of bivalent cations. *Biochem. Biophys. Res. Commun.* **45**, 119–124 (1971).
- 19 Brock, N. L., Ravella, S. R., Schulz, S. & Dickschat, J. S. A detailed view of 2-methylisoborneol biosynthesis. *Angew. Chem. Int. Ed. Engl.* **52**, 2100–2104 (2013).
- 20 Citron, C. A., Barra, L., Wink, J. & Dickschat, J. S. Volatiles from nineteen recently genome sequenced actinomycetes. *Org. Biomol. Chem.* **13**, 2673–2683 (2015).
- 21 Croteau, R. & Satterwhite, D. M. Biosynthesis of monoterpenes. Stereochemical implications of acyclic and monocyclic olefin formation by (+)- and (–)-pinene cyclases from sage. *J. Biol. Chem.* **264**, 15309–15315 (1989).
- 22 Bradford, M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248–254 (1976).
- 23 Laemmli, U. K. Cleavage of structural proteins during the assembly of the head of Bacteriophage T4. *Nature (London)* **227**, 680–685 (1970).
- 24 Davisson, V. J. *et al.* Phosphorylation of isoprenoid alcohols. *J. Org. Chem.* **51**, 4768–4779 (1986).

Supplementary Information accompanies the paper on The Journal of Antibiotics website (<http://www.nature.com/ja>).