

Synthesis and Structures of Mononuclear Co(III) and Cu(II) Complexes Derived from N-[2-(2-Hydroxy-3-methoxybenzylideneamino)cyclohexyl]-2-hydroxybenzamide

HUI-YU WANG, YUN-HUI LI, YU-QIN MA, YING GAO, GUI-MIAO YU and XIU-YUN YANG*

School of Chemistry and Environmental Engineering, Changchun University of Science and Technology, Changchun 130022, P.R. China

*Corresponding author: Fax: +86 0431 84727678; Tel: +86 0431 85582361; E-mail: liyh@cust.edu.cn; whyu@ciac.jl.cn

Received: 18 March 2013;

Accepted: 2 September 2013;

Published online: 30 January 2014;

AJC-14624

Two mononuclear complexes, $[\text{Co}(\text{HL})_2(\text{C}_8\text{H}_7\text{O}_3)] \cdot (\text{CH}_3\text{OH})_2$ (**1**) and $[\text{Cu}(\text{HL})_2]$ (**2**) where $\text{H}_2\text{L} = \text{N}$ -[2-(2-hydroxy-3-methoxybenzylideneamino)cyclohexyl]-2-hydroxybenzamide have been synthesized and characterized by IR, elemental analysis and crystal structure analysis. In complexes (**1**) and (**2**) the core ion has the six-coordinated N_2O_4 environment with a distorted octahedral geometry and four-coordinated N_2O_2 environment with a distorted square-planar geometry, respectively.

Keywords: Cobalt(II), Copper(II), Complexes, Crystal structure.

INTRODUCTION

The novel transition-metal complexes with oxygen and nitrogen donor Schiff bases have received attention not only because of their fascinating self-assembled structures but also because of their important catalytic, magnetic and biological properties¹⁻³. *o*-Vanillin has allowed the construction of polymers with defined geometry and special properties;^{4,5} and the application of transition-metal,^{6,7} lanthanide, or a mixed transition-metal/lanthanide⁸ strategy has yielded many structurally and magnetically interesting systems in recent years⁹. *o*-Vanillin as multidentate ligand displays a variety of bonding geometries, such as chelating, monodentate/bidentate bridging and chelating bridging^{1,10}. Therefore, methods based on such bridging ligands (e.g., methoxide, hydroxide, alkoxide, oxide, carboxylate) have led to the formation of some interesting complexes¹¹.

From the structural point of view, we selected the multidentate Schiff ligand, N-[2-(2-hydroxy-3-methoxybenzylideneamino)cyclohexyl]-2-hydroxybenzamide (H_2L) formed by *o*-vanillin and N-(2-aminocyclohexyl)-2-hydroxybenzamide, to construct coordination complexes. The Schiff base ligand (H_2L) contains adequate number of oxygen and nitrogen donors and these donors with suitable relative positions can chelate one or more metal centers. Accordingly, the design of such ligands is highly effective for establishing simple syntheses for new metal-assembled complexes¹². In this contribution, we describe the synthesis and crystal structures of two mononuclear cobalt(III) and copper(II) complexes with N-[2-(2-hydroxy-3-methoxybenzylideneamino)cyclohexyl]-2-

hydroxybenzamide (H_2L) ligand, $[\text{Co}(\text{HL})_2(\text{C}_8\text{H}_7\text{O}_3)] \cdot (\text{CH}_3\text{OH})_2$ (**1**) and $[\text{Cu}(\text{HL})_2]$ (**2**).

EXPERIMENTAL

All the reagents and solvents employed were commercially available and used as received without further purification. Elemental analysis for C, N and H were performed with a Perkin-Elmer 2400 II analyzer.

Preparation of H_2L : N-(2-Aminocyclohexyl)-2-hydroxybenzamide was synthesized by the condensation of the 1,2-diaminocyclohexane (22.9 g, 0.2 mol) and methyl 2-hydroxybenzoate (15.2 g, 0.1 mol) stirring at 80-100 °C for 13 h as reported method¹³. Then the white powders were collected by filtration, washed with cooled methanol and dried at room temperature. Yield: 10.9 g, 28.6 %. Elemental analysis (%) calcd. for $\text{C}_{13}\text{H}_{18}\text{N}_2\text{O}_2$: C, 66.64, H, 7.74, N, 11.96; found: C, 67.23, H, 7.21, N, 12.06.

The ligand ($\text{H}_2\text{L} = \text{N}$ -[2-(2-Hydroxy-3-methoxybenzylideneamino)cyclohexyl]-2-hydroxybenzamide), was prepared in a similar manner to the reported procedures^{14,15}. A methanol (20 mL) solution of *o*-vanillin (0.1 mol) was drop-wise added to a methanol solution (20 mL) of N-(2-aminocyclohexyl)-2-hydroxybenzamide (0.1 mol) and the mixture was stirred for 6 h. Then the yellow precipitates of the H_2L ligand was separated and filtered off, washed with 5 mL of cooled methanol and then dried in air. Yield was 65 %. Elemental analysis (%) calcd. for $\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_4$: C, 68.46; H, 6.57; N, 7.60; found C, 68.71; H, 6.26; N, 7.52.

Preparation of complex 1: One pot method was used to synthesize complex **1**, in which the Schiff base formed by the

in situ condensation of N-(2-aminocyclohexyl)-2-hydroxybenzamide (0.1 mmol) and *o*-vanillin (0.2 mmol) was utilized directly to react with $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.1 mmol) in the presence of NEt_3 (0.3 mmol) in 20 mL of methanol solution. The resulting mixture was stirred for 3 h at room temperature to afford brown solution, which was left unperturbed to allow the slow evaporation of the solvent. Black single crystals suitable for X-ray diffraction analysis were formed after about two weeks. Yield was 29.7 % (based on Co). Elemental analysis (%) calcd. for $\text{C}_{52}\text{H}_{61}\text{N}_4\text{O}_{13}\text{Co}$: C, 61.63; H, 6.01; N, 5.48; found C, 61.89; H, 6.09; N, 5.55.

Preparation of complex 2: Complex **2** was obtained by adding $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (0.2 mmol) to a solution of H_2L (0.2 mmol) in 20 mL of methanol and the resulting mixture was stirred for about 3 h to afford a green solution. Two week later, green crystals of the title compound were isolated from the solution. Yield was 30.1 % (based on Cu). Elemental analysis (%) calcd. for $\text{C}_{42}\text{H}_{46}\text{N}_4\text{O}_8\text{Cu}$: C, 63.18; H, 5.81; N, 7.02; found C, 63.34; H, 5.56; N, 7.04.

TABLE-1
CRYSTALLOGRAPHIC DATA AND STRUCTURE
REFINEMENT FOR COMPLEXES **1** AND **2**

Parameters	1	2
Formula	$\text{C}_{52}\text{H}_{61}\text{N}_4\text{O}_{13}\text{Co}$	$\text{C}_{42}\text{H}_{46}\text{N}_4\text{O}_8\text{Cu}$
Formula weight	1044.45	798.38
<i>T</i> /K	185(2)	185(2) K
Wavelength (Å)	0.71073	0.71073
Crystal system	Triclinic	Monoclinic
Space group	$P\bar{1}$	Cc
<i>a</i> (Å)	12.3512 (8)	21.0001 (12)
<i>b</i> (Å)	13.4602 (8)	13.1356 (7)
<i>c</i> (Å)	17.3239 (16)	15.0708 (9)
α (°)	107.411 (2)	90
β (°)	93.1790 (10)	114.3960 (10)
γ (°)	112.8120 (10)	90
<i>V</i> (Å ³)	2484.6 (3)	3786.1 (4)
<i>Z</i>	2	4
Calculated density	1.349	1.397
μ (mm ⁻¹)	0.414	0.637
<i>F</i> (000)	1064	1668
Limiting indices	$-14 \leq h \leq 14$, $-16 \leq k \leq 15$, $-17 \leq l \leq 20$	$-25 \leq h \leq 23$, $-13 \leq k \leq 15$, $-17 \leq l \leq 17$
Data/restraints / parameters	8707/420/640	5753/2/501
Goodness-of-fit on <i>F</i> ²	1.059	1.035
<i>R</i> ₁ , <i>wR</i> ₂ [<i>I</i> > 2 σ (<i>I</i>)] ^{a,b}	<i>R</i> ₁ = 0.0676 <i>wR</i> ₂ = 0.1879	<i>R</i> ₁ = 0.0339 <i>wR</i> ₂ = 0.0766
<i>R</i> indices (all data)	<i>R</i> ₁ = 0.1044 <i>wR</i> ₂ = 0.2141	<i>R</i> ₁ = 0.0432 <i>wR</i> ₂ = 0.0819
Largest diff. peak and hole	2.409 and -0.517 e.Å ⁻³	0.244 and -0.393 e.Å ⁻³

$$^a R = \sum ||F_o| - |F_c|| / \sum |F_o|$$

$$^b wR_2 = [\sum w(F_o^2 - F_c^2)^2 / \sum w(F_o^2)^2]^{1/2}$$

X-Ray crystallography: Crystallographic data were collected on a Bruker Apex II CCD diffractometer with graphite monochromated MoK_α radiation ($\lambda = 0.71073$ Å) at a temperature of 185 (±2) K, for **1** and **2**. Data processing was accomplished with the SAINT processing program. These structures

were solved by the direct methods and refined on *F*² by full-matrix least squares using SHELXTL97^{16,17}. The location of Co/Cu atom was easily determined and O, N and C atoms were subsequently determined from the difference Fourier maps. All non-hydrogen atoms were refined with anisotropic thermal parameters. Crystallographic and refinement details of **1** and **2** are listed in Table-1. CCDC- 915373 (**1**) and 915374 (**2**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre *via* www.ccdc.cam.ac.uk/data_request/cif. [or from the Cambridge Crystallographic Data Centre (CCDC), 12 Union Road, Cambridge CB2 1EZ, UK; fax: (44) 1223-336033; email: deposit@ccdc.cam.ac.uk].

RESULTS AND DISCUSSION

The molecular structure of **1** is displayed in Fig. 1 and all the relevant bond lengths and bond angles are listed in Table-2. The complex **1**, $[\text{Co}(\text{HL})_2(\text{C}_8\text{H}_7\text{O}_3)] \cdot (\text{CH}_3\text{OH})_2$, contains of two HL⁻ ligand, one Co(III) ion, one *o*-vanillin anion and two methanol molecules. The geometry around the Co(III) ion centre can be described as a distorted octahedral geometry composed of the N₂O₄ set of atoms, the HL⁻ ligand is chelated to the Co(III) ion in an O,N-bidentate manner and the *o*-vanillin anion coordinates through the phenolate oxygen atom and aldehyde oxygen. The Co1-O and Co1-N distances are similar to the corresponding distances observed in some similar Co

TABLE-2
BOND LENGTHS (Å) AND ANGLES (°) FOR COMPLEX **1**

Bond lengths			
Co1-O1	1.897(3)	Co1-N1	1.955(4)
Co1-O3	1.903(4)	Co1-N4	1.951(3)
Co1-O8	1.880(3)	Co1-O9	1.893(3)
Bond angles			
O8-Co1-O9	88.03(12)	O8-Co1-O3	94.46(15)
O8-Co1-O1	85.92(12)	O9-Co1-O3	87.23(14)
O9-Co1-O1	171.95(13)	O1-Co1-O3	87.97(14)
O8-Co1-N4	176.68(15)	O9-Co1-N4	92.19(13)
O1-Co1-N4	94.16(13)	O3-Co1-N4	88.86(16)
O8-Co1-N1	84.42(14)	O3-Co1-N1	94.01(14)
O1-Co1-N1	90.67(14)	O3-Co1-N1	178.30(14)
N4-Co1-N1	92.26(15)		




Fig. 1. A view of the complex **1**, showing the atomic numbering scheme, methanol molecules are omitted for clarity

complexes^{18,19}. The structure shows distortion from perfect octahedral geometry with respect to bond angles. The trans base angles [N1-Co1-O3 = 178.30 (14)°, N4-Co1-O8 = 176.68 (15)°] are all significantly less than the ideal value of 180°. The axial angle O1-Co1-O9 [171.95(13)°] deviates significantly from the ideal 180°. This angle of distortion can be attributed to the ligand structure.

The present complex **2** of [Cu(HL)₂], together with the molecular structure is displayed in Fig. 2. Selected bond lengths and angles are given in Table-3. The title complex **2** contains of two HL⁻ ligand and one Cu(II) ion. The geometry around the Cu(II) centre can be described as a distorted square-planar composed of the N₂O₂ set of atoms, every HL⁻ ligand is chelated to the Cu atom in an O,N-bidentate manner. The Cu1-O and Cu1-N distances are similar to the corresponding distances observed in some similar Cu complexes²⁰⁻²². There is slightly deviation of the metal centre from the N₂O₂ equatorial plane. The maximum displacements from the least-squares plane through N2, N4, O4 and O8 are 0.5447 (22) Å and 0.3598 (18) Å for atoms N4 and O8, respectively, Cu1 is 0.1208 (2) Å below this plane.




Fig. 2. A view of the complex **2**, showing the atomic numbering scheme

TABLE-3
BOND LENGTHS (Å) AND ANGLES (°) FOR COMPLEX **2**

Bond lengths			
Cu1-O8	1.886 (6)	Cu1-N2	1.969 (7)
Cu1-O4	1.891 (5)	Cu1-N4	1.998 (7)
Bond angles			
O8-Cu1-O4	151.94 (7)	O8-Cu1-N2	92.4 (3)
O4-Cu1-N2	92.3 (3)	O8-Cu1-N4	93.2 (3)
O4-Cu1-N4	91.2 (3)	N2-Cu1-N4	91.2 (3)

The ligand H₂L is a potential multidentate ligand with N₂O₄ donor sites but in this case it acts as a bidentate chelate. In complex **1** and complex **2**, the HL⁻ ligand is chelated to the metal ions only in an O,N-bidentate manner, which from the *o*-vanillin group. 2-Hydroxybenzamide group is not effected the coordination metal cation, thus the angle of both aromatic rings in each ligand is similarly. In complex **1**, both aromatic rings in each ligand surrounding the metal ions form a dihedral

angle that is 78.40 (16)°. The angle is similar to complex **2** of 78.13 (9)°.

Conclusion

Reactions of a multidentate N-[2-(2-Hydroxy-3-methoxybenzylideneamino)cyclohexyl]-2-hydroxy-benzamide (H₂L) with CoCl₂·6H₂O and Cu(NO₃)₂·3H₂O in methanol solvent results in the formation of two mononuclear complexes, [Co(HL)₂(C₈H₇O₃)]·(CH₃OH)₂ (**1**) and [Cu(HL)₂] (**2**). Both complexes are characterized by elemental analysis and crystal structure analysis. The metal centers of complexes **1** and **2** have exhibit a distorted octahedral geometry and square-planar geometry, respectively. The multidentate HL⁻ ligand is chelated to the metal ions only in an O,N-bidentate manner, which from the *o*-vanillin group, thus it is a pity that no synthetic polynuclear metal complexes.

ACKNOWLEDGEMENTS

The authors thank the project supported by the Development and Reform Commission of Jilin Province, the Science and Technology Burrau of Changchun City, the Science and Technology Department of Jilin Province, China (2011174) and Changchun University of Science and Technology for their financial support.

REFERENCES

- S.H. Li, S.K. Gao, S.X. Liu and Y.N. Guo, *Cryst. Growth Des.*, **10**, 495 (2010).
- H. Yin and S.X. Liu, *J. Mol. Struct.*, **918**, 165 (2009).
- L.F. Zou, L. Zhao, Y.N. Guo, J.K. Tang, Q.L. Wang and Y.H. Li, *Inorg. Chim. Acta*, **382**, 65 (2012).
- J.P. Costes, F. Dahan and F. Nicodeme, *Inorg. Chem.*, **40**, 5285 (2001).
- J. Tang, I. Hewitt, N.T. Madhu, G. Chastanet, W. Wernsdorfer, C.E. Anson, C. Benelli, R. Sessoli and A.K. Powell, *Angew. Chem. Int. Ed.*, **45**, 1729 (2006).
- Y. Elerman, A. Elmali and I. Svoboda, *Z Naturforsch B*, **57**, 519 (2002).
- C.D. Papadopoulos, A.G. Hatzidimitriou, G.P. Voutsas and M. Lalia-Kantouri, *Polyhedron*, **26**, 1077 (2007).
- T. Hamamatsu, K. Yabe, M. Towatari, N. Matsumoto, N. Re, A. Pochaba and J. Mrozinski, *Bull. Chem. Soc. Jpn.*, **80**, 523 (2007).
- G.M. Yu, L. Zhao, L.F. Zou, Y.N. Guo, G.F. Xu, Y.H. Li and J.K. Tang, *J. Chem. Crystallogr.*, **46**, 606 (2011).
- E.A. Velcheva, B.A. Stamboliyska and P.J. Boyadjieva, *J. Mol. Struct.*, **963**, 57 (2010).
- L.K. Thompson, C.J. Matthews, L. Zhao, Z.Q. Xu, D.O. Miller, C. Wilson, M.A. Leech, J.A.K. Howard, S.L. Heath, A.G. Whittaker and R.E.P. Winpenny, *J. Solid State Chem.*, **159**, 308 (2001).
- M. Koikawa, H. Yamashita and T. Tokii, *Inorg. Chim. Acta*, **357**, 2635 (2004).
- R. Mitsuhashi, T. Suzuki, Y. Sunatsuki and M. Kojima, *Chem. Lett.*, **40**, 696698 (2011).
- Y. Qiao, X.P. Ju, Z.Q. Gao and L.Q. Kong, *Acta Cryst. E*, **66**, O95 (2009).
- O. Pouralimardan, A.-C. Chamayou, C. Janiak and H. Hosseini-Monfared, *Inorg. Chim. Acta*, **360**, 1599 (2007).
- G.M. Sheldrick, SHELXS-97, Program for Crystal Structure Solution, University of Göttingen, Germany (1997).
- G.M. Sheldrick, SHELXL-97, Program for Crystal Structure Refinement, University of Göttingen, Germany (1997).
- M. Fleck, D. Karmakar, M. Ghosh, A. Ghosh, R. Saha and D. Bandyopadhyay, *Polyhedron*, **34**, 157 (2012).
- A. Ray, S. Banerjee, G.M. Rosair, V. Gramlich and S. Mitra, *Struct. Chem.*, **19**, 459 (2008).
- H. Ali, N.A. Khamis, W.J. Basirun and B.M. Yamin, *Acta Cryst. E*, **60**, m982 (2004).
- A. Elmali, Y. Elerman and I. Svoboda, *Acta Cryst.*, **C56**, 423 (2000).
- L.F. Zou, X.Y. Yang, Y. Gao, H.B. Yao and Y.H. Li, *Acta Cryst. E*, **67**, m511 (2011).