View Article Online

ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: T. Morofuji, T. Yoshida, R. Tsutsumi, M. Yamanaka and N. Kano, *Chem. Commun.*, 2020, DOI: 10.1039/D0CC05830K.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the Information for Authors.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



rsc.li/chemcomm

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 13 October 2020. Downloaded on 10/14/2020 11:57:36 AM.

Arylation of aryllithiums with S-arylphenothiazinium ions for biaryl synthesis

Received 00th January 20xx. Accepted 00th January 20xx

Tatsuya Morofuji,*a Tatsuki Yoshida,a Ryosuke Tsutsumi,^b Masahiro Yamanaka^b and Naokazu Kano*^a

DOI: 10.1039/x0xx00000x

Aryllithiums are one of the most common and important aryl nucleophiles; nevertheless, methods for arylation of aryllithums to produce biaryls have been limited. Herein, we report arylation of aryllithiums with S-arylphenothiazinium ions through selective ligand coupling of intermediary sulfuranes. Various unsymmetrical biaryls could be obtained without transition-metal catalysis.

Aryllithiums are one of the most common and important aryl nucleophiles in organic chemistry. Aryllithiums are known to react with various electrophiles, such as alkyl halides, carbonyl compounds, chlorosilanes and trimethoxyborane.¹ However, methods for anylation of anyllithums to produce biaryls are very limited. Arylation of aryllithiums based on nucleophilic aromatic substitution² or nucleophilic attack to aryne³ was reported, but there were few options regarding the applicable substrates. Although a few palladium-catalyzed methods have been reported,⁴ development of electrophilic arylation reagents that react with aryllithiums in the absence of transition metals is a simple but challenging problem (Figure 1a).⁵

In sulfur chemistry. the ligand coupling of tetraarylsulfuranes (IUPAC nomenclature: tetraaryl-λ⁴sulfanes) has been reported.⁶ Reaction of triarylsulfoniums with arylmetal reagents including aryllithiums affords the tetraarylsulfurane intermediate, which decomposes to the corresponding biaryls and diaryl sulfides without transitionmetal catalysis (Figure 1b).⁷ However, these processes have very limited applications because the selectivity of the resulting products depends on aromatic substituents on the sulfur atom.7b,f One of the reasons why control of the selectivity of the ligand coupling is difficult is that conformation of sulfurane is not fixed through Berry's pseudorotation.7ª To obtain unsymmetrical biaryls, previous ligand coupling was limited to

use electron-poor heteroaromatic rings such as pyridine, azole, or quinoline as coupling partner, which promote desired ligand coupling.7i,8 In contrast, the selective synthesis of unsymmetrical biphenyl derivatives from sulfonium salts and aryl metals is a long-standing problem.

Herein, we report that S-arylphenothiazinium ions can solve the problem and be used as versatile arylation reagents for aryllithiums. The reaction of aryllithiums 1 with Sarylphenothiazinium ions 2 selectively afforded the corresponding biaryls 3 and phenothiazine (Figure 1c). In contrast to the transition-metal-catalyzed approach, bromo-

a) blueprint: development of arylation reagents for aryllithiums









Selectivity depends on substrates





Figure 1. Arylation of aryllithiums. (a) Our blueprint: development of arylation reagents for aryllithiums. (b) Previous work: Reactivity of sulfurane and its limitation for the synthesis of biaryl. (c) This work: S-arylphenothiazinium ions used as electrophilic arylation reagents for synthesizing unsymmetrical biaryls.

^{a.} Department of Chemistry, Faculty of Science, Gakushuin University, 1-5-1 Mejiro, Toshima-ku, Tokyo 171-8588, Japan.

^{b.} Department of Chemistry, Faculty of Science, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan.

⁺ Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: Experimental procedures, DFT calculations, and NMR data. See DOI: 10.1039/x0xx00000x

Journal Name

and iodo-substituted biaryls, which are attractive synthetic intermediates, could be synthesized using the present method. A computational study was performed to reveal the reaction mechanism of this sulfurane-mediated biaryl-formation.

First, we examined the reaction of some cyclic sulfonium ions⁹ with phenyllithium (Table 1). Reaction of Sphenylphenothiazinium hexafluorophosphate (2a) with phenyllithium (1a) in THF at -78 °C and 23 °C afforded biphenyl (3a) and N-methylphenothiazine (4) in good yields (entry 1). When the amount of 1a used was reduced, the conversion of 2a decreased (entry 2). In contrast, the reaction of Sphenylphenoxathiinium 5 with phenyllithium (1a) afforded a mixture of biphenyl (3a), phenoxathiin (6), and ring-opening product 7a (entry 3). As previously reported, the reaction of Sphenylthianthrenium 8 with phenyllithium (1a) under the same conditions afforded biphenyl (3a) in only 8% yield, while dibenzothiophene (9) (55%), diphenyl sulfide (10) (51%), and ring-opening product 7b (31%) were obtained in higher yields 4).10 lt was also reported (entrv that Sphenyldibenzothiophenium (11) afforded ring-opening product 7c in a quantitative yield (entry 5).¹¹ It should be noted that ring opening product and N-methylcarbazole were not observed by ¹H NMR analysis when 2a was used. These results suggest that the phenothiazine moiety is essential for successful formation

 Table 1. Reaction of phenyllithium (1a) with sulfonium salts.

This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

Open Access Article. Published on 13 October 2020. Downloaded on 10/14/2020 11:57:36 AM.



Phenyllithium (**1a**) (0.6 mol) was added to sulfonium salt (0.2 mmol) in THF (5 mL) at -78 °C. The reaction mixture was stirred at -78 °C for 30 min and 23 °C for 1 h. ^{*a*} Determined by ¹H NMR spectroscopy. ^{*b*} 0.4 mmol of **1a** was used. ^{*c*} Isolated yield. ^{*d*} Result from ref **11**.

The scope of the arylation of aryllithiums using Sarylphenothiazinium 2 investigated in the present study is shown in Scheme 1. To perform desymmetrizative monoarylation of inexpensive 1,4-dibromobenzene, it was lithiated with *n*-BuLi and reacted with various S-arylphenothiaziniums successively to obtain the desired bromo-substituted biaryls (3b-3g) in good yields. Significantly, the reaction of S-(4bromophenyl)phenothiazinium with aryllithiums bearing a bromo-substituent also afforded mono-arylated products bearing two bromo groups, 3h, 3i, and 3j. Other halogensubstituted biaryls could also be synthesized by the present method (3k-3o). The resulting bromo- or iodo-substituted biaryls are synthetically attractive intermediates for further derivatization by transition-metal-catalyzed cross-coupling reactions. It should be noted that bromo- or iodo-substituted biaryls have never been synthesized by Murahashi coupling because C-C bond formation takes place at the halogensubstituted position.⁴ Furthermore, various unsymmetrical biaryls were synthesized using the present method (3p-3z). Reactions using *p*-chlorophenyllithium produced the desired products in good yields (3p-3r), whereas the use of aryllithiums bearing a methylchalcogeno group slightly decreased the yields of the product (3s, 3t). Notably, the reaction efficiently produced 3p on the gram scale. Ortho- or meta-methylsubstituted aryllithium was applicable to the present reaction system (**3u**, **3v**). Reactions of π -extended aryllithiums with Sarylphenothiaziniums also afforded the corresponding unsymmetrical biaryls in good yields (3w-3z).

To gain insight into the selective formation of unsymmetrical biaryls from the S-arylphenothiazinium ions, density functional theory (DFT) calculations were conducted at the B3LYP/6-311G+(d,p) level. Possible reaction pathways of phenyllithium(THF)₃ (**1a**) with S-phenylphenothiazinium (**2a**) via the corresponding sulfurane were explored. The relative free energy profiles of the formation of biphenyl (path A), ringopening reaction (path B), and formation of diphenyl sulfide (path C) from tetracoordinated sulfuranes are summarized in Figure 3a. Three types of sulfurane intermediates (IM A, IM B, and IM C) afforded the corresponding transition states (TS A, TS B, and TS C). Each sulfurane intermediate has a different pseudo-trigonal bipyramidal structure and similar free energy. Interconversion between IM A, IM B, and IM C occurs easily through pseudorotation.^{7a} Therefore, the selectivity of the reaction products is determined by the TS, not by the relative stability of IM. TS A is energetically more stable than TS B and TS C, which is in good agreement with the experimental results, as described in Table 1.

DFT calculations using S-phenylphenoxathiinium (4), Sphenylthianthrenium (6), or S-phenyldibenzothiophenium (7) instead of 2a indicated that biphenyl formation was not energetically preferred to ring opening-reaction and formation of diphenyl sulfide when 4, 6, or 7 was used (see SI for details). These computational results are consistent with the experimental results. To understand the major factor in the preference of **path A** for S-phenylphenothiazinium ion, we



Scheme 1. Scope of cross-coupling of aryllithiums and S-arylphenothiazinium ions for synthesizing biaryls.^{*a*} S-Arylphenothiazinium 2 (0.2 mmol) was added to a solution of aryllithium 1 (0.6 mmol) in THF (5 mL) at -78 °C. The reaction mixture was stirred at -78 °C for 30 min and 23 °C for 1 h. See SI for details.^{*a*} Isolated yield. ^{*b*} The reaction was performed on a gram scale.

compared the structural differences between **IM A** and **TS A** with their analogs (Figure 3b). The bond lengths of apical bonds III and IV in each transition state are almost the same. In contrast, apical bonds I and II in **IM A** are longer and shorter, respectively, than those in **IM D**, **IM G**, and **IM J**. **IM A** has a

COMMUNICATION

relatively close structure to **TS A**. This indicates that binned formation from the *S*-arylphenothiazinium long (path A) requires a smaller structural distortion with a lower energy loss than that from other sulfonium ions. Such a structural difference is due to π -electron donation by the lone pair on the nitrogen atom, which would strengthen and shorten apical C–S bond II (Figure 3c). In general, when one apical bond is strengthened, the other apical bond becomes weaker and longer.¹² Therefore, apical bond I in IM A is longer than that in others. This structural difference of IM A can be favorable for biaryl formation from phenyllithium (1a) and *S*-phenylphenothiazinium ion 2a.

In summary, we developed *S*-arylphenothiazinium ions as efficient arylation reagents for aryllithiums. The reaction can provide various unsymmetrical biaryls without transition-metal catalysis. Notably, transformable halogen substituents such as bromo- and iodo-groups are tolerated. The resulting halogensubstituted biaryls are attractive synthetic intermediates. This result contrasts sharply with that obtained through palladiumcatalyzed Murahashi coupling. In contrast to other similar sulfonium salts, *S*-arylphenothiazinium ions are specifically suitable for biaryl synthesis through selective ligand coupling of a sulfurane intermediate, which is consistent with theoretical calculations. We are currently developing new transition-metalfree arylation reactions using *S*-arylphenothiazinium ions; the results will be presented in due course.

This research was partially supported by the MEXTsupported program for the Strategic Research Foundation at Private Universities, and the Society of Iodine Science. DFT calculations were performed using Research Center for Computational Science, Okazaki, Japan.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 The Chemistry of Organolithium Compounds. Elsevier: Pergamon, 1974.
- (a) N. Koike, T. Hattori, A. Takeda, Y. Okaishi, S. Miyano, *Chem. Lett.*, 1997, **26**, 641; (b) J.-M. Becht, A. Gissot, A. Wagner, C. Mioskowski, *Chem. Eur. J.*, 2003, **9**, 3209.
- (a) J.-A. García-López, M. F. Greaney, *Chem. Soc. Rev.*, 2016,
 45, 6766; (b) F. Leroux, M. Schlosser, *Angew. Chem. Int. Ed.*,
 2002, 41, 4272; (c) T. Truong, M. Mesgar, K. K. A. Le, O. Daugulis, *J. Am. Chem. Soc.*, 2014, 136, 8568.
- 4 (a) S. Murahashi, M. Yamamura, K. Yanagisawa, N. Mita, K. Kondo, J. Org. Chem., 1979, 44, 2408; (b) A. Nagaki, A. Kenmoku, Y. Moriwaki, A. Hayashi, J.-I. Yoshida, Angew. Chem. Int. Ed., 2010, 49, 7543; (c) M. Giannerini, M. Fananas-Mastral, B. L. Feringa, Nat. Chem., 2013, 5, 667; (d) E. B. Pinxterhuis, M. Giannerini, V. Hornillos, B. L. Feringa, Nat. Commun., 2016, 7, 11698; (e) D. Heijnen, F. Tosi, C. Vila, M. C. A. Stuart, P. H. Elsinga, W. Szymanski, B. L. Feringa, Angew. Chem. Int. Ed., 2017, 56, 3354.
- 5 For selected examples of transition-metal-free biarylsynthesis, see (a) C.-L. Sun, Z.-J. Shi, *Chem. Rev.*, 2014, **114**, 9219; (b) T. Dohi, M. Ito, K. Morimoto, M. Iwata, Y. Kita, *Angew. Chem. Int. Ed.*, 2008, **47**, 1301; (c) T. Morofuji, A. Shimizu, J.-I. Yoshida, *Angew. Chem. Int. Ed.*, 2012, **51**, 7259;

COMMUNICATION



Figure 3. DFT calculations of reaction of phenyllithium (1a) and S-phenylphenothiazinium ion (2a) at the B3LYP/6-311G+(d,p) level. (a) Relative free energy profiles. (b) Selected calculated bond lengths (Å) or interatomic distances (Å) of sulfuranes and transition states for biphenyl formation. (c) A plausible reason for the structural difference in sulfurane IM A.

- (d) E. Shirakawa, Y. Hayashi, K.-I. Itoh, R. Watabe, N. Uchiyama, W. Konagaya, S. Masui, T. Hayashi, Angew. Chem. Int. Ed., 2012, 51, 218; (e) M. Ito, H. Kubo, I. Itani, K. Morimoto, T. Dohi, Y. Kita, J. Am. Chem. Soc., 2013, 135, 14078; (f) T. Yanagi, S. Otsuka, Y. Kasuga, K. Fujimoto, K. Murakami, K. Nogi, H. Yorimitsu, A. Osuka, J. Am. Chem. Soc., 2016, 138, 14582; (g) M. C. Hilton, X. Zhang, B. T. Boyle, J. V. Alegre-Requena, R. S. Paton, A. McNally, Science, 2018, 362, 799; (h) F. Kloss, T. Neuwirth, V. G. Haensch, C. Hertweck, Angew. Chem. Int. Ed., 2018, 57, 14476; (i) A. Music, A. N. Baumann, P. Spieß, A. Plantefol, T. C. Jagau, D. Didier, J. Am. Chem. Soc., 2020, 142, 4341.
- 6 (a) W. A. Sheppard, J. Am. Chem. Soc., 1971, 93, 5597; (b) S. Ogawa, Y. Matsunaga, S. Sato, I. Iida, N. Furukawa, J. Chem. Soc., Chem. Commun., 1992, 1141.
- 7 (a) Ligand Coupling Reactions with Heteroatomic Compounds. Elsevier: Pergamon, 1998; (b) Y. H. Khim, S. Oae, Bull. Chem. Soc. Jpn., 1969, 42, 1968; (c) B. Trost, R. LaRochelle, R. Atkins, J. Am. Chem. Soc., 1969, 91, 2175; (d) R. W. LaRochelle, B. M. Trost, J. Am. Chem. Soc., 1971, 93, 6077; (e) B. M. Trost, H. C. Arndt, J. Am. Chem. Soc., 1973, 95, 5288; (f) S. Oae, Croatica Chemica Acta, 1986, 59, 129; (g) M. Hori, T. Kataoka, H. Shimizu, M. Ikemori, Y. Aoyama, J. Chem. Soc., Perkin Trans. 1, 1988, 1209; (h) S. Oae, Y. Uchida, Acc. Chem. Res., 1991, 24,

202; (i) S. Oae, H. Ishihara, M. Yoshihara, *Chem. Heterocycl. Compd.*, 1995, **31**, 917; (j) S. Sato, O. Takahashi, N. Furukawa, *Coord. Chem. Rev.*, 1998, **176**, 483; (k) W. M. Dean, c M. Šiaučiulis, T. E. Storr, W. Lewis, R. A. Stockman, *Angew. Chem. Int. Ed.*, 2016, **55**, 10013; (I) D. L. Chen, Y. Sun, M. Chen, X. Li, L. Zhang, X. Huang, Y. Bai, F. Luo, B. Peng, *Org. Lett.*, 2019, **21**, 3986.

- 8 (a) T. Kawai, N. Furukawa, S. Oae, *Tetrahedron Lett.*, 1984, 25, 2549; (b) N. Furukawa, T. Shibutani, H. Fujihara, *Tetrahedron Lett.*, 1987, 28, 5845; (c) M. Zhou, J. Tsien, T. Qin, *Angew. Chem. Int. Ed.*, 2020, 59, 7372
- 9 L. Racicot, T. Kasahara, M. A. Ciufolini, *Org. Lett.*, 2014, **16**, 6382.
- 10 B. Boduszek, H. J. Shine, T. K. Venkatachalam, *J. Org. Chem.*, 1989, **54**, 1616.
- 11 S. Sato, N. Furukawa, Tetrahedron Lett., 1995, 36, 2803.
- 12 L. J. Adzima, E. N. Duesler, J. C. Martin, *J. Org. Chem.*, 1977, **42**, 4001.

Journal Name