

# Communication

# **Biomimetic Chirality Sensing with Pyridoxal-5'-phosphate**

Samantha L. Pilicer, Pegah R. Bakhshi, Keith W Bentley, and Christian Wolf J. Am. Chem. Soc., Just Accepted Manuscript • Publication Date (Web): 27 Jan 2017 Downloaded from http://pubs.acs.org on January 27, 2017

# **Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

# Biomimetic Chirality Sensing with Pyridoxal-5'-phosphate

Samantha L. Pilicer, Pegah R. Bakhshi, Keith W. Bentley and Christian Wolf\*

Department of Chemistry, Georgetown University, Washington, DC 20057, USA

Supporting Information Placeholder

**ABSTRACT:** Pyridoxal-5'-phosphate (PLP) is introduced to a biomimetic indicator displacement assay for simultaneous determination of the absolute configuration, enantiomeric composition and concentration of unprotected amino acids, amino alcohols and amines. The chiroptical assay is based on fast imine metathesis with a PLP aryl imine probe to capture the target compound for circular dichroism and fluorescence sensing analysis. The substrate binding yields characteristic Cotton effects that provide information about the target compound ee and the synchronous release of the indicator results in a nonenantioselective off-on fluorescence response that is independent of the enantiomeric sample composition and readily correlated to the total analyte concentration.

Pyridoxal-5'-phosphate (PLP, the active form of Vitamin  $B_6$ ) is an indispensable and ubiquitous enzyme cofactor that plays a central role in transamination, racemization, elimination and many other reactions.1 The remarkable predominance and versatility of PLP in nature has inspired the development of intriguing examples of biomimetic catalysis,<sup>2</sup> continuous flow chemistry,<sup>3</sup> combinatorial biosynthesis of complex target compounds,<sup>4</sup> biomolecule labeling or protein conjugation<sup>5</sup> and other applications.<sup>6</sup> Despite the myriad of pyridoxal-5'-phosphate-dependent enzymes and the variety of biological processes, the propensity of the PLP aldehyde group toward Schiff base formation with primary amino functions is a strikingly unifying leitmotif and the PLP-imine moiety is a widely encountered resting state in natural and biomimetic catalytic reactions. Despite its wide utilization and general understanding of its properties and functions, PLP has not been exploited for molecular recognition and quantitative chemosensing of chiral compounds.

The general importance of chiral amino acids, amines and amino alcohols in biological processes and asymmetric synthesis has directed increasing attention to optical methods that are capable of high-throughput screening (HTS) of the enantiomeric composition of small sample amounts.<sup>7</sup> In this regard, time-efficiency, cost, and compatibility with automation and microscale analysis are among the most important considerations whereas error margins of up to 10% ee have been considered acceptable for HTS applications.<sup>8</sup> The concept of covalent substrate fixation with a carefully designed aldehyde or ketone probe and subsequent enantiomeric excess (ee) determination of the bound chiral amine by UV, fluorescence or circular dichroism measurements has been successfully exploited by Kim, Hong and Chin,<sup>9</sup> Anslyn,<sup>10</sup> Pu,<sup>11</sup> Joyce<sup>12</sup> and our group.<sup>13</sup> Many of the chemosensors introduced to date, however, have a limited application scope and require elaborate synthesis. More importantly, the chiroptical determination of both enantiopurity *and* concentration of a chiral target compound remains a difficult task.<sup>14</sup>

PLP Biological functions and bioorganic applications:



Figure 1. PLP applications and illustration of the chirality sensing concept.

We now introduce a cost-effective approach to quantitative chirality sensing using readily available PLP imines as probe (Figure 1). At the onset of this study, we rationalized that condensation of PLP with an aromatic amine would afford a fast and broadly useful chirality sensor carrying an integrated indicator unit. This biomimetic sensing strategy relies on covalent binding of a chiral amino acid or another target with a primary amino group and concomitant chirality amplification onto the PLP imine scaffold to generate a circular dichroism signal that can be correlated to the ee and absolute configuration of the substrate. The binding event interconverts the chemosensor, an aromatic PLP aldimine, to a more stable aliphatic imine - a reaction that is thermodynamically favored. Because Schiff base formation and the imine exchange reaction are accelerated by the presence of the phenol and phosphate groups,<sup>15</sup> the use of PLP as chirality probe also addresses the need for time-efficiency in HTS applications. The imine metathesis was therefore expected to be fast, quantitative and a stoichiometric process, that is, for each bound target compound one aryl amine molecule, which we refer to as the indicator, is released. Importantly, this displacement reaction is independent of the chirality of the substrate and it consequently allows concentration analysis based on the fluorescence signal of the free indicator regardless of the enantiopurity of the target compound as previously shown by Yu and Pu.<sup>nc</sup> Altogether, this biomimetic sensing approach enables one to determine both the enantiomeric excess and the total amount of amino acids, amino alcohols and amines by a simple indicator displacement assay (IDA) and fast CD and fluorescence measurements.<sup>16</sup>



**Figure 2.** Structures of the PLP imine sensors **1-3** and nine-teen amino acids, amino alcohols and amines tested.

It has been reported that free PLP and pyridoxal Schiff bases exist as complex equilibria of multiple tautomeric forms with often indistinguishable UV absorptions that are highly sensitive to the solvent polarity, pH, ionic strength, the presence of counterions, solvation effects and other environmental factors.<sup>17</sup> This renders efforts to use UV spectroscopy for reliable quantification of the imine metathesis impractical. Because both PLP and PLP imines typically have low quantum yields it was ruled out to accomplish this by direct fluorescence analysis and we therefore resorted to an indicator displacement assay.<sup>18</sup> With this in mind, we began our study with the synthesis of PLP Schiff bases exhibiting an aromatic imine moiety that would (a) be irreversibly displaced by the more nucleophilic aliphatic sensing target and (b) generate a characteristic fluorescence signal suitable for quantitative analysis upon cleavage from the PLP sensor scaffold (Figure 2 and SI).



Figure 3. Chirality sensing of 4 (600  $\mu$ M in MeOH), 7 (480  $\mu$ M), 15 (450  $\mu$ M) and 21 (360  $\mu$ M) with PLP imine 1.

Initially, we screened solvents and bases to dissolve the PLP imine sensors and the targeted amino acids and to achieve homogeneous conditions for the imine metathesis at millimolar concentrations. We found that the desired indicator displacement occurs within a few minutes in the presence of KOH using methanol as solvent. The reaction mixtures obtained from sensor 1 and the amino acids 4-12 were then diluted with methanol to submillimolar concentrations and directly subjected to CD analysis. We were pleased to find characteristic Cotton effects in all cases including aliphatic substrates such as methionine (Figure 3 and SI). Essentially the same results were obtained with the amino alcohols 13-18 and amines 19-21.

The imine metathesis was further investigated by <sup>1</sup>H NMR and CD spectroscopy (SI). Mixing of sensor 1 and cyclohexylethylamine, 19, in methanolic solution in the presence of stoichiometric amounts of TBAOH showed smooth displacement of the aniline moiety by the substrate within 45 seconds (Figure 4). The reaction coincides with a clear upfield shift of the formylimine proton from 9.39 to 9.03 ppm and of the aromatic PLP proton from 7.99 to 7.82 ppm. As expected, the cleaved indicator has upfield shifted NMR signals that are identical with free aniline and the PLP methylene and methyl groups shift from 5.12 to 4.99 ppm and 2.50 to 2.43 ppm, respectively. The PLP binding of 19 results in a down field shift of the methine proton attached to the chiral center from 2.58 to 3.50 ppm. Importantly, the imine metathesis occurs quantitatively and without hydrolysis toward free PLP which would be clearly visible by the appearance of the PLP aldehyde signal at approximately 10.2 ppm. As a result, the indicator displacement is a perfectly stoichiometric process, i.e. the quantity of the released aniline corresponds exactly to that of the bound amine target. The generation of equimolar indicator amounts thus sets the stage for determination of the concentration of the sensing target independent of the enantiomeric composition, vide infra.



**Figure 4.** Proton NMR analysis of the metathesis using PLP sensor 1 and amine 19. 1) Cyclohexylethylamine, 19; 2) free aniline; 3) PLP-bound cyclohexylethylamine after mixing of 1 and 19 for 45 seconds; 4) PLP sensor 1. All NMR spectra were recorded at 23 mM (both 1 and 19) after addition of 1 equivalent of TBAOH (1 M in MeOH) using CD<sub>3</sub>OD as solvent. •MeOH; •TBAOH; \*TBAOH-impurity.

58

59 60

Having established conceptual proof for the proposed imine metathesis and chirality sensing of all 19 target compounds with the prototype sensor 1 we continued with evaluating the possibility of quantitative ee and concentration analysis using the aminonaphthalene analogues 2 and 3. Nonracemic tryptophan samples were applied to the indicator displacement assay with each sensor and the CD amplitudes measured at 416 nm were plotted against the enantiomeric excess. As expected, we obtained linear responses for both sensors (Figure 5 and SI). The known CD response of PLP sensor 3 was then used to analyze nine Trp samples with random enantiomeric composition. Four of these samples contained the D-isomer in 87%, 78%, 54% and 30% ee. We were pleased to find that the chiroptical analysis based on the indicator displacement assay with 3 gave 88%, 80%, 58% and 32% ee, respectively. The remaining five samples had the L-isomer of Trp in 14%, 38%, 44%, 56% and 85% ee and our sensing results were 20%, 41%, 49%, 56% and 86% ee. The results show that the sign of the Cotton effects provide the absolute configuration of the major Trp enantiomer while the CD amplitudes measured allow calculation of accurate ee values.



**Figure 5.** Top: CD effects observed with nonracemic samples of tryptophan and sensor **3** in MeOH at 480  $\mu$ M (left). Linear relationship between the CD response of **3** and Trp %ee (right). The CD signals observed for Trp samples of random enantiomeric composition are shown in red. Bottom: Off-on fluorescence response to the imine metathesis and indicator release using **3** and Trp. The fluorescence spectra were obtained at 24  $\mu$ M in MeOH,  $\lambda_{exc.}$  325 nm.

As demonstrated above, the binding of enantiomeric substrates to PLP generates characteristic Cotton effects of opposite sign at high wavelengths that can be used to identify the absolute configuration and – when nonracemic samples are investigated – the intensity of the CD response yields the enantiomeric composition. While the CD sensor response is consequently enantioselective the release of the aminonaphthalene indicator is not and therefore suitable for total concentration analysis irrespective of the sample ee. We noticed that the PLP imines exhibit very low fluorescence signals whereas excitation of the free indicator at 325 nm gave a strong fluorescence signal at approximately 430 nm (Figure 5). The insignificant background generated by the PLP imine

sensor thus results in an off-on fluorescence response to the presence of a substrate. To test if the CD and fluorescence responses can be exploited for simultaneous ee and concentration analysis we prepared several Trp samples of varying enantiomeric composition and amount. The samples were then employed in our indicator displacement assay and analyzed as described above using the equations from Figure 5 (Table 1). Comparison of the actual with the experimentally determined values show that the error margins are within the typical range that is acceptable for high-throughput screening purposes. For example, chiroptical sensing of a sample containing a 5.16 mM solution of D-Trp in 93% ee correctly confirmed that the D-form was the major enantiomer and present in 90% ee at 5.35 mM (entry 5). The PLP sensing assay can also be used for ee analysis of analyte mixtures which is demonstrated with nonracemic 5 and 16 (SI).

# Table 1. Comprehensive chirality sensing of tryptophan samples.<sup>a</sup>



<sup>a</sup>Circular dichroism and fluorescence sensing were performed in MeOH at submillimolar concentrations after mixing the sensor and the Trp samples for 15-20 minutes as described in Figure 5.

In summary, we have introduced a biomimetic indicator displacement assay that exploits readily available pyridoxal phosphate derived aryl imines for simultaneous determination of the absolute configuration, enantiomeric composition and concentration of unprotected amino acids, amino alcohols and amines. The chiroptical assay is based on irreversible imine metathesis and facile circular dichroism and fluorescence measurements. The binding of the substrate by the PLP receptor generates characteristic Cotton effects that identify the major enantiomer and the sample ee. The concomitant indicator displacement results in a nonenantioselective off-on fluorescence response that is independent of the enantiomeric sample composition and readily correlated to the total analyte concentration. The sensing assay is broadly applicable - 19 chiral compounds were successfully tested - and it gives results with sufficient accuracy for HTS purposes.

# ASSOCIATED CONTENT

#### Supporting Information

Experimental procedures, compound characterization and sensing details. This material is available free of charge via the Internet at http://pubs.acs.org.

### AUTHOR INFORMATION

# **Corresponding Author**

Cw27@georgetown.edu

## Notes

1

The authors declare no competing financial interests.

## ACKNOWLEDGMENT

This material is based upon work supported by the NSF under CHE-1464547. PRB thanks Georgetown University for a Lisa J. Raines Research Fellowship.

# REFERENCES

1 a) Metzler, D. E. I. M.; Snell, E. E. J. Am. Chem. Soc. 1954, 76, 648-652. b) Griswold, W. R.; Toney, M. D. J. Am. Chem. Soc. 2011, 133, 14823-14830. c) Lai, R.-Yi; Huang, S.; Fenwick, M. K.; Hazra, A.; Zhang, Y.; Rajashankar, K.; Philmus, B.; Kinsland, C.; Sanders, J. M.; Ealick, S. E.; Begley. T. P. J. Am. Chem. Soc. 2012, 134, 9157-9159. d) Chan-Huot, M. Dos, A.; Zander, R.; Sharif, S.; Tolstoy, P. M.; Compton, S.; Fogle, E.; Toney, M. D.; Shenderovich, I.; Denisov, G. S.; Limbach, H.-H. J. Am. Chem. Soc. 2013, 135, 18160-18175. e) Shoji, M.; Hanaoka, K.; Ujiie, Y.; Tanaka, W.; Kondo, D.; Umeda, H.; Kamoshida, Y.; Kayanuma, M.; Kamiya, K.; Shiraishi, K.; Machida, Y.; Murakawa, T.; Hayashi, H. J. Am. Chem. Soc. 2014, 136, 4525-4533. f) Nodwell, M. B.; Koch, M. F.; Alte, F.; Schneider, S.; Sieber, S. A. J. Am. Chem. Soc. 2014, 136, 4992-4999. g) Caulkins, B. G.; Bastin, B.; Yang, C.; Neubauer, T. J.; Young, R. P.; Hilario, E.; Huang, Y.-M. M.; Chang, C.-E. A.; Fan, L.; Dunn, M. F.; Marsella, M. J.; Mueller, L. J. J. Am. Chem. Soc. 2014, 136, 12824-12827.

2 a) Seebeck, F. P.; Guainazzi, A.; Amoreira, C.; Baldridge, K. K.; Hilvert, D. Angew. Chem. Int. Ed. Engl. 2006, 45, 6824-6826. b) Toscano, M. D.; Müller, M. M.; Hilvert, D. Angew. Chem. Int. Ed. Engl. 2007, 46, 4468-4470. c) Fesko, K.; Uhl, M.; Steinreiber, J.; Gruber, K.; Griengl, H. Angew. Chem. Int. Ed. Engl. 2010, 49, 121-124. d) Sattler, J. H.; Fuchs, M.; Mutti, F. G.; Grischek, B.; Engel, P.; Pfeffer, J.; Woodley, J. M.; Kroutil, W. Angew. Chem. Int. Ed. Engl. 2014, 53, 14153-14157. e) Maity, A. N.; Lin, H.-H.; Chiang, H.-S.; Lo, H.-H.; Ke, S.-C. ACS Catal. 2015, 5, 3093-3099. f) Shi, L.; Tao, C.; Yang, Q.; Liu, Y. E.; Chen, J.; Chen, J.; Tian, J.; Liu, F.; Li, B.; Du, Y. Zhao, B. Org. Lett. 2015, 17, 5784-5787.

3 Andrade, L. H.; Kroutil, W.; Jamison, T. F. Org. Lett. 2014, 6092-6095.

4 a) Zhang, W.; Bolla, M. L.; Kahne, D.; Walsh, C. T. *J. Am. Chem.* Soc. 2010, 132, 6402-6411. b) Wang, M.; Zhao, Q.; Zhang, Q.; Liu, W. *J. Am. Chem. Soc.* 2016, 138, 6348-6351.

5 a) Scheck, R. A.; Francis, M. B. ACS Chem. Biol. 2007, 4, 247-251. b) Witus, L. S.; Moore, T.; Thuronyi, B. J.; Esser-Kahn, A. P.; Scheck, R. A.; Iavarone, A. T.; Francis, M. B. J. Am. Chem. Soc. 2010, 132, 16812-16817. c) Wang, X.; Canary, J. W. Bioconjugate Chem. 2012, 23, 2329-2334.

6 a) Wilson, D.; Branda, N. R. Angew. Chem. Int. Ed. Engl. 2012, 51, 5431-5434. b) Thorson, M. K.; Majtan, T.; Kraus, J. P.; Barrios, A. M. Angew. Chem. Int. Ed. Engl. 2013, 52, 4641-4644. c) Xu, Y.; Zhang, Z.; Ali, M. M.; Sauder, J.; Deng, X.; Giang, K.; Aguirre, S. D.; Pelton, R.; Li, Y.; Filipe, C. D. M. Angew. Chem. Int. Ed. Engl. 2014, 53, 2620-2622. d) Li, Q.; Qin, X.; Liu, J.; Gui, C.; Wang, B.; Li, J.; Ju, J. J. Am. Chem. Soc. 2016, 138, 408-415.

7 a) Leung, D.; Kang, S. O.; Anslyn, E. V. Chem. Soc. Rev. 2012, 41, 448-479. b) Wolf, C.; Bentley, K. W. Chem. Soc. Rev. 2013, 42, 5408-5424.

8 Metola, P.; Nichols, S. M.; Kahr, B.; Anslyn, E. V. Chem. Sci. 2014, 5, 4278-4282.

9 a) Kim, H.; So, S. M.; Yen, C. P.-H.; Vinhato, E.; Lough, A. J.; Hong, J.-I.; Kim, H.-J.; Chin, J. *Angew. Chem. Int. Ed.* **2008**, *47*, 8657-8660. b) Seo, M.-S.; Lee, A.; Kim, H. *Org. Lett.* **2014**, *16*, 2950-2953.

10 a) Nieto, S.; Dragna, J. M.; Anslyn, E. V. *Chem. Eur. J.* **2010**, *16*, 227–232. b) Dragna, J. M.; Pescitelli, G.; Tran, L.; Lynch, V. M.; Anslyn, E. V.; Di Bari, L. *J. Am. Chem. Soc.* **2012**, *134*, 4398-4407.

12 Joyce, L. A.; Sherer, E. C.; Welch C. J. Chem. Sci. 2014, 5, 2855-2861.

13 a) Ghosn, M. W.; Wolf, C. J. Am. Chem. Soc. 2009, 131, 16360-16361. b) Ghosn, M. W.; Wolf, C. Tetrahedron 2010, 66, 3989-3994. c) Iwaniuk, D. P.; Wolf, C. J. Am. Chem. Soc. 2011, 133, 2414-2417. d) Ghosn, M. W.; Wolf, C. J. Org. Chem. 2011, 76, 3888-3897. e) Iwaniuk, D. P.; Wolf, C. Org. Lett. 2011, 13, 2602-2605. f) Iwaniuk, D. P.; Bentley, K. W.; Wolf, C. Chirality 2012, 24, 584-589. g) Bentley, K. W., Wolf, C. J. Am. Chem. Soc. 2013, 135, 12200-12203. h) Bentley, K. W.; Wolf, C. J. Org. Chem. 2014, 79, 6517-6531. i) De los Santos, Z. A.; Ding, R.; Wolf, C. Org. Biomol. Chem. 2016, 14, 1934-1939. j) De los Santos, Z. A.; Wolf, C. J. Am. Chem. Soc. 2016, 138, 13517-13520.

14 Selected examples of optical chirality sensing: a) Huang, X.; Fujioka, N.; Pescitelli, G.; Koehn, F. E.; Williamson, R. T.; Nakanishi, K.; Berova, N. J. Am. Chem. Soc. 2002, 124, 10320-10335. b) Zhao, J.; Fyles, T. M.; James, T. D. Angew. Chem. Int. Ed. 2004, 43, 3461-3464. c) Mei, X.; Wolf, C. J. Am. Chem. Soc. 2004, 126, 14736-14737. d) Zhu, L.; Zhong, Z.; Anslyn, E. V. J. Am. Chem. Soc. 2005, 127, 4260-4269. e) Folmer-Andersen, J. F.; Lynch, V. M. Anslyn, E. V. J. Am. Chem. Soc. 2005, 127, 7986-7987. f) Mei, X.; Wolf, C. J. Am. Chem. Soc. 2006, 128, 13326-13327. g) Berova, N.; Pescitelli, G.; Petrovic, A. G.; Proni, G. Chem. Commun. 2009, 5958-5980. h) Dutot, L.; Wright, K.; Gaucher, A.; Wakselman, M.; Mazaleyrat, J.-P.; De Zotti, M.; Peggion, C.; Formaggio, F.; Toniolo, C. C. i) Sciebura, J.; Gawronski, J. Chem. Eur. J. 2011, 17, 13138-13141. j) Wezenberg, S. J.; Salassa, G.; Escudero-Adan, E. C.; Benet-Buchholz, J.; Kleij, A. W. Angew. Chem., Int. Ed. 2011, 50, 713-716. k) Li, X.; Burrell, C. E.; Staples, R. J.; Borhan, B. J. Am. Chem. Soc. 2012, 134, 9026-9029. l) Anyika, M.; Gholami, H.; Ashtekar, K. D.; Acho, R.; Borhan, B. J. Am. Chem. Soc. 2014, 136, 550-553. m) Akdeniz, A.; Mosca, L.; Minami, T.; Anzenbacher Jr., P. Chem. Commun. 2015, 51, 5770-5773. n) Shcherbakova, E. G.; Minami, T.; Brega, V.; James, T. D.; Anzenbacher, Jr., P. Angew. Chem. Int. Ed. 2015, 54, 7130-7133. o) Huang, Z.; Yu, S.; Zhao, X.; Wen, K.; Xu, Y.; Yu, X.; Xu, Y.; Pu, L. J. Am. Chem. Soc. 2015, 137, 16458-16461. p) Akdeniz, A.; Minami, T.; Watanabe, S.; Yokoyama, M.; Ema, T.; Anzenbacher Jr., P. Chem. Sci. 2016, 7, 2016-2022. For a review: 0) Pu, L. Chem. Rev. 2004, 104, 1687-1716.

15 a) Cordes, E. H.; Jencks, W. P. *Biochemistry* **1962**, *1*, 773-778. b) Vazquez, M. A.; Munoz, F.; Donoso, J.; Garcia Blanco, F. *Amino Acids* **1992**, *3*, 81-94. c) Salva, A.; Donoso, J.; Frau, J.; Munoz, F. *THEOCHEM* **2002**, 577, 229-238.

16 For a review on indicator displacement assays: a) Nguyen, B. T.; Anslyn, E. V. Coord. Chem. Rev. **2006**, 250, 3118-3127. Chirality sensing examples: b) Folmer-Andersen, J. F.; Kitamura, M.; Anslyn, E. V. J. Am. Chem. Soc. **2006**, 128, 5652-5653. c) Shabbir, S. H.; Joyce, L. A.; da Cruz, G. M.; Lynch, V. M.; Sorey, S.; Anslyn, E. V. J. Am. Chem. Soc. **2009**, 131, 13125-13131. d) Shabbir, S. H.; Regan, C. J.; Anslyn, E. V. Proc. Natl. Acad. Sci. **2009**, 106, 10487-10492. e) Wen, K., Yu, S., Huang, Z., Chen, L., Xiao, M., Yu, X., Pu, L. J. Am. Chem. Soc. **2015**, 137, 4517-4524.

17 a) Metzler, C. M.; Harris, A. G.; Metzler, D. E. Biochemistry 1988, 27, 4923-4933. b) Kiruba, G. S. M.; Wong, M. W. J. Org. Chem. 2003, 68, 2874-2881. c) Sanz, D.; Perona, A.; Claramunt, R. M.; Elguero, J. Tetrahedron 2005, 61, 145-154. d) Golubev, N. S.; Smirnov, S. N.; Tolstoy, P. M.; Sharif, S.; Toney, M. D.; Denisov, G. S.; Limbach, H.-H. J. Mol. Struct. 2007, 844-845, 319-327. e) Sharif, S.; Schlagen, D.; Toney, M. D.; Limbach, H.-H. J. Am. Chem. Soc. 2007, 129, 4440-4455. f) Sharif, S.; Denisov, G. S.; Toney, M. D.; Limbach, H.-H. J. Am. Chem. Soc. 2007, 129, 6313-6327. g) Hill, M. P.; Carroll, C.; Toney, M. D.; Larsen, D. S. J. Phys. Chem. B. 2008, 112, 5867-5873. h) Crugeiras, J.; Rios, A.; Riveiros, E.; Richard, J. P. J. Am. Chem. Soc. 2009, 131, 15815-15824. i) Chan-Huot, Sharif, S.; Tolstoy, P. M.; Toney, M. D.; Limbach, H.-H. Biochemistry 2010, 49, 10818-10830.

18 Cambron, G.; Sevilla, J. M.; Pineda, T.; Blazquez, M. J. Fluoresc. **1996**, *6*, 1-6.

