Base-Mediated Cyclocondensation of Salicylaldehydes and 2-Bromoallyl Sulfones for the Synthesis of 3-Sulfonylchromene Derivatives and Their Regioselective Friedel–Crafts Heteroarylation Reactions

Anand Kumar, Srinivas Thadkapally, and Rajeev S. Menon*

Medicinal Chemistry and Pharmacology Division, CSIR-Indian Institute of Chemical Technology, Hyderabad 500 007, India

Supporting Information

ABSTRACT: Cesium carbonate-mediated reaction of 2hydroxybenzaldehydes and 2-bromoallyl sulfones afforded 2*H*- and 4*H*-chromenol derivatives endowed with a 3arylsulfonyl group. 2-Bromoallyl sulfones functioned as synthetic equivalents of allenyl sulfones under these conditions. The 2*H*- and 4*H*-chromenol derivatives underwent



regioselective Friedel-Crafts reactions with heteroarenes in the presence of *p*-toluenesulfonic acid to afford 4-heteroaryl-4*H*-chromene derivatives in excellent yields.

INTRODUCTION

Unsaturated 1-benzopyran derivatives, commonly known as chromenes, form the central unit of many biologically active natural products as well as synthetic therapeutic agents.¹ The chromene nucleus is, therefore, generally regarded as a privileged scaffold in medicinal chemistry. It exists in two isomeric forms, 2*H*-chromene **1** and 4*H*-chromene **2**, that can be distinguished by the site of unsaturation (Figure 1).





Consequently, method development for chromene synthesis has become an important and fertile area of investigation.^{2,3} Although a number of methods are available for their construction, there is considerable demand for general approaches that deliver substituted chromenes from readily available precursors.

Salicyclic aldehydes constitute a group of useful building blocks for the synthesis of 1-benzopyran derivatives. Recent investigations by Shi, Bräse, and others have established that the base-mediated cyclocondensation of salicylic aldehydes and α,β -unsaturated carbonyl compounds is a versatile method for accessing dihydrobenzopyran derivatives.⁴ More pertinent to the work reported here is the use of *o*-hydroxybenzaldehydes and electron deficient allenes as building blocks for chromene derivatives. DBU-catalyzed cyclocondensation of allenyl phophonates **3** and salicylaldehyde **4a** was developed by Kumara Swamy (Scheme 1a).⁵ Shi reported that allenyl esters **5** react with **4a** in the presence of catalytic quantities of base to Scheme 1. Base-Mediated Cyclocondensations of Salicyclic Aldehydes and Electron Deficient Allenes for the Synthesis of Chromene Derivatives



afford 2*H*- and 4*H*-chromenes (Scheme 1b).⁶ Our interest in the cyclocondensation reactions of unsaturated sulfones⁷ prompted us to explore their utility in the synthesis of sulfone-bearing chromenes. In the process, it was revealed that 3-sulfonylchromenes can be readily synthesized from salicyclic aldehydes and bromoallyl sulfones **6** via a base-mediated annulation (Scheme 1c). Importantly, the direct use of sensitive allenyl sulfones⁸ was avoided by devising and exploiting a base-mediated vinylic bromide displacement on bromoallyl sulfone **6**.

Received: October 7, 2015

RESULTS AND DISCUSSION

The results described in this article originated from our observation that bromoallyl sulfone $6a^9$ readily undergoes basemediated vinylic substitution reactions with nucleophiles. For example, the treatment of 6a with 4-chlorophenol 7 in the presence of 2 equiv of cesium carbonate afforded enol ether 8 in 84% yield (eq 1). Evidently, 8 is formed via a formal vinylic

$$CI \xrightarrow{F} OH = Br \xrightarrow{SO_2Ph} Cs_2CO_3 \xrightarrow{CI} OG = SO_2Ph (1)$$
7 6a 8 (84%)

substitution that usually requires the use of transition metal catalysts. Analogous vinylic displacement reactions of sulfonamide nucleophiles and subsequent gold-catalyzed cycloisomerization reactions were recently developed in our laboratory.¹⁰

We surmised that under the basic conditions of vinylic substitution, the sulfonylmethylene moiety of the enol ether product (such as 8) may be deprotonated and engaged in an annulation reaction with a tethered electrophile. Although deprotonation adjacent to sulfonyl group typically requires a strong base such as butyl lithium, our recent investigations revealed that milder bases like cesium carbonate and DBU are effective in promoting intramolecular condensations of sulfones and aldehydes.⁷ Thus, salicylaldehyde 4a was employed as the phenol component in the vinylic substitution, and in the event, 2*H*-chromene derivative 9aa was obtained as the product in 69% yield (eq 2).



Further experiments to explore the scope and generality of this annulation reaction revealed an interesting pattern of reactivity. Salicylaldehyde as well as *o*-hydroxybenzaldehydes carrying electron-donating substituents generally afforded 2*H*-chromene derivatives **9** preferentially (Table 1, entries 1–6). On the other hand, *o*-hydroxybenzaldehydes endowed with one or more electron-withdrawing substituents furnished isomeric 4*H*-chromene derivatives **10** as the sole or major product (Table 1, entries 8 and 10–14). Halogen-bearing salicylaldehydes deviated from this general trend to some extent (entries 7 and 9). It may be noted that formation of 2*H*- and 4*H*-chromenes was observed in the base-catalyzed reactions of salicylaldehydes and allenoates by Shi; however, no clear trend in reactivity was discernible.^{6a}

To gain insights into the mechanistic underpinnings of this transformation, the reaction of bromoallyl sulfones with cesium carbonate was conducted. An inseparable mixture of allenyl sulfone 11^{11} and propargyl sulfone 12^{12} was obtained after workup and column chromatography. Further treatment of the mixture of 11 and 12 with salicylaldehyde 4a and base afforded 2*H*-chromene 9aa in 22% yield (eq 3). The use of bromoallyl



sulfone **6** in this transformation is, therefore, advantageous in view of the improved pot-economy as well as overall efficiency. Moreover, allenyl sulfones (such as **11**) are base-sensitive and tend to undergo nucleophile-triggered oligomerization reactions.⁸ The relative scarcity of studies involving allenyl sulfones vis-à-vis analogous allenoates may be attributed to these factors. Here, the utility of 2-bromoallyl sulfones **6a** and **6b** as stable and readily available surrogates of allenyl sulfones is demonstrated.

Additionally, it was observed that 4*H*-chromene **10eb** readily isomerized to furnish the corresponding 2*H*-isomer **9eb** in near quantitative yield upon exposure to catalytic PTSA in DMSO (eq 4).

ОН

Table 1. Scope of the Cyclocondensation of Salicylic Aldehydes and 2-Bromoallyl Sulfones^a

$\begin{array}{c} R^{1} \\ R^{2} \\ R^{3} \end{array} \xrightarrow{CHO} \\ R^{3} \end{array} \xrightarrow{H} \\ R^{3} \end{array} \xrightarrow{SO_{2}R^{4}} \\ R^{4} \\ CH_{3}CN \\ 25 \ ^{\circ}C \\ R^{3} \end{array} \xrightarrow{R^{1} \\ CH_{3}CN \\ R^{2} \\ R^{3} \end{array} \xrightarrow{SO_{2}R^{4}} \\ R^{1} \\ OH \\ R^{2} \\ R^{3} \\ R^{$							
	4 6	9	10				
entry	aldehyde	bromoallyl sulfone	time (h)	product(s)	yield ^b (%)		
1	4a , $R^1 = R^2 = R^3 = H$	$6a, R^4 = Ph$	1	9aa	69		
2	4a , $R^1 = R^2 = R^3 = H$	6b , $R^4 = p$ -tolyl	2	9ab	65		
3	4b , $R^1 = OMe$, $R^2 = R^3 = H$	6a, R4 = Ph	3	9ba	61		
4	4b , $R^1 = OMe$, $R^2 = R^3 = H$	6b , $R^4 = p$ -tolyl	4	9bb	75		
5	4c , $R^1 = H$, $R^2 = OMe$, $R^3 = H$	6a, R4 = Ph	2	9ca	55		
6	4c , $R^1 = H$, $R^2 = OMe$, $R^3 = H$	6b , $\mathbb{R}^4 = p$ -tolyl	2	9cb	60		
7	4d , $R^1 = Cl$, $R^2 = R^3 = H$	6a, R4 = Ph	2	9da	52		
8	4d , $R^1 = Cl$, $R^2 = R^3 = H$	6b , $R^4 = p$ -tolyl	2	10db	67		
9	4e , $R^1 = Br$, $R^2 = R^3 = H$	6a, R4 = Ph	4	9ea and 10ea ^c	53		
10	4e , $R^1 = Br$, $R^2 = R^3 = H$	6b , $\mathbb{R}^4 = p$ -tolyl	4	10eb	62		
11	4f , $R^1 = Cl$, $R^2 = H$, $R^3 = Br$	6a, R4 = Ph	4	10fa	66		
12	4f , $R^1 = Cl$, $R^2 = H$, $R^3 = Br$	6b , $\mathbb{R}^4 = p$ -tolyl	4	9fb and 10fb ^d	77		
13	4g , $R^1 = NO_2$, $R^2 = H$, $R^3 = Br$	6a, R4 = Ph	4	10ga	55		
14	4g, $R^1 = NO_2$, $R^2 = H$, $R^3 = Br$	6b , $R^4 = p$ -tolyl	4	10gb	54		

^{*a*}Reaction conditions: **4a**–**g** (1.0 mmol), **6a** or **6b** (1.5 mmol), Cs₂CO₃ (2.5 mmol), and CH₃CN (8 mL). ^{*b*}Isolated yield. ^{*c*}A 10:3 inseparable mixture. ^{*d*}A 4:10 inseparable mixture.



In view of these observations (eq 3), allenyl sulfone 11 is deemed to be a very likely intermediate in the reaction of salicylaldehyde with bromoallyl sulfones 6. A mechanistic rationale for the formation of chromene derivatives is presented in Scheme 2. Conjugate addition of the phenoxide ion 13 to the





sulfonyl allene **11** generates an α -sulfonyl carbanion **14**. Intramolecular nucleophilic addition of the α -sulfonyl carbanion to the aldehyde and subsequent double-bond isomerization affords the 4*H*-chromene derivative **10**. Isomerization of the latter to the 2*H*-chromene derivative **9** may proceed via the benzopyrilium ion **15**. A similar isomerization was observed by Shi in the base-catalyzed reaction of salicylaldehydes with allenyl carboxylates.^{6a} The preferential formation of the 2*H*-chromene derivatives **9** from *o*-hydroxybenzaldehydes carrying electron-releasing groups on the ring may be explained by the enhanced stability of the corresponding benzopyrilium ions **15**.

The apparent involvement of benzopyrilium ions 15 in the formation of 9 prompted us to explore the possibility of its reactions with electron rich aromatic compounds such as indole with an eye on generating heteroaryl chromene derivatives.¹³ The reaction of 9aa with indole was attempted in the presence of various acid catalysts in ethanol and dichloromethane, and the results are summarized in Table 2. In almost all cases, facile conversion of 9aa to 4H-chromene 16a was observed. The 4-(3-indolyl)-4H-chromene derivative 16a was characterized by spectroscopic analysis. The benzylic proton of 16a appeared as a singlet at δ 5.53, and the corresponding carbon resonated at δ 33.5 in the ¹H and ¹³C NMR spectra. All other signals were in agreement with the assigned structure. The reactions in ethanol consistently afforded higher yields of 16a, albeit with varying amounts of ethyl ketal Et-9aa. The use of p-toluenesulfonic acid as the mediator (50 mol %), however, produced only the 4Hchromene derivative 16a in 72% yield (entry 8). The quantitative yield of 16a was obtained when the loading of p-TSA was increased to 70 mol % (entry 10).

It is conceivable that the corresponding benzopyrilium ion 15 underwent a regioselective Friedel–Crafts reaction with indole at the more accessible secondary benzylic carbon. The scope of this reaction was examined subsequently, and it was

Table 2. Optimization of Conditions for the Acid-Mediated Reaction of 9aa and Indole^a

9aa	SO ₂ Ph OH Cataly Me Solver 25 °C,	N H st 2h	HN SO ₂ Ph SO ₂ Ph to Me 16a (when	SO ₂ Ph OEt Me Et-9aa solvent is EtOH)			
entry	catalyst (mol %)	solvent	products	yields ^b (%)			
1	BF ₃ -OEt ₂ (50)	CH_2Cl_2	16a	42			
		EtOH	16a and Et-9aa	51 and 23			
2	$Bi(OTf)_3$ (50)	CH_2Cl_2	16a	48			
		EtOH	16a and Et-9aa	57 and 13			
3	$Yb(OTf)_3(50)$	CH_2Cl_2	16a	44			
		EtOH	16a and Et-9aa	52 and 27			
4	$AlCl_3$ (50)	EtOH	16a and Et-9aa	40 and 15			
5	$\operatorname{FeCl}_{3}(50)$	EtOH	16a and Et-9aa	49 and 24			
6	$SnCl_4$ (50)	EtOH	16a and Et-9aa	51 and 28			
7	$ZnCl_2$ (50)	EtOH	16a and Et-9aa	44 and 41			
8	<i>p</i> -TSA (50)	EtOH	16a	72			
9	<i>p</i> -TSA (60)	EtOH	16a	84			
10	p-TSA (70)	EtOH	16a	99			
^a Reaction conditions: indole (0.15 mmol) , 9aa (0.2 mmol) , acid							

mediator (50–70 mol %), solvent (2 mL), 25 °C, 2 h. ^bIsolated yields.

found that a variety of indoles react in an analogous fashion to afford the corresponding 4-(3-indolyl)-4H-chromenes 16-22 in excellent yields (Table 3). Other heterocyclic aromatics such as pyrrole and furan also furnished the corresponding 4hetroaryl-4H-chromenes 23-26. It is important to note that 4heteroaryl-4H-chromenes were formed from isomeric chromenol 9 or 10. This is illustrated by the formation of products 22 and 24 from 4H-chromenol 10eb as well as the formation of 25 from 10fa. This observation is consistent with the intermediacy of the purported benzopyrilium ion. Additionally, attempted reactions of aromatic compounds such as thiophene, toluene, and benzene led to the formation of ethanol-derived ketal Et-9aa. The latter was isolated in 71% yield when the reaction was run without any heteroarene. A variety of 4-aryl-4H-chromenes have been shown to possess excellent anticancer activities,¹⁴ and this protocol allows for the facile construction of hitherto unknown sulfonyl derivatives of 4-heteroaryl-4H-chromenes. It may be mentioned here that our efforts to reductively remove or substitute the sulfonyl groups of 4-heteroaryl-4H-chromenes were not successful.

The base-mediated cyclocondensation reaction initiated by the vinylic displacement of the bromoallyl sulfone 6a can also be adapted for the construction of other related heterocycles (eq 5). In light of our recent findings about the addition of



sulfonamide nucleophiles to bromoallyl sulfones,¹⁰ we surmised that the β -(aminosulfonyl)aldehyde 27^{15} could partake in a vinylic displacement-cyclization reaction with bromoallyl sulfone **6a** to afford a six-membered nitrogen heterocycle. The widespread occurrence of piperidines and piperidones in





^{*a*}Reaction conditions: heteroarene (0.30 mmol), 2*H*-chromene (0.40 mmol), *p*-TSA (0.21 mmol), EtOH (2 mL), 25 °C, 2 h. Isolated yields given. ^{*b*}Reaction conducted under the conditions described in footnote *a* but in the absence of heteroarene.

natural alkaloids and synthetic compounds of medicinal importance provided an additional impetus for this preliminary study.¹⁶ Gratifyingly, the reaction of **2**7 with **6**a under the conditions of vinylic displacement afforded the substituted piperidine **28** in 76% yield. Oxidation of the latter by pyridinium chlorochromate (PCC) furnished the 4-piperidone derivative **29** in 86% yield. It may be noted that the oxidation was accompanied by the migration of the exocyclic double bond of **28** to an endocyclic position in conjugation with the carbonyl group (eq 5).

CONCLUSION

The following important developments have emerged from this study. (i) A transition metal-free, vinylic bromide displacement reaction of bromoallyl sulfones **6** with phenolate nucleophiles was discovered. (ii) Under these conditions, **6** functioned as a synthetic equivalent of allenyl sulfones and underwent a smooth cyclocondensation with salicylic aldehydes. (iii) The sulfonyl chromenols thus formed underwent acid-mediated regioselective Friedel–Crafts reactions to afford 4-heteroaryl-4H-chromenes. Additionally, in view of the biological activities displayed by chromenes in general and 4-aryl-4H-chromenes in particular, the synthetic methods described here have potential for applications in medicinal chemistry programs. The success of *p*-TSA-mediated addition of heteroaromatics to the chromenols bodes well for the development of asymmetric versions of the reaction by using chiral Brønsted acids.¹⁷

Preliminary results suggest that the vinylic bromide displacement method can also be applied for the construction of sixmembered nitrogen heterocycles.

EXPERIMENTAL SECTION

All reactions were performed in flame-dried round-bottom flasks, fitted with rubber septa under a positive pressure of nitrogen. Analytical thin-layer chromatography (TLC) was performed using aluminumbacked UVF254 precoated silica gel flexible plates. Removal of solvent under reduced pressure refers to distillation with a rotary evaporator attached to a vacuum pump (~3 mmHg). Technical grade ethyl acetate and petroleum ether used for column chromatography were distilled prior to use. Column chromatography was conducted using silica gel (60-120 mesh and 100-200 mesh) packed in glass columns. Melting points were obtained in open capillary tubes using a micromelting point apparatus and are uncorrected. NMR spectra were recorded on a 300, 400, or 500 MHz nuclear magnetic resonance spectrometer. The proton resonances are annotated as chemical shifts (δ) relative to tetramethylsilane $(\delta 0.0)$ using the residual solvent signal as an internal standard or tetramethylsilane itself: chloroform-d $(\delta$ 7.26, singlet), multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad), coupling constant (J, in hertz), and the number of protons for a given resonance indicated by nH. The chemical shifts of ¹³C NMR are reported in parts per million relative to the central line of the triplet at 77.0 ppm for CDCl₃. IR spectra were recorded on an FT-IR spectrometer, and wavenumbers of maximal absorption peaks are presented in inverse centimeters. High-resolution mass analyses (HRMS) were performed on a mass spectrometer using ESI-TOF techniques. Acetonitrile (LR grade), dimethyl sulfoxide (LR

grade), ethanol (95%), cesium carbonate (AR), and *p*-toluenesulfonic acid were purchased from local suppliers and used as received. 2-Hydroxybenzaldehydes 4a-g, 2,3-dibromopropene, sodium benzenesulfinate, sodium *p*-toluenesulfinate, 3-amino-1-propanol, pyridinium chlorochromate (PCC), and the heteroarenes used for the Friedel– Crafts reactions were used as received from suppliers. 2-(Bromoallylsulfonyl)benzene 6a,⁹ allenyl sulfone 11,¹¹ and propargyl sulfone 12^{12} are known compounds and were identified by comparison of their spectroscopic data with the reported values. *N*-(3-Oxopropyl)benzenesulfonamide 27 was prepared from 3-amino-1propanol as reported previously.¹⁵

General Procedure for the Preparation of 1-(2-Bromoallyl)sulfonylbenzene 6a or 1-(2-Bromoallyl)sulfonyl-4-methylbenzene 6b. Sodium benzenesulfinate or sodium *p*-toluenesulfinate (10 mmol) was added to DMF (10 mL) in a round-bottom flask and left to stir for 20 min. To this suspension was added 2,3-dibromopropene (technical grade 80%, 1.22 mL, 10 mmol), and the mixture was stirred at room temperature under nitrogen for 12 h. After the completion of the reaction, ice and ethyl acetate (20 mL) were added and the mixture was stirred for 15 min. Water (20 mL) was added, and the organic layer was separated in a separatory funnel. The aqueous layer was extracted with ethyl acetate (2 × 15 mL); the organic layers were combined, and the solvent was removed on a rotavapor. The residue obtained was purified by column chromatography on silica gel using ethyl acetate/hexane mixtures as the eluent to obtain pure samples of 6a or 6b.

(2-Bromoally/sulfony/)benzene (**6a**).⁹ White solid (1.51 g, 58%): R_f = 0.6 (30% ethyl acetate in hexanes); mp 77–79 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.94 (d, J = 8.5 Hz, 2H), 7.71–7.68 (m, 1H), 7.60–7.57 (m, 2H), 5.84 (d, J = 2.1 Hz, 1H), 5.77 (d, J = 2.1 Hz, 1H), 4.16 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 137.7, 134.1, 129.1, 128.7, 126.5, 117.1, 66.2; IR (KBr) ν_{max} 3423, 2981, 2919, 1622, 1584, 1447, 1393, 1307, 1256, 1141, 907 cm⁻¹.

1-(2-Bromoallylsulfonyl)-4-methylbenzene (**6b**). White solid (1.56 g, 57%): $R_f = 0.6$ (30% ethyl acetate in hexanes); mp 89–91 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.81 (d, J = 8.2 Hz, 2H), 7.37 (d, J = 8.2 Hz, 2H), 5.83 (d, J = 2.1 Hz, 1H), 5.77 (d, J = 2.1 Hz, 1H), 4.14 (s, 2H), 2.46 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 145.2, 134.8, 129.7, 128.7, 126.3, 117.3, 66.3, 21.6; IR (KBr) ν_{max} 2920, 1623, 1467, 1393, 1308, 1256, 1141, 1084, 908 cm⁻¹; HRMS (ESI) calcd for C₁₀H₁₁BrO₂SNa (M + Na) 296.9561, found 296.9568.

General Procedure for the Cesium Carbonate-Mediated Reaction of Phenolic Substrates (4a-g and 7) with 2-Bomoallyl Sulfones 6a and 6b. 2-Bromoallyl sulfonylbenzene 6a or 2-bromoallyl sulfonyl-4-methylbenzene 6b (1.5 mmol) and the phenolic substrate (1.0 mmol) were dissolved in acetonitrile (8 mL) at room temperature in a 25 mL round-bottom flask. Cesium carbonate (815 mg, 2.5 mmol) was added, and the reaction mixture was stirred under nitrogen for the required amount of time. Upon completion, the solvent was evaporated on a rotavapor, water (20 mL) was added, and the resulting solution was extracted with dichloromethane (3 × 20 mL). The combined organic extracts were dried over anhydrous sodium sulfate and evaporated on a rotavapor, and the residue obtained was subjected to column chromatography on silica gel using an ethyl acetate/hexane mixture as the eluent.

1-Chloro-4-[3-(phenylsulfonyl)prop-1-en-2-yloxy]benzene (8). White solid (259 mg, 84%): $R_f = 0.6$ (30% ethyl acetate in hexanes); mp 78–80 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.97 (d, J = 7.5 Hz, 2H), 7.69–7.66 (m, 1H), 7.59–7.56 (m, 2H), 7.28–7.26 (m, 2H), 6.79 (d, J = 8.5 Hz, 2H), 4.42 (d, J = 2.1 Hz, 1H), 4.20 (d, J = 2.1 Hz, 1H), 4.04 (s, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 152.5, 151.6, 138.8, 133.9, 130.2, 129.8, 129.08, 128.6, 122.3, 96.1, 61.3; IR (KBr) ν_{max} 2926, 2358, 1643, 1485, 1318, 1290, 1218, 1149, 1083, 968 cm⁻¹; HRMS (ESI) calcd for C₁₅H₁₃ClO₃SNa (M + Na) 331.0172, found 331.0166.

2-Methyl-3-(phenylsulfonyl)-2H-chromen-2-ol (**9aa**). Yellow solid (208 mg, 69%): $R_f = 0.5$ (30% ethyl acetate in hexanes); mp 143–145 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.97 (d, J = 8.1 Hz, 2H), 7.83 (s, 1H), 7.66–7.62 (m, 1H), 7.57–7.54 (m, 2H), 7.38–7.35 (m, 2H), 7.06 (t, J = 7.4 Hz, 1H), 6.97 (d, J = 8.1 Hz, 1H), 3.52 (s, 1H), 1.62 (s,

3H); ¹³C NMR (75 MHz, CDCl₃) δ 152.7, 140.4, 135.2, 134.1, 133.5, 133.2, 129.5, 129.1, 128.1, 122.2, 118.3, 117.0, 97.4, 26.6; IR (KBr) $\nu_{\rm max}$ 3438, 3062, 2924, 1688, 1613, 1481, 1303, 1149, 1113, 941, 862, 751, 727, 682, 641 cm⁻¹; HRMS (ESI) calcd for C₁₆H₁₄O₄SNa (M + Na) 325.0510, found 325.0508.

2-Methyl-3-tosyl-2H-chromen-2-ol (**9ab**). Yellow solid (205 mg, 65%): $R_f = 0.4$ (30% ethyl acetate in hexanes); mp 161–163 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.84 (d, J = 8.1 Hz, 2H), 7.79 (s, 1H), 7.38–7.33 (m, 4H), 7.07–7.02 (m, 1H), 6.97 (d, J = 8.1 Hz, 1H), 3.51 (s, 1H), 2.44 (s, 3H), 1.60 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 152.7, 144.6, 137.4, 134.7, 134.4, 133.1, 129.8, 129.4, 128.2, 122.2, 118.4, 117.0, 97.5, 26.5, 21.6; IR (KBr) ν_{max} 3447, 3063, 2952, 1618, 1304, 1146, 943, 754, 676 cm⁻¹; HRMS (ESI) calcd for C₁₇H₁₆O₄SNa (M + Na) 339.0667, found 339.0670.

6-Methoxy-2-methyl-3-(phenylsulfonyl)-2H-chromen-2-ol (**9ba**). Yellow solid (202 mg, 61%): $R_f = 0.5$ (30% ethyl acetate in hexanes); mp 155–157 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.97–7.95 (m, 2H), 7.78 (s, 1H), 7.65–7.62 (m, 1H), 7.57–7.53 (m, 2H), 6.94 (dd, J = 2.8, 8.9 Hz, 1H), 6.90 (d, J = 8.9 Hz, 1H), 6.86 (d, J = 2.8 Hz, 1H), 3.80 (s, 3H), 3.47 (s, 1H), 1.59 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 154.5, 146.8, 140.4, 135.2, 134.7, 133.5, 129.1, 128.2, 119.9, 118.6, 117.9, 112.7, 97.3, 55.8, 26.3; IR (KBr) ν_{max} 3457, 3105, 3009, 2945, 1572, 1491, 1301, 1148, 1083, 940, 725, 690 cm⁻¹; HRMS (ESI) calcd for C₁₇H₁₆O₃SNa (M + Na) 355.0616, found 355.0620.

6-Methoxy-2-methyl-3-tosyl-2H-chromen-2-ol (**9bb**). Pale yellow solid (260 mg, 75%): $R_f = 0.3$ (30% ethyl acetate in hexanes); mp 148–150 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.83 (d, J = 7.8 Hz, 2H), 7.74 (s, 1H), 7.34 (d, J = 7.8 Hz, 2H), 6.94–6.89 (m, 2H), 6.85 (s, 1H), 3.80 (s, 3H), 3.47 (s, 1H), 2.44 (s, 3H), 1.59 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 154.5, 146.8, 144.6, 137.3, 135.0, 134.7, 129.8, 128.2, 119.8, 118.7, 117.9, 112.6, 97.3, 55.8, 26.2, 21.6; IR (KBr) ν_{max} 3407, 3083, 2951, 1569, 1488, 1305, 1267, 1149, 1095, 1027, 822, 687 cm⁻¹; HRMS (ESI) calcd for C₁₈H₁₈O₅SNa (M + Na) 369.0773, found 369.0778.

7-Methoxy-2-methyl-3-(phenylsulfonyl)-2H-chromen-2-ol (9ca). Pale yellow solid (183 mg, 55%): $R_f = 0.4$ (30% ethyl acetate in hexanes); mp 124–126 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.96 (d, J = 7.2 Hz, 2H), 7.79 (s, 1H), 7.65–7.60 (m, 1H), 7.56–7.52 (m, 2H), 7.26 (d, J = 8.5 Hz, 1H), 6.61 (dd, J = 2.2, 8.5 Hz, 1H), 6.50 (d, J = 2.2 Hz, 1H), 3.81 (s, 3H), 3.57 (s, 1H), 1.58 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 164.0, 154.6, 140.8, 135.4, 133.4, 130.7, 130.4, 129.1, 128.0, 111.6, 109.6, 101.7, 97.8, 55.6, 26.4; IR (KBr) ν_{max} 3447, 3064, 2930, 1616, 1297, 1147, 1093, 730, 622 cm⁻¹; HRMS (ESI) calcd for C₁₇H₁₆O₅SNa (M + Na) 355.0616, found 355.0619.

7-Methoxy-2-methyl-3-tosyl-2H-chromen-2-ol (9cb). Pale yellow solid (208 mg, 60%): $R_f = 0.4$ (30% ethyl acetate in hexanes); mp 136–138 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.84 (d, J = 7.6 Hz, 2H), 7.77 (s, 1H), 7.35–7.25 (m, 3H), 6.61 (d, J = 8.3 Hz, 1H), 6.51 (s, 1H), 3.82 (s, 3H), 3.67 (s, 1H), 2.45 (s, 3H), 1.59 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 163.8, 154.5, 144.3, 137.8, 134.9, 130.8, 130.6, 129.7, 128.1, 111.7, 109.5, 101.7, 97.8, 55.5, 26.4, 21.6; IR (KBr) ν_{max} 3447, 3000, 2944, 1609, 1285, 1141, 1091, 673 cm⁻¹; HRMS (ESI) calcd for C₁₈H₁₈O₅SNa (M + Na) 369.0773, found 369.0770.

6-Chloro-2-methyl-3-(phenylsulfonyl)-2H-chromen-2-ol (**9da**). Pale yellow solid (175 mg, 52%): $R_f = 0.6$ (30% ethyl acetate in hexanes); mp 145–147 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.96 (d, J = 7.5 Hz, 2H), 7.73 (s, 1H), 7.67–7.64 (m, 1H), 7.58–7.55 (m, 2H), 7.33–7.29 (m, 2H), 6.91 (d, J = 8.5 Hz, 1H), 3.51 (s, 1H), 1.61 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 151.2, 140.0, 135.6, 133.9, 133.8, 132.8, 129.2, 128.6, 128.3, 127.2, 119.5, 118.5, 97.8, 26.6; IR (KBr) ν_{max} 3436, 3061, 3006, 2950, 1619, 1476, 1308, 1156, 1087, 950, 866, 730, 685 cm⁻¹; HRMS (ESI) calcd for C₁₆H₁₃ClO₄SNa (M + Na) 359.0121, found 359.0127.

6-Chloro-2-methyl-3-tosyl-4H-chromen-4-ol (10db). Pale yellow solid (235 mg, 67%): $R_f = 0.6$ (30% ethyl acetate in hexanes); mp 181–183 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.88 (d, J = 8.3 Hz, 2H), 7.48 (d, J = 2.3 Hz, 1H), 7.37–7.28 (m, 3H), 7.02 (d, J = 8.3 Hz, 1H), 5.70 (d, J = 3.8 Hz, 1H), 3.70 (d, J = 3.8 Hz, 1H), 2.45 (s, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 160.5, 147.9, 144.4, 138.7, 130.1, 129.9, 129.8, 129.5, 127.2, 122.6, 117.9, 116.1, 60.8, 21.6, 18.5; IR (KBr) ν_{max}

3486, 2914, 1635, 1480, 1303, 1248, 1146, 1088, 983, 819, 677, 604 cm⁻¹; HRMS (ESI) calcd for $C_{17}H_{15}ClO_4SNa$ (M + Na) 373.0277, found 373.0278.

6-Bromo-2-methyl-3-(phenylsulfonyl)-2H-chromen-2-ol (**9ea**) and 6-Bromo-2-methyl-3-(phenylsulfonyl)-4H-chromen-4-ol (**10ea**) (1:0.3 inseparable mixture). Pale yellow solid (201 mg, 53%): $R_f = 0.5$ (30% ethyl acetate in hexanes); ¹H NMR (300 MHz, CDCl₃) δ 8.00–7.94 (m, 2.6H), 7.72 (s, 1H), 7.65–7.43 (m, 6.8H), 6.96 (d, J = 8.9 Hz, 0.3H), 6.86 (d, J = 8.5 Hz, 1H), 5.71 (d, J = 2.6Hz, 0.3H), 3.67 (d, J = 2.6 Hz, 0.3H), 3.55 (s, 1H), 2.43 (s, 1H), 1.61 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 151.8, 140.0, 135.7, 133.8, 133.7, 133.4, 132.7, 132.5, 131.6, 129.3, 129.2, 128.3, 127.9, 127.1, 120.0, 118.9, 118.2, 114.3, 97.7, 60.7, 26.6, 18.6; IR (KBr) ν_{max} 3486, 3441, 3058, 1477, 1306, 1247, 1150, 1105, 1084, 881, 719, 648 cm⁻¹; HRMS (ESI) calcd for C₁₆H₁₃BrO₄SNa (M + Na) 402.9616, found 402.9608.

6-Bromo-2-methyl-3-tosyl-4H-chromen-4-ol (**10eb**). Yellow solid (244 mg, 62%): $R_f = 0.6$ (30% ethyl acetate in hexanes); mp 206–208 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.86 (d, J = 8.1 Hz, 2H), 7.62 (d, J = 2.3 Hz, 1H), 7.42 (dd, J = 2.3, 8.9 Hz, 1H), 7.34 (d, J = 8.1 Hz, 2H), 6.95 (d, J = 8.9 Hz, 1H), 5.68 (d, J = 3.2 Hz, 1H), 3.70 (d, J = 3.2 Hz, 1H), 2.44 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 160.4, 148.4, 144.5, 138.7, 132.7, 132.5, 129.9, 127.2, 123.0, 118.2, 117.5, 116.2, 60.7, 21.6, 18.5; IR (KBr) ν_{max} 3486, 3065, 2910, 1635, 1477, 1299, 1247, 1145, 1088, 980, 817, 674 cm⁻¹; HRMS (ESI) calcd for C₁₇H₁₅BrO₄SNa (M + Na) 416.9772, found 416.9777.

8-Bromo-6-chloro-2-methyl-3-(phenylsulfonyl)-4H-chromen-4-ol (**10fa**). Pale yellow solid (273 mg, 66%): $R_f = 0.7$ (30% ethyl acetate in hexanes); mp 176–178 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.99 (d, J = 8.5 Hz, 2H), 7.65–7.62 (m, 1H), 7.58–7.55 (m, 3H), 7.43 (s, 1H), 5.70 (d, J = 3.8 Hz, 1H), 3.66 (s, 1H), 2.50 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 160.7, 145.2, 141.4, 133.6, 133.0, 130.3, 129.4, 128.8, 127.2, 123.6, 116.5, 111.0, 61.1, 18.5; IR (KBr) ν_{max} 3460, 3070, 2922, 1639, 1455, 1326, 1253, 1149, 1089, 1020, 726, 628 cm⁻¹; HRMS (ESI) calcd for C₁₆H₁₂BrClO₄SNa (M + Na) 436.9226, found 436.9231.

8-Bromo-6-chloro-2-methyl-3-tosyl-2H-chromen-2-ol (9fb) and 8-Bromo-6-chloro-2-methyl-3-tosyl-4H-chromen-4-ol (10fb) (0.4:1 inseparable mixture). Pale yellow solid (330 mg, 77%): $R_f = 0.6$ (30% ethyl acetate in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.86 (d, J =8.2 Hz, 2H), 7.82 (d, J = 8.4 Hz, 0.8H), 7.66 (s, 0.4H), 7.56 (d, J = 2.4Hz, 0.4H), 7.54 (d, J = 2.4 Hz, 1H), 7.42 (d, J = 2.4 Hz, 1H), 7.35 (d, J =8.2 Hz, 2.8H), 7.28 (d, J = 2.4 Hz, 0.4H), 5.67 (d, J = 3.5 Hz, 1H), 3.72 (d, J = 3.5 Hz, 1H), 3.69 (s, 0.4H), 2.50 (s, 3H), 2.45 (s, 1.2H), 2.44 (s, 3H), 1.69 (s, 1.2H); ¹³C NMR (75 MHz, CDCl₃) δ 160.4, 148.5, 145.2, 145.1, 145.0, 144.6, 138.4, 137.0, 136.8, 135.2, 132.9, 130.2, 129.9, 128.7, 128.4, 127.8, 127.2, 123.7, 120.4, 116.8, 111.5, 111.0, 98.9, 61.0, 26.6, 21.7, 21.6, 18.4; IR (KBr) $ν_{max}$ 3447, 2927, 1639, 1454, 1325, 1293, 1251, 1143, 1089, 1042, 680 cm⁻¹; HRMS (ESI) calcd for C₁₇H₁₄BrClO₄SNa (M + Na) 450.9382, found 450.9386.

2-Methyl-6-nitro-3-(phenylsulfonyl)-4H-chromen-4-ol (**10ga**). Pale yellow solid (191 mg, 55%): $R_f = 0.5$ (30% ethyl acetate in hexanes); mp 212–214 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.44 (s, 1H), 8.21 (d, J = 8.9 Hz, 1H), 8.01 (d, J = 7.6 Hz, 2H), 7.65–7.58 (m, 3H), 7.20 (d, J = 8.9 Hz, 1H), 5.80 (s, 1H), 3.87 (s, 1H), 2.49 (s, 3H). Because of the poor solubility of **10ga** in CDCl₃, ¹³C NMR was measured in DMSO- d_6 . During measurement, **10ga** isomerized to generate a mixture of 2H-chromenes **9ga** and **10ga**: ¹³C NMR (125 MHz, DMSO- d_6) δ 160.4, 157.6, 153.1, 144.0, 142.3, 141.5, 140.6, 136.5, 133.8, 133.6, 133.4, 129.4, 129.3, 127.8, 127.4, 126.0, 125.7, 125.1, 124.6, 118.7, 117.7, 117.5, 98.6, 79.2, 58.7, 26.8, 18.3; IR (KBr) ν_{max} 3509, 3086, 2925, 1642, 1526, 1342, 1294, 1242, 1139, 1086, 985, 756, 640 cm⁻¹; HRMS (ESI) calcd for C₁₆H₁₃NO₆SNa (M + Na) 370.0361, found 370.0365.

2-Methyl-6-nitro-3-tosyl-4H-chromen-4-ol (**10gb**). Pale yellow solid (195 mg, 54%): $R_f = 0.6$ (30% ethyl acetate in hexanes); mp 185–187 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.43 (s, 1H), 8.20 (d, J = 8.7 Hz, 1H), 7.88 (d, J = 7.0 Hz, 2H), 7.36 (d, J = 7.0 Hz, 2H), 7.19 (d, J = 8.7 Hz, 1H), 5.76 (s, 1H), 3.85 (s, 1H), 2.49 (s, 3H), 2.44 (s,

3H); ¹³C NMR (75 MHz, CDCl₃) δ 159.9, 153.3, 144.8, 144.6, 138.2, 130.0, 127.3, 126.4, 125.1, 122.1, 117.6, 117.2, 60.5, 21.6, 18.4; IR (KBr) ν_{max} 3477, 3097, 2924, 2854, 1642, 1535, 1343, 1298, 1247, 1146, 1091, 681 cm⁻¹; HRMS (ESI) calcd for C₁₇H₁₅NO₆SNa (M + Na) 384.0518, found 384.0526.

CS₂CO₃-Mediated Conversion of 2-Bromoallyl Sulfones 6a to Allenyl Sulfone 11 and Propargyl Sulfone 12. Cesium carbonate (391 mg, 1.2 mmol) was added to a solution of bromoallyl sulfone 6a (260 mg, 1.0 mmol) in acetonitrile (2 mL) at room temperature and stirred for 2h. The solvent was removed on a rotavapor; deionized water (10 mL) was added, and the aqueous solution was extracted with ethyl acetate (3×10 mL). The combined organic extracts was washed with brine, dried over anhydrous sodium sulfate, and concentrated on a rotavapor. The residue on column chromatography on silica gel using a petroleum ether/ethyl acetate eluent afforded 130 mg of an oil that was identified as a 5:1 mixture of allenyl sulfone 11¹¹ and propargyl sulfone 12.¹² The reaction of the isolated mixture of 11 and 12 with salicylaldehyde was conducted according to the general procedure to obtain 9aa in 22% isolated yield.

NMR Data for the Mixture of **11** and **12**. ¹H NMR (500 MHz, CDCl₃) δ 8.00–7.98 (m, 0.4H), 7.93–7.91 (m, 2H), 7.72–7.69 (m, 0.2H), 7.66–7.62 (m, 1H), 7.61–7.59 (m, 0.4H), 7.58–7.54 (m, 2H), 6.26 (t, *J* = 6.4 Hz, 1H), 5.45 (d, *J* = 6.4 Hz, 2H), 3.97 (d, *J* = 2.7 Hz, 0.4H), 2.39 (t, *J* = 0.2H); ¹³C NMR (125 MHz, CDCl₃) δ 209.3, 141.1, 134.3, 133.5, 129.2, 129.1, 128.8 (2), 127.5, 127.3, 100.9, 84.1, 76.2, 71.5, 48.3.

PTSA-Catalyzed Isomerization of 4H-Chromene 10eb to 2H-Chromene 9eb. To a solution of 4H-chromene 10eb (39 mg, 0.1 mmol) in dimethyl sulfoxide (0.5 mL) was added PTSA (0.4 mg, 20 mol %), and the reaction mixture was allowed to sir at 25 overnight. Water (5 mL) was added, and the solution was extracted with ethyl acetate $(3 \times 5 \text{ mL})$. The organic extracts were combined and washed with a saturated sodium bicarbonate solution (10 mL) and brine. After evaporation of the solvent on a rotavapor, the residue was chromatographed on a silica column using hexane/ethyl acetate mixtures as the eluent to obtain 6-bromo-2-methyl-3-tosyl-2Hchromen-2-ol **9eb** as a white solid (39 mg, 99%): $R_f = 0.5$ (30%) ethyl acetate in hexanes); mp 160-162 °Č; ¹H NMR (400 MHz, $CDCl_3$) δ 7.83 (d, J = 8.2 Hz, 2H), 7.67 (s, 1H), 7.46 (d, J = 2.3 Hz, 1H), 7.43 (dd, J = 2.3, 8.6 Hz, 1H), 7.35 (d, J = 8.2 Hz, 2H), 6.86 (d, J = 8.6 Hz, 1H), 3.53 (s, 1H), 2.45 (s, 3H), 1.60 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 151.8, 144.9, 137.2, 136.2, 135.6, 133.1, 131.6, 129.9, 128.5, 120.2, 118.9, 114.3, 97.9, 26.6, 21.6; IR (KBr) ν_{max} 3450, 2920, 1618, 1475, 1397, 1312, 1267, 1247, 1170, 1151, 1095, 945 cm^{-1} ; HRMS (ESI) calcd for $C_{17}H_{15}BrO_4SNa$ (M + Na) 416.9772, found 416.9780.

Optimized General Procedure for the PTSA-Mediated Reaction of Chromenes (9 and 10) with Heteroarenes. Chromene derivative 9 or 10 (0.40 mmol) and the (hetero)arene (0.30 mmol) were dissolved in ethanol (2 mL) at room temperature. PTSA (36 mg, 0.21 mmol) was added, and the reaction mixture was stirred for 2 h. Water (10 mL) was added, and the solution was extracted with ethyl acetate (3×10 mL). The organic extracts were combined and washed with a saturated sodium bicarbonate solution (10 mL) and brine. After evaporation of the solvent on a rotavapor, the residue was chromatographed on a silica column using hexane/ ethyl acetate mixtures as the eluent to obtain analytically pure samples of products 16–27.

 $\overline{3}$ -[2-Methyl-3-(phenylsulfonyl)-4H-chromen-4-yl]-1H-indole (16a). Pale yellow solid (119 mg, 99%): $R_f = 0.5$ (40% ethyl acetate in hexanes); mp 187–189 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.90 (s, 1H), 7.26–7.23 (m, 3H), 7.19–7.09 (m, 5H), 7.06–6.95 (m, 5H), 6.90 (t, J = 7.2 Hz, 1H), 5.53 (s, 1H), 2.52 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 159.4, 148.9, 142.1, 136.3, 131.8, 129.1, 127.9, 127.6, 126.3, 125.3, 124.9, 124.2, 122.7, 121.9, 119.7, 119.0, 118.6, 116.0, 115.5, 111.1, 33.5, 19.0; IR (KBr) ν_{max} 3389, 3060, 2886, 1638, 1487, 1452, 1318, 1286, 1237, 1141, 1089, 757, 727, 630 cm⁻¹; HRMS (ESI) calcd for C₂₄H₁₉NO₃SNa (M + Na) 424.0983, found 424.0972.

3-(2-Methyl-3-tosyl-4H-chromen-4-yl)-1H-indole (16b). Colorless solid (114 mg, 91%): $R_f = 0.6$ (40% ethyl acetate in hexanes); mp

173–175 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.93 (s, 1H), 7.23 (d, J = 8.1 Hz, 1H), 7.18 (d, J = 8.2 Hz, 1H), 7.14–7.09 (m, 5H), 7.06–6.99 (m, 2H), 6.96 (dt, J = 1.1, 7.5 Hz, 1H), 6.91–6.88 (m, 1H), 6.77 (d, J = 8.1 Hz, 2H), 5.51 (s, 1H), 2.51 (s, 3H), 2.17 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 159.0, 148.9, 142.7, 139.2, 136.4, 129.1, 128.5, 127.6, 126.6, 126.4, 125.3, 124.9, 124.3, 122.7, 121.8, 119.6, 119.0, 118.6, 115.9, 111.1, 33.6, 21.3, 18.9; IR (KBr) ν_{max} 3370, 3052, 2920, 1636, 1489, 1316, 1287, 1238, 1143, 1088, 742, 671, 608 cm⁻¹; HRMS (ESI) calcd for C₂₅H₂₁NO₃SNa (M + Na) 438.1140, found 438.1123.

5-Methoxy-3-[2-methyl-3-(phenylsulfonyl)-4H-chromen-4-yl]-1Hindole (17a). Colorless solid (122 mg, 94%): $R_f = 0.5$ (40% ethyl acetate in hexanes); mp 110–112 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.80 (s, 1H), 7.27–7.25 (m, 2H), 7.22–7.19 (m, 1H), 7.15–7.11 (m, 2H), 7.08–7.06 (m, 2H), 7.03–6.97 (m, 4H), 6.72–6.70 (m, 2H), 5.50 (s, 1H), 3.70 (s, 3H), 2.51 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 159.3, 153.9, 149.0, 142.1, 131.9, 131.4, 129.2, 127.9, 127.7, 126.3, 125.7, 125.0, 124.1, 123.4, 118.7, 115.8, 115.5, 112.1, 111.8, 100.4, 55.6, 33.5, 18.9; IR (KBr) ν_{max} 3388, 3061, 2927, 1636, 1583, 1485, 1452, 1299, 1230, 1146, 1088, 755, 629 cm⁻¹; HRMS (ESI) calcd for C₂₃H₂₁NO₄SNa (M + Na) 454.1089, found 454.1084.

5-Methoxy-3-(2-methyl-3-tosyl-4H-chromen-4-yl)-1H-indole (17b). Pale yellow solid (123 mg, 92%): $R_f = 0.5$ (40% ethyl acetate in hexanes); mp 190–192 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.86 (s, 1H), 7.14–7.07 (m, 6H), 7.02–6.96 (m, 2H), 6.80 (d, J = 7.7 Hz, 2H), 6.71–6.68 (m, 2H), 5.48 (s, 1H), 3.70 (s, 3H), 2.49 (s, 3H), 2.19 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 158.9, 153.9, 149.1, 142.7, 139.2, 131.5, 129.2, 128.6, 127.7, 126.4, 125.8, 125.0, 124.2, 123.4, 118.8, 115.8, 112.0, 111.7, 100.6, 55.6, 33.5, 21.3, 18.9; IR (KBr) ν_{max} 3340, 2922, 1639, 1485, 1453, 1297, 1234, 1144, 1087, 1039, 751, 667 cm⁻¹; HRMS (ESI) calcd for C₂₆H₂₃NO₄SNa (M + Na) 468.1245, found 468.1240.

5-Bromo-3-[2-methyl-3-(phenylsulfonyl)-4H-chromen-4-yl]-1Hindole (18a). Yellow solid (137 mg, 95%): $R_f = 0.4$ (40% ethyl acetate in hexanes); mp 188–190 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.04 (s, 1H), 7.33–7.31 (m, 3H), 7.21 (t, J = 7.3 Hz, 1H), 7.16–7.09 (m, 3H), 7.07–7.02 (m, 5H), 6.98 (dt, J = 1.2, 7.6 Hz, 1H), 5.46 (s, 1H), 2.56 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 159.5, 148.7, 141.9, 134.9, 132.1, 128.9, 128.1, 127.9, 126.9, 126.3, 125.0, 124.7, 124.0, 123.8, 121.1, 118.7, 116.2, 115.2, 112.9, 112.7, 33.4, 19.0; IR (KBr) ν_{max} 3331, 3661, 2922, 1638, 1451, 1322, 1289, 1231, 1140, 1087, 764, 721, 628 cm⁻¹; HRMS (ESI) calcd for C₂₄H₁₈BrNO₃SNa (M + Na) 502.0088, found 502.0010.

5-Bromo-3-(2-methyl-3-tosyl-4H-chromen-4-yl)-1H-indole (18b). Yellow solid (145 mg, 98%): $R_f = 0.4$ (40% ethyl acetate in hexanes); mp 178–180 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.20 (s, 1H), 7.29 (s, 1H), 7.20 (d, J = 8.2 Hz, 2H), 7.14–7.08 (m, 3H), 7.04–6.95 (m, 4H), 6.84 (d, J = 8.2 Hz, 2H), 5.42 (s, 1H), 2.55 (s, 3H), 2.19 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 159.1, 148.6, 143.0, 138.8, 134.9, 128.85, 128.7, 127.8, 126.9, 126.35, 124.9, 124.5, 124.0, 123.8, 121.1, 118.6, 116.1, 115.5, 112.8, 112.6, 33.4, 21.3, 18.9; IR (KBr) ν_{max} 3400, 2922, 1635, 1456, 1289, 1233, 1145, 1090, 752, 674 cm⁻¹; HRMS (ESI) calcd for C₂₅H₂₀BrNO₃SNa (M + Na) 518.0224, found 518.0218.

1-*E*thyl-3-[2-methyl-3-(phenylsulfonyl)-4H-chromen-4-yl]-1H-indole (**19a**). Pale yellow solid (127 mg, 99%): $R_f = 0.7$ (40% ethyl acetate in hexanes); mp 170–172 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.24 (d, J = 8.1 Hz, 1H), 7.21 (dd, J = 1.1, 8.4 Hz, 2H), 7.17–7.09 (m, 4H), 7.07–7.04 (m, 1H), 7.02 (dd, J = 0.9, 8.1 Hz, 1H), 6.98–6.93 (m, 4H), 6.90–6.87 (m, 1H), 5.51 (s, 1H), 3.97 (q, J = 7.3 Hz, 2H), 2.54 (s, 3H), 1.37 (t, J = 7.3 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 159.1, 148.8, 142.2, 136.0, 131.6, 129.1, 127.7, 127.5, 126.1, 126.0, 125.6, 124.9, 124.4, 121.3, 119.1, 118.8, 117.4, 115.9, 115.6, 109.1, 40.6, 33.4, 19.0, 15.3; IR (KBr) ν_{max} 3448, 3123, 3054, 2962, 2927, 1637, 1321, 1297, 1235, 1149, 745, 719, 628 cm⁻¹; HRMS (ESI) calcd for C₂₆H₂₃NO₃SNa (M + Na) 452.1296, found 452.1291.

1-Ethyl-3-(2-methyl-3-tosyl-4H-chromen-4-yl)-1H-indole (19b). Colorless solid (132 mg, 99%): $R_f = 0.7$ (40% ethyl acetate in hexanes); mp 144–146 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.24 (d, J = 7.9 Hz, 1H), 7.14–7.04 (m, 6H), 7.01 (dd, J = 1.1, 8.2 Hz, 1H), 6.97–6.93 (m, 2H), 6.89 (dt, J = 0.9, 7.9 Hz, 1H), 6.73 (d, J = 8.1 Hz, Article

2H), 5.49 (s, 1H), 3.99 (q, J = 7.3 Hz, 2H), 2.53 (s, 3H), 2.16 (s, 3H), 1.38 (t, J = 7.3 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 158.8, 148.8, 142.5, 139.3, 136.0, 129.0, 128.4, 127.5, 126.3, 126.0, 125.6, 124.8, 124.5, 121.2, 119.1, 118.9, 117.4, 115.9, 109.1, 40.6, 33.4, 21.2, 19.0, 15.4; IR (KBr) ν_{max} 3431, 3048, 2972, 2927, 1636, 1590, 1463, 1321, 1292, 1234, 1146, 765, 736, 696, 661 cm⁻¹; HRMS (ESI) calcd for C₂₇H₂₅NO₃SNa (M + Na) 466.1453, found 466.1449.

2-Methyl-3-[2-methyl-3-(phenylsulfonyl)-4H-chromen-4-yl]-6nitro-1H-indole (**20**). Yellow solid (123 mg, 89%): $R_f = 0.2$ (40% ethyl acetate in hexanes); mp 254–256 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.06 (s, 1H), 8.03 (s, 1H), 7.86 (dd, J = 2.1, 8.9 Hz, 1H), 7.23 (d, J = 7.4 Hz, 2H), 7.15–7.12 (m, 2H), 7.08 (d, J = 8.2 Hz, 1H), 7.06 (d, J = 8.9 Hz, 1H), 6.98–6.95 (m, 4H), 5.56 (s, 1H), 2.65 (s, 3H), 2.56 (s, 3H); ¹³C NMR [75 MHz, (CD₃)₂SO + CDCl₃] δ 158.6, 147.7, 141.0, 140.0, 137.9, 135.6, 131.2, 128.2, 127.2, 127.1, 125.1, 124.9, 124.3, 122.8, 115.4, 115.2, 113.5, 109.7, 31.1, 18.1, 11.0; IR (KBr) ν_{max} 3376, 3051, 2924, 2362, 1643, 1479, 1326, 1231, 1151, 756, 719, 626 cm⁻¹; HRMS (ESI) calcd for C₂₅H₂₀N₂O₅SNa (M + Na) 483.0991, found 483.0998.

3-[6-Methoxy-2-methyl-3-(phenylsulfonyl)-4H-chromen-4-yl]-1Hindole (21). Orange solid (128 mg, 99%): $R_f = 0.4$ (40% ethyl acetate in hexanes); mp 172–174 °C (CH₂Cl₂/hexane); ¹H NMR (400 MHz, CDCl₃) δ 7.90 (s, 1H), 7.26–7.22 (m, 3H), 7.19–7.15 (m, 2H), 7.10 (d, J = 2.3 Hz, 1H), 7.05 (t, J = 7.1 Hz, 1H), 7.00–6.95 (m, 3H), 6.92–6.88 (m, 1H), 6.66 (dd, J = 2.9, 8.9 Hz, 1H), 6.61 (d, J = 2.9 Hz, 1H), 5.50 (s, 1H), 3.64 (s, 3H), 2.50 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 159.6, 156.5, 143.1, 142.2, 136.4, 131.8, 127.9, 126.2, 125.4, 124.9, 122.7, 121.9, 119.7, 118.8, 118.6, 116.9, 114.4, 113.8, 113.1, 111.1, 55.5, 34.0, 19.0; IR (KBr) ν_{max} 3350, 3055, 2919, 2845, 1625, 1497, 1285, 1227, 1143, 1089, 744, 622 cm⁻¹; HRMS (ESI) calcd for C₂₅H₂₁NO₄SNa (M + Na) 454.1089, found 454.1080.

3-(6-Bromo-2-methyl-3-tosyl-4H-chromen-4-yl)-1H-indole (22). Pale yellow solid (105 mg, 71%): $R_f = 0.6$ (40% ethyl acetate in hexanes); mp 160–162 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.00 (s, 1H), 7.22–7.19 (m, 3H), 7.16 (d, J = 8.0 Hz, 1H), 7.12–7.04 (m, 4H), 6.91–6.88 (m, 2H), 6.76 (d, J = 8.1 Hz, 2H), 5.45 (s, 1H), 2.51 (s, 3H), 2.17 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 158.7, 148.0, 142.8, 138.9, 136.4, 131.8, 130.7, 128.6, 126.4, 126.3, 125.2, 122.9, 122.0, 119.8, 118.4, 118.3, 117.8, 117.1, 115.8, 111.2, 33.5, 21.3, 18.9; IR (KBr) ν_{max} 3442, 3125, 3060, 2922, 1638, 1477, 1318, 1292, 1230, 1146, 1089, 814, 741, 672 cm⁻¹; HRMS (ESI) calcd for C₂₅H₂₀BrNO₃SNa (M + Na) 518.0224, found 518.0229.

2-[2-Methyl-3-(phenylsulfonyl)-4H-chromen-4-yl]-1H-pyrrole (**23a**). Colorless solid (88 mg, 84%): $R_f = 0.7$ (40% ethyl acetate in hexanes); mp 193–195 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.92 (s, 1H), 7.55–7.50 (m, 3H), 7.43–7.39 (m, 2H), 7.26–7.22 (m, 1H), 7.18 (dd, J = 1.2, 7.6 Hz, 1H), 7.13–7.09 (m, 1H), 7.04 (d, J = 8.2 Hz, 1H), 6.67–6.65 (m, 1H), 6.03–6.01 (m, 1H), 5.47–5.45 (m, 1H), 5.26 (s, 1H), 2.29 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 160.6, 150.0, 142.4, 134.0, 132.8, 129.3, 129.0, 128.2, 126.6, 125.1, 122.6, 117.5, 116.1, 115.9, 108.2, 106.9, 34.9, 18.8; IR (KBr) ν_{max} 3370, 2924, 1642, 1289, 1242, 1146, 1089, 760, 724, 629 cm⁻¹; HRMS (ESI) calcd for C₂₀H₁₇NO₃SNa (M + Na) 374.0827, found 374.0834.

2-(2-Methyl-3-tosyl-4H-chromen-4-yl)-1H-pyrrole (**23b**). Colorless solid (107 mg, 98%): $R_f = 0.7$ (40% ethyl acetate in hexanes); mp 186–188 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.98 (s, 1H), 7.42 (d, J = 8.3 Hz, 2H), 7.26–7.17 (m, 4H), 7.12–7.09 (m, 1H), 7.03 (d, J = 8.1 Hz, 1H), 6.68–6.66 (m, 1H), 6.02 (dd, J = 2.8, 5.8 Hz, 1H), 5.44 (s, 1H), 5.23 (s, 1H), 2.38 (s, 3H), 2.28 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 160.3, 150.1, 143.8, 139.5, 134.1, 129.6, 129.3, 128.2, 126.7, 125.0, 122.7, 117.5, 116.2, 116.0, 108.1, 106.8, 34.9, 21.5, 18.7; IR (KBr) ν_{max} 3366, 2927, 1637, 1322, 1297, 1236, 1145, 753, 730, 667 cm⁻¹; HRMS (ESI) calcd for C₂₁H₁₉NO₃SNa (M + Na) 388.0983, found 388.0984.

2-(6-Bromo-2-methyl-3-tosyl-4H-chromen-4-yl)-1H-pyrrole (24). Colorless solid (130 mg, 98%): $R_f = 0.7$ (40% ethyl acetate in hexanes); mp 191–193 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.98 (s, 1H), 7.40 (d, J = 8.1 Hz, 2H), 7.35–7.31 (m, 2H), 7.22 (d, J = 8.1 Hz, 2H), 6.92 (d, J = 8.5 Hz, 1H), 6.69 (s, 1H), 6.05–6.04 (m, 1H), 5.47 (s, 1H), 5.18 (s, 1H), 2.39 (s, 3H), 2.28 (s, 3H); ¹³C NMR (125 MHz,

CDCl₃) δ 160.0, 149.2, 144.0, 139.2, 133.3, 131.8, 131.3, 129.7, 126.7, 124.8, 117.9, 117.8, 117.3, 116.3, 108.3, 107.1, 34.7, 21.5, 18.7; IR (KBr) ν_{max} 3376, 2925, 1635, 1478, 1317, 1291, 1235, 1145, 1123, 1086, 814, 727, 674 cm⁻¹; HRMS (ESI) calcd for C₂₁H₁₈BrNO₃SNa (M + Na) 466.0088, found 466.0083.

2-[8-Bromo-6-chloro-2-methyl-3-(phenylsulfonyl)-4H-chromen-4-yl]-1H-pyrrole (**25**). Pale yellow solid (138 mg, 99%): $R_f = 0.7$ (40% ethyl acetate in hexanes); mp 213-215 °C (CH₂Cl₂/hexane); ¹H NMR (500 MHz, CDCl₃) δ 8.82 (s, 1H), 7.56–7.50 (m, 3H), 7.46–7.41 (m, 3H), 7.11 (d, J = 2.0 Hz, 1H), 6.67 (s, 1H), 6.05 (d, J = 2.6 Hz, 1H), 5.55 (s, 1H), 5.22 (s, 1H), 2.36 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 160.2, 145.8, 141.8, 133.2, 132.6, 131.7, 130.1, 129.1, 128.1, 126.7, 125.5, 118.0, 116.8, 110.9, 108.6, 107.4, 35.4, 18.6; IR (KBr) ν_{max} 3393, 3071, 2932, 1644, 1452, 1284, 1245, 1136, 1089, 726, 687 cm⁻¹; HRMS (ESI) calcd for C₂₀H₁₅BrClNO₃SNa (M + Na) 485.9542, found 485.9530.

4-(Furan-2-yl)-2-methyl-3-tosyl-4H-chromene (**26**). White solid (49 mg, 45%): $R_f = 0.7$ (40% ethyl acetate in hexanes); mp 163–165 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.41 (d, J = 8.1 Hz, 2H), 7.21–7.16 (m, 4H), 7.12 (s, 1H), 7.09–7.06 (m, 1H), 7.01 (d, J = 8.0 Hz, 1H), 6.19–6.18 (m, 2H), 5.36 (s, 1H), 2.43 (s, 3H), 2.37 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 160.7, 155.3, 149.4, 143.4, 141.9, 139.4, 129.4, 128.7, 128.2, 127.0, 125.0, 121.8, 116.3, 114.2, 110.3, 106.6, 35.6, 21.5, 19.0; IR (KBr) ν_{max} 3308, 3064, 2963, 2926, 2854, 1716, 1623, 1557, 1278, 1229, 1183, 1127, 879, 768, 727 cm⁻¹; HRMS (ESI) calcd for C₂₁H₁₈O₄SNa (M + Na) 389.0823, found 389.0832.

2-Ethoxy-2-methyl-3-(phenylsulfonyl)-2H-chromene (Et-9aa). Pale yellow solid (70 mg, 71%): $R_f = 0.7$ (40% ethyl acetate in hexanes); mp 118–120 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.96–7.93 (m, 3H), 7.62–7.59 (m, 1H), 7.54–7.51 (m, 2H), 7.36–7.33 (m, 2H), 7.03–7.00 (m, 1H), 6.92 (d, J = 8.5 Hz, 1H), 3.27–3.21 (m, 1H), 3.14–3.08 (m, 1H), 1.87 (s, 3H), 0.69 (t, J = 7.0 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 153.6, 141.2, 136.3, 133.2, 133.1, 132.8, 129.5, 128.9, 128.2, 121.8, 117.8, 115.9, 99.9, 58.0, 25.8, 14.6; IR (KBr) ν_{max} 3425, 3061, 3016, 2972, 2928, 2886, 2363, 1623, 1447, 1307, 1154, 1114, 1084, 1047, 765, 725, 691, 630 cm⁻¹; HRMS (ESI) calcd for C₁₈H₁₈O₄SNa (M + Na) 353.0823, found 353.0809.

Synthesis of 2-Methylene-1,3-bis(phenylsulfonyl)piperidin-4-ol (28). Cesium carbonate (408 mg, 1.25 mmol) was added to a solution of N-(3-oxopropyl)benzenesulfonamide¹⁵ 27 (107 mg, 0.50 mmol) and 2-bromoallyl sulfonylbenzene 6a (195 mg, 0.75 mmol) in acetonitrile (5 mL) kept at room temperature. The reaction mixture was stirred for 3 h, and the solvent was evaporated on a rotavapor. Water (20 mL) was added, and the resulting solution was extracted with dichloromethane $(3 \times 10 \text{ mL})$. The combined organic extracts were dried over anhydrous sodium sulfate and evaporated on a rotavapor, and the residue obtained was subjected to column chromatography on silica gel using an ethyl acetate/hexane mixture as the eluent to afford 2-methylene-1,3-bis(phenylsulfonyl)piperidin-4ol 28 (149 mg, 76%) as a pale yellow viscous liquid: $R_f = 0.4$ (50%) ethyl acetate in hexanes); ¹H NMR (500 MHz, $CDCl_3$) δ 7.97 (d, J = 7.3 Hz, 2H), 7.74 (d, J = 7.2 Hz, 2H), 7.65 (td, J = 1.2, 7.5 Hz, 2H), 7.58-7.55 (m, 2H), 7.51 (t, J = 7.85 Hz, 2H), 5.25 (d, J = 1.2 Hz, 1H), 4.76-4.74 (m, 1H), 4.20 (d, J = 1.2 Hz, 1H), 3.69-3.64 (m, 2H), 3.57-3.52 (m, 1H), 2.66 (d, J = 3.1 Hz, 1H), 2.38-2.32 (m, 1H), 1.87–1.82 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 138.7, 136.1, 134.1, 133.1, 132.0, 129.5, 129.0, 128.8, 128.3, 113.7, 73.4, 63.6, 42.0, 29.3; IR (KBr) $\nu_{\rm max}$: 3452, 2926, 2855, 1714, 1630, 1446, 1310, 1153, 1084, 762, 570 cm⁻¹; HRMS (ESI) calcd for $C_{18}H_{20}NO_5S_2$ (M + H) 394.0783, found 394.0780.

PCC Oxidation of 28. Pyridinium chlorochromate (66 mg, 0.30 mmol) was added to a solution of alcohol **28** (100 mg, 0.25 mmol) in dichloromethane (5 mL) at room temperature and stirred for 5 h. The reaction mixture was then poured into a separating funnel containing water (20 mL). The organic layer was separated, and the aqueous phase was extracted with dichloromethane (3×10 mL). The combined organic extracts were dried over anhydrous sodium sulfate and evaporated on a rotavapor, and the residue obtained was subjected to column chromatography on silica gel using an ethyl acetate/hexane mixture as the eluent to afford 6-methyl-1,5-bis(phenylsulfonyl)-2,3-

dihydropyridin-4(1*H*)-one **29** as a brown solid (86 mg, 86%): $R_f = 0.6$ (50% ethyl acetate in hexanes); mp 196–198 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.00 (d, J = 7.4 Hz, 2H), 7.87 (d, J = 7.4 Hz, 2H), 7.75–7.73 (m, 1H), 7.65–7.50 (m, 5H), 4.23–4.21 (m, 2H), 2.89 (s, 3H), 2.57–2.55 (m, 2H); ¹³C NMR (125 MHz, CDCl₃) δ 186.7, 164.5, 142.4, 139.3, 134.7, 133.1, 130.1, 128.7, 127.8, 126.9, 124.4, 47.1, 37.9, 19.0; IR (KBr) ν_{max} 3424, 3063, 2922, 1666, 1508, 1368, 1327, 1300, 1152, 1078, 919, 735, 686, 533 cm⁻¹; HRMS (ESI) calcd for C₁₈H₁₈NO₅S₂ (M + H) 392.0626, found 392.0619.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.5b02324.

¹H and ¹³C NMR spectra for all new compounds (PDF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: rajeevmenon@iict.res.in or srajeevmenon@gmail.com.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The Department of Science and Technology (DST), India, is acknowledged for the award of a Ramanujan fellowship and a Start-up Research Grant (young scientists) to R.S.M. Financial support in part from the XIIth five year plan project "Affordable Cancer Therapeutics (ACT)" from the Council of Scientific and Industrial Research (CSIR), India, is also acknowledged.

REFERENCES

 (a) Pratap, R.; Ram, V. J. Chem. Rev. 2014, 114, 10476. (b) Ellis,
 G. P.; Lockhart, I. M. The Chemistry of Heterocyclic Compounds: Chromenes, Chromanones, and Chromones; Ellis, G. P., Ed.; Wiley-VCH: New York, 2009; Vol. 31, pp 1–1196. (c) Patil, S. A.; Wang, J.; Li, X.
 S.; Chen, J.; Jones, T. S.; Hosni-Ahmed, A.; Patil, R.; Seibel, W. L.; Li,
 W.; Miller, D. D. Bioorg. Med. Chem. Lett. 2012, 22, 4458.
 (d) Kemnitzer, W.; Jiang, S.; Wang, Y.; Kasibhatla, S.; Crogan-Grundy, C.; Bubenik, M.; Labrecque, D.; Denis, R.; Lamothe, S.; Attardo, G.; Gourdeau, H.; Tseng, B.; Drewe, J.; Cai, S. X. Bioorg. Med. Chem. Lett. 2008, 18, 603. (e) Elomri, A.; Mitaku, S.; Michel, S.; Skaltsounis, A.-L.; Tillequin, F.; Koch, M.; Pierré, A.; Guilbaud, N.; Léonce, S.; Kraus-Berthier, L.; Rolland, Y.; Atassi, G. J. Med. Chem. 1996, 39, 4762. (f) Gunatilaka, A. A. L.; Kingston, D. G. I.; Wijeratne, E. M. K.; Bandara, B. M. R.; Hofmann, G. A.; Johnson, R. K. J. Nat. Prod. 1994, 57, 518.

(2) For 2H-chromene synthesis, see: (a) Majumdar, N.; Paul, N. D.; Mandal, S.; de Bruin, B.; Wulff, W. D. ACS Catal. 2015, 5, 2329.
(b) Zeng, B. S.; Yu, X.; Siu, P. W.; Scheidt, K. A. Chem. Sci. 2014, 5, 2277.
(c) Bera, K.; Sarkar, S.; Biswas, S.; Maiti, S.; Jana, U. J. Org. Chem. 2011, 76, 3539.
(d) Chandrasekhara Rao, L.; Meshram, H. M.; Satish Kumar, N.; Nageswara Rao, N.; Jagadeesh Babu, N. Tetrahedron Lett. 2014, 55, 1127.
(e) Menon, R. S.; Findlay, A. D.; Bissember, A. C.; Banwell, M. G. J. Org. Chem. 2009, 74, 8901.
(f) Nicolaou, K. C.; Pfefferkorn, J. A.; Cao, G.-Q. Angew. Chem., Int. Ed. 2000, 39, 734.
(g) Chang, S.; Grubbs, R. H. J. Org. Chem. 1998, 63, 864.

(3) For 4H-chromene synthesis, see: (a) Shestopalov, A. M.; Litvinov, Y. M.; Rodinovskaya, L. A.; Malyshev, O. R.; Semenova, M. N.; Semenov, V. V. ACS Comb. Sci. 2012, 14, 484. (b) Malakar, C. C.; Schmidt, D.; Conrad, J.; Beifuss, U. Org. Lett. 2011, 13, 1972. (c) Dong, Z.; Liu, X.; Feng, J.; Wang, M.; Lin, L.; Feng, X. Eur. J. Org. Chem. 2011, 2011, 137. (d) Liu, Y.; Qian, J.; Lou, S.; Zhu, J.; Xu, Z. J. Org. Chem. 2010, 75, 1309. (e) Gao, S.; Tsai, C. H.; Tseng, C.; Yao, C.-F. Tetrahedron 2008, 64, 9143. (f) van Otterlo, W. A. L.; Ngidi, E. L.; Kuzvidza, S.; Morgans, G. L.; Moleele, S. S.; de Koning, C. B. Tetrahedron 2005, 61, 9996.

(4) (a) Shi, Y. L.; Shi, M. Org. Biomol. Chem. 2007, 5, 1499 and references cited therein. (b) Lesch, B.; Torang, J.; Vanderheiden, S.; Bräse, S. Adv. Synth. Catal. 2005, 347, 555. (c) Lesch, B.; Bräse, S. Angew. Chem., Int. Ed. 2004, 43, 115. (d) Lee, K. Y.; Kim, J. M.; Kim, J. N. Bull. Korean Chem. Soc. 2003, 24, 17.

(5) Kumar, N. N. B.; Reddy, M. N.; Swamy, K. C. K. J. Org. Chem. 2009, 74, 5395.

(6) (a) Dai, L.-Z.; Shi, Y.-L.; Zhao, G.-L.; Shi, M. Chem. - Eur. J. 2007,
13, 3701. (b) Shi, Y.-L.; Shi, M. Chem. - Eur. J. 2006, 12, 3374.
(c) Zhao, G.-L.; Shi, Y.-L.; Shi, M. Org. Lett. 2005, 7, 4527.

(7) (a) Joshi, P. R.; Undeela, S.; Reddy, D. D.; Singarapu, K. K.; Menon, R. S. Org. Lett. **2015**, 17, 1449. (b) Reddy, S.; Thadkapally, S.; Mamidyala, M.; Nanubolu, J. B.; Menon, R. S. RSC Adv. **2015**, 5, 8199.

(8) (a) Núñez, A., Jr.; Martín, M. R.; Fraile, A.; García Ruano, J. L. G. Chem. - Eur. J. 2010, 16, 5443. (b) Lu, C.; Lu, X. Tetrahedron 2004, 60, 6575.

(9) Ma, S.; Wei, Q. J. Org. Chem. 1999, 64, 1026.

(10) Undeela, S.; Thadkapally, S.; Nanubolu, J. B.; Singarapu, K. K.; Menon, R. S. *Chem. Commun.* **2015**, *51*, 13748 (The conditions for the vinylic displacement reaction were adapted from this work. A detailed study of the scope of this vinylic displacement reaction is under progress and will be reported in due course).

(11) Watterson, S. H.; Ni, Z.; Murphree, S. S.; Padwa, A. Org. Synth. **1997**, 74, 115.

(12) Achard, T.; Lepronier, V.; Gimbert, Y.; Clavier, H.; Giordano, L.; Tenaglia, A.; Buono, G. Angew. Chem., Int. Ed. **2011**, 50, 3552.

(13) (a) Sun, S.; Bai, R.; Gu, Y. Chem. - Eur. J. 2014, 20, 549.
(b) Chandrasekhara Rao, L.; Satish Kumar, N.; Jagadeesh Babu, N.; Meshram, H. M. Tetrahedron Lett. 2014, 55, 5342. (c) Li, M.; Zhang, B.; Gu, Y. Green Chem. 2012, 14, 2421.

(14) (a) Kemnitzer, W.; Drewe, J.; Jiang, S.; Zhang, H.; Wang, Y.; Zhao, J.; Jia, S.; Herich, J.; Labreque, D.; Storer, R.; Meerovitch, K.; Bouffard, D.; Rej, R.; Denis, R.; Blais, C.; Lamothe, S.; Attardo, G.; Gourdeau, H.; Tseng, B.; Kasibhatla, S.; Cai, S. X. J. Med. Chem. 2004, 47, 6299. (b) Kemnitzer, W.; Drewe, J.; Jiang, S.; Zhang, H.; Zhao, J.; Crogan-Grundy, C.; Xu, L.; Lamothe, S.; Gourdeau, H.; Denis, R.; Tseng, B.; Kasibhatla, S.; Cai, S. X. J. Med. Chem. 2007, 50, 2858 Also see ref 1d..

(15) MaGee, D. I.; Ramaseshan, M.; Leach, J. D. Can. J. Chem. 1995, 73, 2111.

(16) Weintraub, P. M.; Sabol, J. S.; Kane, J. M.; Borcherding, D. R. *Tetrahedron* **2003**, *59*, 2953.

(17) For a review, see: (a) You, S.-L.; Cai, Q.; Zeng, M. Chem. Soc. Rev. 2009, 38, 2190. For selected recent reports, see: (b) Saha, S.; Alamsetti, S. K.; Schneider, C. Chem. Commun. 2015, 51, 1461.
(c) Zhao, W.; Wang, Z.; Chu, B.; Sun, J. Angew. Chem., Int. Ed. 2015, 54, 1910.